

第3部 身体状況調査の結果

第20表 身長・体重の平均値及び標準偏差（性・年齢階級別）

年齢(歳)	男 性						女 性					
	身 長(cm)			体 重(kg)			身 長(cm)			体 重(kg)		
	人 数	平均値	標準偏差	人 数	平均値	標準偏差	人 数	平均値	標準偏差	人 数	平均値	標準偏差
総 数	3,644	161.3	17.7	3,633	60.2	16.8	4,337	150.3	14.0	4,303	49.9	12.2
1	22	81.0	3.7	22	10.7	1.1	31	79.1	4.2	31	10.1	1.3
2	20	90.0	3.7	20	12.7	1.3	29	87.5	4.0	29	12.2	1.3
3	36	96.1	4.3	36	14.1	1.5	22	93.0	6.5	22	13.9	1.8
4	36	103.7	4.9	36	16.5	2.0	29	101.2	3.2	29	15.9	1.7
5	40	110.5	4.1	40	19.4	3.4	43	108.0	5.8	43	17.7	2.2
6	36	116.6	5.0	36	21.4	3.0	34	115.5	4.9	34	20.4	2.8
7	41	121.6	6.6	41	23.8	4.1	38	121.6	5.2	38	22.9	2.9
8	47	127.2	6.4	47	27.2	5.6	49	127.9	6.7	49	26.9	5.6
9	33	131.6	4.8	33	28.8	4.9	33	130.4	7.0	33	27.9	5.8
10	35	138.1	6.6	35	33.7	7.6	40	140.5	7.1	40	35.3	7.4
11	34	143.1	6.8	35	36.0	9.1	32	148.0	6.0	32	40.6	7.1
12	28	150.2	9.2	28	41.3	9.6	27	152.0	5.8	26	45.6	10.5
13	42	158.5	6.3	42	49.2	11.0	33	152.8	4.4	33	43.9	5.9
14	31	165.2	6.3	31	53.3	10.1	31	157.0	6.1	30	48.8	6.3
15	30	166.4	7.3	30	58.0	10.7	32	157.4	4.6	32	49.5	6.1
16	26	169.6	6.0	26	58.6	10.4	32	157.9	4.5	31	51.9	7.9
17	37	171.1	6.6	36	62.2	12.6	29	158.9	5.5	29	50.7	5.3
18	23	172.1	6.7	23	62.8	10.3	24	158.1	5.3	24	50.8	7.1
19	25	171.8	6.8	24	60.3	6.9	15	158.7	3.7	15	51.8	6.2
20	32	170.3	5.1	32	59.8	8.6	34	157.4	6.6	34	51.5	14.4
21	22	173.2	6.6	22	67.5	15.8	17	157.0	4.6	17	47.8	6.3
22	18	171.9	5.7	18	62.6	7.5	26	160.0	4.1	24	51.3	8.6
23	36	170.9	6.6	36	67.3	11.8	32	157.7	5.6	32	52.7	9.2
24	28	173.5	5.1	28	65.4	11.2	24	158.6	4.6	22	50.1	5.9
25	23	172.1	5.5	23	63.7	6.6	31	157.7	4.2	31	50.6	6.1
26-29	102	171.3	6.2	101	66.0	10.9	129	158.5	5.7	124	53.0	9.6
30-39	397	171.0	6.2	396	69.3	11.8	510	158.0	5.7	499	53.1	8.8
40-49	396	170.5	5.4	395	70.3	10.8	481	157.7	5.1	477	55.1	8.7
50-59	547	168.1	5.8	547	67.5	10.0	690	155.0	5.4	687	54.3	8.4
60-69	682	164.8	5.8	678	64.2	9.4	801	152.1	5.3	799	53.3	8.3
70歳以上	739	160.9	6.4	736	60.1	8.7	959	147.1	6.5	957	49.8	8.7
(再掲)												
20歳以上	3,022	166.6	7.2	3,012	65.3	10.7	3,734	153.4	7.1	3,703	52.7	8.8
20-29歳	261	171.6	6.0	260	65.0	10.9	293	158.3	5.4	284	51.9	9.5
60-64	336	165.3	5.8	333	65.1	9.2	412	152.8	5.3	411	53.6	8.3
65-69	346	164.3	5.7	345	63.5	9.5	389	151.3	5.1	388	52.9	8.2
70-74	306	162.5	6.0	304	62.7	8.4	358	149.4	5.6	358	52.5	8.4
75-79	225	160.6	6.4	225	59.4	8.7	284	147.3	5.8	283	50.4	8.3
80歳以上	208	158.8	6.2	207	56.9	8.2	317	144.4	7.0	316	46.3	8.1

注) 体重は妊婦除外

第21表 BMIの平均値及び標準偏差（性・年齢階級別）

年齢(歳)	男 性			女 性		
	人 数	平均値	標準偏差	人 数	平均値	標準偏差
総 数	3,151	23.39	3.29	3,832	22.35	3.53
15-19歳	139	20.83	3.03	131	20.32	2.33
20-29	260	22.08	3.60	284	20.71	3.61
30-39	396	23.65	3.55	499	21.29	3.45
40-49	395	24.18	3.40	477	22.15	3.41
50-59	547	23.88	3.21	687	22.60	3.29
60-69	678	23.63	2.98	799	23.03	3.47
70歳以上	736	23.18	2.95	955	23.02	3.58
(再掲)						
20歳以上	3,012	23.51	3.25	3,701	22.42	3.55
20-69歳	2,276	23.61	3.34	2,746	22.21	3.51
40-69	1,620	23.85	3.17	1,963	22.67	3.41
65-69	345	23.47	2.98	388	23.11	3.55
70-74	304	23.72	2.76	358	23.51	3.58
75-79	225	23.01	2.90	283	23.21	3.51
80-84	154	22.80	2.93	196	22.30	3.40
80歳以上	207	22.57	3.13	314	22.29	3.54
85歳以上	53	21.89	3.60	118	22.26	3.77

注) 妊婦除外

第22表 BMIの分布（15歳以上）

	総 数		男 性		女 性	
	人 数	%	人 数	%	人 数	%
総 数	6,983	100	3,151	100	3,832	100
15未満	15	0.2	1	0.0	14	0.4
15-16未満	36	0.5	10	0.3	26	0.7
16-17未満	77	1.1	21	0.7	56	1.5
17-18未満	269	3.9	80	2.5	189	4.9
18-19未満	432	6.2	119	3.8	313	8.2
19-20未満	632	9.1	229	7.3	403	10.5
20-21未満	784	11.2	285	9.0	499	13.0
21-22未満	846	12.1	358	11.4	488	12.7
22-23未満	862	12.3	411	13.0	451	11.8
23-24未満	725	10.4	390	12.4	335	8.7
24-25未満	662	9.5	372	11.8	290	7.6
25-26未満	495	7.1	272	8.6	223	5.8
26-27未満	375	5.4	205	6.5	170	4.4
27-28未満	267	3.8	147	4.7	120	3.1
28-29未満	157	2.2	85	2.7	72	1.9
29-30未満	117	1.7	61	1.9	56	1.5
30-31未満	87	1.2	43	1.4	44	1.1
31-32未満	47	0.7	21	0.7	26	0.7
32-33未満	38	0.5	16	0.5	22	0.6
33-34未満	17	0.2	6	0.2	11	0.3
34-35未満	16	0.2	7	0.2	9	0.2
35以上	27	0.4	12	0.4	15	0.4

注) 妊婦除外

第23表 BMIの状況（性・年齢階級別）

		総数		肥満区分					
				やせ		普通		肥満	
		人数	%	人数	%	人数	%	人数	%
男性	総数	3,151	100.0	161	5.1	2,115	67.1	875	27.8
	15-19歳	139	100.0	31	22.3	94	67.6	14	10.1
	20-29歳	260	100.0	25	9.6	197	75.8	38	14.6
	30-39歳	396	100.0	9	2.3	270	68.2	117	29.5
	40-49歳	395	100.0	10	2.5	243	61.5	142	35.9
	50-59歳	547	100.0	18	3.3	352	64.4	177	32.4
	60-69歳	678	100.0	29	4.3	450	66.4	199	29.4
	70歳以上	736	100.0	39	5.3	509	69.2	188	25.5
	(再掲)20歳以上	3,012	100.0	130	4.3	2,021	67.1	861	28.6
	(再掲)20-69歳	2,276	100.0	91	4.0	1,512	66.4	673	29.6
	(再掲)40-69歳	1,620	100.0	57	3.5	1,045	64.5	518	32.0
	(再掲)65-69歳	345	100.0	17	4.9	231	67.0	97	28.1
	(再掲)70-74歳	304	100.0	4	1.3	203	66.8	97	31.9
	(再掲)75-79歳	225	100.0	17	7.6	158	70.2	50	22.2
	(再掲)80-84歳	154	100.0	11	7.1	108	70.1	35	22.7
(再掲)80歳以上	207	100.0	18	8.7	148	71.5	41	19.8	
(再掲)85歳以上	53	100.0	7	13.2	40	75.5	6	11.3	
女性	総数	3,832	100.0	427	11.1	2,637	68.8	768	20.0
	15-19歳	131	100.0	29	22.1	97	74.0	5	3.8
	20-29歳	284	100.0	64	22.5	198	69.7	22	7.7
	30-39歳	499	100.0	84	16.8	356	71.3	59	11.8
	40-49歳	477	100.0	50	10.5	341	71.5	86	18.0
	50-59歳	687	100.0	56	8.2	486	70.7	145	21.1
	60-69歳	799	100.0	56	7.0	548	68.6	195	24.4
	70歳以上	955	100.0	88	9.2	611	64.0	256	26.8
	(再掲)20歳以上	3,701	100.0	398	10.8	2,540	68.6	763	20.6
	(再掲)20-69歳	2,746	100.0	310	11.3	1,929	70.2	507	18.5
	(再掲)40-69歳	1,963	100.0	162	8.3	1,375	70.0	426	21.7
	(再掲)65-69歳	388	100.0	32	8.2	255	65.7	101	26.0
	(再掲)70-74歳	358	100.0	23	6.4	222	62.0	113	31.6
	(再掲)75-79歳	283	100.0	25	8.8	183	64.7	75	26.5
	(再掲)80-84歳	196	100.0	23	11.7	131	66.8	42	21.4
(再掲)80歳以上	314	100.0	40	12.7	206	65.6	68	21.7	
(再掲)85歳以上	118	100.0	17	14.4	75	63.6	26	22.0	

注) 妊婦除外 BMIによる肥満の判定はp.17参照

第24表 日比式による肥満区分（6-14歳，性・年齢階級別）

	総数	6-8歳		9-11歳		12-14歳			
		人数	%	人数	%	人数	%		
総数	総数	642	100	245	100	207	100	190	100
	-30%以上～-20%未満	18	2.8	1	0.4	6	2.9	11	5.8
	-20%以上～-10%未満	97	15.1	26	10.6	26	12.6	45	23.7
	-10%以上～0%未満	222	34.6	89	36.3	76	36.7	57	30.0
	0%以上～+10%未満	176	27.4	80	32.7	56	27.1	40	21.1
	+10%以上～+20%未満	71	11.1	30	12.2	21	10.1	20	10.5
	+20%以上～+30%未満	30	4.7	12	4.9	13	6.3	5	2.6
	+30%以上～+40%未満	14	2.2	6	2.4	2	1.0	6	3.2
	+40%以上	14	2.2	1	0.4	7	3.4	6	3.2
男性	総数	327	100	124	100	102	100	101	100
	-30%以上～-20%未満	14	4.3	0	0	4	3.9	10	9.9
	-20%以上～-10%未満	55	16.8	16	12.9	13	12.7	26	25.7
	-10%以上～0%未満	120	36.7	42	33.9	44	43.1	34	33.7
	0%以上～+10%未満	72	22.0	40	32.3	20	19.6	12	11.9
	+10%以上～+20%未満	34	10.4	16	12.9	10	9.8	8	7.9
	+20%以上～+30%未満	15	4.6	6	4.8	5	4.9	4	4.0
	+30%以上～+40%未満	8	2.4	4	3.2	1	1.0	3	3.0
	+40%以上	9	2.8	0	0	5	4.9	4	4.0
女性	総数	315	100	121	100	105	100	89	100
	-30%以上～-20%未満	4	1.3	1	0.8	2	1.9	1	1.1
	-20%以上～-10%未満	42	13.3	10	8.3	13	12.4	19	21.3
	-10%以上～0%未満	102	32.4	47	38.8	32	30.5	23	25.8
	0%以上～+10%未満	104	33.0	40	33.1	36	34.3	28	31.5
	+10%以上～+20%未満	37	11.7	14	11.6	11	10.5	12	13.5
	+20%以上～+30%未満	15	4.8	6	5.0	8	7.6	1	1.1
	+30%以上～+40%未満	6	1.9	2	1.7	1	1.0	3	3.4
	+40%以上	5	1.6	1	0.8	2	1.9	2	2.2

注) 日比式による区分は p.17 参照

第25表 学校保健統計調査方式による肥満区分（6-14歳，性・年齢階級別）

	総数	6-8歳		9-11歳		12-14歳			
		人数	%	人数	%	人数	%		
総数	総数	642	100	245	100	207	100	190	100
	-30%未満～-20%以下	14	2.2	1	0.4	8	3.9	5	2.6
	-20%未満～-10%以上	125	19.5	42	17.1	33	15.9	50	26.3
	-10%未満～0%未満	236	36.8	96	39.2	82	39.6	58	30.5
	0%以上～+10%未満	154	24.0	69	28.2	44	21.3	41	21.6
	+10%以上～+20%未満	58	9.0	23	9.4	18	8.7	17	8.9
	+20%以上～+30%未満	35	5.5	12	4.9	14	6.8	9	4.7
	+30%以上～+40%未満	8	1.2	2	0.8	3	1.4	3	1.6
	+40%以上～+50%未満	8	1.2	0	0	3	1.4	5	2.6
	+50%以上	4	0.6	0	0	2	1.0	2	1.1
男性	総数	327	100	124	100	102	100	101	100
	-30%未満～-20%以下	8	2.4	0	0	5	4.9	3	3.0
	-20%未満～-10%以上	66	20.2	21	16.9	19	18.6	26	25.7
	-10%未満～0%未満	118	36.1	42	33.9	44	43.1	32	31.7
	0%以上～+10%未満	74	22.6	40	32.3	15	14.7	19	18.8
	+10%以上～+20%未満	27	8.3	12	9.7	8	7.8	7	6.9
	+20%以上～+30%未満	21	6.4	8	6.5	6	5.9	7	6.9
	+30%以上～+40%未満	5	1.5	1	0.8	2	2.0	2	2.0
	+40%以上～+50%未満	6	1.8	0	0	2	2.0	4	4.0
	+50%以上	2	0.6	0	0	1	1.0	1	1.0
女性	総数	315	100	121	100	105	100	89	100
	-30%未満～-20%以下	6	1.9	1	0.8	3	2.9	2	2.2
	-20%未満～-10%以上	59	18.7	21	17.4	14	13.3	24	27.0
	-10%未満～0%未満	118	37.5	54	44.6	38	36.2	26	29.2
	0%以上～+10%未満	80	25.4	29	24.0	29	27.6	22	24.7
	+10%以上～+20%未満	31	9.8	11	9.1	10	9.5	10	11.2
	+20%以上～+30%未満	14	4.4	4	3.3	8	7.6	2	2.2
	+30%以上～+40%未満	3	1.0	1	0.8	1	1.0	1	1.1
	+40%以上～+50%未満	2	0.6	0	0	1	1.0	1	1.1
	+50%以上	2	0.6	0	0	1	1.0	1	1.1

注) 学校保健統計調査方式による区分は p.18 参照

第26表 腹囲の分布（6歳以上、性・年齢階級別）

総数	総数		6-8歳		9-11歳		12-14歳		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 20歳以上		(再掲) 40-74歳						
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%			
総数	7,102	100	225	100	175	100	100	100	230	100	486	100	827	100	811	100	1,153	100	1,409	100	1,595	100	6,281	100	4,008	100	0	0			
腹囲(cm)																															
<50	23	0.3	21	9.3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
50<55	123	1.7	88	39.1	4	2.3	0	0	0	0	6	1.2	6	0.7	2	0.2	6	0.5	0	0	3	0.2	4	0.1	1	0.0	1	0.0	0		
55<60	189	2.7	77	34.2	18	10.3	0	0	2	0.9	6	1.2	6	0.7	2	0.2	6	0.5	0	0	7	0.4	27	0.4	10	0.2	10	0.2	0		
60<65	315	4.4	22	9.8	51	29.1	0	0	35	15.2	38	7.8	43	5.2	19	2.3	16	1.4	16	1.1	21	1.3	133	2.4	56	1.4	56	1.4	0		
65<70	528	7.4	9	4.0	49	28.0	0	0	60	26.1	89	18.3	88	10.6	60	7.4	59	5.1	47	3.3	51	3.2	394	6.3	179	4.5	179	4.5	0		
70<75	880	12.4	7	3.1	27	15.4	0	0	62	27.0	117	24.1	176	21.3	125	15.4	115	10.0	107	7.6	131	8.2	771	12.3	394	9.8	394	9.8	0		
75<80	1,092	15.4	1	0.4	10	5.7	0	0	41	17.8	98	20.2	160	19.3	155	19.1	206	17.9	208	14.8	207	13.0	1,094	16.5	637	15.9	637	15.9	0		
80<85	1,263	17.8	0	0	7	4.0	0	0	14	6.1	67	13.8	135	16.3	175	21.6	242	21.0	310	22.0	310	19.4	1,239	19.7	855	21.3	855	21.3	0		
85<90	1,227	17.3	0	0	5	2.9	0	0	6	2.6	33	6.8	100	12.1	119	14.7	256	22.2	324	23.0	381	23.9	1,213	19.3	853	21.3	853	21.3	0		
90<95	736	10.4	0	0	2	1.1	0	0	4	1.7	11	2.3	62	7.5	71	8.8	137	11.9	205	14.5	243	15.2	729	11.6	521	13.0	521	13.0	0		
95<100	430	6.1	0	0	0	0	0	0	3	1.3	12	2.5	27	3.3	46	5.7	61	5.3	117	8.3	163	10.2	426	6.8	298	7.4	298	7.4	0		
100<105	175	2.5	0	0	1	0.6	0	0	1	0.4	7	1.4	19	2.3	23	2.8	35	3.0	43	3.1	46	2.9	173	2.8	124	3.1	124	3.1	0		
105<110	64	0.9	0	0	0	0	0	0	2	0.9	5	1.0	4	0.5	4	0.5	11	1.0	15	1.1	22	1.4	61	1.0	40	1.0	40	1.0	0		
110<115	30	0.4	0	0	0	0	0	0	0	0	1	0.2	4	0.5	1	0.1	5	0.4	11	0.8	8	0.5	30	0.5	20	0.5	20	0.5	0		
115<120	15	0.2	0	0	0	0	0	0	0	0	1	0.2	1	0.1	5	0.6	3	0.3	4	0.3	1	0.1	15	0.2	12	0.3	12	0.3	0		
>=120	12	0.2	0	0	0	0	0	0	0	0	1	0.2	2	0.2	2	0.2	1	0.1	2	0.1	1	0.1	12	0.2	8	0.2	8	0.2	0		
男性	3,206	100	113	100	91	100	114	100	234	100	358	100	358	100	356	100	499	100	644	100	701	100	2,792	100	1,792	100	1,792	100	0	0	
腹囲(cm)																															
<50	8	0.2	8	7.1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
50<55	48	1.5	36	31.9	2	2.2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
55<60	104	3.2	49	43.4	10	11.0	0	0	18	7.9	3	1.3	7	0.8	0	0	1	0.2	17	2.6	3	0.4	14	0.5	5	0.3	5	0.3	0	0	
60<65	84	2.6	12	10.6	24	26.4	2	2	9	3.9	28	12.0	51	14.2	4	1.1	24	4.8	26	4.0	11	1.6	74	2.7	28	1.6	28	1.6	0	0	
65<70	139	4.3	3	2.7	29	31.9	2	2	26	11.3	56	23.9	68	19.0	7	2.0	74	14.4	48	7.0	43	6.1	223	8.0	90	5.0	90	5.0	0	0	
70<75	282	8.8	4	3.5	11	12.1	4	4	35	15.8	56	23.9	68	19.0	23	6.5	64	12.8	81	12.6	88	12.6	44	1.6	148	8.4	148	8.4	0	0	
75<80	438	13.7	1	0.9	3	3.3	5	5.5	18	7.9	42	17.9	75	20.9	92	25.8	111	22.2	154	22.7	159	22.7	633	22.7	431	24.1	431	24.1	0	0	
80<85	647	20.2	0	0	5	5.5	4	4.4	8	3.5	23	9.8	69	19.3	73	20.5	141	28.3	159	24.7	179	25.5	644	23.1	445	24.8	445	24.8	0	0	
85<90	652	20.3	0	0	4	4.4	2	2.1	2	1.8	8	3.4	48	13.4	45	12.6	82	16.4	107	16.6	120	17.1	445	14.7	285	15.9	285	15.9	0	0	
90<95	415	12.9	0	0	1	1.1	1	1.1	3	1.3	7	3.0	17	4.7	33	9.3	42	8.4	64	9.9	70	10.0	233	8.3	178	9.9	178	9.9	0	0	
95<100	237	7.4	0	0	0	0	0	0	1	0.4	5	2.1	13	3.6	18	5.1	18	3.6	20	3.1	20	2.9	94	3.4	65	3.6	65	3.6	0	0	
100<105	96	3.0	0	0	1	1.1	1	1.1	2	0.9	4	1.7	3	0.8	4	1.1	4	0.8	3	0.5	8	1.1	26	0.9	15	0.8	15	0.8	0	0	
105<110	29	0.9	0	0	0	0	0	0	0	0	0	0	2	0.6	1	0.3	3	0.6	6	0.9	0	0	13	0.5	10	0.6	10	0.6	0	0	
110<115	13	0.4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0.2	3	0.5	0	0	7	0.3	7	0.4	7	0.4	0	0	
115<120	7	0.2	0	0	0	0	0	0	0	0	1	0.4	2	0.6	3	0.8	1	0.2	3	0.5	0	0	7	0.3	7	0.4	7	0.4	0	0	
>=120	7	0.2	0	0	0	0	0	0	0	0	1	0.4	2	0.6	3	0.8	1	0.2	0	0	0	0	7	0.3	7	0.4	7	0.4	0	0	
女性	3,896	100	112	100	84	100	116	100	252	100	469	100	469	100	455	100	654	100	765	100	894	100	3,489	100	2,216	100	2,216	100	0	0	
腹囲(cm)																															
<50	15	0.4	13	11.6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
50<55	75	1.9	52	46.4	2	2.4	0	0	0	0	6	2.4	6	1.3	2	0.4	6	0.9	0	0	3	0.3	4	0.1	1	0.0	1	0.0	1	0.0	0
55<60	85	2.2	28	25.0	8	9.5	0	0	26	10.3	35	13.9	40	8.5	19	4.2	52	7.8	12	1.6	7	0.8	27	0.8	10	0.5	10	0.5	0	0	
60<65	231	5.9	10	8.9	27	32.1	0	0	34	13.3	61	24.2	81	17.3	56	12.4	152	23.3	30	4.0	18	2.0	139	4.0	51	2.3	51	2.3	0	0	
65<70	389	10.0	6	5.4	20	23.8	0	0	22	8.8	61	24.2	125	26.7	102	22.4	91	13.9	81	10.6	88	9.8	548	15.7	304	13.7	304	13.7	0	0	
70<75	598	15.3	3	2.7	16	19.0	22	19.0	22	8.8	42	16.7	92	19.6	98	21.5	142	21.7	127	16.6	119	13.3	620	17.8	408	18.4	408	18.4	0	0	
75<80	654	16.8	0	0	7	8.3	23	19.8	23	9.1	42	16.7	92	19.6	98	21.5	142	21.7	127	16.6	119	13.3	620	17.8	408	18.4	408	18.4	0	0	
80<85	616	15.8	0	0	2	2.4	6	5.2	25	9.9	30	11.5	60	12.8	83	18.2	131	20.0	156	20.4	151	16.9	606	17.4	424	19.1	424	19.1	0	0	
85<90	575																														

第27表 腹囲の区分(性・年齢階級別)

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	6,281	100	486	100	827	100	811	100	1,153	100	1,409	100	1,595	100	4,008	100
85cm未満(男)/90cm未満(女)	4,191	66.7	425	87.4	639	77.3	583	71.9	759	65.8	853	60.5	932	58.4	2,540	63.4
85cm以上(男)/90cm以上(女)	2,090	33.3	61	12.6	188	22.7	228	28.1	394	34.2	556	39.5	663	41.6	1,468	36.6
男性	2,792	100	234	100	358	100	356	100	499	100	644	100	701	100	1,792	100
	1,358	48.6	185	79.1	204	57.0	176	49.4	207	41.5	282	43.8	304	43.4	783	43.7
	1,434	51.4	49	20.9	154	43.0	180	50.6	292	58.5	362	56.2	397	56.6	1,009	56.3
女性	3,489	100	252	100	469	100	455	100	654	100	765	100	894	100	2,216	100
	2,833	81.2	240	95.2	435	92.8	407	89.5	552	84.4	571	74.6	628	70.2	1,757	79.3
	656	18.8	12	4.8	34	7.2	48	10.5	102	15.6	194	25.4	266	29.8	459	20.7

注) 妊婦除外

第28表 BMI・腹囲の区分(性・年齢階級別)

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	6,266	100	484	100	824	100	811	100	1,151	100	1,405	100	1,591	100	4,000	100
BMI<25, 腹囲<85/90cm	3,872	61.8	412	85.1	602	73.1	525	64.7	690	59.9	788	56.1	855	53.7	2,317	57.9
BMI≥25, 腹囲<85/90cm	306	4.9	12	2.5	35	4.2	58	7.2	67	5.8	61	4.3	73	4.6	215	5.4
BMI<25, 腹囲≥85/90cm	872	13.9	19	3.9	63	7.6	76	9.4	162	14.1	240	17.1	312	19.6	596	14.9
BMI≥25, 腹囲≥85/90cm	1,216	19.4	41	8.5	124	15.0	152	18.7	232	20.2	316	22.5	351	22.1	872	21.8
男性	2,784	100	233	100	356	100	356	100	499	100	641	100	699	100	1,787	100
	1,296	46.6	180	77.3	190	53.4	160	44.9	198	39.7	271	42.3	297	42.5	742	41.5
	56	2.0	5	2.1	13	3.7	16	4.5	9	1.8	8	1.2	5	0.7	36	2.0
	683	24.5	19	8.2	59	16.6	68	19.1	139	27.9	180	28.1	218	31.2	471	26.4
	749	26.9	29	12.4	94	26.4	112	31.5	153	30.7	182	28.4	179	25.6	538	30.1
女性	3,482	100	251	100	468	100	455	100	652	100	764	100	892	100	2,213	100
	2,576	74.0	232	92.4	412	88.0	365	80.2	492	75.5	517	67.7	558	62.6	1,575	71.2
	250	7.2	7	2.8	22	4.7	42	9.2	58	8.9	53	6.9	68	7.6	179	8.1
	189	5.4	0	0	4	0.9	8	1.8	23	3.5	60	7.9	94	10.5	125	5.6
	467	13.4	12	4.8	30	6.4	40	8.8	79	12.1	134	17.5	172	19.3	334	15.1

注) 妊婦除外

第29表の1 血圧を下げる薬の服用状況
(性・年齢階級別)

		総数		使用あり		使用無し	
		人数	%	人数	%	人数	%
総数	総数	5,089	100.0	1,320	25.9	3,769	74.1
	20-29歳	322	100.0	3	0.9	319	99.1
	30-39歳	592	100.0	10	1.7	582	98.3
	40-49歳	592	100.0	30	5.1	562	94.9
	50-59歳	916	100.0	167	18.2	749	81.8
	60-69歳	1,224	100.0	352	28.8	872	71.2
	70歳以上	1,443	100.0	758	52.5	685	47.5
男性	総数	2,150	100.0	595	27.7	1,555	72.3
	20-29歳	143	100.0	2	1.4	141	98.6
	30-39歳	223	100.0	7	3.1	216	96.9
	40-49歳	237	100.0	18	7.6	219	92.4
	50-59歳	357	100.0	74	20.7	283	79.3
	60-69歳	544	100.0	172	31.6	372	68.4
	70歳以上	646	100.0	322	49.8	324	50.2
女性	総数	2,939	100.0	725	24.7	2,214	75.3
	20-29歳	179	100.0	1	0.6	178	99.4
	30-39歳	369	100.0	3	0.8	366	99.2
	40-49歳	355	100.0	12	3.4	343	96.6
	50-59歳	559	100.0	93	16.6	466	83.4
	60-69歳	680	100.0	180	26.5	500	73.5
	70歳以上	797	100.0	436	54.7	361	45.3

注) 妊婦除外

第29表の2 脈の乱れを治す薬の服用状況
(性・年齢階級別)

		総数		使用あり		使用無し	
		人数	%	人数	%	人数	%
総数	総数	5,088	100.0	184	3.6	4,904	96.4
	20-29歳	322	100.0	0	0	322	100.0
	30-39歳	592	100.0	3	0.5	589	99.5
	40-49歳	592	100.0	5	0.8	587	99.2
	50-59歳	916	100.0	10	1.1	906	98.9
	60-69歳	1,224	100.0	38	3.1	1,186	96.9
	70歳以上	1,442	100.0	128	8.9	1,314	91.1
男性	総数	2,150	100.0	88	4.1	2,062	95.9
	20-29歳	143	100.0	0	0	143	100.0
	30-39歳	223	100.0	1	0.4	222	99.6
	40-49歳	237	100.0	2	0.8	235	99.2
	50-59歳	357	100.0	5	1.4	352	98.6
	60-69歳	544	100.0	19	3.5	525	96.5
	70歳以上	646	100.0	61	9.4	585	90.6
女性	総数	2,938	100.0	96	3.3	2,842	96.7
	20-29歳	179	100.0	0	0	179	100.0
	30-39歳	369	100.0	2	0.5	367	99.5
	40-49歳	355	100.0	3	0.8	352	99.2
	50-59歳	559	100.0	5	0.9	554	99.1
	60-69歳	680	100.0	19	2.8	661	97.2
	70歳以上	796	100.0	67	8.4	729	91.6

注) 妊婦除外

第29表の3 インスリン注射または血糖を下げる薬の服用状況 (性・年齢階級別)

		総数		使用あり		使用無し	
		人数	%	人数	%	人数	%
総数	総数	5,089	100.0	273	5.4	4,816	94.6
	20-29歳	322	100.0	1	0.3	321	99.7
	30-39歳	592	100.0	2	0.3	590	99.7
	40-49歳	592	100.0	8	1.4	584	98.6
	50-59歳	916	100.0	36	3.9	880	96.1
	60-69歳	1,224	100.0	87	7.1	1,137	92.9
	70歳以上	1,443	100.0	139	9.6	1,304	90.4
男性	総数	2,150	100.0	149	6.9	2,001	93.1
	20-29歳	143	100.0	0	0	143	100.0
	30-39歳	223	100.0	1	0.4	222	99.6
	40-49歳	237	100.0	6	2.5	231	97.5
	50-59歳	357	100.0	20	5.6	337	94.4
	60-69歳	544	100.0	47	8.6	497	91.4
	70歳以上	646	100.0	75	11.6	571	88.4
女性	総数	2,939	100.0	124	4.2	2,815	95.8
	20-29歳	179	100.0	1	0.6	178	99.4
	30-39歳	369	100.0	1	0.3	368	99.7
	40-49歳	355	100.0	2	0.6	353	99.4
	50-59歳	559	100.0	16	2.9	543	97.1
	60-69歳	680	100.0	40	5.9	640	94.1
	70歳以上	797	100.0	64	8.0	733	92.0

注) 妊婦除外

第29表の4 コレステロールを下げる薬の服用状況 (性・年齢階級別)

		総数		使用あり		使用無し	
		人数	%	人数	%	人数	%
総数	総数	5,088	100.0	595	11.7	4,493	88.3
	20-29歳	322	100.0	0	0	322	100.0
	30-39歳	592	100.0	4	0.7	588	99.3
	40-49歳	592	100.0	22	3.7	570	96.3
	50-59歳	916	100.0	83	9.1	833	90.9
	60-69歳	1,224	100.0	212	17.3	1,012	82.7
	70歳以上	1,442	100.0	274	19.0	1,168	81.0
男性	総数	2,149	100.0	191	8.9	1,958	91.1
	20-29歳	143	100.0	0	0	143	100.0
	30-39歳	223	100.0	4	1.8	219	98.2
	40-49歳	237	100.0	14	5.9	223	94.1
	50-59歳	357	100.0	25	7.0	332	93.0
	60-69歳	544	100.0	70	12.9	474	87.1
	70歳以上	645	100.0	78	12.1	567	87.9
女性	総数	2,939	100.0	404	13.7	2,535	86.3
	20-29歳	179	100.0	0	0	179	100.0
	30-39歳	369	100.0	0	0	369	100.0
	40-49歳	355	100.0	8	2.3	347	97.7
	50-59歳	559	100.0	58	10.4	501	89.6
	60-69歳	680	100.0	142	20.9	538	79.1
	70歳以上	797	100.0	196	24.6	601	75.4

注) 妊婦除外

第29表の5 中性脂肪（トリグリセライド）を下げる薬の服用状況（性・年齢階級別）

		総数		使用あり		使用無し	
		人数	%	人数	%	人数	%
総数	総数	5,086	100.0	193	3.8	4,893	96.2
	20-29歳	322	100.0	0	0	322	100.0
	30-39歳	592	100.0	3	0.5	589	99.5
	40-49歳	592	100.0	6	1.0	586	99.0
	50-59歳	915	100.0	33	3.6	882	96.4
	60-69歳	1,224	100.0	59	4.8	1,165	95.2
	70歳以上	1,441	100.0	92	6.4	1,349	93.6
	男性	総数	2,148	100.0	94	4.4	2,054
	20-29歳	143	100.0	0	0	143	100.0
	30-39歳	223	100.0	1	0.4	222	99.6
	40-49歳	237	100.0	5	2.1	232	97.9
	50-59歳	356	100.0	21	5.9	335	94.1
	60-69歳	544	100.0	30	5.5	514	94.5
	70歳以上	645	100.0	37	5.7	608	94.3
女性	総数	2,938	100.0	99	3.4	2,839	96.6
	20-29歳	179	100.0	0	0	179	100.0
	30-39歳	369	100.0	2	0.5	367	99.5
	40-49歳	355	100.0	1	0.3	354	99.7
	50-59歳	559	100.0	12	2.1	547	97.9
	60-69歳	680	100.0	29	4.3	651	95.7
	70歳以上	796	100.0	55	6.9	741	93.1

注) 妊婦除外

第29表の6 貧血治療のための薬（鉄剤）の服用状況（性・年齢階級別）

		総数		使用あり		使用無し	
		人数	%	人数	%	人数	%
総数	総数	5,089	100.0	73	1.4	5,016	98.6
	20-29歳	322	100.0	0	0	322	100.0
	30-39歳	592	100.0	5	0.8	587	99.2
	40-49歳	592	100.0	10	1.7	582	98.3
	50-59歳	916	100.0	12	1.3	904	98.7
	60-69歳	1,224	100.0	12	1.0	1,212	99.0
	70歳以上	1,443	100.0	34	2.4	1,409	97.6
	男性	総数	2,150	100.0	23	1.1	2,127
20-29歳		143	100.0	0	0	143	100.0
30-39歳		223	100.0	1	0.4	222	99.6
40-49歳		237	100.0	0	0	237	100.0
50-59歳		357	100.0	3	0.8	354	99.2
60-69歳		544	100.0	4	0.7	540	99.3
70歳以上		646	100.0	15	2.3	631	97.7
女性		総数	2,939	100.0	50	1.7	2,889
	20-29歳	179	100.0	0	0	179	100.0
	30-39歳	369	100.0	4	1.1	365	98.9
	40-49歳	355	100.0	10	2.8	345	97.2
	50-59歳	559	100.0	9	1.6	550	98.4
	60-69歳	680	100.0	8	1.2	672	98.8
	70歳以上	797	100.0	19	2.4	778	97.6

注) 妊婦除外

第30表の1 血圧の状況（性・年齢階級別）

総数		総数		高血圧学会（2回測定値の平均値）												（再掲） 収縮期高血圧	
				至適血圧		正常血圧		正常高値血圧		Ⅰ度高血圧		Ⅱ度高血圧		Ⅲ度高血圧			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%*
総数	総数	4,783	100.0	1,222	25.5	861	18.0	972	20.3	1,225	25.6	403	8.4	100	2.1	896	18.7
	15-19歳	117	100.0	91	77.8	19	16.2	5	4.3	2	1.7	0	0	0	0	2	1.7
	20-29歳	253	100.0	167	66.0	51	20.2	21	8.3	12	4.7	2	0.8	0	0	5	2.0
	30-39歳	529	100.0	316	59.7	103	19.5	58	11.0	39	7.4	12	2.3	1	0.2	7	1.3
	40-49歳	524	100.0	201	38.4	105	20.0	98	18.7	82	15.6	25	4.8	13	2.5	18	3.4
	50-59歳	833	100.0	152	18.2	170	20.4	194	23.3	226	27.1	74	8.9	17	2.0	102	12.2
	60-69歳	1,171	100.0	176	15.0	206	17.6	258	22.0	372	31.8	125	10.7	34	2.9	266	22.7
	70歳以上	1,356	100.0	119	8.8	207	15.3	338	24.9	492	36.3	165	12.2	35	2.6	496	36.6
	（再掲）65-74歳	1,150	100.0	142	12.3	179	15.6	263	22.9	395	34.3	141	12.3	30	2.6	337	29.3
	（再掲）75歳以上	800	100.0	62	7.8	129	16.1	194	24.3	299	37.4	95	11.9	21	2.6	314	39.3
男性	総数	1,969	100.0	331	16.8	344	17.5	443	22.5	590	30.0	211	10.7	50	2.5	390	19.8
	15-19歳	48	100.0	28	58.3	15	31.3	3	6.3	2	4.2	0	0	0	0	2	4.2
	20-29歳	103	100.0	48	46.6	27	26.2	17	16.5	9	8.7	2	1.9	0	0	5	4.9
	30-39歳	182	100.0	62	34.1	50	27.5	38	20.9	22	12.1	10	5.5	0	0	3	1.6
	40-49歳	197	100.0	44	22.3	41	20.8	39	19.8	50	25.4	15	7.6	8	4.1	6	3.0
	50-59歳	313	100.0	35	11.2	57	18.2	76	24.3	105	33.5	34	10.9	6	1.9	40	12.8
	60-69歳	516	100.0	61	11.8	67	13.0	119	23.1	174	33.7	77	14.9	18	3.5	119	23.1
	70歳以上	610	100.0	53	8.7	87	14.3	151	24.8	228	37.4	73	12.0	18	3.0	215	35.2
	（再掲）65-74歳	528	100.0	59	11.2	75	14.2	119	22.5	184	34.8	75	14.2	16	3.0	145	27.5
	（再掲）75歳以上	352	100.0	25	7.1	54	15.3	83	23.6	139	39.5	41	11.6	10	2.8	138	39.2
女性	総数	2,814	100.0	891	31.7	517	18.4	529	18.8	635	22.6	192	6.8	50	1.8	506	18.0
	15-19歳	69	100.0	63	91.3	4	5.8	2	2.9	0	0	0	0	0	0	0	0
	20-29歳	150	100.0	119	79.3	24	16.0	4	2.7	3	2.0	0	0	0	0	0	0
	30-39歳	347	100.0	254	73.2	53	15.3	20	5.8	17	4.9	2	0.6	1	0.3	4	1.2
	40-49歳	327	100.0	157	48.0	64	19.6	59	18.0	32	9.8	10	3.1	5	1.5	12	3.7
	50-59歳	520	100.0	117	22.5	113	21.7	118	22.7	121	23.3	40	7.7	11	2.1	62	11.9
	60-69歳	655	100.0	115	17.6	139	21.2	139	21.2	198	30.2	48	7.3	16	2.4	147	22.4
	70歳以上	746	100.0	66	8.8	120	16.1	187	25.1	264	35.4	92	12.3	17	2.3	281	37.7
	（再掲）65-74歳	622	100.0	83	13.3	104	16.7	144	23.2	211	33.9	66	10.6	14	2.3	192	30.9
	（再掲）75歳以上	448	100.0	37	8.3	75	16.7	111	24.8	160	35.7	54	12.1	11	2.5	176	39.3

注) 日本高血圧学会による血圧の分類は p. 16 参照

※ 総数に対する割合

第30表の2 血圧の状況（性・年齢階級別）

総数		総数		高血圧学会(2回測定値の平均値)												(再掲)	
				至適血圧		正常血圧		正常高値血圧		Ⅰ度高血圧		Ⅱ度高血圧		Ⅲ度高血圧		収縮期高血圧	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%*
総数	総数	3,545	100.0	1,148	32.4	713	20.1	671	18.9	723	20.4	221	6.2	69	1.9	457	12.9
	15-19歳	117	100.0	91	77.8	19	16.2	5	4.3	2	1.7	0	0	0	0	2	1.7
	20-29歳	246	100.0	163	66.3	50	20.3	21	8.5	10	4.1	2	0.8	0	0	5	2.0
	30-39歳	512	100.0	307	60.0	102	19.9	56	10.9	36	7.0	10	2.0	1	0.2	6	1.2
	40-49歳	496	100.0	199	40.1	104	21.0	87	17.5	76	15.3	20	4.0	10	2.0	17	3.4
	50-59歳	683	100.0	147	21.5	149	21.8	155	22.7	162	23.7	57	8.3	13	1.9	79	11.6
	60-69歳	839	100.0	157	18.7	175	20.9	182	21.7	231	27.5	67	8.0	27	3.2	158	18.8
	70歳以上	652	100.0	84	12.9	114	17.5	165	25.3	206	31.6	65	10.0	18	2.8	190	29.1
	(再掲)65-74歳	705	100.0	117	16.6	137	19.4	161	22.8	205	29.1	66	9.4	19	2.7	160	22.7
	(再掲)75歳以上	340	100.0	41	12.1	58	17.1	83	24.4	116	34.1	33	9.7	9	2.6	113	33.2
男性	総数	1,414	100.0	305	21.6	279	19.7	306	21.6	363	25.7	122	8.6	39	2.8	201	14.2
	15-19歳	48	100.0	28	58.3	15	31.3	3	6.3	2	4.2	0	0	0	0	2	4.2
	20-29歳	101	100.0	48	47.5	26	25.7	17	16.8	8	7.9	2	2.0	0	0	5	5.0
	30-39歳	175	100.0	62	35.4	49	28.0	36	20.6	19	10.9	9	5.1	0	0	2	1.1
	40-49歳	181	100.0	43	23.8	41	22.7	33	18.2	46	25.4	13	7.2	5	2.8	6	3.3
	50-59歳	249	100.0	34	13.7	47	18.9	60	24.1	78	31.3	25	10.0	5	2.0	34	13.7
	60-69歳	353	100.0	52	14.7	53	15.0	85	24.1	107	30.3	40	11.3	16	4.5	64	18.1
	70歳以上	307	100.0	38	12.4	48	15.6	72	23.5	103	33.6	33	10.7	13	4.2	88	28.7
	(再掲)65-74歳	319	100.0	47	14.7	56	17.6	76	23.8	92	28.8	35	11.0	13	4.1	59	18.5
	(再掲)75歳以上	159	100.0	18	11.3	24	15.1	30	18.9	63	39.6	18	11.3	6	3.8	59	37.1
女性	総数	2,131	100.0	843	39.6	434	20.4	365	17.1	360	16.9	99	4.6	30	1.4	256	12.0
	15-19歳	69	100.0	63	91.3	4	5.8	2	2.9	0	0	0	0	0	0	0	0
	20-29歳	145	100.0	115	79.3	24	16.6	4	2.8	2	1.4	0	0	0	0	0	0
	30-39歳	337	100.0	245	72.7	53	15.7	20	5.9	17	5.0	1	0.3	1	0.3	4	1.2
	40-49歳	315	100.0	156	49.5	63	20.0	54	17.1	30	9.5	7	2.2	5	1.6	11	3.5
	50-59歳	434	100.0	113	26.0	102	23.5	95	21.9	84	19.4	32	7.4	8	1.8	45	10.4
	60-69歳	486	100.0	105	21.6	122	25.1	97	20.0	124	25.5	27	5.6	11	2.3	94	19.3
	70歳以上	345	100.0	46	13.3	66	19.1	93	27.0	103	29.9	32	9.3	5	1.4	102	29.6
	(再掲)65-74歳	386	100.0	70	18.1	81	21.0	85	22.0	113	29.3	31	8.0	6	1.6	101	26.2
	(再掲)75歳以上	181	100.0	23	12.7	34	18.8	53	29.3	53	29.3	15	8.3	3	1.7	54	29.8

注) 妊婦・血圧を下げる薬服用者除外

※ 総数に対する割合

第31表の1-1 最低血圧の分布（性・年齢階級別）

	総数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上			
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%		
男性	総数		1,969	100	48	100	103	100	182	100	197	100	313	100	516	100	610	100
	2回の平均値(mmHg)																	
	-39	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	40-49	3	0.2	0	0	0	0	0	0	0	0	0	0	1	0.2	2	0.3	
	50-59	45	2.3	9	18.8	4	3.9	2	1.1	4	2.0	1	0.3	6	1.2	19	3.1	
	60-69	213	10.8	22	45.8	32	31.1	21	11.5	17	8.6	12	3.8	36	7.0	73	12.0	
	70-79	551	28.0	14	29.2	38	36.9	65	35.7	50	25.4	54	17.3	145	28.1	185	30.3	
	80-89	696	35.3	3	6.3	23	22.3	65	35.7	59	29.9	141	45.0	178	34.5	227	37.2	
	90-99	328	16.7	0	0	5	4.9	19	10.4	46	23.4	75	24.0	107	20.7	76	12.5	
	100-109	108	5.5	0	0	1	1.0	10	5.5	14	7.1	26	8.3	35	6.8	22	3.6	
	110-119	16	0.8	0	0	0	0	0	0	1	0.5	3	1.0	7	1.4	5	0.8	
	120-129	7	0.4	0	0	0	0	0	0	4	2.0	1	0.3	1	0.2	1	0.2	
	130-139	2	0.1	0	0	0	0	0	0	2	1.0	0	0	0	0	0	0	
	140-	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
女性	総数		2,814	100	69	100	150	100	347	100	327	100	520	100	655	100	746	100
	2回の平均値(mmHg)																	
	-39	1	0.0	0	0	0	0	1	0.3	0	0	0	0	0	0	0	0	0
	40-49	13	0.5	3	4.3	2	1.3	1	0.3	0	0	0	0	1	0.2	6	0.8	
	50-59	142	5.0	23	33.3	20	13.3	32	9.2	12	3.7	12	2.3	7	1.1	36	4.8	
	60-69	545	19.4	27	39.1	62	41.3	123	35.4	74	22.6	55	10.6	82	12.5	122	16.4	
	70-79	953	33.9	12	17.4	49	32.7	131	37.8	120	36.7	157	30.2	222	33.9	262	35.1	
	80-89	789	28.0	4	5.8	14	9.3	43	12.4	86	26.3	186	35.8	228	34.8	228	30.6	
	90-99	287	10.2	0	0	3	2.0	13	3.7	26	8.0	77	14.8	96	14.7	72	9.7	
	100-109	65	2.3	0	0	0	0	2	0.6	5	1.5	28	5.4	13	2.0	17	2.3	
	110-119	12	0.4	0	0	0	0	0	0	1	0.3	5	1.0	3	0.5	3	0.4	
	120-129	7	0.2	0	0	0	0	1	0.3	3	0.9	0	0	3	0.5	0	0	
	130-139	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	140-	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

第31表の1-1 最低血圧の平均値及び標準偏差（性・年齢階級別）

	総数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	81.7	11.7	66.2	8.3	74.2	9.8	80.0	10.0	84.7	13.3	86.2	10.0	83.5	11.2	79.9	11.4
女性	77.0	11.5	63.0	9.2	68.1	8.8	71.1	10.0	76.6	11.7	81.4	10.9	80.3	10.2	77.2	11.0

注) 2回測定値の平均値

第31表の1-2 最高血圧の分布（性・年齢階級別）

	総数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上			
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%		
男性	総数		1,969	100	48	100	103	100	182	100	197	100	313	100	516	100	610	100
	2回の平均値(mmHg)																	
	-89	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	90-99	19	1.0	4	8.3	2	1.9	5	2.7	3	1.5	0	0	3	0.6	2	0.3	
	100-109	116	5.9	10	20.8	18	17.5	23	12.6	17	8.6	14	4.5	17	3.3	17	2.8	
	110-119	257	13.1	15	31.3	30	29.1	52	28.6	31	15.7	37	11.8	48	9.3	44	7.2	
	120-129	399	20.3	15	31.3	28	27.2	54	29.7	57	28.9	66	21.1	85	16.5	94	15.4	
	130-139	445	22.6	2	4.2	17	16.5	33	18.1	37	18.8	79	25.2	123	23.8	154	25.2	
	140-149	332	16.9	2	4.2	5	4.9	12	6.6	26	13.2	55	17.6	101	19.6	131	21.5	
	150-159	203	10.3	0	0	2	1.9	1	0.5	14	7.1	39	12.5	59	11.4	88	14.4	
	160-169	105	5.3	0	0	1	1.0	2	1.1	6	3.0	11	3.5	43	8.3	42	6.9	
	170-179	56	2.8	0	0	0	0	0	0	2	1.0	7	2.2	23	4.5	24	3.9	
	180-	37	1.9	0	0	0	0	0	0	4	2.0	5	1.6	14	2.7	14	2.3	
女性	総数		2,814	100	69	100	150	100	347	100	327	100	520	100	655	100	746	100
	2回の平均値(mmHg)																	
	-89	14	0.5	3	4.3	2	1.3	6	1.7	0	0	2	0.4	0	0	1	0.1	
	90-99	136	4.8	22	31.9	26	17.3	53	15.3	19	5.8	7	1.3	6	0.9	3	0.4	
	100-109	378	13.4	24	34.8	59	39.3	121	34.9	66	20.2	43	8.3	43	6.6	22	2.9	
	110-119	460	16.3	17	24.6	40	26.7	94	27.1	90	27.5	83	16.0	88	13.4	48	6.4	
	120-129	529	18.8	1	1.4	20	13.3	47	13.5	70	21.4	125	24.0	140	21.4	126	16.9	
	130-139	509	18.1	2	2.9	2	1.3	14	4.0	50	15.3	120	23.1	135	20.6	186	24.9	
	140-149	382	13.6	0	0	1	0.7	7	2.0	15	4.6	77	14.8	118	18.0	164	22.0	
	150-159	207	7.4	0	0	0	0	3	0.9	7	2.1	31	6.0	68	10.4	98	13.1	
	160-169	107	3.8	0	0	0	0	1	0.3	5	1.5	18	3.5	30	4.6	53	7.1	
	170-179	52	1.8	0	0	0	0	1	0.3	3	0.9	8	1.5	12	1.8	28	3.8	
	180-	40	1.4	0	0	0	0	0	0	2	0.6	6	1.2	15	2.3	17	2.3	

第31表の1-2 最高血圧の平均値及び標準偏差（性・年齢階級別）

	総数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	135.1	18.7	115.5	11.2	121.0	13.0	122.0	12.4	130.6	18.5	135.9	17.7	139.7	19.0	140.2	17.5
女性	128.6	20.2	103.7	10.5	108.2	10.3	110.9	12.8	120.7	16.4	130.5	17.3	134.5	18.6	140.0	17.9

注) 2回測定値の平均値

第31表の2-1 最低血圧の分布（性・年齢階級別）

	総数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
男性	総数																
	1,414	100	48	100	101	100	175	100	181	100	249	100	353	100	307	100	
	40mmHg未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	40-49	2	0.1	0	0	0	0	0	0	0	0	0	1	0.3	1	0.3	
	50-59	34	2.4	9	18.8	4	4.0	2	1.1	4	2.2	0	0	6	1.7	9	2.9
	60-69	161	11.4	22	45.8	32	31.7	20	11.4	16	8.8	11	4.4	27	7.6	33	10.7
	70-79	413	29.2	14	29.2	37	36.6	64	36.6	50	27.6	50	20.1	109	30.9	89	29.0
	80-89	481	34.0	3	6.3	23	22.8	63	36.0	53	29.3	114	45.8	111	31.4	114	37.1
	90-99	226	16.0	0	0	4	4.0	17	9.7	42	23.2	51	20.5	71	20.1	41	13.4
	100-109	77	5.4	0	0	1	1.0	9	5.1	12	6.6	20	8.0	21	5.9	14	4.6
	110-119	14	1.0	0	0	0	0	0	0	1	0.6	2	0.8	6	1.7	5	1.6
	120-129	4	0.3	0	0	0	0	0	0	1	0.6	1	0.4	1	0.3	1	0.3
	130-139	2	0.1	0	0	0	0	0	0	2	1.1	0	0	0	0	0	0
	140mmHg以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
女性	総数																
	2,131	100	69	100	145	100	337	100	315	100	434	100	486	100	345	100	
	40mmHg未満	1	0.0	0	0	0	0	1	0.3	0	0	0	0	0	0	0	
	40-49	7	0.3	3	4.3	1	0.7	1	0.3	0	0	0	0	1	0.2	1	0.3
	50-59	112	5.3	23	33.3	17	11.7	30	8.9	12	3.8	12	2.8	6	1.2	12	3.5
	60-69	462	21.7	27	39.1	62	42.8	117	34.7	74	23.5	51	11.8	69	14.2	62	18.0
	70-79	755	35.4	12	17.4	49	33.8	130	38.6	118	37.5	140	32.3	179	36.8	127	36.8
	80-89	561	26.3	4	5.8	14	9.7	43	12.8	80	25.4	152	35.0	163	33.5	105	30.4
	90-99	172	8.1	0	0	2	1.4	13	3.9	24	7.6	51	11.8	54	11.1	28	8.1
	100-109	46	2.2	0	0	0	0	1	0.3	3	1.0	24	5.5	10	2.1	8	2.3
	110-119	9	0.4	0	0	0	0	0	0	1	0.3	4	0.9	2	0.4	2	0.6
	120-129	6	0.3	0	0	0	0	1	0.3	3	1.0	0	0	2	0.4	0	0
	130-139	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	140mmHg以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

注) 妊婦・血圧を下げる薬服用者除外

第31表の2-1 最低血圧の平均値及び標準偏差（性・年齢階級別）

	総数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	81.5	11.8	66.2	8.3	74.0	9.8	79.8	9.9	83.7	12.7	85.5	10.0	82.8	11.6	81.0	11.9
女性	76.2	11.2	63.0	9.2	68.4	8.4	71.2	9.9	76.1	11.5	80.6	11.0	79.1	9.8	77.3	10.4

注) 妊婦・血圧を下げる薬服用者除外 2回測定値の平均値

第31表の2-2 最高血圧の分布（性・年齢階級別）

	総数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
男性	総数																
	1,414	100	48	100	101	100	175	100	181	100	249	100	353	100	307	100	
	90mmHg未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	90-99	18	1.3	4	8.3	2	2.0	5	2.9	3	1.7	0	0	2	0.6	2	0.7
	100-109	108	7.6	10	20.8	18	17.8	23	13.1	17	9.4	14	5.6	15	4.2	11	3.6
	110-119	228	16.1	15	31.3	30	29.7	52	29.7	29	16.0	30	12.0	39	11.0	33	10.7
	120-129	320	22.6	15	31.3	27	26.7	51	29.1	55	30.4	57	22.9	66	18.7	49	16.0
	130-139	303	21.4	2	4.2	17	16.8	30	17.1	30	16.6	60	24.1	91	25.8	73	23.8
	140-149	208	14.7	2	4.2	4	4.0	11	6.3	26	14.4	42	16.9	61	17.3	62	20.2
	150-159	113	8.0	0	0	2	2.0	1	0.6	12	6.6	27	10.8	32	9.1	39	12.7
	160-169	53	3.7	0	0	1	1.0	2	1.1	5	2.8	9	3.6	21	5.9	15	4.9
	170-179	35	2.5	0	0	0	0	0	0	2	1.1	5	2.0	14	4.0	14	4.6
	180mmHg以上	28	2.0	0	0	0	0	0	0	2	1.1	5	2.0	12	3.4	9	2.9
女性	総数																
	2,131	100	69	100	145	100	337	100	315	100	434	100	486	100	345	100	
	90mmHg未満	14	0.7	3	4.3	2	1.4	6	1.8	0	0	2	0.5	0	0	1	0.3
	90-99	133	6.2	22	31.9	26	17.9	52	15.4	19	6.0	7	1.6	6	1.2	1	0.3
	100-109	361	16.9	24	34.8	55	37.9	115	34.1	66	21.0	42	9.7	41	8.4	18	5.2
	110-119	424	19.9	17	24.6	40	27.6	92	27.3	89	28.3	80	18.4	74	15.2	32	9.3
	120-129	438	20.6	1	1.4	19	13.1	47	13.9	68	21.6	112	25.8	123	25.3	68	19.7
	130-139	336	15.8	2	2.9	2	1.4	14	4.2	43	13.7	92	21.2	94	19.3	89	25.8
	140-149	227	10.7	0	0	1	0.7	7	2.1	15	4.8	59	13.6	73	15.0	72	20.9
	150-159	101	4.7	0	0	0	0	2	0.6	6	1.9	18	4.1	42	8.6	33	9.6
	160-169	50	2.3	0	0	0	0	1	0.3	5	1.6	13	3.0	15	3.1	16	4.6
	170-179	26	1.2	0	0	0	0	1	0.3	2	0.6	5	1.2	8	1.6	10	2.9
	180mmHg以上	21	1.0	0	0	0	0	0	0	2	0.6	4	0.9	10	2.1	5	1.4

注) 妊婦・血圧を下げる薬服用者除外

第31表の2-2 最高血圧の平均値及び標準偏差（性・年齢階級別）

	総数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	132.4	19.0	115.5	11.2	120.8	12.9	121.6	12.4	129.3	17.3	135.0	18.4	137.7	19.8	138.7	19.0
女性	124.2	19.0	103.7	10.5	108.2	10.4	110.9	12.7	120.2	16.1	128.4	16.9	131.7	18.4	136.1	17.9

注) 妊婦・血圧を下げる薬服用者除外 2回測定値の平均値

第32表の1 血色素量の分布（性・年齢階級別）

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%		
男性	総数	1,812	100	101	100	178	100	185	100	292	100	483	100	573	100	258	100	241	100	181	100	118	100	33	100
	10g/dl未満	8	0.4	0	0	0	0	0	0	1	0.2	7	1.2	0	0	2	0.8	1	0.6	3	2.5	1	3.0		
	10-11未満	20	1.1	0	0	0	0	1	0.3	5	1.0	14	2.4	3	1.2	2	0.8	3	1.7	6	5.1	3	9.1		
	11-12	48	2.6	0	0	1	0.6	0	0	1	0.3	12	2.5	34	5.9	8	3.1	6	2.5	14	7.7	8	6.8		
	12-13	107	5.9	1	1.0	0	0	2	1.1	11	3.8	18	3.7	75	13.1	10	3.9	29	12.0	19	10.5	21	17.8		
	13-14	286	15.8	5	5.0	7	3.9	12	6.5	27	9.2	99	20.5	136	23.7	53	20.5	56	23.2	42	23.2	32	27.1		
	14-15	527	29.1	23	22.8	46	25.8	59	31.9	101	34.6	150	31.1	148	25.8	80	31.0	55	22.8	56	30.9	29	24.6		
	15-16	521	28.8	33	32.7	75	42.1	69	37.3	106	36.3	128	26.5	110	19.2	65	25.2	64	26.6	31	17.1	12	10.2		
	16-17	246	13.6	33	32.7	43	24.2	37	20.0	32	11.0	57	11.8	44	7.7	30	11.6	25	10.4	13	7.2	6	5.1		
	17-18	41	2.3	6	5.9	6	3.4	4	2.2	11	3.8	9	1.9	5	0.9	6	2.3	2	0.8	2	1.1	1	0.8		
	18g/dl以上	8	0.4	0	0	0	0	2	1.1	2	0.7	4	0.8	0	0	3	1.2	0	0	0	0	0	0		
女性	総数	2,617	100	140	100	337	100	316	100	500	100	620	100	704	100	297	100	284	100	228	100	127	100	65	100
	10g/dl未満	64	2.4	0	0	12	3.6	17	5.4	9	1.8	4	0.6	22	3.1	2	0.7	2	0.7	7	3.1	7	5.5		
	10-11未満	98	3.7	6	4.3	19	5.6	12	3.8	13	2.6	8	1.3	40	5.7	7	2.4	5	1.8	16	7.0	12	9.4		
	11-12	290	11.1	11	7.9	39	11.6	34	10.8	31	6.2	52	8.4	123	17.5	38	12.8	34	12.0	38	16.7	32	25.2		
	12-13	748	28.6	45	32.1	100	29.7	86	27.2	132	26.4	180	29.0	205	29.1	76	25.6	87	30.6	69	30.3	36	28.3		
	13-14	943	36.0	56	40.0	123	36.5	120	38.0	196	39.2	240	38.7	208	29.5	107	36.0	105	37.0	64	28.1	27	21.3		
	14-15	385	14.7	17	12.1	37	11.0	43	13.6	96	19.2	107	17.3	85	12.1	52	17.5	40	14.1	28	12.3	11	8.7		
	15-16	79	3.0	5	3.6	6	1.8	4	1.3	21	4.2	24	3.9	19	2.7	12	4.0	9	3.2	6	2.6	2	1.6		
	16-17	10	0.4	0	0	1	0.3	0	0	2	0.4	5	0.8	2	0.3	3	1.0	2	0.7	0	0	0	0		
	17-18	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	18g/dl以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		

第32表の1 血色素量の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	14.7	1.4	15.6	1.0	15.4	0.9	15.3	1.0	15.0	1.1	14.6	1.3	14.0	1.6	14.6	1.4	14.3	1.5	14.0	1.5	13.5	1.7	12.9	1.5
女性	12.9	1.2	13.0	1.0	12.8	1.3	12.8	1.3	13.2	1.2	13.2	1.1	12.7	1.3	13.1	1.2	13.1	1.1	12.6	1.3	12.3	1.4	12.2	1.5

第32表の2 血色素量の分布（性・年齢階級別）

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%		
男性	総数	1,794	100	101	100	177	100	185	100	289	100	482	100	560	100	258	100	237	100	177	100	114	100	32	100
	10g/dl未満	7	0.4	0	0	0	0	0	0	1	0.2	6	1.1	0	0	2	0.8	1	0.6	2	1.8	1	3.1		
	10-11未満	19	1.1	0	0	0	0	0	1	0.3	5	1.0	13	2.3	3	1.2	2	0.8	3	1.7	6	5.3	2	6.3	
	11-12	47	2.6	0	0	1	0.6	0	1	0.3	12	2.5	33	5.9	8	3.1	6	2.5	13	7.3	8	7.0	6	18.8	
	12-13	104	5.8	1	1.0	0	0	2	1.1	11	3.8	18	3.7	72	12.9	10	3.9	27	11.4	19	10.7	20	17.5	6	18.8
	13-14	281	15.7	5	5.0	7	4.0	12	6.5	26	9.0	99	20.5	132	23.6	53	20.5	54	22.8	41	23.2	31	27.2	6	18.8
	14-15	524	29.2	23	22.8	46	26.0	59	31.9	100	34.6	150	31.1	146	26.1	80	31.0	55	23.2	54	30.5	29	25.4	8	25.0
	15-16	518	28.9	33	32.7	74	41.8	69	37.3	105	36.3	127	26.3	110	19.6	65	25.2	64	27.0	31	17.5	12	10.5	3	9.4
	16-17	245	13.7	33	32.7	43	24.3	37	20.0	32	11.1	57	11.8	43	7.7	30	11.6	25	10.5	13	7.3	5	4.4	0	0
	17-18	41	2.3	6	5.9	6	3.4	4	2.2	11	3.8	9	1.9	5	0.9	6	2.3	2	0.8	2	1.1	1	0.9	0	0
	18g/dl以上	8	0.4	0	0	0	0	2	1.1	2	0.7	4	0.8	0	0	3	1.2	0	0	0	0	0	0	0	0
女性	総数	2,575	100	140	100	333	100	307	100	491	100	615	100	689	100	295	100	280	100	223	100	125	100	61	100
	10g/dl未満	59	2.3	0	0	10	3.0	16	5.2	9	1.8	4	0.7	20	2.9	2	0.7	2	0.7	5	2.2	7	5.6	6	9.8
	10-11未満	91	3.5	6	4.3	19	5.7	12	3.9	9	1.8	7	1.1	38	5.5	6	2.0	5	1.8	16	7.2	11	8.8	6	9.8
	11-12	283	11.0	11	7.9	38	11.4	33	10.7	29	5.9	52	8.5	120	17.4	38	12.9	34	12.1	38	17.0	31	24.8	17	27.9
	12-13	741	28.8	45	32.1	100	30.0	84	27.4	131	26.7	178	28.9	203	29.5	76	25.8	86	30.7	69	30.9	36	28.8	12	19.7
	13-14	928	36.0	56	40.0	122	36.6	116	37.8	194	39.5	238	38.7	202	29.3	106	35.9	102	36.4	61	27.4	27	21.6	12	19.7
	14-15	384	14.9	17	12.1	37	11.1	42	13.7	96	19.6	107	17.4	85	12.3	52	17.6	40	14.3	28	12.6	11	8.8	6	9.8
	15-16	79	3.1	5	3.6	6	1.8	4	1.3	21	4.3	24	3.9	19	2.8	12	4.1	9	3.2	6	2.7	2	1.6	2	3.3
	16-17	10	0.4	0	0	1	0.3	0	0	2	0.4	5	0.8	2	0.3	3	1.0	2	0.7	0	0	0	0	0	0
	17-18	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	18g/dl以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

注) 貧血治療のための薬服用者除外

第32表の2 血色素量の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	14.7	1.4	15.6	1.0	15.4	0.9	15.3	1.0	15.0	1.1	14.6	1.3	14.0	1.6	14.6	1.4	14.3	1.5	14.0	1.5	13.5	1.7	13.0	1.5
女性	13.0	1.2	13.0	1.0	12.8	1.2	12.8	1.3	13.2	1.2	13.2	1.1	12.7	1.3	13.1	1.2	13.1	1.1	12.7	1.3	12.3	1.4	12.2	1.5

注) 貧血治療のための薬服用者除外

第33表 赤血球数の分布（性・年齢階級別）

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%		
男性	総数	1,812	100	101	100	178	100	185	100	292	100	483	100	573	100	258	100	241	100	181	100	118	100	33	100
	400万个/μl未満	130	7.2	0	0	2	1.1	0	0	5	1.7	26	5.4	97	16.9	14	5.4	24	10.0	32	17.7	29	24.6	12	36.4
	400-450未満	419	23.1	3	3.0	5	2.8	21	11.4	53	18.2	134	27.7	203	35.4	81	31.4	86	35.7	57	31.5	47	39.8	13	39.4
	450-500	769	42.4	29	28.7	65	36.5	87	47.0	160	54.8	225	46.6	203	35.4	110	42.6	88	36.5	71	39.2	36	30.5	8	24.2
	500-550	427	23.6	52	51.5	87	48.9	72	38.9	69	23.6	82	17.0	65	11.3	45	17.4	38	15.8	21	11.6	6	5.1	0	0
	550-600	63	3.5	16	15.8	19	10.7	4	2.2	5	1.7	14	2.9	5	0.9	7	2.7	5	2.1	0	0	0	0	0	0
	600-650	4	0.2	1	1.0	0	0	1	0.5	0	0	2	0.4	0	0	1	0.4	0	0	0	0	0	0	0	
	650万个/μl以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
女性	総数	2,617	100	140	100	337	100	316	100	500	100	620	100	704	100	297	100	284	100	228	100	127	100	65	100
	400万个/μl未満	412	15.7	6	4.3	34	10.1	28	8.9	41	8.2	85	13.7	218	31.0	57	19.2	56	19.7	76	33.3	53	41.7	33	50.8
	400-450未満	1,324	50.6	78	55.7	170	50.4	171	54.1	247	49.4	327	52.7	331	47.0	141	47.5	144	50.7	107	46.9	57	44.9	23	35.4
	450-500	788	30.1	52	37.1	120	35.6	106	33.5	182	36.4	188	30.3	140	19.9	88	29.6	80	28.2	39	17.1	14	11.0	7	10.8
	500-550	91	3.5	4	2.9	13	3.9	11	3.5	29	5.8	19	3.1	15	2.1	10	3.4	4	1.4	6	2.6	3	2.4	2	3.1
	550-600	2	0.1	0	0	0	0	0	0	1	0.2	1	0.2	0	0	1	0.3	0	0	0	0	0	0	0	0
	600-650	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	650万个/μl以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

第33表 赤血球数の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	470.7	47.8	513.5	35.6	507.5	34.2	490.2	34.4	478.4	37.2	466.5	44.2	445.1	48.6	464.6	44.1	456.3	45.3	445.1	48.4	430.3	50.7	417.1	40.5
女性	434.3	37.1	443.8	28.5	440.7	33.6	439.8	30.1	444.5	33.9	436.2	34.7	418.1	41.6	432.5	38.5	429.5	34.7	416.1	42.0	405.1	44.1	400.5	48.8

第34表 ヘマトクリット値の分布（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総数	1,812	100	101	100	178	100	185	100	292	100	483	100	573	100	258	100	241	100	181	100	118	100	33	100		
	25%未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	25-30未満	5	0.3	0	0	0	0	0	0	0	0	0	0	5	0.9	0	0	2	0.8	1	0.6	2	1.7	0	0	0	0
	30-35	26	1.4	0	0	0	0	0	0	1	0.3	6	1.2	19	3.3	4	1.6	3	1.2	4	2.2	9	7.6	3	9.1		
	35-40	128	7.1	0	0	2	1.1	0	0	9	3.1	21	4.3	96	16.8	13	5.0	32	13.3	29	16.0	23	19.5	12	36.4		
	40-45	639	35.3	20	19.8	28	15.7	54	29.2	107	36.6	190	39.3	240	41.9	101	39.1	96	39.8	80	44.2	51	43.2	13	39.4		
	45-50	840	46.4	62	61.4	119	66.9	107	57.8	151	51.7	223	46.2	178	31.1	115	44.6	87	36.1	57	31.5	29	24.6	5	15.2		
	50-55	167	9.2	18	17.8	28	15.7	23	12.4	24	8.2	40	8.3	34	5.9	22	8.5	21	8.7	9	5.0	4	3.4	0	0		
	55%以上	7	0.4	1	1.0	1	0.6	1	0.5	0	0	3	0.6	1	0.2	3	1.2	0	0	1	0.6	0	0	0	0		
女性	総数	2,617	100	140	100	337	100	316	100	500	100	620	100	704	100	297	100	284	100	228	100	127	100	65	100		
	25%未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	25-30未満	15	0.6	0	0	3	0.9	1	0.3	2	0.4	1	0.2	8	1.1	0	0	0	0	4	1.8	2	1.6	2	3.1		
	30-35	138	5.3	3	2.1	20	5.9	26	8.2	18	3.6	14	2.3	57	8.1	13	4.4	9	3.2	17	7.5	21	16.5	10	15.4		
	35-40	805	30.8	52	37.1	119	35.3	96	30.4	114	22.8	166	26.8	258	36.6	84	28.3	96	33.8	84	36.8	50	39.4	28	43.1		
	40-45	1,412	54.0	72	51.4	176	52.2	176	55.7	303	60.6	369	59.5	316	44.9	164	55.2	147	51.8	100	43.9	50	39.4	19	29.2		
	45-50	240	9.2	13	9.3	18	5.3	17	5.4	62	12.4	69	11.1	61	8.7	35	11.8	30	10.6	21	9.2	4	3.1	6	9.2		
	50-55	7	0.3	0	0	1	0.3	0	0	1	0.2	1	0.2	4	0.6	1	0.3	2	0.7	2	0.9	0	0	0	0		
	55%以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		

第34表 ヘマトクリット値の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	45.2	4.0	47.7	3.0	47.2	2.7	46.7	2.7	45.7	3.1	45.2	3.7	43.3	4.5	45.2	3.9	44.2	4.2	43.4	4.3	42.0	4.9	40.7	4.4
女性	40.7	3.4	40.8	2.9	40.2	3.4	40.3	3.4	41.4	3.2	41.4	3.1	40.1	3.8	41.3	3.4	41.0	3.2	40.0	4.0	38.8	3.9	38.6	4.3

第35表 平均赤血球容積 (MCV) の分布 (性・年齢階級別)

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総数	1,812	100	101	100	178	100	185	100	292	100	483	100	573	100	258	100	241	100	181	100	118	100	33	100
	70歳未満	1	0.1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	70-75	1	0.1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	75-80	5	0.3	1	1.0	1	0.6	0	0	0	0	1	0.2	2	0.3	0	0	0	0	0	0	1	0.8	1	3.0
	80-85	15	0.8	0	0	2	1.1	2	1.1	2	0.7	4	0.8	5	0.9	1	0.4	2	0.8	1	0.6	1	0.8	1	3.0
	85-90	139	7.7	19	18.8	33	18.5	22	11.9	21	7.2	22	4.6	22	3.8	12	4.7	10	4.1	7	3.9	5	4.2	0	0
	90-95	594	32.8	51	50.5	88	49.4	70	37.8	109	37.3	137	28.4	139	24.3	73	28.3	70	29.0	41	22.7	25	21.2	3	9.1
	95-100	647	35.7	25	24.8	42	23.6	57	30.8	110	37.7	187	38.7	226	39.4	95	36.8	84	34.9	80	44.2	45	38.1	17	51.5
	100-105	319	17.6	5	5.0	10	5.6	28	15.1	42	14.4	98	20.3	136	23.7	55	21.3	59	24.5	37	20.4	30	25.4	10	30.3
	105-110	67	3.7	0	0	2	1.1	5	2.7	6	2.1	25	5.2	29	5.1	16	6.2	8	3.3	11	6.1	9	7.6	1	3.0
	110-115	20	1.1	0	0	0	0	1	0.5	1	0.3	8	1.7	10	1.7	6	2.3	5	2.1	4	2.2	1	0.8	0	0
	115-120	1	0.1	0	0	0	0	0	0	0	0	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0
	120歳以上	3	0.2	0	0	0	0	0	0	1	0.3	1	0.2	1	0.2	0	0	0	0	0	0	1	0.8	0	0
女性	総数	2,617	100	140	100	337	100	316	100	500	100	620	100	704	100	297	100	284	100	228	100	127	100	65	100
	70歳未満	13	0.5	0	0	5	1.5	4	1.3	4	0.8	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	70-75	17	0.6	2	1.4	4	1.2	9	2.8	2	0.4	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	75-80	27	1.0	1	0.7	9	2.7	9	2.8	5	1.0	3	0.5	0	0	1	0.3	0	0	0	0	0	0	0	0
	80-85	78	3.0	8	5.7	22	6.5	21	6.6	10	2.0	4	0.6	13	1.8	3	1.0	5	1.8	1	0.4	5	3.9	2	3.1
	85-90	300	11.5	27	19.3	70	20.8	46	14.6	73	14.6	48	7.7	36	5.1	24	8.1	15	5.3	14	6.1	6	4.7	1	1.5
	90-95	1,031	39.4	62	44.3	139	41.2	116	36.7	218	43.6	255	41.1	241	34.2	102	34.3	109	38.4	79	34.6	36	28.3	17	26.2
	95-100	875	33.4	33	23.6	69	20.5	90	28.5	156	31.2	240	38.7	287	40.8	126	42.4	111	39.1	89	39.0	54	42.5	33	50.8
	100-105	240	9.2	6	4.3	16	4.7	19	6.0	30	6.0	58	9.4	111	15.8	32	10.8	36	12.7	39	17.1	25	19.7	11	16.9
	105-110	31	1.2	1	0.7	3	0.9	1	0.3	2	0.4	11	1.8	13	1.8	8	2.7	7	2.5	4	1.8	1	0.8	1	1.5
	110-115	2	0.1	0	0	0	0	0	0	0	0	1	0.2	1	0.1	1	0.3	0	0	1	0.4	0	0	0	0
	115-120	3	0.1	0	0	0	0	1	0.3	0	0	0	0	2	0.3	0	0	1	0.4	1	0.4	0	0	0	0
	120歳以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

第35表 平均赤血球容積 (MCV) の平均値及び標準偏差 (性・年齢階級別)

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	96.3	5.3	93.1	4.0	93.2	4.3	95.4	4.9	95.8	4.8	97.2	5.3	97.5	5.4	97.6	5.3	97.0	5.6	97.8	4.9	97.9	5.8	97.5	5.5
女性	94.0	5.6	92.0	5.1	91.4	6.5	91.8	7.1	93.3	5.4	95.2	4.2	96.0	4.5	95.6	4.4	95.7	4.4	96.3	4.4	96.0	4.7	96.6	4.3

第36表 平均赤血球血色素量（MCH）の分布（性・年齢階級別）

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総数	1,812	100	101	100	178	100	185	100	292	100	483	100	573	100	258	100	241	100	181	100	118	100	33	100
	20pg未満	2	0.1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0.4	0	0	1	0.8	0	0
	20-25未満	7	0.4	0	0	1	0.6	0	0	1	0.3	2	0.4	3	0.5	0	0	2	0.8	0	0	0	0	1	3.0
	25-27	20	1.1	2	2.0	2	1.1	0	0	2	0.7	6	1.2	8	1.4	2	0.8	3	1.2	1	0.6	2	1.7	2	6.1
	27-29	109	6.0	8	7.9	19	10.7	12	6.5	12	4.1	27	5.6	31	5.4	16	6.2	10	4.1	11	6.1	10	8.5	0	0
	29-31	679	37.5	66	65.3	98	55.1	85	45.9	115	39.4	146	30.2	169	29.5	77	29.8	74	30.7	52	28.7	31	26.3	12	36.4
	31-33	751	41.4	25	24.8	54	30.3	62	33.5	125	42.8	213	44.1	272	47.5	111	43.0	113	46.9	85	47.0	59	50.0	15	45.5
	33-35	212	11.7	0	0	4	2.2	23	12.4	30	10.3	79	16.4	76	13.3	46	17.8	32	13.3	29	16.0	12	10.2	3	9.1
	35-40	32	1.8	0	0	0	0	3	1.6	7	2.4	10	2.1	12	2.1	6	2.3	6	2.5	3	1.7	3	2.5	0	0
	40pg以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
女性	総数	2,617	100	140	100	337	100	316	100	500	100	620	100	704	100	297	100	284	100	228	100	127	100	65	100
	20pg未満	17	0.6	0	0	5	1.5	7	2.2	5	1.0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	20-25未満	76	2.9	7	5.0	22	6.5	24	7.6	11	2.2	4	0.6	8	1.1	2	0.7	3	1.1	1	0.4	3	2.4	1	1.5
	25-27	80	3.1	7	5.0	22	6.5	22	7.0	10	2.0	7	1.1	12	1.7	5	1.7	3	1.1	4	1.8	4	3.1	1	1.5
	27-29	400	15.3	25	17.9	84	24.9	55	17.4	92	18.4	70	11.3	74	10.5	34	11.4	23	8.1	31	13.6	13	10.2	7	10.8
	29-31	1,371	52.4	83	59.3	148	43.9	137	43.4	284	56.8	355	57.3	364	51.7	166	55.9	160	56.3	115	50.4	59	46.5	30	46.2
	31-33	621	23.7	16	11.4	50	14.8	67	21.2	89	17.8	174	28.1	225	32.0	82	27.6	87	30.6	72	31.6	40	31.5	26	40.0
	33-35	46	1.8	2	1.4	6	1.8	2	0.6	9	1.8	9	1.5	18	2.6	7	2.4	7	2.5	4	1.8	7	5.5	0	0
	35-40	6	0.2	0	0	0	0	2	0.6	0	0	1	0.2	3	0.4	1	0.3	1	0.4	1	0.4	1	0.8	0	0
	40pg以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

第36表 平均赤血球血色素量（MCH）の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	31.2	1.8	30.3	1.2	30.4	1.3	31.2	1.5	31.3	1.7	31.5	1.8	31.4	2.0	31.5	1.7	31.4	2.1	31.5	1.7	31.4	2.2	30.9	2.2
女性	29.9	2.1	29.4	2.0	29.0	2.7	29.1	2.9	29.7	2.1	30.3	1.5	30.4	1.5	30.4	1.6	30.5	1.5	30.4	1.5	30.4	1.9	30.4	1.5

第 37 表 平均赤血球血色素濃度 (MCHC) の分布 (性・年齢階級別)

	総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%		
男 性	総 数		1,812	100	101	100	178	100	185	100	292	100	483	100	573	100	258	100	241	100	181	100	118	100	33	100
	22%未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	22-24未満	1	0.1	0	0	0	0	0	0	0	0	0	0	1	0.2	0	0	0	0	0	0	1	0.8	0	0	
	24-26	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	26-28	3	0.2	0	0	0	0	0	0	0	0	1	0.2	2	0.3	0	0	1	0.4	0	0	0	0	1	3.0	
	28-30	29	1.6	1	1.0	0	0	1	0.5	4	1.4	11	2.3	12	2.1	6	2.3	5	2.1	4	2.2	3	2.5	0	0	
	30-32	504	27.8	24	23.8	47	26.4	36	19.5	54	18.5	142	29.4	201	35.1	88	34.1	64	26.6	70	38.7	51	43.2	16	48.5	
	32-34	1,185	65.4	66	65.3	122	68.5	135	73.0	211	72.3	311	64.4	340	59.3	155	60.1	162	67.2	100	55.2	62	52.5	16	48.5	
	34%以上	90	5.0	10	9.9	9	5.1	13	7.0	23	7.9	18	3.7	17	3.0	9	3.5	9	3.7	7	3.9	1	0.8	0	0	
女 性	総 数		2,617	100	140	100	337	100	316	100	500	100	620	100	704	100	297	100	284	100	228	100	127	100	65	100
	22%未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	22-24未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	24-26	4	0.2	0	0	0	0	1	0.3	3	0.6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	26-28	17	0.6	1	0.7	5	1.5	4	1.3	3	0.6	2	0.3	2	0.3	1	0.3	0	0	2	0.9	0	0	0	0	
	28-30	155	5.9	7	5.0	30	8.9	36	11.4	22	4.4	28	4.5	32	4.5	16	5.4	11	3.9	10	4.4	8	6.3	3	4.6	
	30-32	1,218	46.5	60	42.9	142	42.1	137	43.4	228	45.6	279	45.0	372	52.8	142	47.8	132	46.5	131	57.5	70	55.1	39	60.0	
	32-34	1,204	46.0	71	50.7	156	46.3	136	43.0	239	47.8	306	49.4	296	42.0	138	46.5	140	49.3	84	36.8	49	38.6	23	35.4	
	34%以上	19	0.7	1	0.7	4	1.2	2	0.6	5	1.0	5	0.8	2	0.3	0	0	1	0.4	1	0.4	0	0	0	0	

第 37 表 平均赤血球血色素濃度 (MCHC) の平均値及び標準偏差 (性・年齢階級別)

	総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男 性	32.5	1.1	32.6	1.1	32.6	1.0	32.7	0.9	32.7	1.0	32.4	1.1	32.2	1.1	32.3	1.1	32.4	1.1	32.2	1.0	32.0	1.2	31.7	1.2
女 性	31.8	1.1	31.9	1.1	31.7	1.3	31.6	1.4	31.8	1.2	31.9	1.0	31.7	1.0	31.8	1.0	31.9	0.9	31.6	1.0	31.6	1.0	31.5	1.0

第38表の1 フェリチンの分布（性・年齢階級別）

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総数	1,819	100	101	100	178	100	185	100	292	100	485	100	578	100	259	100	243	100	183	100	119	100	33	100
	1.1ng/ml未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1.1-5未満	10	0.5	0	0	1	0.6	0	0	1	0.3	4	0.8	4	0.7	2	0.8	2	0.8	0	0	2	1.7	0	0
	5-10	42	2.3	2	2.0	0	0	0	0	10	3.4	12	2.5	18	3.1	7	2.7	8	3.3	4	2.2	4	3.4	2	6.1
	10-15	35	1.9	0	0	0	0	1	0.5	2	0.7	13	2.7	19	3.3	9	3.5	4	1.6	7	3.8	8	6.7	0	0
	15-20	34	1.9	1	1.0	2	1.1	1	0.5	5	1.7	6	1.2	19	3.3	3	1.2	8	3.3	4	2.2	6	5.0	1	3.0
	20-25	43	2.4	2	2.0	2	1.1	3	1.6	1	0.3	11	2.3	24	4.2	8	3.1	10	4.1	7	3.8	3	2.5	4	12.1
	25-30	43	2.4	0	0	0	0	4	2.2	11	3.8	8	1.6	20	3.5	4	1.5	6	2.5	8	4.4	5	4.2	1	3.0
	30-35	36	2.0	0	0	1	0.6	1	0.5	8	2.7	10	2.1	16	2.8	7	2.7	6	2.5	7	3.8	3	2.5	0	0
	35-40	49	2.7	4	4.0	2	1.1	7	3.8	4	1.4	11	2.3	21	3.6	6	2.3	10	4.1	6	3.3	4	3.4	1	3.0
	40-45	42	2.3	3	3.0	3	1.7	1	0.5	7	2.4	11	2.3	17	2.9	7	2.7	8	3.3	4	2.2	3	2.5	2	6.1
	45-50	45	2.5	6	5.9	5	2.8	1	0.5	4	1.4	11	2.3	18	3.1	6	2.3	6	2.5	6	3.2	4	3.4	2	6.1
	50-60	100	5.5	5	5.0	7	3.9	10	5.4	17	5.8	24	4.9	37	6.4	11	4.2	16	6.6	11	6.0	7	5.9	3	9.1
	60-70	110	6.0	11	10.9	3	1.7	13	7.0	14	4.8	29	6.0	40	6.9	16	6.2	18	7.4	12	6.6	8	6.7	2	6.1
	70-80	111	6.1	10	9.9	19	10.7	16	8.6	12	4.1	34	7.0	20	3.5	17	6.6	9	3.7	5	2.7	5	4.2	1	3.0
	80-90	85	4.7	8	7.9	16	9.0	8	4.3	16	5.5	16	3.3	21	3.6	8	3.1	6	2.5	5	2.7	7	5.9	3	9.1
	90-100	106	5.8	5	5.0	12	6.7	14	7.6	21	7.2	26	5.4	28	4.8	17	6.6	10	4.1	10	5.5	7	5.9	1	3.0
	100-150	381	20.9	20	19.8	45	25.3	42	22.7	66	22.6	106	21.9	102	17.6	59	22.8	43	17.7	35	19.1	20	16.8	4	12.1
	150-200	245	13.5	17	16.8	31	17.4	22	11.9	34	11.6	69	14.2	72	12.5	35	13.5	34	14.0	22	12.0	11	9.2	5	15.2
	200ng/ml以上	302	16.6	7	6.9	29	16.3	41	22.2	59	20.2	84	17.3	82	14.2	37	14.3	39	16.0	30	16.4	12	10.1	1	3.0
女性	総数	2,631	100	144	100	338	100	317	100	502	100	622	100	708	100	300	100	285	100	230	100	128	100	65	100
	1.1ng/ml未満	1	0.0	0	0	0	0	1	0.3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1.1-5未満	165	6.3	16	11.1	58	17.2	59	18.6	21	4.2	4	0.6	7	1.0	2	0.7	3	1.1	2	0.9	1	0.8	1	1.5
	5-10	201	7.6	18	12.5	55	16.3	53	16.7	28	5.6	11	1.8	36	5.1	7	2.3	13	4.6	7	3.0	12	9.4	4	6.2
	10-15	160	6.1	23	16.0	40	11.8	37	11.7	16	3.2	10	1.6	34	4.8	4	1.3	8	2.8	12	5.2	9	7.0	5	7.7
	15-20	135	5.1	15	10.4	36	10.7	20	6.3	19	3.8	9	1.4	36	5.1	5	1.7	9	3.2	13	5.7	7	5.5	7	10.8
	20-25	134	5.1	10	6.9	21	6.2	28	8.8	17	3.4	18	2.9	40	5.6	9	3.0	11	3.9	15	6.5	11	8.6	3	4.6
	25-30	151	5.7	14	9.7	32	9.5	25	7.9	28	5.6	20	3.2	32	4.5	8	2.7	14	4.9	13	5.7	3	2.3	2	3.1
	30-35	129	4.9	2	1.4	17	5.0	18	5.7	21	4.2	31	5.0	40	5.6	17	5.7	14	4.9	15	6.5	7	5.5	4	6.2
	35-40	114	4.3	8	5.6	13	3.8	13	4.1	32	6.4	24	3.9	24	3.4	10	3.3	8	2.8	8	3.5	8	6.3	0	0
	40-45	124	4.7	7	4.9	19	5.6	13	4.1	22	4.4	37	5.9	26	3.7	17	5.7	10	3.5	9	3.9	3	2.3	4	6.2
	45-50	110	4.2	8	5.6	13	3.8	7	2.2	26	5.2	23	3.7	33	4.7	12	4.0	9	3.2	10	4.3	9	7.0	5	7.7
	50-60	220	8.4	11	7.6	13	3.8	18	5.7	55	11.0	61	9.8	62	8.8	29	9.7	27	9.5	21	9.1	13	10.2	1	1.5
	60-70	171	6.5	5	3.5	3	0.9	11	3.5	45	9.0	51	8.2	56	7.9	25	8.3	19	6.7	20	8.7	9	7.0	8	12.3
	70-80	161	6.1	5	3.5	5	1.5	5	1.6	37	7.4	65	10.5	44	6.2	29	9.7	18	6.3	10	4.3	9	7.0	7	10.8
	80-90	110	4.2	2	1.4	5	1.5	2	0.6	25	5.0	41	6.6	35	4.9	16	5.3	16	5.6	11	4.8	3	2.3	5	7.7
	90-100	93	3.5	0	0	3	0.9	1	0.3	17	3.4	40	6.4	32	4.5	18	6.0	21	7.4	10	4.3	1	0.8	0	0
	100-150	290	11.0	0	0	3	0.9	4	1.3	64	12.7	109	17.5	110	15.5	51	17.0	54	18.9	38	16.5	14	10.9	4	6.2
	150-200	86	3.3	0	0	1	0.3	1	0.3	16	3.2	42	6.8	26	3.7	28	9.3	14	4.9	9	3.9	2	1.6	1	1.5
	200ng/ml以上	76	2.9	0	0	1	0.3	1	0.3	13	2.6	26	4.2	35	4.9	13	4.3	17	6.0	7	3.0	7	5.5	4	6.2

第38表の1 フェリチンの平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	127.4	114.4	108.6	70.4	134.8	82.7	142.4	117.6	143.2	160.4	129.6	106.0	113.8	104.9	123.8	105.2	118.9	105.8	122.8	115.1	98.7	93.2	80.9	60.7
女性	60.5	79.3	26.5	20.9	24.5	25.8	25.1	30.2	64.0	51.7	83.4	57.6	77.9	124.8	85.7	62.1	81.3	61.5	70.5	61.6	90.9	262.9	63.2	61.8

第38表の2 フェリチンの分布（性・年齢階級別）

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総数	1,801	100	101	100	177	100	185	100	289	100	484	100	565	100	259	100	239	100	179	100	115	100	32	100
	1.1ng/ml未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1.1-5未満	10	0.6	0	0	1	0.6	0	0	1	0.3	4	0.8	4	0.7	2	0.8	2	0.8	0	0	2	1.7	0	0
	5-10	41	2.3	2	2.0	0	0	0	0	10	3.5	12	2.5	17	3.0	7	2.7	7	2.9	4	2.2	4	3.5	2	6.3
	10-15	35	1.9	0	0	0	0	1	0.5	2	0.7	13	2.7	19	3.4	9	3.5	4	1.7	7	3.9	8	7.0	0	0
	15-20	34	1.9	1	1.0	2	1.1	1	0.5	5	1.7	6	1.2	19	3.4	3	1.2	8	3.3	4	2.2	6	5.2	1	3.1
	20-25	43	2.4	2	2.0	2	1.1	3	1.6	1	0.3	11	2.3	24	4.2	8	3.1	10	4.2	7	3.9	3	2.6	4	12.5
	25-30	40	2.2	0	0	0	0	4	2.2	11	3.8	8	1.7	17	3.0	4	1.5	6	2.5	6	3.4	5	4.3	0	0
	30-35	35	1.9	0	0	1	0.6	1	0.5	7	2.4	10	2.1	16	2.8	7	2.7	6	2.5	7	3.9	3	2.6	0	0
	35-40	49	2.7	4	4.0	2	1.1	7	3.8	4	1.4	11	2.3	21	3.7	6	2.3	10	4.2	6	3.4	4	3.5	1	3.1
	40-45	41	2.3	3	3.0	3	1.7	1	0.5	7	2.4	11	2.3	16	2.8	7	2.7	8	3.3	3	1.7	3	2.6	2	6.3
	45-50	45	2.5	6	5.9	5	2.8	1	0.5	4	1.4	11	2.3	18	3.2	6	2.3	6	2.5	6	3.4	4	3.5	2	6.3
	50-60	100	5.6	5	5.0	7	4.0	10	5.4	17	5.9	24	5.0	37	6.5	11	4.2	16	6.7	11	6.1	7	6.1	3	9.4
	60-70	110	6.1	11	10.9	3	1.7	13	7.0	14	4.8	29	6.0	40	7.1	16	6.2	18	7.5	12	6.7	8	7.0	2	6.3
	70-80	111	6.2	10	9.9	19	10.7	16	8.6	12	4.2	34	7.0	20	3.5	17	6.6	9	3.8	5	2.8	5	4.3	1	3.1
	80-90	83	4.6	8	7.9	16	9.0	8	4.3	16	5.5	16	3.3	19	3.4	8	3.1	6	2.5	4	2.2	6	5.2	3	9.4
	90-100	106	5.9	5	5.0	12	6.8	14	7.6	21	7.3	26	5.4	28	5.0	17	6.6	10	4.2	10	5.6	7	6.1	1	3.1
	100-150	379	21.0	20	19.8	45	25.4	42	22.7	66	22.8	106	21.9	100	17.7	59	22.8	42	17.6	35	19.6	19	16.5	4	12.5
	150-200	241	13.4	17	16.8	31	17.5	22	11.9	33	11.4	68	14.0	70	12.4	35	13.5	33	13.8	22	12.3	10	8.7	5	15.6
	200ng/ml以上	298	16.5	7	6.9	28	15.8	41	22.2	58	20.1	84	17.4	80	14.2	37	14.3	38	15.9	30	16.8	11	9.6	1	3.1
女性	総数	2,578	100	140	100	327	100	308	100	493	100	617	100	693	100	298	100	281	100	225	100	126	100	61	100
	1.1ng/ml未満	1	0.0	0	0	0	0	1	0.3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1.1-5未満	159	6.2	16	11.4	55	16.8	58	18.8	19	3.9	4	0.6	7	1.0	2	0.7	3	1.1	2	0.9	1	0.8	1	1.6
	5-10	192	7.4	17	12.1	52	15.9	51	16.6	25	5.1	11	1.8	36	5.2	7	2.3	13	4.6	7	3.1	12	9.5	4	6.6
	10-15	157	6.1	23	16.4	38	11.6	37	12.0	16	3.2	10	1.6	33	4.8	4	1.3	7	2.5	12	5.3	9	7.1	5	8.2
	15-20	130	5.0	15	10.7	36	11.0	19	6.2	18	3.7	9	1.5	33	4.8	5	1.7	9	3.2	12	5.3	7	5.6	5	8.2
	20-25	131	5.1	10	7.1	21	6.4	26	8.4	17	3.4	18	2.9	39	5.6	9	3.0	11	3.9	15	6.7	11	8.7	2	3.3
	25-30	148	5.7	13	9.3	32	9.8	24	7.8	28	5.7	20	3.2	31	4.5	8	2.7	14	5.0	12	5.3	3	2.4	2	3.3
	30-35	127	4.9	2	1.4	17	5.2	17	5.5	21	4.3	31	5.0	39	5.6	17	5.7	13	4.6	15	6.7	7	5.6	4	6.6
	35-40	113	4.4	8	5.7	13	4.0	12	3.9	32	6.5	24	3.9	24	3.5	10	3.4	8	2.8	8	3.6	8	6.3	0	0
	40-45	120	4.7	6	4.3	19	5.8	13	4.2	22	4.5	37	6.0	23	3.3	17	5.7	10	3.6	7	3.1	2	1.6	4	6.6
	45-50	109	4.2	8	5.7	13	4.0	7	2.3	26	5.3	23	3.7	32	4.6	12	4.0	9	3.2	10	4.4	8	6.3	5	8.2
	50-60	216	8.4	11	7.9	12	3.7	18	5.8	54	11.0	61	9.9	60	8.7	29	9.7	26	9.3	20	8.9	13	10.3	1	1.6
	60-70	170	6.6	5	3.6	3	0.9	11	3.6	45	9.1	50	8.1	56	8.1	24	8.1	19	6.8	20	8.9	9	7.1	8	13.1
	70-80	157	6.1	4	2.9	4	1.2	5	1.6	37	7.5	64	10.4	43	6.2	29	9.7	18	6.4	10	4.4	9	7.1	6	9.8
	80-90	110	4.3	2	1.4	5	1.5	2	0.6	25	5.1	41	6.6	35	5.1	16	5.4	16	5.7	11	4.9	3	2.4	5	8.2
	90-100	91	3.5	0	0	3	0.9	1	0.3	16	3.2	39	6.3	32	4.6	18	6.0	21	7.5	10	4.4	1	0.8	0	0
	100-150	286	11.1	0	0	3	0.9	4	1.3	63	12.8	107	17.3	109	15.7	50	16.8	53	18.9	38	16.9	14	11.1	4	6.6
	150-200	86	3.3	0	0	1	0.3	1	0.3	16	3.2	42	6.8	26	3.8	28	9.4	14	5.0	9	4.0	2	1.6	1	1.6
	200ng/ml以上	75	2.9	0	0	0	0	1	0.3	13	2.6	26	4.2	35	5.1	13	4.4	17	6.0	7	3.1	7	5.6	4	6.6

注) 妊婦・貧血治療のための薬服用者除外

第38表の2 フェリチンの平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	127.5	114.7	108.6	70.4	134.4	82.8	142.4	117.6	143.3	161.1	129.5	106.1	114.1	105.5	123.8	105.2	118.7	106.1	124.5	115.8	97.1	93.9	82.6	60.9
女性	60.9	79.8	26.3	20.8	24.0	23.2	25.2	30.6	64.5	51.6	83.3	57.8	78.6	126.0	85.7	62.2	81.7	61.7	71.2	62.0	91.7	264.9	65.1	63.0

注) 妊婦・貧血治療のための薬服用者除外

第39表 総たんぱく質値の分布（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総数	1,818	100	100	100	178	100	185	100	292	100	485	100	578	100	259	100	243	100	183	100	119	100	33	100		
	6.0g/dl未満	4	0.2	0	0	0	0	0	0	0	0	0	0	4	0.7	0	0	1	0.4	1	0.5	2	1.7	0	0		
	6.0-6.5未満	21	1.2	0	0	0	0	2	1.1	3	1.0	5	1.0	11	1.9	4	1.5	4	1.6	3	1.6	4	3.4	0	0		
	6.5-7.0	214	11.8	4	4.0	9	5.1	19	10.3	33	11.3	57	11.8	92	15.9	26	10.0	29	11.9	29	15.8	26	21.8	8	24.2		
	7.0-7.5	676	37.2	27	27.0	45	25.3	68	36.8	119	40.8	195	40.2	222	38.4	105	40.5	100	41.2	70	38.3	38	31.9	14	42.4		
	7.5-8.0	635	34.9	42	42.0	78	43.8	72	38.9	99	33.9	169	34.8	175	30.3	95	36.7	79	32.5	53	29.0	35	29.4	8	24.2		
	8.0-8.5	234	12.9	22	22.0	39	21.9	23	12.4	33	11.3	48	9.9	69	11.9	22	8.5	28	11.5	24	13.1	14	11.8	3	9.1		
	8.5g/dl以上	34	1.9	5	5.0	7	3.9	1	0.5	5	1.7	11	2.3	5	0.9	7	2.7	2	0.8	3	1.6	0	0	0	0		
女性	総数	2,631	100	144	100	338	100	317	100	502	100	622	100	708	100	300	100	285	100	230	100	128	100	65	100		
	6.0g/dl未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	6.0-6.5未満	18	0.7	1	0.7	4	1.2	1	0.3	1	0.2	6	1.0	5	0.7	4	1.3	1	0.4	0	0	2	1.6	2	3.1		
	6.5-7.0	213	8.1	8	5.6	18	5.3	30	9.5	35	7.0	45	7.2	77	10.9	20	6.7	32	11.2	25	10.9	14	10.9	6	9.2		
	7.0-7.5	986	37.5	44	30.6	133	39.3	132	41.6	151	30.1	231	37.1	295	41.7	111	37.0	114	40.0	91	39.6	55	43.0	35	53.8		
	7.5-8.0	1,013	38.5	68	47.2	131	38.8	119	37.5	206	41.0	248	39.9	241	34.0	118	39.3	102	35.8	88	38.3	36	28.1	15	23.1		
	8.0-8.5	352	13.4	18	12.5	47	13.9	33	10.4	99	19.7	83	13.3	72	10.2	43	14.3	27	9.5	23	10.0	18	14.1	4	6.2		
	8.5g/dl以上	49	1.9	5	3.5	5	1.5	2	0.6	10	2.0	9	1.4	18	2.5	4	1.3	9	3.2	3	1.3	3	2.3	3	4.6		

第39表 総たんぱく質値の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	7.5	0.5	7.7	0.4	7.7	0.4	7.5	0.4	7.4	0.4	7.4	0.5	7.4	0.5	7.5	0.5	7.4	0.5	7.4	0.5	7.3	0.5	7.3	0.4
女性	7.5	0.4	7.6	0.4	7.5	0.4	7.4	0.4	7.6	0.4	7.5	0.4	7.4	0.5	7.5	0.4	7.5	0.5	7.5	0.4	7.4	0.5	7.4	0.5

第40表 アルブミン値の分布（性・年齢階級別）

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総数	1,818	100	100	100	178	100	185	100	292	100	485	100	578	100	259	100	243	100	183	100	119	100	33	100
	3.0g/dl未満	2	0.1	0	0	0	0	0	0	0	0	0	0	2	0.3	0	0	0	0	1	0.5	1	0.8	0	0
	3.0-3.2未満	4	0.2	0	0	0	0	0	0	0	0	0	4	0.7	0	0	3	1.2	1	0.5	0	0	0	0	
	3.2-3.4	3	0.2	0	0	0	0	0	0	0	0	1	0.2	2	0.3	0	0	1	0.4	1	0.5	0	0	0	
	3.4-3.6	9	0.5	0	0	0	0	0	0	1	0.3	0	0	8	1.4	0	0	1	0.4	3	1.6	2	1.7	2	6.1
	3.6-3.8	17	0.9	0	0	1	0.6	0	0	2	0.7	2	0.4	12	2.1	2	0.8	3	1.2	3	1.6	3	2.5	3	9.1
	3.8-4.0	68	3.7	0	0	0	0	1	0.5	1	0.3	17	3.5	49	8.5	14	5.4	11	4.5	11	6.0	19	16.0	8	24.2
	4.0-4.2	194	10.7	1	1.0	3	1.7	5	2.7	16	5.5	50	10.3	119	20.6	29	11.2	32	13.2	45	24.6	33	27.7	9	27.3
	4.2-4.4	365	20.1	2	2.0	3	1.7	20	10.8	48	16.4	131	27.0	161	27.9	68	26.3	74	30.5	56	30.6	25	21.0	6	18.2
	4.4-4.6	473	26.0	14	14.0	22	12.4	44	23.8	95	32.5	150	30.9	148	25.6	79	30.5	82	33.7	38	20.8	25	21.0	3	9.1
	4.6-4.8	396	21.8	24	24.0	71	39.9	72	38.9	79	27.1	92	19.0	58	10.0	46	17.8	25	10.3	22	12.0	9	7.6	2	6.1
	4.8-5.0	208	11.4	36	36.0	49	27.5	32	17.3	41	14.0	37	7.6	13	2.2	19	7.3	9	3.7	2	1.1	2	1.7	0	0
	5.0-5.2	62	3.4	15	15.0	24	13.5	9	4.9	9	3.1	3	0.6	2	0.3	1	0.4	2	0.8	0	0	0	0	0	0
	5.2g/dl以上	17	0.9	8	8.0	5	2.8	2	1.1	0	0	2	0.4	0	0	1	0.4	0	0	0	0	0	0	0	0
女性	総数	2,631	100	144	100	338	100	317	100	502	100	622	100	708	100	300	100	285	100	230	100	128	100	65	100
	3.0g/dl未満	1	0.0	0	0	0	0	0	0	0	0	0	0	0	0	1	0.3	0	0	0	0	0	0	0	0
	3.0-3.2未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	3.2-3.4	1	0.0	1	0.7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	3.4-3.6	8	0.3	0	0	3	0.9	0	0	1	0.2	0	0	4	0.6	0	0	1	0.4	2	0.9	1	0.8	0	0
	3.6-3.8	17	0.6	0	0	3	0.9	0	0	1	0.2	4	0.6	9	1.3	2	0.7	2	0.7	3	1.3	1	0.8	3	4.6
	3.8-4.0	82	3.1	2	1.4	6	1.8	10	3.2	5	1.0	13	2.1	46	6.5	8	2.7	11	3.9	17	7.4	10	7.8	8	12.3
	4.0-4.2	299	11.4	7	4.9	20	5.9	26	8.2	21	4.2	78	12.5	147	20.8	50	16.7	51	17.9	45	19.6	30	23.4	21	32.3
	4.2-4.4	620	23.6	17	11.8	67	19.8	81	25.6	93	18.5	147	23.6	215	30.4	76	25.3	87	30.5	69	30.0	42	32.8	17	26.2
	4.4-4.6	829	31.5	38	26.4	113	33.4	107	33.8	185	36.9	199	32.0	187	26.4	83	27.7	83	29.1	58	25.2	35	27.3	11	16.9
	4.6-4.8	524	19.9	45	31.3	91	26.9	71	22.4	120	23.9	125	20.1	72	10.2	56	18.7	34	11.9	25	10.9	8	6.3	5	7.7
	4.8-5.0	210	8.0	27	18.8	30	8.9	19	6.0	63	12.5	46	7.4	25	3.5	20	6.7	13	4.6	11	4.8	1	0.8	0	0
	5.0-5.2	34	1.3	4	2.8	4	1.2	3	0.9	11	2.2	9	1.4	3	0.4	4	1.3	3	1.1	0	0	0	0	0	0
	5.2g/dl以上	6	0.2	3	2.1	1	0.3	0	0	2	0.4	0	0	0	0	0	0	0	0	0	0	0	0	0	0

第40表 アルブミン値の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	4.44	0.32	4.79	0.25	4.73	0.24	4.59	0.23	4.51	0.25	4.40	0.26	4.24	0.31	4.38	0.26	4.31	0.29	4.23	0.30	4.17	0.31	4.05	0.29
女性	4.41	0.27	4.56	0.28	4.46	0.26	4.42	0.23	4.49	0.23	4.41	0.26	4.29	0.26	4.38	0.28	4.33	0.25	4.29	0.26	4.25	0.23	4.18	0.25

第41表の1 ヘモグロビンA1cの分布(性・年齢階級別)

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	3.9%以下	6	0.3	1	1.0	0	0	1	0.5	2	0.7	2	0.4	0	0	1	0.4	0	0	0	0	0	0	0	0
	4.0-4.1	3	0.2	1	1.0	0	0	0	0	0	0	0	0.3	0	0	0	0	0	2	1.1	0	0	0	0	
	4.2-4.3	4	0.2	0	0	2	1.1	0	0	0	0	1	0.2	1	0.4	0	0	0	0	0	1	0.8	0	0	
	4.4-4.5	34	1.9	14	13.9	6	3.4	3	1.6	4	1.4	4	0.8	3	0.5	0	0	3	1.2	0	0	0	0	0	
	4.6-4.7	121	6.7	18	17.8	23	12.9	12	6.5	18	6.2	31	6.4	19	3.3	18	7.0	8	3.3	5	2.8	4	3.4	2	6.1
	4.8-4.9	252	13.9	28	27.7	49	27.5	42	22.7	41	14.0	40	8.3	52	9.1	21	8.1	28	11.6	13	7.2	7	5.9	4	12.1
	5.0-5.1	298	16.4	21	20.8	41	23.0	38	20.5	44	15.1	73	15.1	81	14.1	30	11.6	29	12.0	26	14.4	19	16.1	7	21.2
	5.2	184	10.1	7	6.9	20	11.2	26	14.1	35	12.0	51	10.5	45	7.9	28	10.9	21	8.7	17	9.4	5	4.2	2	6.1
	5.3	184	10.1	7	6.9	18	10.1	19	10.3	39	13.4	42	8.7	59	10.3	21	8.1	26	10.8	23	12.7	7	5.9	3	9.1
	5.4	132	7.3	1	1.0	6	3.4	10	5.4	16	5.5	47	9.7	52	9.1	29	11.2	25	10.4	17	9.4	10	8.5	0	0
	5.5	111	6.1	1	1.0	3	1.7	9	4.9	18	6.2	31	6.4	49	8.6	19	7.4	20	8.3	13	7.2	13	11.0	3	9.1
	5.6	94	5.2	1	1.0	5	2.8	7	3.8	9	3.1	36	7.4	36	6.3	23	8.9	13	5.4	10	5.5	11	9.3	2	6.1
	5.7	64	3.5	0	0	2	1.1	2	1.1	10	3.4	23	4.8	27	4.7	11	4.3	11	4.6	10	5.5	6	5.1	0	0
	5.8	51	2.8	0	0	0	0	1	0.5	12	4.1	18	3.7	20	3.5	9	3.5	5	2.1	6	3.3	7	5.9	2	6.1
	5.9	35	1.9	0	0	1	0.6	4	2.2	4	1.4	10	2.1	16	2.8	5	1.9	4	1.7	8	4.4	3	2.5	1	3.0
	6.0	29	1.6	1	1.0	0	0	2	1.1	6	2.1	8	1.7	12	2.1	5	1.9	7	2.9	2	1.1	2	1.7	1	3.0
	6.1	23	1.3	0	0	0	0	0	0	3	1.0	8	1.7	12	2.1	4	1.6	2	0.8	5	2.8	4	3.4	1	3.0
	6.2-6.3	39	2.2	0	0	1	0.6	1	0.5	3	1.0	17	3.5	17	3.0	8	3.1	5	2.1	6	3.3	4	3.4	2	6.1
	6.4-6.5	23	1.3	0	0	0	0	1	0.5	2	0.7	5	1.0	15	2.6	2	0.8	7	2.9	5	2.8	3	2.5	0	0
	6.6-6.7	26	1.4	0	0	0	0	0	0	6	2.1	8	1.7	12	2.1	4	1.6	5	2.1	3	1.7	2	1.7	2	6.1
	6.8-6.9	15	0.8	0	0	0	0	0	0	3	1.0	4	0.8	8	1.4	2	0.8	3	1.2	2	1.1	3	2.5	0	0
	7.0-7.1	17	0.9	0	0	0	0	0	0	4	1.4	6	1.2	7	1.2	5	1.9	4	1.7	1	0.6	2	1.7	0	0
	7.2-7.3	6	0.3	0	0	0	0	0	0	1	0.3	0	0	5	0.9	0	0	3	1.2	1	0.6	1	0.8	0	0
	7.4-7.5	14	0.8	0	0	0	0	4	2.2	1	0.3	3	0.6	6	1.0	2	0.8	2	0.8	3	1.7	0	0	1	3.0
	7.6-7.7	9	0.5	0	0	0	0	0	0	2	0.7	2	0.4	5	0.9	0	0	2	0.8	1	0.6	2	1.7	0	0
	7.8-7.9	6	0.3	0	0	0	0	1	0.5	0	0	3	0.6	2	0.3	3	1.2	1	0.4	0	0	1	0.8	0	0
	8.0-8.1	4	0.2	0	0	0	0	0	0	1	0.3	1	0.2	2	0.3	0	0	2	0.8	0	0	0	0	0	0
	8.2-8.3	3	0.2	0	0	0	0	0	0	0	0	2	0.4	1	0.2	1	0.4	0	0	1	0.6	0	0	0	0
	8.4-8.5	4	0.2	0	0	0	0	0	0	1	0.3	1	0.2	2	0.3	0	0	2	0.8	0	0	0	0	0	0
	8.6-8.7	1	0.1	0	0	0	0	0	0	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0	0	0
	8.8-8.9	1	0.1	0	0	0	0	0	0	0	0	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0
	9.0%以上	20	1.1	0	0	1	0.6	2	1.1	7	2.4	6	1.2	4	0.7	5	1.9	2	0.8	1	0.6	1	0.8	0	0
	(再掲) 6.1%以上	211	11.6	0	0	2	1.1	9	4.9	34	11.6	67	13.8	99	17.3	37	14.3	41	17.0	29	16.0	23	19.5	6	18.2
	総数	1,813	100	101	100	178	100	185	100	292	100	484	100	573	100	258	100	241	100	181	100	118	100	33	100

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上		
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数
女性	3.9%以下	2	0.1	1	0.7	1	0.3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	4.0-4.1	3	0.1	0	0	1	0.3	0	0	2	0.4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	4.2-4.3	11	0.4	2	1.4	1	0.3	1	0.3	1	0.2	3	0.5	3	0.4	2	0.7	0	0	2	0.9	0	0	1	1.5	0
	4.4-4.5	53	2.0	11	7.8	11	3.3	7	2.2	10	2.0	8	1.3	6	0.8	2	0.7	0	0	4	1.8	2	1.6	0	0	0
	4.6-4.7	172	6.6	24	17.0	47	13.9	31	9.8	24	4.8	19	3.1	27	3.8	10	3.4	8	2.8	10	4.4	8	6.3	1	1.5	0
	4.8-4.9	359	13.7	35	24.8	84	24.9	67	21.2	58	11.6	54	8.7	61	8.6	24	8.1	25	8.5	15	6.6	13	10.2	8	12.3	0
	5.0-5.1	451	17.2	26	18.4	93	27.5	66	20.9	88	17.6	97	15.6	81	11.5	44	14.8	27	9.5	32	14.0	15	11.7	7	10.8	0
	5.2	278	10.6	9	6.4	30	8.9	39	12.3	47	9.4	71	11.5	82	11.6	34	11.4	31	10.9	31	13.6	13	10.2	7	10.8	0
	5.3	291	11.1	17	12.1	26	7.7	28	8.9	61	12.2	74	11.9	85	12.0	31	10.4	39	13.7	30	13.2	10	7.8	6	9.2	0
	5.4	257	9.8	5	3.5	17	5.0	22	7.0	59	11.8	76	12.3	78	11.0	35	11.8	34	11.9	22	9.6	13	10.2	9	13.8	0
	5.5	201	7.7	8	5.7	10	3.0	18	5.7	45	9.0	57	9.2	63	8.9	21	7.1	24	8.4	16	7.0	17	13.3	6	9.2	0
	5.6	149	5.7	1	0.7	11	3.3	12	3.8	36	7.2	37	6.0	52	7.4	22	7.4	21	7.4	16	7.0	10	7.8	5	7.7	0
	5.7	85	3.2	1	0.7	1	0.3	10	3.2	19	3.8	20	3.2	34	4.8	12	4.0	12	4.2	11	4.8	5	3.9	6	9.2	0
	5.8	65	2.5	0	0	1	0.3	4	1.3	9	1.8	23	3.7	28	4.0	12	4.0	12	4.2	3	1.3	10	7.8	3	4.6	0
	5.9	49	1.9	0	0	1	0.3	3	0.9	9	1.8	9	1.5	27	3.8	6	2.0	7	2.5	14	6.1	4	3.1	2	3.1	0
	6.0	27	1.0	0	0	0	0	1	0.3	2	0.4	10	1.6	14	2.0	7	2.4	8	2.8	5	2.2	1	0.8	0	0	0
	6.1	32	1.2	0	0	0	0	2	0.6	6	1.2	10	1.6	14	2.0	2	0.7	3	1.1	8	3.5	1	0.8	2	3.1	0
	6.2-6.3	33	1.3	0	0	1	0.3	2	0.6	3	0.6	13	2.1	14	2.0	7	2.4	9	3.2	4	1.8	1	0.8	0	0	0
	6.4-6.5	22	0.8	0	0	0	0	0	0	8	1.6	6	1.0	8	1.1	4	1.3	5	1.8	1	0.4	2	1.6	0	0	0
	6.6-6.7	18	0.7	1	0.7	0	0	0	0	1	0.2	10	1.6	6	0.8	7	2.4	3	1.1	1	0.4	0	0	2	3.1	0
	6.8-6.9	10	0.4	0	0	0	0	1	0.3	0	0	6	1.0	3	0.4	3	1.0	2	0.7	0	0	1	0.8	0	0	0
	7.0-7.1	8	0.3	0	0	0	0	2	0.6	0	0	3	0.5	3	0.4	1	0.3	1	0.4	1	0.4	1	0.8	0	0	0
	7.2-7.3	7	0.3	0	0	0	0	0	0	3	0.6	0	0	4	0.6	0	0	4	1.4	0	0	0	0	0	0	0
	7.4-7.5	3	0.1	0	0	0	0	0	0	1	0.2	1	0.2	1	0.1	1	0.3	1	0.4	0	0	0	0	0	0	0
	7.6-7.7	4	0.2	0	0	0	0	0	0	1	0.2	1	0.2	2	0.3	1	0.3	1	0.4	1	0.4	0	0	0	0	0
	7.8-7.9	5	0.2	0	0	0	0	0	0	0	0	3	0.5	2	0.3	3	1.0	1	0.4	0	0	1	0.8	0	0	0
	8.0-8.1	4	0.2	0	0	0	0	0	0	0	0	1	0.2	3	0.4	1	0.3	2	0.7	1	0.4	0	0	0	0	0
	8.2-8.3	5	0.2	0	0	1	0.3	0	0	0	0	4	0.6	0	0	4	1.3	0	0	0	0	0	0	0	0	0
	8.4-8.5	2	0.1	0	0	0	0	0	0	1	0.2	0	0	1	0.1	0	0	1	0.4	0	0	0	0	0	0	0
	8.6-8.7	1	0.0	0	0	0	0	0	0	0	0	1	0.2	0	0	0	0	0	0	0	0	0	0	0	0	0
	8.8-8.9	5	0.2	0	0	0	0	0	0	3	0.6	0	0	2	0.3	0	0	2	0.7	0	0	0	0	0	0	0
	9.0%以上	9	0.3	0	0	1	0.3	0	0	3	0.6	3	0.5	2	0.3	1	0.3	2	0.7	0	0	0	0	0	0	0
	(再掲) 6.1%以上	168	6.4	1	0.7	3	0.9	7	2.2	30	6.0	62	10.0	65	9.2	35	11.8	37	13.0	17	7.5	7	5.5	4	6.2	0
	総数	2,621	100	141	100	338	100	316	100	500	100	620	100	706	100	297	100	285	100	228	100	128	100	65	100	0

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上				
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
総数	3.9%以下	8	0.2	2	0.8	1	0.2	1	0.2	2	0.3	2	0.2	0	0	1	0.2	0	0	0	0	0	0	0	0	0	0	0
	4.0-4.1	6	0.1	1	0.4	1	0.2	0	0	2	0.3	0	0	2	0.2	0	0	0	0	2	0.5	0	0	0	0	0	0	0
	4.2-4.3	15	0.3	2	0.8	3	0.6	1	0.2	1	0.1	4	0.4	4	0.3	3	0.5	0	0	2	0.5	1	0.4	1	1.0	1	1.0	
	4.4-4.5	87	2.0	25	10.3	17	3.3	10	2.0	14	1.8	12	1.1	9	0.7	2	0.4	3	0.6	4	1.0	2	0.8	0	0	0	0	0
	4.6-4.7	293	6.6	42	17.4	70	13.6	43	8.6	42	5.3	50	4.5	46	3.6	28	5.0	16	3.0	15	3.7	12	4.9	3	3.1	3	3.1	
	4.8-4.9	611	13.8	63	26.0	133	25.8	109	21.8	99	12.5	94	8.5	113	8.8	45	8.1	53	10.1	28	6.8	20	8.1	12	12.2	12	12.2	
	5.0-5.1	749	16.9	47	19.4	134	26.0	104	20.8	132	16.7	170	15.4	162	12.7	74	13.3	56	10.6	58	14.2	34	13.8	14	14.3	14	14.3	
	5.2	462	10.4	16	6.6	50	9.7	65	13.0	82	10.4	122	11.1	127	9.9	62	11.2	52	9.9	48	11.7	18	7.3	9	9.2	9	9.2	
	5.3	475	10.7	24	9.9	44	8.5	47	9.4	100	12.6	116	10.5	144	11.3	52	9.4	65	12.4	53	13.0	17	6.9	9	9.2	9	9.2	
	5.4	389	8.8	6	2.5	23	4.5	32	6.4	75	9.5	123	11.1	130	10.2	64	11.5	59	11.2	39	9.5	23	9.3	9	9.2	9	9.2	
	5.5	312	7.0	9	3.7	13	2.5	27	5.4	63	8.0	88	8.0	112	8.8	40	7.2	44	8.4	29	7.1	30	12.2	9	9.2	9	9.2	
	5.6	243	5.5	2	0.8	16	3.1	19	3.8	45	5.7	73	6.6	88	6.9	45	8.1	34	6.5	26	6.4	21	8.5	7	7.1	7	7.1	
	5.7	149	3.4	1	0.4	3	0.6	12	2.4	29	3.7	43	3.9	61	4.8	23	4.1	23	4.4	21	5.1	11	4.5	6	6.1	6	6.1	
	5.8	116	2.6	0	0	1	0.2	5	1.0	21	2.7	41	3.7	48	3.8	21	3.8	17	3.2	9	2.2	17	6.9	5	5.1	5	5.1	
	5.9	84	1.9	0	0	2	0.4	7	1.4	13	1.6	19	1.7	43	3.4	11	2.0	11	2.1	22	5.4	7	2.8	3	3.1	3	3.1	
	6.0	56	1.3	1	0.4	0	0	3	0.6	8	1.0	18	1.6	26	2.0	12	2.2	15	2.9	7	1.7	3	1.2	1	1.0	1	1.0	
	6.1	55	1.2	0	0	0	0	2	0.4	9	1.1	18	1.6	26	2.0	6	1.1	5	1.0	13	3.2	5	2.0	3	3.1	3	3.1	
	6.2-6.3	72	1.6	0	0	2	0.4	3	0.6	6	0.8	30	2.7	31	2.4	15	2.7	14	2.7	10	2.4	5	2.0	2	2.0	2	2.0	
	6.4-6.5	45	1.0	0	0	0	0	1	0.2	10	1.3	11	1.0	23	1.8	6	1.1	12	2.3	6	1.5	5	2.0	0	0	0	0	
	6.6-6.7	44	1.0	1	0.4	0	0	0	0	7	0.9	18	1.6	18	1.4	11	2.0	8	1.5	4	1.0	2	0.8	4	4.1	4	4.1	
	6.8-6.9	25	0.6	0	0	0	0	1	0.2	3	0.4	10	0.9	11	0.9	5	0.9	5	1.0	2	0.5	4	1.6	0	0	0	0	
	7.0-7.1	25	0.6	0	0	0	0	2	0.4	4	0.5	9	0.8	10	0.8	6	1.1	5	1.0	2	0.5	3	1.2	0	0	0	0	
	7.2-7.3	13	0.3	0	0	0	0	0	0	4	0.5	0	0	9	0.7	0	0	7	1.3	1	0.2	1	0.4	0	0	0	0	
7.4-7.5	17	0.4	0	0	0	0	4	0.8	2	0.3	4	0.4	7	0.5	3	0.5	3	0.6	3	0.7	0	0	1	1.0	1	1.0		
7.6-7.7	13	0.3	0	0	0	0	0	0	3	0.4	3	0.3	7	0.5	1	0.2	3	0.6	2	0.5	2	0.8	0	0	0	0		
7.8-7.9	11	0.2	0	0	0	0	1	0.2	0	0	6	0.5	4	0.3	6	1.1	2	0.4	0	0	2	0.8	0	0	0	0		
8.0-8.1	8	0.2	0	0	0	0	0	0	1	0.1	2	0.2	5	0.4	1	0.2	4	0.8	1	0.2	0	0	0	0	0	0		
8.2-8.3	8	0.2	0	0	1	0.2	0	0	0	0	6	0.5	1	0.1	5	0.9	0	0	1	0.2	0	0	0	0	0	0		
8.4-8.5	6	0.1	0	0	0	0	0	0	2	0.3	1	0.1	3	0.2	0	0	3	0.6	0	0	0	0	0	0	0	0		
8.6-8.7	2	0.0	0	0	0	0	0	0	0	0	2	0.2	0	0	1	0.2	0	0	0	0	0	0	0	0	0	0		
8.8-8.9	6	0.1	0	0	0	0	0	0	3	0.4	0	0	3	0.2	0	0	3	0.6	0	0	0	0	0	0	0	0		
9.0%以上	29	0.7	0	0	2	0.4	2	0.4	10	1.3	9	0.8	6	0.5	6	1.1	4	0.8	1	0.2	1	0.4	0	0	0	0		
(再掲) 6.1%以上	379	8.5	1	0.4	5	1.0	16	3.2	64	8.1	129	11.7	164	12.8	72	13.0	78	14.8	46	11.2	30	12.2	10	10.2	10	10.2		
総数	4,434	100	242	100	516	100	501	100	792	100	1,104	100	1,279	100	555	100	526	100	409	100	246	100	98	100	98	100		

第41表の1 ヘモグロビンA1cの平均値及び標準偏差(性・年齢階級別)

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	5.4	0.8	4.9	0.3	5.0	0.4	5.2	0.7	5.5	1.0	5.5	0.8	5.6	0.8	5.6	0.9	5.6	0.8	5.5	0.7	5.6	0.8	5.5	0.6
女性	5.3	0.6	5.0	0.4	5.0	0.4	5.1	0.4	5.4	0.7	5.4	0.7	5.5	0.6	5.5	0.7	5.6	0.8	5.4	0.5	5.4	0.5	5.4	0.4
総数	5.4	0.7	4.9	0.3	5.0	0.4	5.2	0.5	5.4	0.8	5.5	0.7	5.5	0.7	5.5	0.8	5.6	0.8	5.4	0.6	5.5	0.7	5.4	0.5

第41表の2 ヘモグロビンA1cの分布（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上				
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
男性	3.9%以下	6	0.4	1	1.0	0	0	1	0.6	2	0.7	2	0.5	0	0	1	0.4	0	0	0	0	0	0	0	0	0	0	0
	4.0-4.1	3	0.2	1	1.0	0	0	0	0	0	0	0	0	2	0.4	0	0	0	0	2	1.2	0	0	0	0	0	0	
	4.2-4.3	4	0.2	0	0	2	1.1	0	0	0	0	1	0.2	1	0.2	1	0.4	0	0	0	0	0	1	1.0	0	0	0	
	4.4-4.5	34	2.0	14	13.9	6	3.4	3	1.7	4	1.4	4	0.9	3	0.6	0	0	3	1.4	0	0	0	0	0	0	0	0	
	4.6-4.7	121	7.2	18	17.8	23	13.0	12	6.6	18	6.5	31	7.0	19	3.7	18	7.7	8	3.8	5	3.0	4	3.8	2	6.5	2	6.5	
	4.8-4.9	252	15.0	28	27.7	49	27.7	42	23.2	41	14.9	40	9.1	52	10.2	21	9.0	28	13.5	13	7.9	7	6.7	4	12.9	4	12.9	
	5.0-5.1	294	17.5	21	20.8	40	22.6	38	21.0	44	15.9	71	16.1	80	15.7	30	12.9	29	13.9	25	15.2	19	18.3	7	22.6	7	22.6	
	5.2	182	10.8	7	6.9	20	11.3	26	14.4	35	12.7	51	11.6	43	8.5	28	12.0	21	10.1	15	9.1	5	4.8	2	6.5	2	6.5	
	5.3	182	10.8	7	6.9	18	10.2	18	9.9	39	14.1	42	9.5	58	11.4	21	9.0	26	12.5	23	13.9	6	5.8	3	9.7	3	9.7	
	5.4	128	7.6	1	1.0	6	3.4	10	5.5	16	5.8	45	10.2	50	9.8	28	12.0	24	11.5	16	9.7	10	9.6	0	0	0	0	
	5.5	109	6.5	1	1.0	3	1.7	9	5.0	18	6.5	31	7.0	47	9.3	19	8.2	19	9.1	12	7.3	13	12.5	3	9.7	3	9.7	
	5.6	93	5.5	1	1.0	5	2.8	7	3.9	9	3.3	35	8.0	36	7.1	22	9.4	13	6.3	10	6.1	11	10.6	2	6.5	2	6.5	
	5.7	61	3.6	0	0	2	1.1	2	1.1	10	3.6	21	4.8	26	5.1	10	4.3	10	4.8	10	6.1	6	5.8	0	0	0	0	
	5.8	49	2.9	0	0	0	0	1	0.6	12	4.3	16	3.6	20	3.9	8	3.4	5	2.4	6	3.6	7	6.7	2	6.5	2	6.5	
	5.9	31	1.8	0	0	1	0.6	4	2.2	4	1.4	9	2.0	13	2.6	5	2.1	4	1.9	5	3.0	3	2.9	1	3.2	1	3.2	
	6.0	22	1.3	1	1.0	0	0	2	1.1	3	1.1	6	1.4	10	2.0	3	1.3	5	2.4	2	1.2	2	1.9	1	3.2	1	3.2	
	6.1	22	1.3	0	0	0	0	0	0	3	1.1	8	1.8	11	2.2	4	1.7	2	1.0	5	3.0	3	2.9	1	3.2	1	3.2	
	6.2-6.3	24	1.4	0	0	1	0.6	1	0.6	1	0.4	10	2.3	11	2.2	5	2.1	2	1.0	5	3.0	2	1.9	2	6.5	2	6.5	
	6.4-6.5	10	0.6	0	0	0	0	1	0.6	1	0.4	1	0.2	7	1.4	0	0	2	1.0	4	2.4	1	1.0	0	0	0	0	
	6.6-6.7	13	0.8	0	0	0	0	0	0	3	1.1	5	1.1	5	1.0	2	0.9	1	0.5	2	1.2	1	1.0	1	3.2	1	3.2	
	6.8-6.9	9	0.5	0	0	0	0	0	0	3	1.1	3	0.7	3	0.6	1	0.4	1	0.5	1	0.6	1	1.0	0	0	0	0	
	7.0-7.1	4	0.2	0	0	0	0	0	0	1	0.4	1	0.2	2	0.4	1	0.4	0	0	1	0.6	1	1.0	0	0	0	0	
	7.2-7.3	4	0.2	0	0	0	0	0	0	1	0.4	0	0	3	0.6	0	0	2	1.0	1	0.6	0	0	0	0	0	0	
	7.4-7.5	6	0.4	0	0	0	0	3	1.7	0	0	1	0.2	2	0.4	1	0.4	1	0.5	1	0.6	0	0	0	0	0	0	
	7.6-7.7	2	0.1	0	0	0	0	0	0	0	0	0	0	2	0.4	0	0	1	0.5	1	0.6	0	0	0	0	0	0	
	7.8-7.9	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	8.0-8.1	2	0.1	0	0	0	0	0	0	1	0.4	1	0.2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	8.2-8.3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	8.4-8.5	3	0.2	0	0	0	0	0	0	1	0.4	1	0.2	1	0.2	0	0	1	0.5	0	0	0	0	0	0	0	0	0
	8.6-8.7	1	0.1	0	0	0	0	0	0	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0	0	0	0	0	0
8.8-8.9	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
9.0%以上	12	0.7	0	0	1	0.6	1	0.6	6	2.2	3	0.7	1	0.2	3	1.3	0	0	0	0	1	1.0	0	0	0	0	0	
	(再掲) 6.1%以上	112	6.7	0	0	2	1.1	6	3.3	21	7.6	35	8.0	48	9.4	18	7.7	13	6.3	21	12.7	10	9.6	4	12.9	4	12.9	
	総数	1,683	100	101	100	177	100	181	100	276	100	440	100	508	100	233	100	208	100	165	100	104	100	31	100	31	100	

注) インスリン注射または血糖を下げる薬服用者除外

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
		女性	3.9%以下	2	0.1	1	0.7	1	0.3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	4.0-4.1	3	0.1	0	0	1	0.3	0	0	2	0.4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	4.2-4.3	11	0.4	2	1.4	1	0.3	1	0.3	1	0.2	3	0.5	3	0.5	2	0.7	0	0	2	0.9	0	0	1	1.7		
	4.4-4.5	53	2.1	11	7.9	11	3.3	7	2.2	10	2.1	8	1.4	6	0.9	2	0.7	0	0	4	1.8	2	1.7	0	0		
	4.6-4.7	172	6.8	24	17.1	47	13.9	31	9.9	24	4.9	19	3.3	27	4.1	10	3.6	8	3.1	10	4.5	8	6.9	1	1.7		
	4.8-4.9	357	14.2	35	25.0	84	24.9	67	21.3	58	12.0	53	9.1	60	9.2	24	8.8	25	9.7	14	6.4	13	11.2	8	13.8		
	5.0-5.1	450	17.9	26	18.6	93	27.6	66	21.0	88	18.1	97	16.6	80	12.3	44	16.1	27	10.5	32	14.5	15	12.9	6	10.3		
	5.2	276	11.0	9	6.4	30	8.9	39	12.4	46	9.5	71	12.2	81	12.4	34	12.4	31	12.0	31	14.1	13	11.2	6	10.3		
	5.3	291	11.6	17	12.1	26	7.7	28	8.9	61	12.6	74	12.7	85	13.0	31	11.3	39	15.1	30	13.6	10	8.6	6	10.3		
	5.4	255	10.2	5	3.6	17	5.0	22	7.0	59	12.2	76	13.0	76	11.7	35	12.8	34	13.2	22	10.0	12	10.3	8	13.8		
	5.5	198	7.9	8	5.7	10	3.0	17	5.4	45	9.3	57	9.8	61	9.4	21	7.7	23	8.9	16	7.3	16	13.8	6	10.3		
	5.6	147	5.9	1	0.7	11	3.3	12	3.8	36	7.4	37	6.3	50	7.7	22	8.0	20	7.8	16	7.3	9	7.8	5	8.6		
	5.7	84	3.3	1	0.7	1	0.3	10	3.2	19	3.9	20	3.4	33	5.1	12	4.4	11	4.3	11	7.0	5	4.3	6	10.3		
	5.8	59	2.3	0	0	1	0.3	4	1.3	9	1.9	21	3.6	24	3.7	11	4.0	12	4.7	2	0.9	9	7.8	1	1.7		
	5.9	45	1.8	0	0	1	0.3	3	1.0	9	1.9	8	1.4	24	3.7	5	1.8	5	1.9	14	6.4	3	2.6	2	3.4		
	6.0	22	0.9	0	0	0	0	1	0.3	2	0.4	8	1.4	11	1.7	5	1.8	8	3.1	3	1.4	0	0	0	0		
	6.1	27	1.1	0	0	0	0	2	0.6	5	1.0	9	1.5	11	1.7	2	0.7	2	0.8	8	3.6	0	0	1	1.7		
	6.2-6.3	20	0.8	0	0	1	0.3	2	0.6	2	0.4	8	1.4	7	1.1	4	1.5	4	1.6	3	1.4	0	0	0	0		
	6.4-6.5	15	0.6	0	0	0	0	0	0	6	1.2	4	0.7	5	0.8	3	1.1	3	1.2	1	0.5	1	0.9	0	0		
	6.6-6.7	6	0.2	0	0	0	0	0	0	0	0	4	0.7	2	0.3	3	1.1	1	0.4	0	0	0	0	1	1.7		
	6.8-6.9	4	0.2	0	0	0	0	1	0.3	0	0	3	0.5	0	0	2	0.7	0	0	0	0	0	0	0	0		
	7.0-7.1	3	0.1	0	0	0	0	1	0.3	0	0	1	0.2	1	0.2	0	0	1	0.4	0	0	0	0	0	0		
	7.2-7.3	2	0.1	0	0	0	0	0	0	1	0.2	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0		
	7.4-7.5	1	0.0	0	0	0	0	0	0	0	0	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0		
	7.6-7.7	1	0.0	0	0	0	0	0	0	0	0	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0		
	7.8-7.9	1	0.0	0	0	0	0	0	0	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0	0	0		
	8.0-8.1	1	0.0	0	0	0	0	0	0	0	0	0	0	1	0.2	0	0	0	0	1	0.5	0	0	0	0		
	8.2-8.3	1	0.0	0	0	0	0	0	0	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0	0	0		
	8.4-8.5	1	0.0	0	0	0	0	0	0	1	0.2	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	8.6-8.7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	8.8-8.9	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	9.0%以上	3	0.1	0	0	1	0.3	0	0	1	0.2	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0	0	
	(再掲) 6.1%以上	86	3.4	0	0	2	0.6	6	1.9	16	3.3	31	5.3	31	4.8	16	5.8	15	5.8	13	5.9	1	0.9	2	3.4		
	総数	2,511	100	140	100	337	100	314	100	485	100	583	100	652	100	274	100	258	100	220	100	116	100	58	100		

注) インスリン注射または血糖を下げる薬服用者除外

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上				
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
総数	3.9%以下	8	0.2	2	0.8	1	0.2	1	0.2	2	0.3	2	0.2	0	0	1	0.2	0	0	0	0	0	0	0	0	0	0	0
	4.0-4.1	6	0.1	1	0.4	1	0.2	0	0	2	0.3	0	0	2	0.2	0	0	0	0	2	0.5	0	0	0	0	0	0	0
	4.2-4.3	15	0.4	2	0.8	3	0.6	1	0.2	1	0.1	4	0.4	4	0.3	3	0.6	0	0	2	0.5	1	0.5	1	1.1	1	1.1	
	4.4-4.5	87	2.1	25	10.4	17	3.3	10	2.0	14	1.8	12	1.2	9	0.8	2	0.4	3	0.6	4	1.0	2	0.9	0	0	0	0	0
	4.6-4.7	293	7.0	42	17.4	70	13.6	43	8.7	42	5.5	50	4.9	46	4.0	28	5.5	16	3.4	15	3.9	12	5.5	3	3.4	3	3.4	
	4.8-4.9	609	14.5	63	26.1	133	25.9	109	22.0	99	13.0	93	9.1	112	9.7	45	8.9	53	11.4	27	7.0	20	9.1	12	13.5	12	13.5	
	5.0-5.1	744	17.7	47	19.5	133	25.9	104	21.0	132	17.3	168	16.4	160	13.8	74	14.6	56	12.0	57	14.8	34	15.5	13	14.6	13	14.6	
	5.2	458	10.9	16	6.6	50	9.7	65	13.1	81	10.6	122	11.9	124	10.7	62	12.2	52	11.2	46	11.9	18	8.2	8	9.0	8	9.0	
	5.3	473	11.3	24	10.0	44	8.6	46	9.3	100	13.1	116	11.3	143	12.3	52	10.3	65	13.9	53	13.8	16	7.3	9	10.1	9	10.1	
	5.4	383	9.1	6	2.5	23	4.5	32	6.5	75	9.9	121	11.8	126	10.9	63	12.4	58	12.4	38	9.9	22	10.0	8	9.0	8	9.0	
	5.5	307	7.3	9	3.7	13	2.5	26	5.3	63	8.3	88	8.6	108	9.3	40	7.9	42	9.0	28	7.3	29	13.2	9	10.1	9	10.1	
	5.6	240	5.7	2	0.8	16	3.1	19	3.8	45	5.9	72	7.0	86	7.4	44	8.7	33	7.1	26	6.8	20	9.1	7	7.9	7	7.9	
	5.7	145	3.5	1	0.4	3	0.6	12	2.4	29	3.8	41	4.0	59	5.1	22	4.3	21	4.5	21	5.5	11	5.0	6	6.7	6	6.7	
	5.8	108	2.6	0	0	1	0.2	5	1.0	21	2.8	37	3.6	44	3.8	19	3.7	17	3.6	8	2.1	16	7.3	3	3.4	3	3.4	
	5.9	76	1.8	0	0	2	0.4	7	1.4	13	1.7	17	1.7	37	3.2	10	2.0	9	1.9	19	4.9	6	2.7	3	3.4	3	3.4	
	6.0	44	1.0	1	0.4	0	0	3	0.6	5	0.7	14	1.4	21	1.8	8	1.6	13	2.8	5	1.3	2	0.9	1	1.1	1	1.1	
	6.1	49	1.2	0	0	0	0	2	0.4	8	1.1	17	1.7	22	1.9	6	1.2	4	0.9	13	3.4	3	1.4	2	2.2	2	2.2	
	6.2-6.3	44	1.0	0	0	2	0.4	3	0.6	3	0.4	18	1.8	18	1.6	9	1.8	6	1.3	8	1.3	2	0.9	2	2.2	2	2.2	
	6.4-6.5	25	0.6	0	0	0	0	1	0.2	7	0.9	5	0.5	12	1.0	3	0.6	5	1.1	5	1.1	2	0.9	0	0	0	0	0
	6.6-6.7	19	0.5	0	0	0	0	0	0	3	0.4	9	0.9	7	0.6	5	1.0	2	0.4	2	0.5	1	0.5	2	2.2	2	2.2	
	6.8-6.9	13	0.3	0	0	0	0	1	0.2	3	0.4	6	0.6	3	0.3	3	0.6	1	0.2	1	0.3	1	0.5	0	0	0	0	0
	7.0-7.1	7	0.2	0	0	0	0	1	0.2	1	0.1	2	0.2	3	0.3	1	0.2	1	0.2	1	0.3	1	0.5	0	0	0	0	0
	7.2-7.3	6	0.1	0	0	0	0	0	0	2	0.3	0	0	4	0.3	0	0	3	0.6	1	0.3	0	0	0	0	0	0	0
	7.4-7.5	7	0.2	0	0	0	0	3	0.6	0	0	1	0.1	3	0.3	1	0.2	2	0.4	1	0.3	0	0	0	0	0	0	0
	7.6-7.7	3	0.1	0	0	0	0	0	0	0	0	0	0	3	0.3	0	0	2	0.4	1	0.3	0	0	0	0	0	0	0
	7.8-7.9	1	0.0	0	0	0	0	0	0	0	0	0	0	1	0.1	0	0	1	0.2	0	0	0	0	0	0	0	0	0
8.0-8.1	3	0.1	0	0	0	0	0	0	1	0.1	1	0.1	1	0.1	0	0	0	0	1	0.3	0	0	0	0	0	0	0	
8.2-8.3	1	0.0	0	0	0	0	0	0	0	0	0	0	1	0.1	0	0	1	0.2	0	0	0	0	0	0	0	0	0	
8.4-8.5	4	0.1	0	0	0	0	0	0	2	0.3	1	0.1	1	0.1	0	0	1	0.2	0	0	0	0	0	0	0	0	0	
8.6-8.7	1	0.0	0	0	0	0	0	0	0	0	0	0	1	0.1	0	0	1	0.2	0	0	0	0	0	0	0	0	0	
8.8-8.9	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
9.0%以上	15	0.4	0	0	2	0.4	1	0.2	7	0.9	3	0.3	2	0.2	3	0.6	1	0.2	0	0	1	0.5	0	0	0	0	0	
(再掲) 6.1%以上	198	4.7	0	0	4	0.8	12	2.4	37	4.9	66	6.5	79	6.8	34	6.7	28	6.0	34	8.8	11	5.0	6	6.7	6	6.7		
総数	4,194	100	241	100	514	100	495	100	761	100	1,023	100	1,160	100	507	100	466	100	385	100	220	100	89	100	89	100		

注) インスリン注射または血糖を下げる薬服用者除外

第41表の2 ヘモグロビンA1cの平均値及び標準偏差(性・年齢階級別)

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	5.3	0.7	4.9	0.3	5.0	0.4	5.2	0.6	5.4	1.0	5.4	0.6	5.4	0.6	5.4	0.7	5.4	0.5	5.4	0.5	5.5	0.7	5.4	0.5
女性	5.2	0.5	5.0	0.3	5.0	0.4	5.1	0.4	5.3	0.5	5.3	0.4	5.4	0.4	5.4	0.5	5.4	0.5	5.3	0.4	5.3	0.4	5.3	0.4
総数	5.3	0.6	4.9	0.3	5.0	0.4	5.2	0.5	5.3	0.7	5.4	0.5	5.4	0.5	5.4	0.6	5.4	0.5	5.4	0.5	5.4	0.5	5.4	0.4

注) インスリン注射または血糖を下げる薬服用者除外

第42表の1 血糖値の分布（食後3時間以上、性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	40mg/dl未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	40-60	3	0.2	1	1.1	0	0	0	0	1	0.4	1	0.2	0	0	1	0.4	0	0	0	0	0	0	0	0	0	0
	60-80	76	4.7	6	6.9	13	8.1	13	8.3	8	3.1	16	3.6	20	3.8	10	4.3	11	5.0	6	3.5	2	1.9	1	3.4	1	3.4
	80-100	865	53.2	70	80.5	126	78.8	105	67.3	147	57.2	227	51.5	190	36.1	106	45.5	87	39.4	63	37.1	31	29.2	9	31.0	9	31.0
	100-120	401	24.6	9	10.3	19	11.9	23	14.7	62	24.1	118	26.8	170	32.3	74	31.8	70	31.7	49	28.8	39	36.8	12	41.4	12	41.4
	120-140	142	8.7	0	0	1	0.6	7	4.5	17	6.6	47	10.7	70	13.3	27	11.6	22	10.0	30	17.6	17	16.0	1	3.4	1	3.4
	140-160	53	3.3	0	0	0	0	3	1.9	8	3.1	11	2.5	31	5.9	5	2.1	11	5.0	12	7.1	6	5.7	2	6.9	2	6.9
	160-180	33	2.0	1	1.1	0	0	2	1.3	3	1.2	6	1.4	21	4.0	2	0.9	8	3.6	4	2.4	5	4.7	4	13.8	4	13.8
	180-200	14	0.9	0	0	0	0	0	0	3	1.2	5	1.1	6	1.1	1	0.4	1	0.5	4	2.4	1	0.9	0	0	0	0
	200-220	12	0.7	0	0	0	0	1	0.6	0	0	2	0.5	9	1.7	0	0	6	2.7	0	0	3	2.8	0	0	0	0
	220-240	6	0.4	0	0	0	0	1	0.6	2	0.8	3	0.7	0	0	2	0.9	0	0	0	0	0	0	0	0	0	0
	240-260	7	0.4	0	0	0	0	0	0	1	0.4	2	0.5	4	0.8	2	0.9	2	0.9	1	0.6	1	0.9	0	0	0	0
	260mg/dl以上	15	0.9	0	0	1	0.6	1	0.6	5	1.9	3	0.7	5	1.0	3	1.3	3	1.4	1	0.6	1	0.9	0	0	0	0
(再掲)110mg/dl以上	391	24.0	6	6.9	5	3.1	19	12.2	52	20.2	111	25.2	198	37.6	62	26.6	80	36.2	59	34.7	48	45.3	11	37.9	11	37.9	
総数	1,627	100	87	100	160	100	156	100	257	100	441	100	526	100	233	100	221	100	170	100	106	100	29	100	29	100	
女 性	40mg/dl未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	40-60	1	0.0	0	0	0	0	0	0	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0	0	0	0	0
	60-80	104	4.5	13	11.1	33	11.4	14	5.5	18	4.0	13	2.4	13	2.1	6	2.2	6	2.4	4	2.0	3	2.7	0	0	0	0
	80-100	1,199	52.4	88	75.2	200	69.2	166	65.1	254	55.9	279	50.7	212	33.9	125	46.1	88	34.8	72	35.6	35	31.3	17	29.3	17	29.3
	100-120	680	29.7	13	11.1	43	14.9	65	25.5	147	32.4	165	30.0	247	39.5	88	32.5	95	37.5	75	37.1	48	42.9	29	50.0	29	50.0
	120-140	177	7.7	2	1.7	8	2.8	8	3.1	19	4.2	50	9.1	90	14.4	23	8.5	38	15.0	32	15.8	12	10.7	8	13.8	8	13.8
	140-160	65	2.8	1	0.9	2	0.7	1	0.4	8	1.8	20	3.6	33	5.3	14	5.2	6	2.4	14	6.9	10	8.9	3	5.2	3	5.2
	160-180	23	1.0	0	0	3	1.0	0	0	2	0.4	7	1.3	11	1.8	5	1.8	6	2.4	3	1.5	1	0.9	1	1.7	1	1.7
	180-200	12	0.5	0	0	0	0	0	0	1	0.2	1	0.2	10	1.6	0	0	8	3.2	1	0.5	1	0.9	0	0	0	0
	200-220	5	0.2	0	0	0	0	0	0	0	0	2	0.4	3	0.5	2	0.7	2	0.8	0	0	1	0.9	0	0	0	0
	220-240	6	0.3	0	0	0	0	1	0.4	0	0	2	0.4	3	0.5	1	0.4	3	1.2	0	0	0	0	0	0	0	0
	240-260	10	0.4	0	0	0	0	0	0	2	0.4	6	1.1	2	0.3	3	1.1	1	0.4	0	0	1	0.9	0	0	0	0
	260mg/dl以上	8	0.3	0	0	0	0	0	0	3	0.7	4	0.7	1	0.2	3	1.1	0	0	1	0.5	0	0	0	0	0	0
(再掲)110mg/dl以上	520	22.7	7	6.0	22	7.6	29	11.4	74	16.3	144	26.2	244	39.0	76	28.0	99	39.1	73	36.1	45	40.2	27	46.6	27	46.6	
総数	2,290	100	117	100	289	100	255	100	454	100	550	100	625	100	271	100	253	100	202	100	112	100	58	100	58	100	

第42表の1 血糖値の平均値及び標準偏差（食後3時間以上、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男 性	105.8	33.6	91.7	11.9	92.1	15.9	98.9	27.4	107.5	44.7	106.1	32.9	113.2	33.6	107.7	37.9	112.8	36.6	111.3	29.8	116.8	34.3	113.3	28.9
女 性	102.8	25.1	91.0	11.5	93.0	14.1	95.3	14.7	100.5	22.3	106.3	30.5	111.2	27.4	108.8	34.7	111.9	29.2	110.1	27.7	112.5	26.6	110.0	18.1

第 42 表の 2 血糖値の分布（食後 3 時間以上，性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上					
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%		
男 性	40mg/dl未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	40-60	3	0.2	1	1.1	0	0	0	0	0	0	1	0.4	1	0.3	0	0	1	0.5	0	0	0	0	0	0	0	0	0	0
	60-80	74	4.9	6	6.9	13	8.2	13	8.6	8	3.3	15	3.8	19	4.1	9	4.3	10	5.2	6	3.9	2	2.2	1	3.7	1	3.7	1	3.7
	80-100	844	56.0	70	80.5	126	79.2	104	68.4	144	59.3	220	55.0	180	38.7	104	49.5	84	44.0	59	38.3	28	30.1	9	33.3	9	33.3	9	33.3
	100-120	380	25.2	9	10.3	18	11.3	22	14.5	59	24.3	110	27.5	162	34.8	69	32.9	67	35.1	45	29.2	38	40.9	12	44.4	12	44.4	12	44.4
	120-140	119	7.9	0	0	1	0.6	7	4.6	13	5.3	37	9.3	61	13.1	20	9.5	19	9.9	28	18.2	13	14.0	1	3.7	1	3.7	1	3.7
	140-160	39	2.6	0	0	0	0	3	2.0	7	2.9	7	1.8	22	4.7	3	1.4	6	3.1	10	6.5	5	5.4	1	3.7	1	3.7	1	3.7
	160-180	19	1.3	1	1.1	0	0	1	0.7	1	0.4	3	0.8	13	2.8	0	0	3	1.6	3	1.9	4	4.3	3	11.1	3	11.1	3	11.1
	180-200	11	0.7	0	0	0	0	0	0	3	1.2	3	0.8	5	1.1	1	0.5	1	0.5	3	1.9	1	1.1	0	0	0	0	0	0
	200-220	3	0.2	0	0	0	0	0	0	0	0	0	0	1	0.3	2	0.4	0	0	0	0	0	0	2	2.2	0	0	0	0
	220-240	3	0.2	0	0	0	0	1	0.7	1	0.4	1	0.4	1	0.3	0	0	1	0.5	0	0	0	0	0	0	0	0	0	0
	240-260	1	0.1	0	0	0	0	0	0	1	0.4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	260mg/dl以上	10	0.7	0	0	1	0.6	1	0.7	5	2.1	2	0.5	1	0.2	2	1.0	1	0.5	0	0	0	0	0	0	0	0	0	0
(再掲)110mg/dl以上	306	20.3	6	6.9	5	3.1	16	10.5	43	17.7	82	20.5	154	33.1	44	21.0	56	29.3	50	32.5	39	41.9	9	33.3	9	33.3	9	33.3	
総 数	1,506	100	87	100	159	100	152	100	243	100	400	100	465	100	210	100	191	100	154	100	93	100	27	100	27	100	27	100	
女 性	40mg/dl未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	40-60	1	0.0	0	0	0	0	0	0	0	0	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0	0	0	0	0
	60-80	102	4.6	13	11.2	33	11.5	14	5.5	18	4.1	13	2.5	11	1.9	6	2.4	4	1.7	4	2.0	3	2.9	0	0	0	0	0	0
	80-100	1,184	53.9	88	75.9	200	69.4	165	65.2	252	57.1	272	52.8	207	35.4	121	48.6	85	36.6	71	36.2	34	33.3	17	31.5	17	31.5		
	100-120	660	30.0	13	11.2	42	14.6	65	25.7	143	32.4	162	31.5	235	40.2	86	34.5	91	39.2	74	37.8	45	44.1	25	46.3	25	46.3		
	120-140	164	7.5	1	0.9	8	2.8	8	3.2	16	3.6	46	8.9	85	14.6	22	8.8	34	14.7	32	16.3	11	10.8	8	14.8	8	14.8		
	140-160	47	2.1	1	0.9	2	0.7	1	0.4	7	1.6	9	1.7	27	4.6	5	2.0	4	1.7	13	6.6	7	6.9	3	5.6	3	5.6		
	160-180	20	0.9	0	0	3	1.0	0	0	2	0.5	6	1.2	9	1.5	5	2.0	6	2.6	1	0.5	1	1.0	1	1.9	1	1.9		
	180-200	9	0.4	0	0	0	0	0	0	1	0.2	1	0.2	7	1.2	0	0	5	2.2	1	0.5	1	1.0	0	0	0	0		
	200-220	2	0.1	0	0	0	0	0	0	0	0	0	0	1	0.2	1	0.2	1	0.4	1	0.4	0	0	0	0	0	0		
	220-240	2	0.1	0	0	0	0	0	0	0	0	0	0	1	0.2	1	0.2	0	0	1	0.4	0	0	0	0	0	0		
	240-260	1	0.0	0	0	0	0	0	0	0	0	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0	0	0		
	260mg/dl以上	5	0.2	0	0	0	0	0	0	2	0.5	3	0.6	0	0	2	0.8	0	0	0	0	0	0	0	0	0	0		
(再掲)110mg/dl以上	454	20.7	6	5.2	21	7.3	28	11.1	65	14.7	117	22.7	217	37.2	59	23.7	86	37.1	69	35.2	37	36.3	25	46.3	25	46.3			
総 数	2,197	100	116	100	288	100	253	100	441	100	515	100	584	100	249	100	232	100	196	100	102	100	54	100	54	100			

注) インスリン注射または血糖を下げる薬服用者除外

第 42 表の 2 血糖値の平均値及び標準偏差（食後 3 時間以上，性・年齢階級別）

	総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男 性	102.3	28.4	91.7	11.9	92.0	15.9	97.6	25.4	106.1	44.6	102.1	25.7	107.6	24.3	103.0	29.1	104.4	23.2	108.3	23.0	112.5	27.1	109.7	26.1
女 性	100.8	20.3	90.7	10.9	93.0	14.1	94.7	11.7	99.2	18.4	102.8	23.9	109.0	21.8	104.2	27.5	109.6	25.5	108.1	18.5	108.9	20.2	109.7	18.7

注) インスリン注射または血糖を下げる薬服用者除外

第43表の1 総コレステロール値の分布（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	120mg/dl未満	19	1.0	3	3.0	1	0.6	2	1.1	1	0.3	2	0.4	10	1.7	2	0.8	1	0.4	6	3.3	2	1.7	1	3.0		
	120-140	55	3.0	5	5.0	3	1.7	1	0.5	3	1.0	11	2.3	32	5.5	6	2.3	7	2.9	10	5.5	11	9.2	4	12.1		
	140-160	154	8.5	17	17.0	6	3.4	12	6.5	12	4.1	44	9.1	63	10.9	25	9.7	24	9.9	15	8.2	17	14.3	7	21.2		
	160-180	352	19.4	26	26.0	41	23.0	31	16.8	46	15.8	82	16.9	126	21.8	46	17.8	52	21.4	42	23.0	27	22.7	5	15.2		
	180-200	425	23.4	28	28.0	37	20.8	48	25.9	57	19.5	118	24.3	137	23.7	61	23.6	58	23.9	49	26.8	21	17.6	9	27.3		
	200-220	390	21.5	12	12.0	36	20.2	38	20.5	78	26.7	119	24.5	107	18.5	57	22.0	46	18.9	36	19.7	22	18.5	3	9.1		
	220-240	236	13.0	3	3.0	33	18.5	31	16.8	46	15.8	61	12.6	62	10.7	37	14.3	34	14.0	14	7.7	12	10.1	2	6.1		
	240-260	120	6.6	5	5.0	10	5.6	11	5.9	30	10.3	29	6.0	35	6.1	18	6.9	18	7.4	9	4.9	6	5.0	2	6.1		
	260-280	44	2.4	1	1.0	9	5.1	7	3.8	11	3.8	12	2.5	4	0.7	4	1.5	2	0.8	2	1.1	0	0	0	0		
	280-300	14	0.8	0	0	1	0.6	1	0.5	5	1.7	5	1.0	2	0.3	2	0.8	1	0.4	0	0	1	0.8	0	0		
	300mg/dl以上	9	0.5	0	0	1	0.6	3	1.6	3	1.0	2	0.4	0	0	1	0.4	0	0	0	0	0	0	0	0		
総数		1,818	100	100	100	178	100	185	100	292	100	485	100	578	100	259	100	243	100	183	100	119	100	33	100		
女 性	120mg/dl未満	4	0.2	0	0	2	0.6	0	0	0	0	0	0	2	0.3	0	0	0	0	1	0.4	0	0	1	1.5		
	120-140	39	1.5	9	6.3	7	2.1	5	1.6	4	0.8	3	0.5	11	1.6	0	0	4	1.4	4	1.7	1	0.8	2	3.1		
	140-160	159	6.0	27	18.8	54	16.0	26	8.2	5	1.0	7	1.1	40	5.6	5	1.7	12	4.2	17	7.4	7	5.5	4	6.2		
	160-180	372	14.1	37	25.7	82	24.3	59	18.6	33	6.6	55	8.8	106	15.0	38	12.7	37	13.0	34	14.8	24	18.8	11	16.9		
	180-200	570	21.7	35	24.3	87	25.7	71	22.4	91	18.1	119	19.1	167	23.6	64	21.3	65	22.8	51	22.2	36	28.1	15	23.1		
	200-220	611	23.2	15	10.4	54	16.0	79	24.9	120	23.9	162	26.0	181	25.6	80	26.7	70	24.6	51	22.2	36	28.1	24	36.9		
	220-240	436	16.6	14	9.7	30	8.9	43	13.6	100	19.9	131	21.1	118	16.7	46	15.3	55	19.3	45	19.6	14	10.9	4	6.2		
	240-260	248	9.4	5	3.5	12	3.6	20	6.3	81	16.1	80	12.9	50	7.1	40	13.3	22	7.7	20	8.7	5	3.9	3	4.6		
	260-280	123	4.7	2	1.4	7	2.1	11	3.5	40	8.0	40	6.4	23	3.2	14	4.7	13	4.6	5	2.2	4	3.1	1	1.5		
	280-300	49	1.9	0	0	2	0.6	2	0.6	18	3.6	18	2.9	9	1.3	9	3.0	6	2.1	2	0.9	1	0.8	0	0		
	300mg/dl以上	20	0.8	0	0	1	0.3	1	0.3	10	2.0	7	1.1	1	0.1	4	1.3	1	0.4	0	0	0	0	0	0		
総数		2,631	100	144	100	338	100	317	100	502	100	622	100	708	100	300	100	285	100	230	100	128	100	65	100		

第43表の1 総コレステロール値の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男 性	196.6	34.9	179.5	31.4	201.4	33.2	203.8	39.1	207.7	34.1	198.0	33.3	188.8	33.4	196.8	34.2	194.4	32.3	186.7	33.0	184.0	34.5	177.4	33.9
女 性	206.7	34.4	182.6	31.4	187.7	31.4	199.9	31.9	222.0	33.4	217.9	31.9	203.0	31.3	214.3	32.9	207.7	32.1	201.9	32.4	198.9	27.7	194.2	27.5

第43表の2 総コレステロール値の分布（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	120mg/dl未満	17	1.0	3	3.0	1	0.6	2	1.2	1	0.4	2	0.5	8	1.6	2	0.9	1	0.5	4	2.5	2	2.0	1	3.3		
	120-140	48	3.0	5	5.0	3	1.7	1	0.6	3	1.1	10	2.4	26	5.3	5	2.3	7	3.4	7	4.4	8	8.2	4	13.3		
	140-160	134	8.3	17	17.0	6	3.4	11	6.4	12	4.6	36	8.7	52	10.5	19	8.6	16	7.7	13	8.2	16	16.3	7	23.3		
	160-180	319	19.7	26	26.0	40	23.0	29	16.8	42	16.0	71	17.1	111	22.4	38	17.1	46	22.1	37	23.3	25	25.5	3	10.0		
	180-200	384	23.7	28	28.0	36	20.7	46	26.6	49	18.7	103	24.8	122	24.6	55	24.8	51	24.5	45	28.3	18	18.4	8	26.7		
	200-220	336	20.7	12	12.0	34	19.5	35	20.2	75	28.6	95	22.8	85	17.2	45	20.3	37	17.8	29	18.2	16	16.3	3	10.0		
	220-240	215	13.3	3	3.0	33	19.0	29	16.8	39	14.9	55	13.2	56	11.3	34	15.3	32	15.4	13	8.2	9	9.2	2	6.7		
	240-260	108	6.7	5	5.0	10	5.7	10	5.8	27	10.3	27	6.5	29	5.9	18	8.1	15	7.2	9	5.7	3	3.1	2	6.7		
	260-280	41	2.5	1	1.0	9	5.2	6	3.5	9	3.4	12	2.9	4	0.8	4	1.8	2	1.0	2	1.3	0	0	0	0		
	280-300	10	0.6	0	0	1	0.6	1	0.6	3	1.1	3	0.7	2	0.4	1	0.5	1	0.5	0	0	1	1.0	0	0		
	300mg/dl以上	8	0.5	0	0	1	0.6	3	1.7	2	0.8	2	0.5	0	0	1	0.5	0	0	0	0	0	0	0	0		
	総数	1,620	100	100	100	174	100	173	100	262	100	416	100	495	100	222	100	208	100	159	100	98	100	30	100		
女 性	120mg/dl未満	4	0.2	0	0	2	0.6	0	0	0	0	0	0	2	0.4	0	0	0	0	1	0.6	0	0	1	2.0		
	120-140	36	1.6	9	6.3	7	2.1	5	1.6	4	0.9	2	0.4	9	1.7	0	0	2	1.0	4	2.3	1	1.1	2	4.0		
	140-160	140	6.3	27	18.8	54	16.1	26	8.4	4	0.9	3	0.6	26	5.0	1	0.4	6	3.0	12	6.9	4	4.3	4	8.0		
	160-180	303	13.5	37	25.7	81	24.1	59	19.1	28	6.3	33	6.8	65	12.5	23	10.1	20	10.0	19	10.9	19	20.2	7	14.0		
	180-200	474	21.2	35	24.3	87	25.9	70	22.7	77	17.4	85	17.5	120	23.1	49	21.6	47	23.4	38	21.7	26	27.7	9	18.0		
	200-220	496	22.2	15	10.4	54	16.1	76	24.6	104	23.5	115	23.6	132	25.4	56	24.7	48	23.9	40	22.9	23	24.5	21	42.0		
	220-240	382	17.1	14	9.7	30	8.9	40	12.9	87	19.6	116	23.8	95	18.3	38	16.7	42	20.9	37	21.1	13	13.8	3	6.0		
	240-260	223	10.0	5	3.5	12	3.6	20	6.5	74	16.7	70	14.4	42	8.1	34	15.0	18	9.0	18	10.3	4	4.3	2	4.0		
	260-280	115	5.1	2	1.4	6	1.8	10	3.2	38	8.6	38	7.8	21	4.0	13	5.7	12	6.0	4	2.3	4	4.3	1	2.0		
	280-300	47	2.1	0	0	2	0.6	2	0.6	17	3.8	18	3.7	8	1.5	9	4.0	6	3.0	2	1.1	0	0	0	0		
	300mg/dl以上	19	0.8	0	0	1	0.3	1	0.3	10	2.3	7	1.4	0	0	4	1.8	0	0	0	0	0	0	0	0		
	総数	2,239	100	144	100	336	100	309	100	443	100	487	100	520	100	227	100	201	100	175	100	94	100	50	100		

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

第43表の2 総コレステロール値の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男 性	196.5	34.80	179.5	31.37	201.6	33.39	203.7	39.41	206.3	33.40	198.3	33.63	189.1	33.27	198.1	34.46	195.2	32.49	188.3	32.21	181.1	33.57	177.5	35.57
女 性	207.6	35.23	182.6	31.39	187.5	31.25	199.2	31.84	223.2	34.01	222.3	32.07	205.5	31.72	218.8	33.27	211.7	31.88	204.4	33.08	200.1	27.68	194.7	28.85

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

第44表の1 中性脂肪（トリグリセライド）値の分布（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上					
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%		
男性	20mg/dl未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	20-50	57	3.1	12	12.0	5	2.8	3	1.6	6	2.1	15	3.1	16	2.8	4	1.5	5	2.1	4	2.2	4	2.2	4	3.4	3	9.1		
	50-80	348	19.1	29	29.0	39	21.9	28	15.1	38	13.0	81	16.7	133	23.0	51	19.7	49	20.2	43	23.5	30	25.2	11	33.3				
	80-110	381	21.0	21	21.0	39	21.9	31	16.8	63	21.6	112	23.1	115	19.9	60	23.2	45	18.5	36	19.7	25	21.0	9	27.3				
	110-140	335	18.4	13	13.0	30	16.9	37	20.0	44	15.1	92	19.0	119	20.6	47	18.1	51	21.0	41	22.4	21	17.6	6	18.2				
	140-170	228	12.5	12	12.0	21	11.8	23	12.4	36	12.3	68	14.0	68	11.8	32	12.4	33	13.6	22	12.0	12	10.1	1	3.0				
	170-200	139	7.6	4	4.0	9	5.1	17	9.2	23	7.9	35	7.2	51	8.8	18	6.9	23	9.5	16	8.7	12	10.1	0	0				
	200mg/dl以上	330	18.2	9	9.0	35	19.7	46	24.9	82	28.1	82	16.9	76	13.1	47	18.1	37	15.2	21	11.5	15	12.6	3	9.1				
	(再掲)150mg/dl以上	614	33.8	20	20.0	58	32.6	82	44.3	128	43.8	162	33.4	164	28.4	85	32.8	77	31.7	50	27.3	34	28.6	3	9.1				
総数	1,818	100	100	100	178	100	185	100	292	100	485	100	578	100	259	100	243	100	183	100	119	100	33	100					
女性	20mg/dl未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	20-50	198	7.5	37	25.7	59	17.5	36	11.4	22	4.4	25	4.0	19	2.7	14	4.7	7	2.5	6	2.6	5	3.9	1	1.5				
	50-80	638	24.2	57	39.6	125	37.0	96	30.3	126	25.1	114	18.3	120	16.9	56	18.7	40	14.0	45	19.6	27	21.1	8	12.3				
	80-110	691	26.3	21	14.6	76	22.5	82	25.9	128	25.5	166	26.7	218	30.8	79	26.3	87	30.5	68	29.6	39	30.5	24	36.9				
	110-140	423	16.1	14	9.7	34	10.1	44	13.9	77	15.3	116	18.6	138	19.5	53	17.7	59	20.7	40	17.4	21	16.4	18	27.7				
	140-170	239	9.1	6	4.2	19	5.6	21	6.6	44	8.8	67	10.8	82	11.6	32	10.7	32	11.2	30	13.0	13	10.2	7	10.8				
	170-200	150	5.7	4	2.8	16	4.7	9	2.8	36	7.2	40	6.4	45	6.4	16	5.3	19	6.7	16	7.0	8	6.3	2	3.1				
	200mg/dl以上	292	11.1	5	3.5	9	2.7	29	9.1	69	13.7	94	15.1	86	12.1	50	16.7	41	14.4	25	10.9	15	11.7	5	7.7				
	(再掲)150mg/dl以上	597	22.7	14	9.7	38	11.2	54	17.0	131	26.1	178	28.6	182	25.7	85	28.3	82	28.8	58	25.2	31	24.2	11	16.9				
総数	2,631	100	144	100	338	100	317	100	502	100	622	100	708	100	300	100	285	100	230	100	128	100	65	100					

第44表の1 中性脂肪（トリグリセライド）値の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	144.3	95.5	113.8	85.4	151.6	109.0	165.6	109.2	167.2	108.8	141.5	92.7	131.2	77.8	139.0	80.7	138.5	78.7	127.2	68.1	131.6	93.7	98.2	44.1
女性	118.0	70.8	81.8	52.0	89.6	52.3	107.8	72.8	125.5	78.5	129.0	67.0	128.6	72.0	131.7	73.6	134.8	75.3	127.7	75.9	121.4	66.0	118.1	49.5

第44表の2 中性脂肪（トリグリセライド）値の分布（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上							
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%				
男 性	20mg/dl未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	20-50	56	3.5	12	12.0	5	2.9	3	1.7	6	2.3	14	3.4	16	3.2	4	1.8	5	2.4	4	2.5	4	4.1	3	10.0						
	50-80	326	20.1	29	29.0	39	22.4	27	15.6	38	14.5	71	17.1	122	24.6	43	19.4	46	22.1	40	25.2	25	25.5	11	36.7						
	80-110	343	21.2	21	21.0	37	21.3	30	17.3	58	22.1	99	23.8	98	19.8	54	24.3	37	17.8	32	20.1	22	22.4	7	23.3						
	110-140	296	18.3	13	13.0	30	17.2	35	20.2	42	16.0	80	19.2	96	19.4	39	17.6	41	19.7	33	20.8	16	16.3	6	20.0						
	140-170	202	12.5	12	12.0	20	11.5	22	12.7	33	12.6	56	13.5	59	11.9	28	12.6	29	13.9	21	13.2	8	8.2	1	3.3						
	170-200	122	7.5	4	4.0	9	5.2	16	9.2	18	6.9	29	7.0	46	9.3	15	6.8	21	10.1	15	9.4	10	10.2	0	0						
	200mg/dl以上	275	17.0	9	9.0	34	19.5	40	23.1	67	25.6	67	16.1	58	11.7	39	17.6	29	13.9	14	8.8	13	13.3	2	6.7						
	(再掲)150mg/dl以上	525	32.4	20	20.0	56	32.2	74	42.8	107	40.8	132	31.7	136	27.5	71	32.0	65	31.3	42	26.4	27	27.6	2	6.7						
総数	1,620	100	100	100	174	100	173	100	262	100	416	100	495	100	222	100	208	100	159	100	98	100	30	100							
女 性	20mg/dl未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	20-50	192	8.6	37	25.7	59	17.6	35	11.3	21	4.7	23	4.7	17	3.3	12	5.3	5	2.5	6	3.4	5	5.3	1	2.0						
	50-80	565	25.2	57	39.6	125	37.2	95	30.7	113	25.5	88	18.1	87	16.7	42	18.5	27	13.4	37	21.1	17	18.1	6	12.0						
	80-110	584	26.1	21	14.6	75	22.3	81	26.2	112	25.3	132	27.1	163	31.3	62	27.3	63	31.3	48	27.4	32	34.0	20	40.0						
	110-140	351	15.7	14	9.7	34	10.1	44	14.2	63	14.2	96	19.7	100	19.2	42	18.5	43	21.4	30	17.1	15	16.0	12	24.0						
	140-170	201	9.0	6	4.2	19	5.7	18	5.8	42	9.5	49	10.1	67	12.9	22	9.7	24	11.9	28	16.0	10	10.6	5	10.0						
	170-200	115	5.1	4	2.8	16	4.8	8	2.6	33	7.4	26	5.3	28	5.4	11	4.8	11	5.5	10	5.7	5	5.3	2	4.0						
	200mg/dl以上	231	10.3	5	3.5	8	2.4	28	9.1	59	13.3	73	15.0	58	11.2	36	15.9	28	13.9	16	9.1	10	10.6	4	8.0						
	(再掲)150mg/dl以上	474	21.2	14	9.7	37	11.0	50	16.2	116	26.2	129	26.5	128	24.6	60	26.4	56	27.9	42	24.0	21	22.3	9	18.0						
総数	2,239	100	144	100	336	100	309	100	443	100	487	100	520	100	227	100	201	100	175	100	94	100	50	100							

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

第44表の2 中性脂肪（トリグリセライド）値の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差		
男 性	140.7	92.0	113.8	85.4	150.2	106.9	162.7	108.4	158.8	99.9	139.5	94.1	126.7	69.0	138.6	82.7	136.3	78.7	121.3	55.6	124.6	70.2	95.3	41.5		
女 性	114.9	69.9	81.8	52.0	88.3	47.1	107.0	72.8	124.0	77.2	126.8	66.4	127.2	73.5	129.1	73.3	133.9	75.4	125.7	78.5	120.1	69.1	118.5	52.6		

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

第45表の1 HDL-コレステロール値の分布（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	30mg/dl未満	15	0.8	1	1.0	1	0.6	0	0	3	1.0	2	0.4	8	1.4	1	0.4	3	1.2	4	2.2	1	0.8	0	0	0	0
	30-40	162	8.9	4	4.0	14	7.9	20	10.8	22	7.5	40	8.2	62	10.7	26	10.0	23	9.5	17	9.3	19	16.0	3	9.1		
	40-50	427	23.5	18	18.0	42	23.6	41	22.2	69	23.6	115	23.7	142	24.6	59	22.8	57	23.5	47	25.7	29	24.4	9	27.3		
	50-60	485	26.7	32	32.0	45	25.3	49	26.5	71	24.3	121	24.9	167	28.9	57	22.0	77	31.7	45	24.6	34	28.6	11	33.3		
	60-70	384	21.1	27	27.0	39	21.9	38	20.5	70	24.0	93	19.2	117	20.2	52	20.1	48	19.8	42	23.0	21	17.6	6	18.2		
	70-80	195	10.7	13	13.0	18	10.1	24	13.0	36	12.3	62	12.8	42	7.3	36	13.9	18	7.4	14	7.7	7	5.9	3	9.1		
	80-90	102	5.6	2	2.0	14	7.9	6	3.2	15	5.1	38	7.8	27	4.7	21	8.1	11	4.5	10	5.5	5	4.2	1	3.0		
	90-100	27	1.5	2	2.0	4	2.2	4	2.2	0	0	9	1.9	8	1.4	5	1.9	4	1.6	4	2.2	0	0	0	0		
	100mg/dl以上	21	1.2	1	1.0	1	0.6	3	1.6	6	2.1	5	1.0	5	0.9	2	0.8	2	0.8	0	0	3	2.5	0	0		
	総数	1,818	100	100	100	178	100	185	100	292	100	485	100	578	100	259	100	243	100	183	100	119	100	33	100		
女性	30mg/dl未満	6	0.2	0	0	0	0	1	0.3	1	0.2	0	0	4	0.6	0	0	2	0.7	2	0.9	0	0	0	0		
	30-40	64	2.4	2	1.4	6	1.8	5	1.6	8	1.6	11	1.8	32	4.5	8	2.7	10	3.5	16	7.0	5	3.9	1	1.5		
	40-50	291	11.1	6	4.2	26	7.7	20	6.3	46	9.2	86	13.8	107	15.1	52	17.3	46	16.1	32	13.9	14	10.9	15	23.1		
	50-60	526	20.0	24	16.7	46	13.6	52	16.4	102	20.3	133	21.4	169	23.9	68	22.7	65	22.8	52	22.6	30	23.4	22	33.8		
	60-70	694	26.4	40	27.8	94	27.8	92	29.0	124	24.7	159	25.6	185	26.1	76	25.3	68	23.9	60	26.1	44	34.4	13	20.0		
	70-80	507	19.3	35	24.3	75	22.2	71	22.4	98	19.5	113	18.2	115	16.2	52	17.3	54	18.9	32	13.9	22	17.2	7	10.8		
	80-90	324	12.3	22	15.3	59	17.5	44	13.9	68	13.5	70	11.3	61	8.6	28	9.3	26	9.1	23	10.0	7	5.5	5	7.7		
	90-100	144	5.5	9	6.3	25	7.4	16	5.0	33	6.6	39	6.3	22	3.1	14	4.7	10	3.5	7	3.0	4	3.1	1	1.5		
	100mg/dl以上	75	2.9	6	4.2	7	2.1	16	5.0	22	4.4	11	1.8	13	1.8	2	0.7	4	1.4	6	2.6	2	1.6	1	1.5		
	総数	2,631	100	144	100	338	100	317	100	502	100	622	100	708	100	300	100	285	100	230	100	128	100	65	100		

第45表の1 HDL-コレステロール値の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男性	57.3	15.1	59.0	13.4	58.2	14.7	57.7	15.6	57.8	14.9	58.6	15.4	55.3	15.0	58.7	15.9	55.8	15.0	55.1	14.7	54.8	16.3	53.9	11.8
女性	66.8	16.0	70.6	14.7	69.9	15.4	70.0	16.3	68.7	16.5	65.9	15.7	62.6	15.4	63.5	15.6	63.2	15.5	62.6	16.4	62.7	13.7	59.9	14.5

第45表の2 HDL-コレステロール値の分布（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	30mg/dl未満	12	0.7	1	1.0	1	0.6	0	0	2	0.8	1	0.2	7	1.4	0	0	3	1.4	3	1.9	1	1.0	0	0	0	0
	30-40	143	8.8	4	4.0	13	7.5	18	10.4	18	6.9	35	8.4	55	11.1	23	10.4	20	9.6	13	8.2	19	19.4	3	10.0		
	40-50	376	23.2	18	18.0	42	24.1	36	20.8	64	24.4	100	24.0	116	23.4	51	23.0	46	22.1	41	25.8	20	20.4	9	30.0		
	50-60	426	26.3	32	32.0	43	24.7	47	27.2	60	22.9	100	24.0	144	29.1	45	20.3	69	33.2	39	24.5	26	26.5	10	33.3		
	60-70	346	21.4	27	27.0	38	21.8	36	20.8	64	24.4	78	18.8	103	20.8	45	20.3	42	20.2	38	23.9	18	18.4	5	16.7		
	70-80	179	11.0	13	13.0	18	10.3	24	13.9	34	13.0	55	13.2	35	7.1	32	14.4	15	7.2	12	7.5	6	6.1	2	6.7		
	80-90	95	5.9	2	2.0	14	8.0	6	3.5	15	5.7	34	8.2	24	4.8	20	9.0	8	3.8	10	6.3	5	5.1	1	3.3		
	90-100	24	1.5	2	2.0	4	2.3	3	1.7	0	0	9	2.2	6	1.2	5	2.3	3	1.4	3	1.9	0	0	0	0		
	100mg/dl以上	19	1.2	1	1.0	1	0.6	3	1.7	5	1.9	4	1.0	5	1.0	1	0.5	2	1.0	0	0	3	3.1	0	0		
	総数	1,620	100	100	100	174	100	173	100	262	100	416	100	495	100	222	100	208	100	159	100	98	100	30	100		
女 性	30mg/dl未満	4	0.2	0	0	0	0	1	0.3	1	0.2	0	0	2	0.4	0	0	0	0	2	1.1	0	0	0	0		
	30-40	51	2.3	2	1.4	6	1.8	5	1.6	6	1.4	9	1.8	23	4.4	8	3.5	9	4.5	10	5.7	3	3.2	1	2.0		
	40-50	235	10.5	6	4.2	25	7.4	18	5.8	39	8.8	66	13.6	81	15.6	38	16.7	30	14.9	26	14.9	12	12.8	13	26.0		
	50-60	445	19.9	24	16.7	45	13.4	51	16.5	94	21.2	97	19.9	134	25.8	47	20.7	51	25.4	40	22.9	26	27.7	17	34.0		
	60-70	597	26.7	40	27.8	94	28.0	91	29.4	110	24.8	129	26.5	133	25.6	58	25.6	48	23.9	46	26.3	31	33.0	8	16.0		
	70-80	430	19.2	35	24.3	75	22.3	69	22.3	85	19.2	86	17.7	80	15.4	42	18.5	37	18.4	23	13.1	15	16.0	5	10.0		
	80-90	279	12.5	22	15.3	59	17.6	43	13.9	58	13.1	58	11.9	39	7.5	22	9.7	16	8.0	17	9.7	2	2.1	4	8.0		
	90-100	130	5.8	9	6.3	25	7.4	16	5.2	31	7.0	32	6.6	17	3.3	10	4.4	7	3.5	6	3.4	3	3.2	1	2.0		
	100mg/dl以上	68	3.0	6	4.2	7	2.1	15	4.9	19	4.3	10	2.1	11	2.1	2	0.9	3	1.5	5	2.9	2	2.1	1	2.0		
	総数	2,239	100	144	100	336	100	309	100	443	100	487	100	520	100	227	100	201	100	175	100	94	100	50	100		

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

第45表の2 HDL-コレステロール値の平均値及び標準偏差（性・年齢階級別）

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上		
		平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	
男 性	57.6	15.2	59.0	13.4	58.3	14.8	58.2	15.5	58.2	14.8	58.9	15.6	55.3	15.2	59.1	15.8	55.6	14.9	55.5	14.6	55.2	17.6	52.8	11.6
女 性	67.2	16.1	70.6	14.7	70.0	15.3	70.1	16.3	68.7	16.3	66.3	16.0	62.4	15.5	63.8	16.2	62.9	15.3	62.9	16.7	61.4	13.6	59.9	15.5

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

第46表の1-1 LDL-コレステロール値（Friedewaldの式）の分布（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	40mg/dl未満	10	0.6	0	0	0	0	1	0.6	1	0.4	3	0.6	5	0.9	2	0.8	0	0	3	1.6	2	1.7	0	0	0	0
	40-50	16	0.9	1	1.0	1	0.6	0	0	3	1.1	5	1.1	6	1.1	3	1.2	2	0.8	4	2.2	0	0	0	0	0	0
	50-60	34	1.9	8	8.1	2	1.2	3	1.7	1	0.4	8	1.7	12	2.1	5	2.0	3	1.3	6	3.3	3	2.6	0	0	0	0
	60-70	74	4.2	7	7.1	4	2.4	10	5.6	6	2.2	20	4.2	27	4.7	13	5.1	12	5.0	7	3.8	6	5.2	2	6.1	2	6.1
	70-80	126	7.1	9	9.1	13	7.8	12	6.7	20	7.2	26	5.5	46	8.1	16	6.3	21	8.8	12	6.6	8	6.9	5	15.2	5	15.2
	80-90	174	9.8	13	13.1	21	12.6	13	7.3	24	8.6	42	8.9	61	10.7	24	9.4	18	7.5	16	8.8	19	16.4	8	24.2	8	24.2
	90-100	211	11.9	16	16.2	21	12.6	21	11.8	28	10.0	50	10.5	75	13.2	25	9.8	25	10.5	25	13.7	23	19.8	2	6.1	2	6.1
	100-110	224	12.7	13	13.1	21	12.6	22	12.4	31	11.1	73	15.4	64	11.2	40	15.7	30	12.6	19	10.4	12	10.3	3	9.1	3	9.1
	110-120	247	14.0	12	12.1	21	12.6	29	16.3	30	10.8	73	15.4	82	14.4	35	13.7	35	14.6	32	17.6	11	9.5	4	12.1	4	12.1
	120-130	205	11.6	6	6.1	15	9.0	18	10.1	33	11.8	60	12.7	73	12.8	32	12.5	33	13.8	24	13.2	12	10.3	4	12.1	4	12.1
	130-140	166	9.4	6	6.1	17	10.2	15	8.4	41	14.7	43	9.1	44	7.7	22	8.6	22	9.2	15	8.2	6	5.2	1	3.0	1	3.0
	140-150	92	5.2	4	4.0	8	4.8	11	6.2	20	7.2	24	5.1	25	4.4	10	3.9	11	4.6	5	2.7	8	6.9	1	3.0	1	3.0
	150-160	86	4.9	2	2.0	9	5.4	11	6.2	22	7.9	21	4.4	21	3.7	14	5.5	12	5.0	7	3.8	1	0.9	1	3.0	1	3.0
	160-170	42	2.4	1	1.0	4	2.4	2	1.1	5	1.8	14	3.0	16	2.8	7	2.7	9	3.8	3	1.6	3	2.6	1	3.0	1	3.0
	170-180	26	1.5	0	0	3	1.8	5	2.8	6	2.2	3	0.6	9	1.6	2	0.8	4	1.7	3	1.6	1	0.9	1	3.0	1	3.0
	180-190	16	0.9	1	1.0	4	2.4	3	1.7	3	1.1	3	0.6	2	0.4	2	0.8	1	0.4	0	0	1	0.9	0	0	0	0
	190-200	9	0.5	0	0	2	1.2	1	0.6	2	0.7	3	0.6	1	0.2	2	0.8	0	0	1	0.5	0	0	0	0	0	0
	200-210	7	0.4	0	0	1	0.6	0	0	3	1.1	2	0.4	1	0.2	0	0	1	0.4	0	0	0	0	0	0	0	0
	210-220	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	220mg/dl以上	2	0.1	0	0	0	0	1	0.6	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0	0	0	0	0
総数	1,767	100	99	100	167	100	178	100	279	100	474	100	570	100	255	100	239	100	182	100	116	100	33	100			
女性	40mg/dl未満	2	0.1	0	0	0	0	2	0.6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	40-50	16	0.6	1	0.7	5	1.5	5	1.6	0	0	1	0.2	4	0.6	0	0	0	0	3	1.3	0	0	1	1.5	1	1.5
	50-60	36	1.4	6	4.2	12	3.6	7	2.2	1	0.2	2	0.3	8	1.1	1	0.3	5	1.8	0	0	2	1.6	1	1.5	1	1.5
	60-70	87	3.3	12	8.3	19	5.6	14	4.5	7	1.4	9	1.4	26	3.7	4	1.3	8	2.8	11	4.9	4	3.1	3	4.6	3	4.6
	70-80	156	6.0	22	15.3	52	15.4	22	7.0	12	2.4	14	2.3	34	4.9	9	3.0	12	4.3	11	4.9	7	5.5	4	6.2	4	6.2
	80-90	216	8.3	29	20.1	45	13.4	39	12.4	16	3.2	30	4.8	57	8.1	14	4.7	20	7.1	23	10.2	11	8.7	3	4.6	3	4.6
	90-100	302	11.6	22	15.3	51	15.1	41	13.1	53	10.7	60	9.7	75	10.7	35	11.7	27	9.6	23	10.2	17	13.4	8	12.3	8	12.3
	100-110	343	13.1	19	13.2	44	13.1	44	14.0	53	10.7	71	11.4	112	16.0	39	13.0	49	17.4	30	13.3	20	15.7	13	20.0	13	20.0
	110-120	339	13.0	12	8.3	39	11.6	32	10.2	64	13.0	85	13.7	107	15.3	39	13.0	39	13.8	35	15.5	22	17.3	11	16.9	11	16.9
	120-130	306	11.7	5	3.5	25	7.4	34	10.8	64	13.0	101	16.3	77	11.0	48	16.1	31	11.0	23	10.2	16	12.6	7	10.8	7	10.8
	130-140	264	10.1	5	3.5	19	5.6	35	11.1	66	13.4	69	11.1	70	10.0	30	10.0	32	11.3	21	9.3	8	6.3	9	13.8	9	13.8
	140-150	188	7.2	5	3.5	11	3.3	12	3.8	48	9.7	59	9.5	53	7.6	24	8.0	21	7.4	20	8.8	11	8.7	1	1.5	1	1.5
	150-160	123	4.7	2	1.4	2	0.6	12	3.8	32	6.5	38	6.1	37	5.3	20	6.7	16	5.7	15	6.6	4	3.1	2	3.1	2	3.1
	160-170	86	3.3	3	2.1	7	2.1	4	1.3	27	5.5	33	5.3	12	1.7	13	4.3	5	1.8	6	2.7	1	0.8	0	0	0	0
	170-180	67	2.6	0	0	1	0.3	5	1.6	26	5.3	21	3.4	14	2.0	10	3.3	10	3.5	1	0.4	2	1.6	1	1.5	1	1.5
	180-190	30	1.1	0	0	1	0.3	1	0.3	10	2.0	12	1.9	6	0.9	5	1.7	2	0.7	2	0.9	1	0.8	1	1.5	1	1.5
	190-200	22	0.8	1	0.7	2	0.6	3	1.0	6	1.2	4	0.6	6	0.9	2	0.7	4	1.4	1	0.4	1	0.8	0	0	0	0
	200-210	11	0.4	0	0	0	0	0	0	4	0.8	6	1.0	1	0.1	1	0.3	1	0.4	0	0	0	0	0	0	0	0
	210-220	10	0.4	0	0	1	0.3	2	0.6	3	0.6	3	0.5	1	0.1	2	0.7	0	0	1	0.4	0	0	0	0	0	0
	220mg/dl以上	6	0.2	0	0	1	0.3	0	0	2	0.4	3	0.5	0	0	3	1.0	0	0	0	0	0	0	0	0	0	0
総数	2,610	100	144	100	337	100	314	100	494	100	621	100	700	100	299	100	282	100	226	100	127	100	65	100			

第46表の1-1 LDL-コレステロール値（Friedewaldの式）の平均値及び標準偏差（性・年齢階級別）

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上		
		平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	
男性	111.3	30.6	98.6	27.7	113.9	31.1	114.4	34.6	118.0	30.9	111.8	29.6	107.9	29.2	111.0	30.8	111.6	29.0	106.5	29.9	103.6	28.2	103.9	28.5
女性	116.4	31.0	95.6	26.2	99.9	28.1	108.6	30.6	128.4	30.0	126.2	29.8	115.0	28.1	124.5	30.5	117.7	29.0	114.5	28.6	112.2	26.1	110.7	25.7

Friedewaldの式 LDLコレステロール=総コレステロール-HDLコレステロール-中性脂肪/5
 (中性脂肪(トリグリセライド)が400mg/dl未満の場合)

第 46 表の 1-2 LDL-コレステロール値 (Friedewald の式) の分布 (性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	40mg/dl未満	9	0.6	0	0	0	0	1	0.6	1	0.4	3	0.7	4	0.8	2	0.9	0	0	2	1.3	2	2.1	0	0	0	0
	40-50	13	0.8	1	1.0	1	0.6	0	0	3	1.2	5	1.2	3	0.6	3	1.4	2	1.0	1	0.6	0	0	0	0	0	0
	50-60	31	2.0	8	8.1	2	1.2	3	1.8	1	0.4	6	1.5	11	2.2	3	1.4	3	1.5	5	3.1	3	3.1	0	0	0	0
	60-70	65	4.1	7	7.1	4	2.4	10	6.0	5	2.0	19	4.7	20	4.1	12	5.5	8	3.9	6	3.8	4	4.1	2	6.7		
	70-80	112	7.1	9	9.1	13	7.9	11	6.6	19	7.5	21	5.2	39	7.9	12	5.5	17	8.3	10	6.3	7	7.2	5	16.7		
	80-90	148	9.4	13	13.1	20	12.2	11	6.6	19	7.5	34	8.4	51	10.4	19	8.7	14	6.8	14	8.8	18	18.6	5	16.7		
	90-100	195	12.3	16	16.2	20	12.2	19	11.4	26	10.3	44	10.8	70	14.3	22	10.1	23	11.2	24	15.1	21	21.6	2	6.7		
	100-110	194	12.3	13	13.1	21	12.8	21	12.6	28	11.1	58	14.3	53	10.8	33	15.1	24	11.7	15	9.4	11	11.3	3	10.0		
	110-120	226	14.3	12	12.1	21	12.8	29	17.4	29	11.5	63	15.5	72	14.7	29	13.3	29	14.1	30	18.9	9	9.3	4	13.3		
	120-130	180	11.4	6	6.1	14	8.5	16	9.6	31	12.3	51	12.5	62	12.6	27	12.4	29	14.1	21	13.2	8	8.2	4	13.3		
	130-140	153	9.7	6	6.1	17	10.4	15	9.0	37	14.6	40	9.8	38	7.7	21	9.6	20	9.8	13	8.2	4	4.1	1	3.3		
	140-150	81	5.1	4	4.0	8	4.9	10	6.0	18	7.1	19	4.7	22	4.5	8	3.7	10	4.9	5	3.1	6	6.2	1	3.3		
	150-160	81	5.1	2	2.0	9	5.5	10	6.0	21	8.3	20	4.9	19	3.9	14	6.4	11	5.4	6	3.8	1	1.0	1	3.3		
	160-170	41	2.6	1	1.0	4	2.4	2	1.2	5	2.0	14	3.4	15	3.1	7	3.2	9	4.4	3	1.9	2	2.1	1	3.3		
	170-180	23	1.5	0	0	3	1.8	5	3.0	4	1.6	3	0.7	8	1.6	2	0.9	4	2.0	3	1.9	0	0	1	3.3		
	180-190	14	0.9	1	1.0	4	2.4	2	1.2	3	1.2	2	0.5	2	0.4	1	0.5	1	0.5	0	0	1	1.0	0	0		
	190-200	8	0.5	0	0	2	1.2	1	0.6	1	0.4	3	0.7	1	0.2	2	0.9	0	0	1	0.6	0	0	0	0		
	200-210	5	0.3	0	0	1	0.6	0	0	2	0.8	1	0.2	1	0.2	0	0	1	0.5	0	0	0	0	0	0		
	210-220	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	220mg/dl以上	2	0.1	0	0	0	0	1	0.6	0	0	1	0.2	0	0	1	0.5	0	0	0	0	0	0	0	0		
総 数		1,581	100	99	100	164	100	167	100	253	100	407	100	491	100	218	100	205	100	159	100	97	100	30	100		
女 性	40mg/dl未満	2	0.1	0	0	0	0	2	0.7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	40-50	15	0.7	1	0.7	5	1.5	5	1.6	0	0	1	0.2	3	0.6	0	0	0	0	2	1.2	0	0	1	2.0		
	50-60	31	1.4	6	4.2	12	3.6	7	2.3	1	0.2	1	0.2	4	0.8	1	0.4	1	0.5	0	0	2	2.2	1	2.0		
	60-70	73	3.3	12	8.3	19	5.7	14	4.6	7	1.6	6	1.2	15	2.9	2	0.9	4	2.0	6	3.5	3	3.2	2	4.0		
	70-80	133	6.0	22	15.3	52	15.5	22	7.2	8	1.8	7	1.4	22	4.3	4	1.8	9	4.5	7	4.1	3	3.2	3	6.0		
	80-90	176	7.9	29	20.1	45	13.4	39	12.7	11	2.5	19	3.9	33	6.4	8	3.5	8	4.0	18	10.5	6	6.5	1	2.0		
	90-100	248	11.2	22	15.3	51	15.2	41	13.4	47	10.8	34	7.0	53	10.3	20	8.8	19	9.5	16	9.4	12	12.9	6	12.0		
	100-110	271	12.2	19	13.2	44	13.1	42	13.7	44	10.1	48	9.9	74	14.4	29	12.8	32	16.1	19	11.1	14	15.1	9	18.0		
	110-120	282	12.7	12	8.3	38	11.3	31	10.1	57	13.1	65	13.4	79	15.4	29	12.8	25	12.6	26	15.2	18	19.4	10	20.0		
	120-130	258	11.6	5	3.5	25	7.4	31	10.1	54	12.4	83	17.1	60	11.7	37	16.4	25	12.6	20	11.7	10	10.8	5	10.0		
	130-140	232	10.4	5	3.5	19	5.7	34	11.1	60	13.8	55	11.3	59	11.5	21	9.3	27	13.6	17	9.9	6	6.5	9	18.0		
	140-150	170	7.7	5	3.5	11	3.3	12	3.9	43	9.9	54	11.1	45	8.8	23	10.2	14	7.0	19	11.1	11	11.8	1	2.0		
	150-160	113	5.1	2	1.4	2	0.6	12	3.9	31	7.1	35	7.2	31	6.0	18	8.0	15	7.5	11	6.4	4	4.3	1	2.0		
	160-170	79	3.6	3	2.1	7	2.1	4	1.3	24	5.5	31	6.4	10	1.9	12	5.3	4	2.0	5	2.9	1	1.1	0	0		
	170-180	62	2.8	0	0	1	0.3	5	1.6	24	5.5	20	4.1	12	2.3	9	4.0	9	4.5	1	0.6	2	2.2	0	0		
	180-190	29	1.3	0	0	1	0.3	1	0.3	10	2.3	11	2.3	6	1.2	5	2.2	2	1.0	2	1.2	1	1.1	1	2.0		
	190-200	20	0.9	1	0.7	2	0.6	2	0.7	6	1.4	4	0.8	5	1.0	2	0.9	4	2.0	1	0.6	0	0	0	0		
	200-210	11	0.5	0	0	0	0	0	0	4	0.9	6	1.2	1	0.2	1	0.4	1	0.5	0	0	0	0	0	0		
	210-220	10	0.5	0	0	1	0.3	2	0.7	3	0.7	3	0.6	1	0.2	2	0.9	0	0	1	0.6	0	0	0	0		
	220mg/dl以上	6	0.3	0	0	1	0.3	0	0	2	0.5	3	0.6	0	0	3	1.3	0	0	0	0	0	0	0	0		
総 数		2,221	100	144	100	336	100	306	100	436	100	486	100	513	100	226	100	199	100	171	100	93	100	50	100		

注) コレステロールを下げる薬または中性脂肪(トリグリセライド)を下げる薬服用者除外

第 46 表の 1-2 LDL-コレステロール値 (Friedewald の式) の平均値及び標準偏差 (性・年齢階級別)

	総 数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上		
		平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	
男 性	111.6	30.6	98.6	27.7	114.1	31.3	114.3	34.7	117.7	30.3	112.2	30.1	108.8	29.0	112.1	31.3	113.1	29.5	108.5	28.8	101.1	27.2	105.7	29.3
女 性	117.6	31.8	95.6	26.2	99.8	28.1	108.1	30.5	129.9	30.3	130.7	30.0	118.4	28.2	129.3	31.3	122.7	28.8	117.4	28.8	115.1	26.6	111.1	25.3

注) コレステロールを下げる薬または中性脂肪(トリグリセライド)を下げる薬服用者除外

〔参考〕第46表の2-1 LDL-コレステロール値（直接測定法）の分布（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	40mg/dl未満	8	0.4	0	0	0	0	0	0	3	1.0	2	0.4	3	0.5	2	0.8	0	0	1	0.5	2	1.7	0	0	0	0
	40-50	7	0.4	0	0	0	0	2	1.1	1	0.3	0	0	4	0.7	0	0	1	0.4	3	1.6	0	0	0	0	0	0
	50-60	27	1.5	2	2.0	3	1.7	1	0.5	1	0.3	10	2.1	10	1.7	4	1.5	2	0.8	5	2.7	3	2.5	0	0	0	0
	60-70	59	3.2	9	9.0	1	0.6	4	2.2	7	2.4	12	2.5	26	4.5	8	3.1	13	5.3	9	4.9	3	2.5	1	3.0	1	3.0
	70-80	91	5.0	8	8.0	5	2.8	7	3.8	12	4.1	22	4.5	37	6.4	14	5.4	14	5.8	8	4.4	10	8.4	5	15.2	5	15.2
	80-90	137	7.5	7	7.0	13	7.3	13	7.0	20	6.8	32	6.6	52	9.0	24	9.3	20	8.2	17	9.3	11	9.2	4	12.1	4	12.1
	90-100	226	12.4	13	13.0	24	13.5	16	8.6	23	7.9	68	14.0	82	14.2	32	12.4	24	9.9	22	12.0	30	25.2	6	18.2	6	18.2
	100-110	224	12.3	16	16.0	25	14.0	25	13.5	31	10.6	61	12.6	66	11.4	33	12.7	25	10.3	24	13.1	11	9.2	6	18.2	6	18.2
	110-120	245	13.5	14	14.0	26	14.6	18	9.7	34	11.6	70	14.4	83	14.4	35	13.5	41	16.9	26	14.2	15	12.6	1	3.0	1	3.0
	120-130	224	12.3	8	8.0	16	9.0	31	16.8	31	10.6	67	13.8	71	12.3	35	13.5	31	12.8	29	15.8	7	5.9	4	12.1	4	12.1
	130-140	208	11.4	9	9.0	20	11.2	26	14.1	45	15.4	44	9.1	64	11.1	20	7.7	33	13.6	17	9.3	11	9.2	3	9.1	3	9.1
	140-150	116	6.4	6	6.0	18	10.1	11	5.9	26	8.9	37	7.6	18	3.1	20	7.7	7	2.9	6	3.3	5	4.2	0	0	0	0
	150-160	102	5.6	4	4.0	9	5.1	12	6.5	22	7.5	25	5.2	30	5.2	13	5.0	16	6.6	6	3.3	6	5.0	2	6.1	2	6.1
	160-170	68	3.7	3	3.0	6	3.4	7	3.8	14	4.8	19	3.9	19	3.3	11	4.2	10	4.1	7	3.8	2	1.7	0	0	0	0
	170-180	36	2.0	0	0	5	2.8	4	2.2	12	4.1	7	1.4	8	1.4	4	1.5	4	1.6	2	1.1	2	1.7	0	0	0	0
	180-190	15	0.8	0	0	3	1.7	4	2.2	3	1.0	2	0.4	3	0.5	0	0	1	0.4	1	0.5	0	0	1	3.0	1	3.0
	190-200	14	0.8	1	1.0	3	1.7	2	1.1	4	1.4	3	0.6	1	0.2	2	0.8	0	0	0	0	1	0.8	0	0	0	0
	200-210	8	0.4	0	0	0	0	1	0.5	3	1.0	3	0.6	1	0.2	1	0.4	1	0.4	0	0	0	0	0	0	0	0
	210-220	1	0.1	0	0	1	0.6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	220mg/dl以上	2	0.1	0	0	0	0	1	0.5	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0	0	0	0	0
	総数		1,818	100	100	100	178	100	185	100	292	100	485	100	578	100	259	100	243	100	183	100	119	100	33	100	
女 性	40mg/dl未満	2	0.1	0	0	0	0	1	0.3	0	0	0	0	1	0.1	0	0	1	0.4	0	0	0	0	0	0	0	0
	40-50	4	0.2	0	0	0	0	1	0.3	0	0	1	0.2	2	0.3	1	0.3	0	0	1	0.4	1	0.8	0	0	0	0
	50-60	21	0.8	2	1.4	9	2.7	4	1.3	1	0.2	0	0	5	0.7	0	0	1	0.4	2	0.9	1	0.8	1	1.5	1	1.5
	60-70	48	1.8	9	6.3	13	3.8	8	2.5	4	0.8	5	0.8	9	1.3	3	1.0	6	2.1	3	1.3	0	0	0	0	0	0
	70-80	107	4.1	10	6.9	28	8.3	19	6.0	5	1.0	13	2.1	32	4.5	7	2.3	8	2.8	11	4.8	8	6.3	5	7.7	5	7.7
	80-90	203	7.7	29	20.1	49	14.5	24	7.6	17	3.4	25	4.0	59	8.3	13	4.3	25	8.8	20	8.7	9	7.0	5	7.7	5	7.7
	90-100	282	10.7	25	17.4	53	15.7	43	13.6	32	6.4	54	8.7	75	10.6	26	8.7	30	10.5	26	11.3	13	10.2	6	9.2	6	9.2
	100-110	337	12.8	19	13.2	52	15.4	47	14.8	57	11.4	65	10.5	97	13.7	38	12.7	29	10.2	32	13.9	27	21.1	9	13.8	9	13.8
	110-120	344	13.1	13	9.0	43	12.7	32	10.1	56	11.2	82	13.2	118	16.7	44	14.7	49	17.2	36	15.7	19	14.8	14	21.5	14	21.5
	120-130	323	12.3	19	13.2	29	8.6	35	11.0	71	14.1	91	14.6	78	11.0	41	13.7	28	9.8	25	10.9	15	11.7	10	15.4	10	15.4
	130-140	286	10.9	4	2.8	19	5.6	31	9.8	66	13.1	85	13.7	81	11.4	36	12.0	32	11.2	26	11.3	13	10.2	10	15.4	10	15.4
	140-150	237	9.0	2	1.4	19	5.6	31	9.8	69	13.7	56	9.0	60	8.5	26	8.7	32	11.2	16	7.0	10	7.8	2	3.1	2	3.1
	150-160	165	6.3	7	4.9	6	1.8	15	4.7	39	7.8	51	8.2	47	6.6	22	7.3	19	6.7	21	9.1	6	4.7	1	1.5	1	1.5
	160-170	99	3.8	0	0	9	2.7	10	3.2	27	5.4	41	6.6	12	1.7	20	6.7	7	2.5	5	2.2	0	0	0	0	0	0
	170-180	74	2.8	3	2.1	4	1.2	8	2.5	25	5.0	20	3.2	14	2.0	9	3.0	7	2.5	3	1.3	3	2.3	1	1.5	1	1.5
	180-190	35	1.3	0	0	0	0	2	0.6	10	2.0	13	2.1	10	1.4	4	1.3	6	2.1	1	0.4	2	1.6	1	1.5	1	1.5
	190-200	34	1.3	1	0.7	1	0.3	3	0.9	16	3.2	7	1.1	6	0.8	3	1.0	4	1.4	1	0.4	1	0.8	0	0	0	0
	200-210	18	0.7	1	0.7	2	0.6	1	0.3	4	0.8	9	1.4	1	0.1	3	1.0	1	0.4	0	0	0	0	0	0	0	0
	210-220	7	0.3	0	0	0	0	1	0.3	3	0.6	2	0.3	1	0.1	2	0.7	0	0	1	0.4	0	0	0	0	0	0
	220mg/dl以上	5	0.2	0	0	2	0.6	1	0.3	0	0	2	0.3	0	0	2	0.7	0	0	0	0	0	0	0	0	0	0
	総数		2,631	100	144	100	338	100	317	100	502	100	622	100	708	100	300	100	285	100	230	100	128	100	65	100	

〔参考〕第46表の2-1 LDL-コレステロール値（直接測定法）の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男 性	116.0	30.2	107.5	28.3	120.7	29.3	121.7	33.3	123.2	31.8	115.7	29.2	110.7	28.5	114.3	30.2	114.6	28.3	109.5	28.6	106.3	28.5	105.4	27.3
女 性	120.8	30.3	103.4	27.3	106.7	28.7	116.1	30.5	132.5	29.1	128.9	29.5	117.6	27.5	126.9	30.3	120.8	28.9	116.6	27.1	114.6	26.5	113.3	23.8

[参考] 第46表の2-2 LDL-コレステロール値（直接測定法）の分布（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	40mg/dl未満	7	0.4	0	0	0	0	0	0	3	1.1	2	0.5	2	0.4	2	0.9	0	0	0	0	2	2.0	0	0	0	0
	40-50	6	0.4	0	0	0	0	2	1.2	1	0.4	0	0	3	0.6	0	0	1	0.5	2	1.3	0	0	0	0	0	0
	50-60	26	1.6	2	2.0	3	1.7	1	0.6	1	0.4	10	2.4	9	1.8	4	1.8	2	1.0	4	2.5	3	3.1	0	0	0	0
	60-70	52	3.2	9	9.0	1	0.6	4	2.3	7	2.7	11	2.6	20	4.0	7	3.2	10	4.8	8	5.0	1	1.0	1	1.0	1	3.3
	70-80	79	4.9	8	8.0	5	2.9	7	4.0	10	3.8	19	4.6	30	6.1	11	5.0	10	4.8	7	4.4	9	9.2	4	4.0	4	13.3
	80-90	122	7.5	7	7.0	13	7.5	13	7.5	18	6.9	27	6.5	44	8.9	20	9.0	18	8.7	12	7.5	10	10.2	4	4.0	4	13.3
	90-100	197	12.2	13	13.0	22	12.6	14	8.1	19	7.3	55	13.2	74	14.9	24	10.8	21	10.1	21	13.2	28	28.6	4	4.0	4	13.3
	100-110	196	12.1	16	16.0	25	14.4	20	11.6	28	10.7	53	12.7	54	10.9	30	13.5	18	8.7	22	13.8	8	8.2	6	6.0	6	20.0
	110-120	222	13.7	14	14.0	25	14.4	18	10.4	32	12.2	58	13.9	75	15.2	28	12.6	39	18.8	23	14.5	12	12.2	1	1.0	1	3.3
	120-130	205	12.7	8	8.0	16	9.2	31	17.9	29	11.1	58	13.9	63	12.7	30	13.5	26	12.5	26	16.4	7	7.1	4	4.0	4	13.3
	130-140	176	10.9	9	9.0	19	10.9	24	13.9	40	15.3	36	8.7	48	9.7	18	8.1	26	12.5	13	8.2	6	6.1	3	3.0	3	10.0
	140-150	106	6.5	6	6.0	18	10.3	11	6.4	22	8.4	32	7.7	17	3.4	18	8.1	7	3.4	6	3.8	4	4.1	0	0	0	0
	150-160	95	5.9	4	4.0	9	5.2	11	6.4	20	7.6	22	5.3	29	5.9	12	5.4	16	7.7	6	3.8	5	5.1	2	2.0	2	6.7
	160-170	64	4.0	3	3.0	6	3.4	6	3.5	14	5.3	19	4.6	16	3.2	11	5.0	8	3.8	6	3.8	2	2.0	0	0	0	0
	170-180	33	2.0	0	0	5	2.9	4	2.3	11	4.2	7	1.7	6	1.2	4	1.8	4	1.9	2	1.3	0	0	0	0	0	0
	180-190	13	0.8	0	0	3	1.7	4	2.3	2	0.8	1	0.2	3	0.6	0	0	1	0.5	1	0.6	0	0	0	0	1	3.3
	190-200	12	0.7	1	1.0	3	1.7	1	0.6	3	1.1	3	0.7	1	0.2	2	0.9	0	0	0	0	1	1.0	0	0	0	0
	200-210	6	0.4	0	0	0	0	1	0.6	2	0.8	2	0.5	1	0.2	0	0	1	0.5	0	0	0	0	0	0	0	0
	210-220	1	0.1	0	0	1	0.6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	220mg/dl以上	2	0.1	0	0	0	0	1	0.6	0	0	1	0.2	0	0	1	0.5	0	0	0	0	0	0	0	0	0	0
総数		1,620	100	100	100	174	100	173	100	262	100	416	100	495	100	222	100	208	100	159	100	98	100	30	100		
女 性	40mg/dl未満	1	0.0	0	0	0	0	1	0.3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	40-50	4	0.2	0	0	0	0	1	0.3	0	0	1	0.2	2	0.4	1	0.4	0	0	1	0.6	1	1.1	0	0	0	0
	50-60	19	0.8	2	1.4	9	2.7	4	1.3	1	0.2	0	0	3	0.6	0	0	0	0	1	0.6	1	1.1	1	2.0	0	0
	60-70	43	1.9	9	6.3	13	3.9	8	2.6	4	0.9	4	0.8	5	1.0	2	0.9	3	1.5	2	1.1	0	0	0	0	0	0
	70-80	95	4.2	10	6.9	28	8.3	19	6.1	5	1.1	10	2.1	23	4.4	5	2.2	6	3.0	8	4.6	5	5.3	4	4.0	4	8.0
	80-90	158	7.1	29	20.1	49	14.6	24	7.8	11	2.5	13	2.7	32	6.2	6	2.6	10	5.0	12	6.9	7	7.4	3	3.0	3	6.0
	90-100	232	10.4	25	17.4	53	15.8	43	13.9	27	6.1	34	7.0	50	9.6	18	7.9	18	9.0	21	12.0	5	5.3	6	6.0	6	12.0
	100-110	275	12.3	19	13.2	52	15.5	47	15.2	46	10.4	43	8.8	68	13.1	25	11.0	22	10.9	22	12.6	20	21.3	4	4.0	4	8.0
	110-120	267	11.9	13	9.0	41	12.2	30	9.7	49	11.1	50	10.3	84	16.2	27	11.9	33	16.4	23	13.1	16	17.0	12	12.0	12	24.0
	120-130	281	12.6	19	13.2	29	8.6	34	11.0	61	13.8	77	15.8	61	11.7	33	14.5	23	11.4	21	12.0	10	10.6	7	7.0	7	14.0
	130-140	247	11.0	4	2.8	19	5.7	28	9.1	60	13.5	72	14.8	64	12.3	28	12.3	23	11.4	23	13.1	8	8.5	10	10.0	10	20.0
	140-150	210	9.4	2	1.4	19	5.7	31	10.0	61	13.8	48	9.9	49	9.4	21	9.3	24	11.9	14	8.0	10	10.6	1	1.0	1	2.0
	150-160	151	6.7	7	4.9	6	1.8	14	4.5	37	8.4	46	9.4	41	7.9	21	9.3	17	8.5	17	9.7	6	6.4	1	1.0	1	2.0
	160-170	90	4.0	0	0	9	2.7	10	3.2	25	5.6	37	7.6	9	1.7	17	7.5	5	2.5	4	2.3	0	0	0	0	0	0
	170-180	70	3.1	3	2.1	4	1.2	7	2.3	23	5.2	20	4.1	13	2.5	9	4.0	7	3.5	3	1.7	3	3.2	0	0	0	0
	180-190	32	1.4	0	0	0	0	2	0.6	10	2.3	12	2.5	8	1.5	4	1.8	5	2.5	1	0.6	1	1.1	1	1.0	1	2.0
	190-200	34	1.5	1	0.7	1	0.3	3	1.0	16	3.6	7	1.4	6	1.2	3	1.3	4	2.0	1	0.6	1	1.1	0	0	0	0
	200-210	18	0.8	1	0.7	2	0.6	1	0.3	4	0.9	9	1.8	1	0.2	3	1.3	1	0.5	0	0	0	0	0	0	0	0
	210-220	7	0.3	0	0	0	0	1	0.3	3	0.7	2	0.4	1	0.2	2	0.9	0	0	1	0.6	0	0	0	0	0	0
	220mg/dl以上	5	0.2	0	0	2	0.6	1	0.3	0	0	2	0.4	0	0	2	0.9	0	0	0	0	0	0	0	0	0	0
総数		2,239	100	144	100	336	100	309	100	443	100	487	100	520	100	227	100	201	100	175	100	94	100	50	100		

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

[参考] 第46表の2-2 LDL-コレステロール値（直接測定法）の平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男 性	116.3	30.3	107.5	28.3	121.0	29.5	121.5	33.5	122.7	31.5	116.0	29.8	111.3	28.4	115.3	30.6	115.6	28.6	110.8	27.8	104.2	27.7	107.1	28.0
女 性	122.1	31.1	103.4	27.3	106.6	28.8	115.6	30.6	134.0	29.4	133.1	30.0	120.6	27.9	131.2	31.4	125.5	28.7	119.0	27.7	116.9	27.4	113.4	23.6

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

第47表の1 クレアチニンの分布（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上				
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
男 性	0.4mg/dl未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	0.4-0.5	1	0.1	0	0	0	0	0	0	0	0	0	0	0	1	0.2	0	0	1	0.4	0	0	0	0	0	0	0	
	0.5-0.6	41	2.3	2	2.0	3	1.7	2	1.1	4	1.4	14	2.9	16	2.8	5	1.9	6	2.5	9	4.9	1	0.8	0	0	0	0	
	0.6-0.7	210	11.6	16	16.0	16	9.0	17	9.2	38	13.0	62	12.8	61	10.6	31	12.0	28	11.5	22	12.0	9	7.6	2	6.1	2	6.1	
	0.7-0.8	491	27.0	28	28.0	58	32.6	58	31.4	76	26.0	140	28.9	131	22.7	77	29.7	54	22.2	41	22.4	30	25.2	6	18.2	6	18.2	
	0.8-0.9	524	28.8	35	35.0	57	32.0	69	37.3	95	32.5	121	24.9	147	25.4	68	26.3	67	27.6	54	29.5	19	16.0	7	21.2	7	21.2	
	0.9-1.0	285	15.7	15	15.0	29	16.3	25	13.5	54	18.5	74	15.3	88	15.2	36	13.9	36	14.8	24	13.1	22	18.5	6	18.2	6	18.2	
	1.0-1.2	201	11.1	4	4.0	14	7.9	12	6.5	18	6.2	58	12.0	95	16.4	33	12.7	40	16.5	22	12.0	25	21.0	8	24.2	8	24.2	
	1.2-1.4	34	1.9	0	0	0	0	1	0.5	4	1.4	9	1.9	20	3.5	3	1.2	9	3.7	5	2.7	5	4.2	1	3.0	1	3.0	
	1.4-1.6	10	0.6	0	0	0	0	0	0	1	0.3	2	0.4	7	1.2	2	0.8	0	0	1	0.5	3	2.5	3	9.1	3	9.1	
	1.6-1.8	8	0.4	0	0	0	0	1	0.5	0	0	3	0.6	4	0.7	2	0.8	2	0.8	1	0.5	1	0.8	0	0	0	0	
	1.8-2.0	4	0.2	0	0	0	0	0	0	0	0	0	0	4	0.7	0	0	0	0	2	1.1	2	1.7	0	0	0	0	
	2.0mg/dl以上	9	0.5	0	0	1	0.6	0	0	2	0.7	2	0.4	4	0.7	2	0.8	0	0	2	1.1	2	1.7	0	0	0	0	
	総数	1,818	100	100	100	178	100	185	100	292	100	485	100	578	100	259	100	243	100	183	100	119	100	33	100	33	100	
女 性	0.4mg/dl未満	14	0.5	2	1.4	3	0.9	1	0.3	2	0.4	2	0.3	4	0.6	1	0.3	3	1.1	0	0	1	0.8	0	0	0	0	
	0.4-0.5	288	10.9	25	17.4	50	14.8	39	12.3	64	12.7	57	9.2	53	7.5	29	9.7	24	8.4	17	7.4	8	6.3	4	6.2	4	6.2	
	0.5-0.6	859	32.6	54	37.5	121	35.8	132	41.6	172	34.3	201	32.3	179	25.3	85	28.3	87	30.5	54	23.5	26	20.3	12	18.5	12	18.5	
	0.6-0.7	905	34.4	50	34.7	119	35.2	99	31.2	185	36.9	235	37.8	217	30.6	119	39.7	92	32.3	74	32.2	38	29.7	13	20.0	13	20.0	
	0.7-0.8	372	14.1	13	9.0	36	10.7	42	13.2	59	11.8	93	15.0	129	18.2	52	17.3	47	16.5	51	22.2	24	18.8	7	10.8	7	10.8	
	0.8-0.9	101	3.8	0	0	8	2.4	3	0.9	17	3.4	26	4.2	47	6.6	9	3.0	17	6.0	13	5.7	10	7.8	7	10.8	7	10.8	
	0.9-1.0	37	1.4	0	0	1	0.3	1	0.3	0	0	4	0.6	31	4.4	2	0.7	9	3.2	9	3.9	6	4.7	7	10.8	7	10.8	
	1.0-1.2	30	1.1	0	0	0	0	0	0	2	0.4	0	0	28	4.0	0	0	5	1.8	6	2.6	11	8.6	6	9.2	6	9.2	
	1.2-1.4	14	0.5	0	0	0	0	0	0	1	0.2	1	0.2	12	1.7	1	0.3	0	0	3	1.3	3	2.3	6	9.2	6	9.2	
	1.4-1.6	5	0.2	0	0	0	0	0	0	0	0	2	0.3	3	0.4	1	0.3	0	0	2	0.9	0	0	1	1.5	1	1.5	
	1.6-1.8	4	0.2	0	0	0	0	0	0	0	0	1	0.2	3	0.4	1	0.3	1	0.4	1	0.4	0	0	1	1.5	1	1.5	
	1.8-2.0	1	0.0	0	0	0	0	0	0	0	0	0	0	1	0.1	0	0	0	0	0	0	0	1	0.8	0	0	0	
	2.0mg/dl以上	1	0.0	0	0	0	0	0	0	0	0	0	0	1	0.1	0	0	0	0	0	0	0	0	0	1	1.5	1	1.5
	総数	2,631	100	144	100	338	100	317	100	502	100	622	100	708	100	300	100	285	100	230	100	128	100	65	100	65	100	

第47表の1 クレアチニンの平均値及び標準偏差（性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上	
	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差	平均値	標準偏差
男 性	0.9	0.4	0.8	0.1	0.8	0.2	0.8	0.1	0.9	0.7	0.9	0.2	0.9	0.3	0.9	0.3	0.9	0.2	0.9	0.3	1.0	0.4	1.0	0.2
女 性	0.6	0.1	0.6	0.1	0.6	0.1	0.6	0.1	0.6	0.1	0.6	0.1	0.7	0.2	0.6	0.1	0.6	0.1	0.7	0.2	0.7	0.2	0.8	0.3

第 47 表の 2 推算 GFR の分布 (性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-69歳		(再掲) 70-74歳		(再掲) 75-79歳		(再掲) 80-84歳		(再掲) 85歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	15ml/min/1.73m ² 未満	5	0.3	0	0	0	0	0	0	2	0.7	1	0.2	2	0.3	1	0.4	0	0	1	0.5	1	0.8	0	0	0	0
	15-29	9	0.5	0	0	1	0.6	0	0	0	0	1	0.2	7	1.2	1	0.4	0	0	3	1.6	4	3.4	0	0	0	0
	30-59	280	15.4	0	0	0	0	5	2.7	14	4.8	78	16.1	183	31.7	46	17.8	68	28.0	44	24.0	53	44.5	18	54.5		
	60-89	1,176	64.7	24	24.0	100	56.2	133	71.9	230	78.8	344	70.9	345	59.7	183	70.7	155	63.8	119	65.0	57	47.9	14	42.4		
	90ml/min/1.73m ² 以上	348	19.1	76	76.0	77	43.3	47	25.4	46	15.8	61	12.6	41	7.1	28	10.8	20	8.2	16	8.7	4	3.4	1	3.0		
	総 数	1,818	100	100	100	178	100	185	100	292	100	485	100	578	100	259	100	243	100	183	100	119	100	33	100		
女 性	15ml/min/1.73m ² 未満	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	15-29	14	0.5	0	0	0	0	0	0	0	0	3	0.5	11	1.6	2	0.7	1	0.4	3	1.3	2	1.6	5	7.7		
	30-59	318	12.1	0	0	1	0.3	3	0.9	30	6.0	72	11.6	212	29.9	42	14.0	60	21.1	71	30.9	50	39.1	31	47.7		
	60-89	1,607	61.1	33	22.9	156	46.2	192	60.6	347	69.1	449	72.2	430	60.7	217	72.3	193	67.7	140	60.9	71	55.5	26	40.0		
	90ml/min/1.73m ² 以上	692	26.3	111	77.1	181	53.6	122	38.5	125	24.9	98	15.8	55	7.8	39	13.0	31	10.9	16	7.0	5	3.9	3	4.6		
	総 数	2,631	100	144	100	338	100	317	100	502	100	622	100	708	100	300	100	285	100	230	100	128	100	65	100		
総 数	15ml/min/1.73m ² 未満	5	0.1	0	0	0	0	0	0	2	0.3	1	0.1	2	0.2	1	0.2	0	0	1	0.2	1	0.4	0	0	0	0
	15-29	23	0.5	0	0	1	0.2	0	0	0	0	4	0.4	18	1.4	3	0.5	1	0.2	6	1.5	6	2.4	5	5.1		
	30-59	598	13.4	0	0	1	0.2	8	1.6	44	5.5	150	13.6	395	30.7	88	15.7	128	24.2	115	27.8	103	41.7	49	50.0		
	60-89	2,783	62.6	57	23.4	256	49.6	325	64.7	577	72.7	793	71.6	775	60.3	400	71.6	348	65.9	259	62.7	128	51.8	40	40.8		
	90ml/min/1.73m ² 以上	1,040	23.4	187	76.6	258	50.0	169	33.7	171	21.5	159	14.4	96	7.5	67	12.0	51	9.7	32	7.7	9	3.6	4	4.1		
	総 数	4,449	100	244	100	516	100	502	100	794	100	1,107	100	1,286	100	559	100	528	100	413	100	247	100	98	100		

「エビデンスに基づく CKD 診療ガイドライン 2009」(日本腎臓学会)

推算 GFR (ml/min/1.73m²) = 194 × 血清クレアチニン値 (mg/dl)^{-1.094} × 年齢 (歳)^{-0.287} (男性)

= 194 × 血清クレアチニン値 (mg/dl)^{-1.094} × 年齢 (歳)^{-0.287} × 0.739 (女性)

第48表の1 メタボリックシンドローム（内臓脂肪症候群）の状況（性・年齢階級別）

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%		
総数	総数	4,406	100	236	100	506	100	500	100	790	100	1,099	100	1,275	100	2,914	100	1,079	100	750	100
	腹囲85cm以上(男), 90cm以上(女)+項目2つ以上該当	733	16.6	6	2.5	20	4.0	46	9.2	115	14.6	229	20.8	317	24.9	528	18.1	273	25.3	179	23.9
	腹囲85cm以上(男), 90cm以上(女)+項目1つ該当	610	13.8	16	6.8	43	8.5	58	11.6	113	14.3	167	15.2	213	16.7	433	14.9	186	17.2	118	15.7
	上記以外	3,063	69.5	214	90.7	443	87.5	396	79.2	562	71.1	703	64.0	745	58.4	1,953	67.0	620	57.5	453	60.4
男性	総数	1,806	100	99	100	177	100	185	100	291	100	482	100	572	100	1,199	100	498	100	331	100
	腹囲85cm以上+項目2つ以上該当	457	25.3	4	4.0	14	7.9	31	16.8	78	26.8	140	29.0	190	33.2	324	27.0	157	31.5	115	34.7
	腹囲85cm以上+項目1つ該当	395	21.9	11	11.1	31	17.5	49	26.5	77	26.5	106	22.0	121	21.2	294	24.5	117	23.5	59	17.8
	上記以外	954	52.8	84	84.8	132	74.6	105	56.8	136	46.7	236	49.0	261	45.6	581	48.5	224	45.0	157	47.4
女性	総数	2,600	100	137	100	329	100	315	100	499	100	617	100	703	100	1,715	100	581	100	419	100
	腹囲90cm以上+項目2つ以上該当	276	10.6	2	1.5	6	1.8	15	4.8	37	7.4	89	14.4	127	18.1	204	11.9	116	20.0	64	15.3
	腹囲90cm以上+項目1つ該当	215	8.3	5	3.6	12	3.6	9	2.9	36	7.2	61	9.9	92	13.1	139	8.1	69	11.9	59	14.1
	上記以外	2,109	81.1	130	94.9	311	94.5	291	92.4	426	85.4	467	75.7	484	68.8	1,372	80.0	396	68.2	296	70.6

注) 妊婦除外, ヘモグロビン A1c \geq 5.5%の場合 項目の基準は p. 81 参照

第48表の2 メタボリックシンドローム（内臓脂肪症候群）の状況（性・年齢階級別）

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%		
総数	総数	4,406	100	236	100	506	100	500	100	790	100	1,099	100	1,275	100	2,914	100	1,079	100	750	100
	腹囲85cm以上(男), 90cm以上(女)+項目2つ以上該当	657	14.9	4	1.7	18	3.6	41	8.2	97	12.3	207	18.8	290	22.7	469	16.1	247	22.9	166	22.1
	腹囲85cm以上(男), 90cm以上(女)+項目1つ該当	674	15.3	16	6.8	44	8.7	63	12.6	129	16.3	185	16.8	237	18.6	483	16.6	208	19.3	131	17.5
	上記以外	3,075	69.8	216	91.5	444	87.7	396	79.2	564	71.4	707	64.3	748	58.7	1,962	67.3	624	57.8	453	60.4
男性	総数	1,806	100	99	100	177	100	185	100	291	100	482	100	572	100	1,199	100	498	100	331	100
	腹囲85cm以上+項目2つ以上該当	411	22.8	3	3.0	13	7.3	27	14.6	68	23.4	126	26.1	174	30.4	288	24.0	140	28.1	107	32.3
	腹囲85cm以上+項目1つ該当	436	24.1	12	12.1	31	17.5	53	28.6	86	29.6	118	24.5	136	23.8	326	27.2	133	26.7	67	20.2
	上記以外	959	53.1	84	84.8	133	75.1	105	56.8	137	47.1	238	49.4	262	45.8	585	48.8	225	45.2	157	47.4
女性	総数	2,600	100	137	100	329	100	315	100	499	100	617	100	703	100	1,715	100	581	100	419	100
	腹囲90cm以上+項目2つ以上該当	246	9.5	1	0.7	5	1.5	14	4.4	29	5.8	81	13.1	116	16.5	181	10.6	107	18.4	59	14.1
	腹囲90cm以上+項目1つ該当	238	9.2	4	2.9	13	4.0	10	3.2	43	8.6	67	10.9	101	14.4	157	9.2	75	12.9	64	15.3
	上記以外	2,116	81.4	132	96.4	311	94.5	291	92.4	427	85.6	469	76.0	486	69.1	1,377	80.3	399	68.7	296	70.6

注) 妊婦除外, ヘモグロビン A1c \geq 5.6%の場合 項目の基準は p. 81 参照

第49表の1 腹囲区分別、血中脂質、血圧、血糖のいずれかのリスクを有する割合（性・年齢階級別）

		総 数		0		1		2		3		
		人数	%	人数	%	人数	%	人数	%	人数	%	
総 数	総 数	85 cm未満(男)/90 cm未満(女)	2,873	100.0	1,056	36.8	1,155	40.2	518	18.0	144	5.0
		85 cm以上(男)/90 cm以上(女)	1,533	100.0	190	12.4	610	39.8	544	35.5	189	12.3
	20-29歳	85 cm未満(男)/90 cm未満(女)	201	100.0	177	88.1	24	11.9	0	0	0	0
		85 cm以上(男)/90 cm以上(女)	35	100.0	13	37.1	16	45.7	6	17.1	0	0
	30-39歳	85 cm未満(男)/90 cm未満(女)	394	100.0	308	78.2	79	20.1	6	1.5	1	0.3
		85 cm以上(男)/90 cm以上(女)	112	100.0	49	43.8	43	38.4	20	17.9	0	0
	40-49歳	85 cm未満(男)/90 cm未満(女)	367	100.0	206	56.1	137	37.3	21	5.7	3	0.8
		85 cm以上(男)/90 cm以上(女)	133	100.0	29	21.8	58	43.6	37	27.8	9	6.8
	50-59歳	85 cm未満(男)/90 cm未満(女)	529	100.0	156	29.5	261	49.3	96	18.1	16	3.0
		85 cm以上(男)/90 cm以上(女)	261	100.0	33	12.6	113	43.3	92	35.2	23	8.8
	60-69歳	85 cm未満(男)/90 cm未満(女)	667	100.0	145	21.7	330	49.5	147	22.0	45	6.7
		85 cm以上(男)/90 cm以上(女)	432	100.0	36	8.3	167	38.7	163	37.7	66	15.3
	70歳以上	85 cm未満(男)/90 cm未満(女)	715	100.0	64	9.0	324	45.3	248	34.7	79	11.0
		85 cm以上(男)/90 cm以上(女)	560	100.0	30	5.4	213	38.0	226	40.4	91	16.3
(再掲)40-74歳	85 cm未満(男)/90 cm未満(女)	1,839	100.0	537	29.2	857	46.6	351	19.1	94	5.1	
	85 cm以上(男)/90 cm以上(女)	1,075	100.0	114	10.6	433	40.3	391	36.4	137	12.7	
(再掲)65-74歳	85 cm未満(男)/90 cm未満(女)	588	100.0	95	16.2	274	46.6	162	27.6	57	9.7	
	85 cm以上(男)/90 cm以上(女)	491	100.0	32	6.5	186	37.9	195	39.7	78	15.9	
(再掲)75歳以上	85 cm未満(男)/90 cm未満(女)	439	100.0	34	7.7	195	44.4	161	36.7	49	11.2	
	85 cm以上(男)/90 cm以上(女)	311	100.0	14	4.5	118	37.9	127	40.8	52	16.7	
男 性	総 数	85 cm未満	834	100.0	250	30.0	380	45.6	169	20.3	35	4.2
		85 cm以上	972	100.0	120	12.3	395	40.6	337	34.7	120	12.3
	20-29歳	85 cm未満	75	100.0	60	80.0	15	20.0	0	0	0	0
		85 cm以上	24	100.0	9	37.5	11	45.8	4	16.7	0	0
	30-39歳	85 cm未満	96	100.0	60	62.5	32	33.3	3	3.1	1	1.0
		85 cm以上	81	100.0	36	44.4	31	38.3	14	17.3	0	0
	40-49歳	85 cm未満	87	100.0	42	48.3	36	41.4	8	9.2	1	1.1
		85 cm以上	98	100.0	18	18.4	49	50.0	24	24.5	7	7.1
	50-59歳	85 cm未満	116	100.0	30	25.9	65	56.0	19	16.4	2	1.7
		85 cm以上	175	100.0	20	11.4	77	44.0	61	34.9	17	9.7
	60-69歳	85 cm未満	217	100.0	37	17.1	117	53.9	50	23.0	13	6.0
		85 cm以上	265	100.0	19	7.2	106	40.0	96	36.2	44	16.6
	70歳以上	85 cm未満	243	100.0	21	8.6	115	47.3	89	36.6	18	7.4
		85 cm以上	329	100.0	18	5.5	121	36.8	138	41.9	52	15.8
(再掲)40-74歳	85 cm未満	512	100.0	117	22.9	264	51.6	109	21.3	22	4.3	
	85 cm以上	687	100.0	69	10.0	294	42.8	237	34.5	87	12.7	
(再掲)65-74歳	85 cm未満	202	100.0	30	14.9	100	49.5	60	29.7	12	5.9	
	85 cm以上	296	100.0	22	7.4	117	39.5	115	38.9	42	14.2	
(再掲)75歳以上	85 cm未満	151	100.0	13	8.6	69	45.7	57	37.7	12	7.9	
	85 cm以上	180	100.0	6	3.3	59	32.8	82	45.6	33	18.3	
女 性	総 数	90 cm未満	2,039	100.0	806	39.5	775	38.0	349	17.1	109	5.3
		90 cm以上	561	100.0	70	12.5	215	38.3	207	36.9	69	12.3
	20-29歳	90 cm未満	126	100.0	117	92.9	9	7.1	0	0	0	0
		90 cm以上	11	100.0	4	36.4	5	45.5	2	18.2	0	0
	30-39歳	90 cm未満	298	100.0	248	83.2	47	15.8	3	1.0	0	0
		90 cm以上	31	100.0	13	41.9	12	38.7	6	19.4	0	0
	40-49歳	90 cm未満	280	100.0	164	58.6	101	36.1	13	4.6	2	0.7
		90 cm以上	35	100.0	11	31.4	9	25.7	13	37.1	2	5.7
	50-59歳	90 cm未満	413	100.0	126	30.5	196	47.5	77	18.6	14	3.4
		90 cm以上	86	100.0	13	15.1	36	41.9	31	36.0	6	7.0
	60-69歳	90 cm未満	450	100.0	108	24.0	213	47.3	97	21.6	32	7.1
		90 cm以上	167	100.0	17	10.2	61	36.5	67	40.1	22	13.2
	70歳以上	90 cm未満	472	100.0	43	9.1	209	44.3	159	33.7	61	12.9
		90 cm以上	231	100.0	12	5.2	92	39.8	88	38.1	39	16.9
(再掲)40-74歳	90 cm未満	1,327	100.0	420	31.7	593	44.7	242	18.2	72	5.4	
	90 cm以上	388	100.0	45	11.6	139	35.8	154	39.7	50	12.9	
(再掲)65-74歳	90 cm未満	386	100.0	65	16.8	174	45.1	102	26.4	45	11.7	
	90 cm以上	195	100.0	10	5.1	69	35.4	80	41.0	36	18.5	
(再掲)75歳以上	90 cm未満	288	100.0	21	7.3	126	43.8	104	36.1	37	12.8	
	90 cm以上	131	100.0	8	6.1	59	45.0	45	34.4	19	14.5	

注) 妊婦除外、ヘモグロビン A1c≧5.5%の場合 項目の基準はp.55参照

第49表の2 腹囲区分別, 血中脂質, 血圧, 血糖のリスクの保有状況 [すべての組み合わせ①血中脂質②血圧③血糖] (性・年齢階級別)

			総数		②有③有④有		②有③有④無		②無③有④有		②有③無④有		②有③無④無		②無③有④無		②無③無④有		②無③無④無	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総数	85cm未満(男)/90cm未満(女)	2,873	100.0	144	5.0	147	5.1	336	11.7	35	1.2	70	2.4	893	31.1	192	6.7	1,056	36.8
		85cm以上(男)/90cm以上(女)	1,533	100.0	189	12.3	157	10.2	363	23.7	24	1.6	25	1.6	516	33.7	69	4.5	190	12.4
20-29歳		85cm未満(男)/90cm未満(女)	201	100.0	0	0	0	0	0	0	0	0	2	1.0	15	7.5	7	3.5	177	88.1
		85cm以上(男)/90cm以上(女)	35	100.0	0	0	3	8.6	3	8.6	0	0	2	5.7	11	31.4	3	8.6	13	37.1
30-39歳		85cm未満(男)/90cm未満(女)	394	100.0	1	0.3	2	0.5	4	1.0	0	0	6	1.5	52	13.2	21	5.3	308	78.2
		85cm以上(男)/90cm以上(女)	112	100.0	0	0	10	8.9	8	7.1	2	1.8	5	4.5	33	29.5	5	4.5	49	43.8
40-49歳		85cm未満(男)/90cm未満(女)	367	100.0	3	0.8	6	1.6	14	3.8	1	0.3	10	2.7	96	26.2	31	8.4	206	56.1
		85cm以上(男)/90cm以上(女)	133	100.0	9	6.8	12	9.0	24	18.0	1	0.8	2	1.5	49	36.8	7	5.3	29	21.8
50-59歳		85cm未満(男)/90cm未満(女)	529	100.0	16	3.0	22	4.2	64	12.1	10	1.9	15	2.8	202	38.2	44	8.3	156	29.5
		85cm以上(男)/90cm以上(女)	261	100.0	23	8.8	24	9.2	65	24.9	3	1.1	1	0.4	94	36.0	18	6.9	33	12.6
60-69歳		85cm未満(男)/90cm未満(女)	667	100.0	45	6.7	41	6.1	93	13.9	13	1.9	22	3.3	254	38.1	54	8.1	145	21.7
		85cm以上(男)/90cm以上(女)	432	100.0	66	15.3	39	9.0	113	26.2	11	2.5	8	1.9	139	32.2	20	4.6	36	8.3
70歳以上		85cm未満(男)/90cm未満(女)	715	100.0	79	11.0	76	10.6	161	22.5	11	1.5	15	2.1	274	38.3	35	4.9	64	9.0
		85cm以上(男)/90cm以上(女)	560	100.0	91	16.3	69	12.3	150	26.8	7	1.3	7	1.3	190	33.9	16	2.9	30	5.4
(再掲)40-74歳		85cm未満(男)/90cm未満(女)	1,839	100.0	94	5.1	96	5.2	226	12.3	29	1.6	56	3.0	659	35.8	142	7.7	537	29.2
		85cm以上(男)/90cm以上(女)	1,075	100.0	137	12.7	104	9.7	266	24.7	21	2.0	15	1.4	363	33.8	55	5.1	114	10.6
(再掲)65-74歳		85cm未満(男)/90cm未満(女)	588	100.0	57	9.7	51	8.7	100	17.0	11	1.9	15	2.6	221	37.6	38	6.5	95	16.2
		85cm以上(男)/90cm以上(女)	491	100.0	78	15.9	54	11.0	127	25.9	14	2.9	7	1.4	157	32.0	22	4.5	32	6.5
(再掲)75歳以上		85cm未満(男)/90cm未満(女)	439	100.0	49	11.2	49	11.2	106	24.1	6	1.4	6	1.4	167	38.0	22	5.0	34	7.7
		85cm以上(男)/90cm以上(女)	311	100.0	52	16.7	40	12.9	86	27.7	1	0.3	3	1.0	109	35.0	6	1.9	14	4.5
男性	総数	85cm未満	834	100.0	35	4.2	34	4.1	124	14.9	11	1.3	19	2.3	320	38.4	41	4.9	250	30.0
		85cm以上	972	100.0	120	12.3	102	10.5	222	22.8	13	1.3	15	1.5	341	35.1	39	4.0	120	12.3
20-29歳		85cm未満	75	100.0	0	0	0	0	0	0	0	0	1	1.3	14	18.7	0	0	60	80.0
		85cm以上	24	100.0	0	0	3	12.5	1	4.2	0	0	1	4.2	9	37.5	1	4.2	9	37.5
30-39歳		85cm未満	96	100.0	1	1.0	0	0	3	3.1	0	0	3	3.1	28	29.2	1	1.0	60	62.5
		85cm以上	81	100.0	0	0	8	9.9	5	6.2	1	1.2	5	6.2	23	28.4	3	3.7	36	44.4
40-49歳		85cm未満	87	100.0	1	1.1	3	3.4	5	5.7	0	0	6	6.9	27	31.0	3	3.4	42	48.3
		85cm以上	98	100.0	7	7.1	11	11.2	13	13.3	0	0	2	2.0	41	41.8	6	6.1	18	18.4
50-59歳		85cm未満	116	100.0	2	1.7	5	4.3	10	8.6	4	3.4	1	0.9	57	49.1	7	6.0	30	25.9
		85cm以上	175	100.0	17	9.7	18	10.3	42	24.0	1	0.6	1	0.6	66	37.7	10	5.7	20	11.4
60-69歳		85cm未満	217	100.0	13	6.0	9	4.1	37	17.1	4	1.8	3	1.4	100	46.1	14	6.5	37	17.1
		85cm以上	265	100.0	44	16.6	22	8.3	69	26.0	5	1.9	4	1.5	91	34.3	11	4.2	19	7.2
70歳以上		85cm未満	243	100.0	18	7.4	17	7.0	69	28.4	3	1.2	5	2.1	94	38.7	16	6.6	21	8.6
		85cm以上	329	100.0	52	15.8	40	12.2	92	28.0	6	1.8	2	0.6	111	33.7	8	2.4	18	5.5
(再掲)40-74歳		85cm未満	512	100.0	22	4.3	22	4.3	77	15.0	10	2.0	13	2.5	222	43.4	29	5.7	117	22.9
		85cm以上	687	100.0	87	12.7	66	9.6	160	23.3	11	1.6	9	1.3	254	37.0	31	4.5	69	10.0
(再掲)65-74歳		85cm未満	202	100.0	12	5.9	11	5.4	44	21.8	5	2.5	4	2.0	83	41.1	13	6.4	30	14.9
		85cm以上	296	100.0	42	14.2	29	9.8	77	26.0	9	3.0	4	1.4	103	34.8	10	3.4	22	7.4
(再掲)75歳以上		85cm未満	151	100.0	12	7.9	12	7.9	44	29.1	1	0.7	2	1.3	56	37.1	11	7.3	13	8.6
		85cm以上	180	100.0	33	18.3	25	13.9	56	31.1	1	0.6	0	0	55	30.6	4	2.2	6	3.3
女性	総数	90cm未満	2,039	100.0	109	5.3	113	5.5	212	10.4	24	1.2	51	2.5	573	28.1	151	7.4	806	39.5
		90cm以上	561	100.0	69	12.3	55	9.8	141	25.1	11	2.0	10	1.8	175	31.2	30	5.3	70	12.5
20-29歳		90cm未満	126	100.0	0	0	0	0	0	0	0	0	1	0.8	1	0.8	7	5.6	117	92.9
		90cm以上	11	100.0	0	0	0	0	2	18.2	0	0	1	9.1	2	18.2	2	18.2	4	36.4
30-39歳		90cm未満	298	100.0	0	0	2	0.7	1	0.3	0	0	3	1.0	24	8.1	20	6.7	248	83.2
		90cm以上	31	100.0	0	0	2	6.5	3	9.7	1	3.2	0	0	10	32.3	2	6.5	13	41.9
40-49歳		90cm未満	280	100.0	2	0.7	3	1.1	9	3.2	1	0.4	4	1.4	69	24.6	28	10.0	164	58.6
		90cm以上	35	100.0	2	5.7	1	2.9	11	31.4	1	2.9	0	0	8	22.9	1	2.9	11	31.4
50-59歳		90cm未満	413	100.0	14	3.4	17	4.1	54	13.1	6	1.5	14	3.4	145	35.1	37	9.0	126	30.5
		90cm以上	86	100.0	6	7.0	6	7.0	23	26.7	2	2.3	0	0	28	32.6	8	9.3	13	15.1
60-69歳		90cm未満	450	100.0	32	7.1	32	7.1	56	12.4	9	2.0	19	4.2	154	34.2	40	8.9	108	24.0
		90cm以上	167	100.0	22	13.2	17	10.2	44	26.3	6	3.6	4	2.4	48	28.7	9	5.4	17	10.2
70歳以上		90cm未満	472	100.0	61	12.9	59	12.5	92	19.5	8	1.7	10	2.1	180	38.1	19	4.0	43	9.1
		90cm以上	231	100.0	39	16.9	29	12.6	58	25.1	1	0.4	5	2.2	79	34.2	8	3.5	12	5.2
(再掲)40-74歳		90cm未満	1,327	100.0	72	5.4	74	5.6	149	11.2	19	1.4	43	3.2	437	32.9	113	8.5	420	31.7
		90cm以上	388	100.0	50	12.9	38	9.8	106	27.3	10	2.6	6	1.5	109	28.1	24	6.2	45	11.6
(再掲)65-74歳		90cm未満	386	100.0	45	11.7	40	10.4	56	14.5	6	1.6	11	2.8	138	35.8	25	6.5	65	16.8
		90cm以上	195	100.0	36	18.5	25	12.8	50	25.6	5	2.6	3	1.5	54	27.7	12	6.2	10	5.1
(再掲)75歳以上		90cm未満	288	100.0	37	12.8	37	12.8	62	21.5	5	1.7	4	1.4	111	38.5	11	3.8	21	7.3
		90cm以上	131	100.0	19	14.5	15	11.5	30	22.9	0	0	3	2.3	54	41.2	2	1.5	8	6.1

注) 妊婦除外, ヘモグロビン A1c \geq 5.5%の場合 リスクの項目の基準は p.55 参照

第50表 「糖尿病が強く疑われる人」及び「糖尿病の可能性を否定できない人」の状況
(性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	糖尿病が強く疑われる人	211	11.6	0	0	2	1.1	9	4.9	34	11.6	67	13.8	99	17.3	151	12.6	78	15.6	58	17.5
	(再掲)糖尿病が強く疑われる 人のうち服薬者*	99	46.9	0	0	0	0	3	33.3	13	38.2	32	47.8	51	51.5	76	50.3	47	60.3	23	39.7
	糖尿病の可能性を否定できない人 上記以外	273	15.1	2	2.0	8	4.5	16	8.6	41	14.0	95	19.6	111	19.4	192	16.0	93	18.6	71	21.4
	総 数	1,329	73.3	99	98.0	168	94.4	160	86.5	217	74.3	322	66.5	363	63.4	859	71.5	328	65.7	203	61.1
女 性	糖尿病が強く疑われる人	168	6.4	1	0.7	3	0.9	7	2.2	30	6.0	62	10.0	65	9.2	136	7.9	72	12.4	28	6.7
	(再掲)糖尿病が強く疑われる 人のうち服薬者*	82	48.8	1	100	1	33.3	1	14.3	14	46.7	31	50.0	34	52.3	68	50.0	41	56.9	12	42.9
	糖尿病の可能性を否定できない人 上記以外	375	14.3	2	1.4	14	4.1	30	9.5	75	15.0	99	16.0	155	22.0	264	15.3	119	20.4	95	22.6
	総 数	2,078	79.3	138	97.9	321	95.0	279	88.3	395	79.0	459	74.0	486	68.8	1,321	76.8	391	67.2	298	70.8
総 数	糖尿病が強く疑われる人	379	8.5	1	0.4	5	1.0	16	3.2	64	8.1	129	11.7	164	12.8	287	9.8	150	13.9	86	11.4
	(再掲)糖尿病が強く疑われる 人のうち服薬者*	181	47.8	1	100	1	20.0	4	25.0	27	42.2	63	48.8	85	51.8	144	50.2	88	58.7	35	40.7
	糖尿病の可能性を否定できない人 上記以外	648	14.6	4	1.7	22	4.3	46	9.2	116	14.6	194	17.6	266	20.8	456	15.6	212	19.6	166	22.0
	総 数	4,434	100	242	100	516	100	501	100	792	100	1,104	100	1,279	100	2,923	100	1,081	100	753	100

注) ヘモグロビン A1c \geq 5.6%の場合

※ 「服薬者」とは、質問票で「インスリン注射または血糖を下げる薬」の使用有と回答した者

判定の基準

- ① 「糖尿病が強く疑われる人」とは、ヘモグロビン A1c の値が 6.1%以上、または、質問票で「現在糖尿病の治療を受けている」と答えた人である。
- ② 「糖尿病の可能性を否定できない人」とは、ヘモグロビン A1c の値が 5.6%以上、6.1%未満で、①以外の人である。

第51表 「糖尿病が強く疑われる人」及び「糖尿病の可能性を否定できない人」の状況
(性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	糖尿病が強く疑われる人	211	11.6	0	0	2	1.1	9	4.9	34	11.6	67	13.8	99	17.3	151	12.6	78	15.6	58	17.5
	(再掲)糖尿病が強く疑われる 人のうち服薬者*	99	46.9	0	0	0	0	3	33.3	13	38.2	32	47.8	51	51.5	76	50.3	47	60.3	23	39.7
	糖尿病の可能性を否定できない人 上記以外	384	21.2	3	3.0	11	6.2	25	13.5	59	20.2	126	26.0	160	27.9	270	22.5	132	26.5	100	30.1
	総 数	1,218	67.2	98	97.0	165	92.7	151	81.6	199	68.2	291	60.1	314	54.8	781	65.0	289	57.9	174	52.4
女 性	糖尿病が強く疑われる人	168	6.4	1	0.7	3	0.9	7	2.2	30	6.0	62	10.0	65	9.2	136	7.9	72	12.4	28	6.7
	(再掲)糖尿病が強く疑われる 人のうち服薬者*	82	48.8	1	100	1	33.3	1	14.3	14	46.7	31	50.0	34	52.3	68	50.0	41	56.9	12	42.9
	糖尿病の可能性を否定できない人 上記以外	576	22.0	10	7.1	24	7.1	48	15.2	120	24.0	156	25.2	218	30.9	408	23.7	164	28.2	134	31.8
	総 数	1,877	71.6	130	92.2	311	92.0	261	82.6	350	70.0	402	64.8	423	59.9	1,177	68.4	346	59.5	259	61.5
総 数	糖尿病が強く疑われる人	379	8.5	1	0.4	5	1.0	16	3.2	64	8.1	129	11.7	164	12.8	287	9.8	150	13.9	86	11.4
	(再掲)糖尿病が強く疑われる 人のうち服薬者*	181	47.8	1	100	1	20.0	4	25.0	27	42.2	63	48.8	85	51.8	144	50.2	88	58.7	35	40.7
	糖尿病の可能性を否定できない人 上記以外	960	21.7	13	5.4	35	6.8	73	14.6	179	22.6	282	25.5	378	29.6	678	23.2	296	27.4	234	31.1
	総 数	3,095	69.8	228	94.2	476	92.2	412	82.2	549	69.3	693	62.8	737	57.6	1,958	67.0	635	58.7	433	57.5

注) ヘモグロビン A1c \geq 5.5%の場合

※ 「服薬者」とは、質問票で「インスリン注射または血糖を下げる薬」の使用有と回答した者

判定の基準

- ① 「糖尿病が強く疑われる人」とは、ヘモグロビン A1c の値が 6.1%以上、または、質問票で「現在糖尿病の治療を受けている」と答えた人である。
- ② 「糖尿病の可能性を否定できない人」とは、ヘモグロビン A1c の値が 5.6%以上、6.1%未満で、①以外の人である。

第52表の1 高血圧症有病者の状況（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	高血圧症有病者	1,073	56.1	11	10.9	35	19.3	79	40.3	171	55.0	325	63.2	452	74.1	747	58.4	349	66.2	280	79.5
	(再掲)高血圧症有病者のうち服薬者*	555	51.7	2	18.2	7	20.0	16	20.3	64	37.4	163	50.2	303	67.0	353	47.3	209	59.9	193	68.9
	正常高値血圧者	302	15.8	17	16.8	35	19.3	33	16.8	60	19.3	85	16.5	72	11.8	220	17.2	76	14.4	30	8.5
	上記以外	538	28.1	73	72.3	111	61.3	84	42.9	80	25.7	104	20.2	86	14.1	312	24.4	102	19.4	42	11.9
	総数	1,913	100	101	100	181	100	196	100	311	100	514	100	610	100	1,279	100	527	100	352	100
女 性	高血圧症有病者	1,160	42.4	3	2.0	22	6.4	54	16.6	210	40.5	331	50.6	540	72.5	798	44.4	386	62.2	337	75.4
	(再掲)高血圧症有病者のうち服薬者*	672	57.9	1	33.3	3	13.6	12	22.2	86	41.0	169	51.1	401	74.3	401	50.3	236	61.1	267	79.2
	正常高値血圧者	363	13.3	4	2.7	20	5.8	54	16.6	95	18.3	97	14.8	93	12.5	286	15.9	85	13.7	53	11.9
	上記以外	1,216	44.4	142	95.3	304	87.9	218	66.9	214	41.2	226	34.6	112	15.0	713	39.7	150	24.2	57	12.8
	総数	2,739	100	149	100	346	100	326	100	519	100	654	100	745	100	1,797	100	621	100	447	100
総 数	高血圧症有病者	2,233	48.0	14	5.6	57	10.8	133	25.5	381	45.9	656	56.2	992	73.2	1,545	50.2	735	64.0	617	77.2
	(再掲)高血圧症有病者のうち服薬者*	1,227	54.9	3	21.4	10	17.5	28	21.1	150	39.4	332	50.6	704	71.0	754	48.8	445	60.5	460	74.6
	正常高値血圧者	665	14.3	21	8.4	55	10.4	87	16.7	155	18.7	182	15.6	165	12.2	506	16.4	161	14.0	83	10.4
	上記以外	1,754	37.7	215	86.0	415	78.7	302	57.9	294	35.4	330	28.3	198	14.6	1,025	33.3	252	22.0	99	12.4
	総数	4,652	100	250	100	527	100	522	100	830	100	1,168	100	1,355	100	3,076	100	1,148	100	799	100

注) 判定の基準は p.16 参照

※ 「服薬者」とは、質問票で「血圧を下げる薬」の使用有と回答した者

第52表の2 高血圧症有病者の状況（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	高血圧症有病者	1,073	56.1	11	10.9	35	19.3	79	40.3	171	55.0	325	63.2	452	74.1	747	58.4	349	66.2	280	79.5
	(再掲)高血圧症有病者のうち服薬者*	555	51.7	2	18.2	7	20.0	16	20.3	64	37.4	163	50.2	303	67.0	353	47.3	209	59.9	193	68.9
	正常高値血圧者	302	15.8	17	16.8	35	19.3	33	16.8	60	19.3	85	16.5	72	11.8	220	17.2	76	14.4	30	8.5
	上記以外	538	28.1	73	72.3	111	61.3	84	42.9	80	25.7	104	20.2	86	14.1	312	24.4	102	19.4	42	11.9
	総数	1,913	100	101	100	181	100	196	100	311	100	514	100	610	100	1,279	100	527	100	352	100
女 性	高血圧症有病者	1,160	42.5	3	2.1	22	6.5	54	16.6	210	40.5	331	50.6	540	72.5	798	44.4	386	62.2	337	75.4
	(再掲)高血圧症有病者のうち服薬者*	672	57.9	1	33.3	3	13.6	12	22.2	86	41.0	169	51.1	401	74.3	401	50.3	236	61.1	267	79.2
	正常高値血圧者	363	13.3	4	2.8	20	5.9	54	16.6	95	18.3	97	14.8	93	12.5	286	15.9	85	13.7	53	11.9
	上記以外	1,205	44.2	138	95.2	297	87.6	218	66.9	214	41.2	226	34.6	112	15.0	713	39.7	150	24.2	57	12.8
	総数	2,728	100	145	100	339	100	326	100	519	100	654	100	745	100	1,797	100	621	100	447	100
総 数	高血圧症有病者	2,233	48.1	14	5.7	57	11.0	133	25.5	381	45.9	656	56.2	992	73.2	1,545	50.2	735	64.0	617	77.2
	(再掲)高血圧症有病者のうち服薬者*	1,227	54.9	3	21.4	10	17.5	28	21.1	150	39.4	332	50.6	704	71.0	754	48.8	445	60.5	460	74.6
	正常高値血圧者	665	14.3	21	8.5	55	10.6	87	16.7	155	18.7	182	15.6	165	12.2	506	16.4	161	14.0	83	10.4
	上記以外	1,743	37.6	211	85.8	408	78.5	302	57.9	294	35.4	330	28.3	198	14.6	1,025	33.3	252	22.0	99	12.4
	総数	4,641	100	246	100	520	100	522	100	830	100	1,168	100	1,355	100	3,076	100	1,148	100	799	100

注) 妊婦除外

※ 「服薬者」とは、質問票で「血圧を下げる薬」の使用有と回答した者

第53表 「脂質異常症が疑われる人」の状況（採血時間によらず、性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	脂質異常症が疑われる人	352	19.4	5	5.1	18	10.2	30	16.2	49	16.8	105	21.6	145	25.1	242	20.1	118	23.5	87	26.0
	(再掲)脂質異常症が疑われる人のうち服薬者*	197	56.0	0	0	4	22.2	12	40.0	29	59.2	69	65.7	83	57.2	145	59.9	72	61.0	48	55.2
	上記以外	1,462	80.6	94	94.9	159	89.8	155	83.8	242	83.2	380	78.4	432	74.9	962	79.9	384	76.5	247	74.0
	総数	1,814	100	99	100	177	100	185	100	291	100	485	100	577	100	1,204	100	502	100	334	100
女 性	脂質異常症が疑われる人	446	17.0	2	1.4	8	2.4	14	4.4	66	13.1	144	23.2	212	30.0	316	18.3	173	29.6	120	28.4
	(再掲)脂質異常症が疑われる人のうち服薬者*	391	87.7	0	0	2	25.0	8	57.1	59	89.4	135	93.8	187	88.2	285	90.2	156	90.2	104	86.7
	上記以外	2,183	83.0	142	98.6	329	97.6	303	95.6	436	86.9	478	76.8	495	70.0	1,409	81.7	411	70.4	303	71.6
	総数	2,629	100	144	100	337	100	317	100	502	100	622	100	707	100	1,725	100	584	100	423	100
総 数	脂質異常症が疑われる人	798	18.0	7	2.9	26	5.1	44	8.8	115	14.5	249	22.5	357	27.8	558	19.1	291	26.8	207	27.3
	(再掲)脂質異常症が疑われる人のうち服薬者*	588	73.7	0	0	6	23.1	20	45.5	88	76.5	204	81.9	270	75.6	430	77.1	228	78.4	152	73.4
	上記以外	3,645	82.0	236	97.1	488	94.9	458	91.2	678	85.5	858	77.5	927	72.2	2,371	80.9	795	73.2	550	72.7
	総数	4,443	100	243	100	514	100	502	100	793	100	1,107	100	1,284	100	2,929	100	1,086	100	757	100

* 「服薬者」とは、質問票で「コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬」の使用有と回答した者

「脂質異常症が疑われる人」の判定

国民健康・栄養調査の血液検査では、空腹時採血が困難であるため、脂質異常症の診断基準項目である中性脂肪による判定は行わず、下記の通りとした。

・「脂質異常症が疑われる人」

HDL コレステロールが 40mg/dl 未満、もしくはコレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬を服用している者。

第54表 「脂質異常症が疑われる人」の状況（採血時間によらず、動脈硬化性疾患予防ガイドライン 2007年版の基準、Friedewald の式、性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	脂質異常症が疑われる人	861	47.5	29	29.3	88	49.7	102	55.1	168	57.7	233	48.0	241	41.8	608	50.5	230	45.8	136	40.7
	(再掲)脂質異常症が疑われる人のうち服薬者*	106	12.3	0	0	2	2.3	8	7.8	23	13.7	35	15.0	38	15.8	80	13.2	31	13.5	24	17.6
	上記以外	953	52.5	70	70.7	89	50.3	83	44.9	123	42.3	252	52.0	336	58.2	596	49.5	272	54.2	198	59.3
	総数	1,814	100	99	100	177	100	185	100	291	100	485	100	577	100	1,204	100	502	100	334	100
女 性	脂質異常症が疑われる人	1,063	40.4	22	15.3	71	21.1	107	33.8	259	51.6	307	49.4	297	42.0	808	46.8	284	48.6	162	38.3
	(再掲)脂質異常症が疑われる人のうち服薬者*	155	14.6	0	0	1	1.4	4	3.7	24	9.3	56	18.2	70	23.6	118	14.6	62	21.8	36	22.2
	上記以外	1,566	59.6	122	84.7	266	78.9	210	66.2	243	48.4	315	50.6	410	58.0	917	53.2	300	51.4	261	61.7
	総数	2,629	100	144	100	337	100	317	100	502	100	622	100	707	100	1,725	100	584	100	423	100
総 数	脂質異常症が疑われる人	1,924	43.3	51	21.0	159	30.9	209	41.6	427	53.8	540	48.8	538	41.9	1,416	48.3	514	47.3	298	39.4
	(再掲)脂質異常症が疑われる人のうち服薬者*	261	13.6	0	0	3	1.9	12	5.7	47	11.0	91	16.9	108	20.1	198	14.0	93	18.1	60	20.1
	上記以外	2,519	56.7	192	79.0	355	69.1	293	58.4	366	46.2	567	51.2	746	58.1	1,513	51.7	572	52.7	459	60.6
	総数	4,443	100	243	100	514	100	502	100	793	100	1,107	100	1,284	100	2,929	100	1,086	100	757	100

* 「服薬者」とは、質問票で「コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬」の使用有と回答した者

「脂質異常症が疑われる人」の判定（定義）「動脈硬化性疾患予防ガイドライン（2007年版）」の基準

LDL コレステロール（Friedewald の式で算出）140mg/dl 以上、または、中性脂肪（トリグリセライド）150mg/dl 以上、または、HDL コレステロール 40mg/dl 未満

Friedewald の式 LDL コレステロール = 総コレステロール - HDL コレステロール - 中性脂肪 / 5
 （中性脂肪（トリグリセライド）が 400mg/dl 未満の場合。）

第55表 運動習慣の状況（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総数	5,057	100	323	100	592	100	584	100	905	100	1,218	100	1,435	100	1,196	100	853	100
	習慣無(健康上の理由)	356	7.0	4	1.2	9	1.5	15	2.6	41	4.5	79	6.5	208	14.5	100	8.4	157	18.4
	習慣無(その他)	3,190	63.1	257	79.6	499	84.3	447	76.3	625	69.1	653	53.6	709	49.4	573	47.9	432	50.6
	運動の習慣有	1,513	29.9	62	19.2	84	14.2	124	21.2	239	26.4	486	39.9	518	36.1	523	43.7	264	30.9
男性	総数	2,125	100	141	100	222	100	230	100	349	100	541	100	642	100	549	100	373	100
	習慣無(健康上の理由)	113	5.3	1	0.7	4	1.8	3	1.3	8	2.3	27	5.0	70	10.9	39	7.1	48	12.9
	習慣無(その他)	1,306	61.4	108	76.6	177	79.7	170	73.3	241	69.1	307	56.7	303	47.2	261	47.5	186	49.9
	運動の習慣有	706	33.3	32	22.7	41	18.5	57	25.4	100	28.7	207	38.3	269	41.9	249	45.4	139	37.3
女性	総数	2,932	100	182	100	370	100	354	100	556	100	677	100	793	100	647	100	480	100
	習慣無(健康上の理由)	243	8.3	3	1.6	5	1.4	12	3.4	33	5.9	52	7.7	138	17.4	61	9.4	109	22.7
	習慣無(その他)	1,884	64.3	149	81.9	322	87.0	277	78.2	384	69.1	346	51.1	406	51.2	312	48.2	246	51.3
	運動の習慣有	805	27.5	30	16.5	43	11.6	65	18.4	139	25.0	279	41.2	249	31.4	274	42.3	125	26.0

第56表 1週間の運動日数（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総数	5,057	100	323	100	592	100	586	100	905	100	1,218	100	1,435	100	1,196	100	853	100
	運動無し	3,546	70.1	261	80.8	508	85.8	462	78.8	666	73.6	732	60.1	917	63.9	673	56.3	589	69.1
	2日/週	336	6.6	12	3.7	31	5.2	41	7.0	76	8.4	96	7.9	80	5.6	84	7.0	38	4.5
	3日/週	250	4.9	16	5.0	23	3.9	21	3.6	41	4.5	76	6.2	73	5.1	79	6.6	40	4.7
	4日/週	153	3.0	7	2.2	9	1.5	12	2.0	20	2.2	55	4.5	50	3.5	55	4.6	25	2.9
	5日/週	195	3.9	14	4.3	11	1.9	13	2.2	32	3.5	67	5.5	58	4.0	62	5.2	30	3.5
	6日/週	66	1.3	5	1.5	3	0.5	5	0.9	10	1.1	21	1.7	22	1.5	22	1.8	11	1.3
	7日/週	511	10.1	8	2.5	7	1.2	30	5.1	60	6.6	171	14.0	235	16.4	221	18.5	120	14.1
男性	総数	2,125	100	141	100	222	100	232	100	349	100	541	100	642	100	549	100	373	100
	運動無し	1,419	66.7	109	77.3	181	81.5	173	74.6	249	71.3	334	61.7	373	58.1	300	54.6	234	62.7
	2日/週	149	7.0	7	5.0	17	7.7	20	8.6	40	11.5	31	5.7	34	5.3	29	5.3	17	4.6
	3日/週	107	5.0	7	5.0	13	5.9	9	3.9	14	4.0	28	5.2	36	5.6	35	6.4	18	4.8
	4日/週	56	2.6	5	3.5	1	0.5	3	1.3	6	1.7	19	3.5	22	3.4	23	4.2	12	3.2
	5日/週	95	4.5	7	5.0	3	1.4	7	3.0	14	4.0	29	5.4	35	5.5	30	5.5	18	4.8
	6日/週	27	1.3	2	1.4	2	0.9	2	0.9	1	0.3	8	1.5	12	1.9	11	2.0	7	1.9
	7日/週	272	12.8	4	2.8	5	2.3	16	6.9	25	7.2	92	17.0	130	20.2	121	22.0	67	18.0
女性	総数	2,932	100	182	100	370	100	354	100	556	100	677	100	793	100	647	100	480	100
	運動無し	2,127	72.5	152	83.5	327	88.4	289	81.6	417	75.0	398	58.8	544	68.6	373	57.7	355	74.0
	2日/週	187	6.4	5	2.7	14	3.8	21	5.9	36	6.5	65	9.6	46	5.8	55	8.5	21	4.4
	3日/週	143	4.9	9	4.9	10	2.7	12	3.4	27	4.9	48	7.1	37	4.7	44	6.8	22	4.6
	4日/週	97	3.3	2	1.1	8	2.2	9	2.5	14	2.5	36	5.3	28	3.5	32	4.9	13	2.7
	5日/週	100	3.4	7	3.8	8	2.2	6	1.7	18	3.2	38	5.6	23	2.9	32	4.9	12	2.5
	6日/週	39	1.3	3	1.6	1	0.3	3	0.8	9	1.6	13	1.9	10	1.3	11	1.7	4	0.8
	7日/週	239	8.2	4	2.2	2	0.5	14	4.0	35	6.3	79	11.7	105	13.2	100	15.5	53	11.0

第57表 運動を行う日の平均運動時間（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総数	5,057	100	323	100	592	100	584	100	905	100	1,218	100	1,435	100	1,196	100	853	100
	運動無し	3,546	70.1	261	80.8	508	85.8	462	78.8	666	73.6	732	60.1	917	63.9	673	56.3	589	69.1
	30-60分未満/日	598	11.8	20	6.2	29	4.9	45	7.7	94	10.4	179	14.7	231	16.1	210	17.6	120	14.1
	60-120分未満/日	584	11.5	28	8.7	36	6.1	49	8.4	101	11.2	199	16.3	171	11.9	196	16.4	93	10.9
	120-180分未満/日	228	4.5	9	2.8	12	2.0	19	3.2	34	3.8	68	5.6	86	6.0	78	6.5	37	4.3
	180分以上/日	101	2.0	5	1.5	7	1.2	9	1.5	10	1.1	40	3.3	30	2.1	39	3.3	14	1.6
男性	総数	2,125	100	141	100	222	100	230	100	349	100	541	100	642	100	549	100	373	100
	運動無し	1,419	66.7	109	77.3	181	81.5	173	74.6	249	71.3	334	61.7	373	58.1	300	54.6	234	62.7
	30-60分未満/日	255	12.0	8	5.7	15	6.8	15	6.5	36	10.3	74	13.7	107	16.7	98	17.9	53	14.2
	60-120分未満/日	278	13.1	16	11.3	17	7.7	23	9.9	44	12.6	84	15.5	94	14.6	95	17.3	51	13.7
	120-180分未満/日	123	5.8	6	4.3	5	2.3	14	6.0	17	4.9	34	6.3	47	7.3	38	6.9	24	6.4
	180分以上/日	50	2.4	2	1.4	4	1.8	5	2.2	3	0.9	15	2.8	21	3.3	18	3.3	11	2.9
女性	総数	2,932	100	182	100	370	100	354	100	556	100	677	100	793	100	647	100	480	100
	運動無し	2,127	72.5	152	83.5	327	88.4	289	81.6	417	75.0	398	58.8	544	68.6	373	57.7	355	74.0
	30-60分未満/日	343	11.7	12	6.6	14	3.8	30	8.5	58	10.4	105	15.5	124	15.6	112	17.3	67	14.0
	60-120分未満/日	306	10.4	12	6.6	19	5.1	26	7.3	57	10.3	115	17.0	77	9.7	101	15.6	42	8.8
	120-180分未満/日	105	3.6	3	1.6	7	1.9	5	1.4	17	3.1	34	5.0	39	4.9	40	6.2	13	2.7
	180分以上/日	51	1.7	3	1.6	3	0.8	4	1.1	7	1.3	25	3.7	9	1.1	21	3.2	3	0.6

第58表 運動強度の状況（性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総数	5,057	100	323	100	592	100	584	100	905	100	1,218	100	1,435	100	1,196	100	853	100
	運動無し	3,546	70.1	261	80.8	508	85.8	462	78.8	666	73.6	732	60.1	917	63.9	673	56.3	589	69.1
	高強度	126	2.5	18	5.6	22	3.7	13	2.2	29	3.2	33	2.7	11	0.8	22	1.8	3	0.4
	中強度	494	9.8	22	6.8	37	6.3	57	9.7	95	10.5	170	14.0	113	7.9	153	12.8	54	6.3
	低強度	891	17.6	22	6.8	25	4.2	52	8.9	115	12.7	283	23.2	394	27.5	348	29.1	207	24.3
男性	総数	2,125	100	141	100	222	100	230	100	349	100	541	100	642	100	549	100	373	100
	運動無し	1,419	66.7	109	77.3	181	81.5	173	74.6	249	71.3	334	61.7	373	58.1	300	54.6	234	62.7
	高強度	76	3.6	10	7.1	14	6.3	9	3.9	17	4.9	18	3.3	8	1.2	12	2.2	2	0.5
	中強度	223	10.5	12	8.5	15	6.8	24	10.3	37	10.6	70	12.9	65	10.1	77	14.0	36	9.7
	低強度	407	19.1	10	7.1	12	5.4	24	10.3	46	13.2	119	22.0	196	30.5	160	29.1	101	27.1
女性	総数	2,932	100	182	100	370	100	354	100	556	100	677	100	793	100	647	100	480	100
	運動無し	2,127	72.5	152	83.5	327	88.4	289	81.6	417	75.0	398	58.8	544	68.6	373	57.7	355	74.0
	高強度	50	1.7	8	4.4	8	2.2	4	1.1	12	2.2	15	2.2	3	0.4	10	1.5	1	0.2
	中強度	271	9.2	10	5.5	22	5.9	33	9.3	58	10.4	100	14.8	48	6.1	76	11.7	18	3.8
	低強度	484	16.5	12	6.6	13	3.5	28	7.9	69	12.4	164	24.2	198	25.0	188	29.1	106	22.1

第59表 運動習慣別栄養素等摂取量

		総 数		男 性		女 性	
		習慣無(その他)	運動習慣有	習慣無(その他)	運動習慣有	習慣無(その他)	運動習慣有
対象者数		3,095	1,475	1,265	685	1,830	790
エネルギー	kcal	1,919	1,937	2,185	2,126	1,735	1,772
たんぱく質	g	69.0	73.0	76.8	78.2	63.7	68.5
うち動物性	g	35.5	37.8	39.9	40.5	32.5	35.4
脂質	g	51.4	51.3	55.6	53.6	48.6	49.3
うち動物性	g	24.9	25.4	27.4	27.3	23.1	23.8
コレステロール	mg	305	304	332	320	286	289
炭水化物	g	276.0	276.5	310.4	300.0	252.3	256.0
食物繊維	g	15.4	17.5	15.9	17.6	15.0	17.4
水溶性	g	3.5	3.9	3.6	3.9	3.4	3.9
不溶性	g	11.3	12.8	11.7	13.0	11.1	12.8
ビタミンA	μgRE ^{*1}	599	680	661	652	557	704
ビタミンD	μg	8.2	9.3	9.0	9.9	7.6	8.8
ビタミンE	mg ^{*2}	9.1	10.7	8.9	9.0	9.2	12.1
ビタミンK	μg	245	288	256	291	237	286
ビタミンB ₁	mg	1.79	1.89	1.75	1.49	1.83	2.25
ビタミンB ₂	mg	1.49	1.63	1.51	1.57	1.48	1.69
ナイアシン	mgNE ^{*3}	15.4	16.1	17.5	17.4	14.0	15.0
ビタミンB ₆	mg	1.99	2.27	1.85	1.77	2.10	2.71
ビタミンB ₁₂	μg	7.0	8.0	8.1	8.4	6.2	7.6
葉酸	μg	320	364	341	365	305	363
パントテン酸	mg	5.39	5.78	5.89	6.06	5.04	5.55
ビタミンC	mg	129	164	115	137	138	187
食塩(ナトリウム×2.54/1,000)	g	11.0	11.4	12.2	12.1	10.2	10.7
カリウム	mg	2,359	2,666	2,479	2,718	2,275	2,622
カルシウム	mg	507	585	519	581	499	588
マグネシウム	mg	253	282	273	293	239	272
リン	mg	982	1,058	1,076	1,117	916	1,007
鉄	mg	8.1	9.0	8.6	9.1	7.7	9.0
亜鉛	mg	8.1	8.4	9.1	9.0	7.4	7.8
銅	mg	1.21	1.29	1.33	1.37	1.12	1.22
脂肪エネルギー比率 ^{*4}	%	23.9	23.6	22.6	22.4	24.9	24.7
炭水化物エネルギー比率 ^{*4}	%	61.5	61.2	63.2	62.8	60.4	59.8
動物性たんぱく質比率 ^{*4}	%	49.4	49.7	49.9	49.7	49.1	49.8

注) 20歳以上 妊婦・「健康上の理由」により運動習慣の無い者除外

*¹ RE: レチノール当量 *² α-トコフェロール量。α-トコフェロール以外のビタミンEは含んでいない *³ NE: ナイアシン当量*⁴ これらの比率は個々の計算値を平均したものである

第 60 表の 1 歩行数の平均値及び標準偏差（性・年齢階級別）

	総 数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 20歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
	平均 値	標準 偏差	平均 値	標準 偏差	平均 値	標準 偏差	平均 値	標準 偏差	平均 値	標準 偏差	平均 値	標準 偏差	平均 値	標準 偏差	平均 値	標準 偏差	平均 値	標準 偏差	平均 値	標準 偏差	平均 値	標準 偏差
総 数	6,491	4,160	7,981	5,070	7,495	4,242	7,057	4,063	7,345	3,935	7,052	3,792	6,711	4,129	4,370	3,662	6,426	4,104	6,039	3,768	3,566	3,502
男 性	7,074	4,560	8,345	5,669	8,370	4,805	7,728	4,678	7,824	4,396	7,317	4,039	7,232	4,471	5,102	4,073	7,011	4,490	6,439	3,879	4,525	4,294
女 性	6,006	3,726	7,593	4,328	6,720	3,500	6,525	3,411	6,931	3,439	6,843	3,575	6,265	3,757	3,790	3,185	5,945	3,688	5,687	3,633	2,854	2,553

第 60 表の 2 歩行数の分布（性・年齢階級別）

	総 数	15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 20歳以上		(再掲) 65-74歳		(再掲) 75歳以上			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%		
総数	総 数	7,459	100	310	100	645	100	957	100	992	100	1,320	100	1,547	100	1,688	100	7,149	100	1,425	100	1,019	100
	2000歩未満	834	11.2	15	4.8	45	7.0	46	4.8	47	4.7	67	5.1	134	8.7	480	28.4	819	11.5	168	11.8	382	37.5
	2000-4000	1,398	18.7	46	14.8	82	12.7	180	18.8	136	13.7	197	14.9	308	19.9	449	26.6	1,352	18.9	317	22.2	292	28.7
	4000-6000	1,572	21.1	55	17.7	122	18.9	214	22.4	220	22.2	337	25.5	322	20.8	302	17.9	1,517	21.2	306	21.5	157	15.4
	6000-8000	1,375	18.4	66	21.3	147	22.8	189	19.7	213	21.5	264	20.0	279	18.0	217	12.9	1,309	18.3	262	18.4	95	9.3
	8000-10000	1,010	13.5	58	18.7	106	16.4	141	14.7	163	16.4	205	15.5	217	14.0	120	7.1	952	13.3	176	12.4	48	4.7
	10000歩以上	1,270	17.0	70	22.6	143	22.2	187	19.5	213	21.5	250	18.9	287	18.6	120	7.1	1,200	16.8	196	13.8	45	4.4
男性	総 数	3,387	100	160	100	303	100	423	100	460	100	581	100	714	100	746	100	3,227	100	666	100	434	100
	2000歩未満	336	9.9	11	6.9	23	7.6	25	5.9	25	5.4	36	6.2	58	8.1	158	21.2	325	10.1	68	10.2	121	27.9
	2000-4000	560	16.5	16	10.0	26	8.6	74	17.5	57	12.4	84	14.5	118	16.5	185	24.8	544	16.9	128	19.2	115	26.5
	4000-6000	668	19.7	32	20.0	50	16.5	72	17.0	92	20.0	136	23.4	134	18.8	152	20.4	636	19.7	137	20.6	85	19.6
	6000-8000	609	18.0	30	18.8	65	21.5	78	18.4	92	20.0	95	16.4	138	19.3	111	14.9	579	17.9	134	20.1	52	12.0
	8000-10000	476	14.1	29	18.1	50	16.5	63	14.9	60	13.0	100	17.2	108	15.1	66	8.8	447	13.9	92	13.8	26	6.0
	10000歩以上	738	21.8	42	26.3	89	29.4	111	26.2	134	29.1	130	22.4	158	22.1	74	9.9	696	21.6	107	16.1	35	8.1
女性	総 数	4,072	100	150	100	342	100	534	100	532	100	739	100	833	100	942	100	3,922	100	759	100	585	100
	2000歩未満	498	12.2	4	2.7	22	6.4	21	3.9	22	4.1	31	4.2	76	9.1	322	34.2	494	12.6	100	13.2	261	44.6
	2000-4000	838	20.6	30	20.0	56	16.4	106	19.9	79	14.8	113	15.3	190	22.8	264	28.0	808	20.6	189	24.9	177	30.3
	4000-6000	904	22.2	23	15.3	72	21.1	142	26.6	128	24.1	201	27.2	188	22.6	150	15.9	881	22.5	169	22.3	72	12.3
	6000-8000	766	18.8	36	24.0	82	24.0	111	20.8	121	22.7	169	22.9	141	16.9	106	11.3	730	18.6	128	16.9	43	7.4
	8000-10000	534	13.1	29	19.3	56	16.4	78	14.6	103	19.4	105	14.2	109	13.1	54	5.7	505	12.9	84	11.1	22	3.8
	10000歩以上	532	13.1	28	18.7	54	15.8	76	14.2	79	14.8	120	16.2	129	15.5	46	4.9	504	12.9	89	11.7	10	1.7

第 61 表 (再掲) 運動習慣の有無別の歩行数の平均値及び標準偏差（性・年齢階級別）

		総 数			20-29歳			30-39歳			40-49歳			50-59歳			60-69歳			70歳以上		
		人数	平均 値	標準 偏差	人数	平均 値	標準 偏差	人数	平均 値	標準 偏差	人数	平均 値	標準 偏差	人数	平均 値	標準 偏差	人数	平均 値	標準 偏差	人数	平均 値	標準 偏差
総数	運動習慣無	3,320	5,679	3,705	241	6,586	3,926	474	6,599	3,540	437	6,988	3,587	636	6,508	3,555	697	5,672	3,539	835	3,583	3,123
	運動習慣有	1,447	7,731	4,292	56	8,027	3,613	80	8,691	5,017	118	8,719	4,256	224	8,382	3,872	479	8,307	4,531	490	6,443	3,888
	総 数	4,767	6,302	4,005	297	6,858	3,904	554	6,901	3,855	555	7,356	3,802	860	6,996	3,730	1,176	6,745	4,177	1,325	4,641	3,692
男性	運動習慣無	1,319	6,049	4,084	97	7,269	4,523	162	6,997	3,887	162	7,531	4,014	235	6,621	3,867	321	6,192	3,804	342	4,023	3,724
	運動習慣有	668	8,159	4,697	26	8,153	3,797	38	9,477	6,390	55	8,736	4,891	90	8,764	4,012	204	8,930	5,173	255	7,007	4,024
	総 数	1,987	6,758	4,413	123	7,456	4,380	200	7,468	4,557	217	7,837	4,274	325	7,215	4,018	525	7,256	4,581	597	5,298	4,125
女性	運動習慣無	2,001	5,435	3,412	144	6,126	3,407	312	6,392	3,334	275	6,668	3,275	401	6,442	3,362	376	5,227	3,235	493	3,278	2,587
	運動習慣有	779	7,365	3,876	30	7,918	3,509	42	7,980	3,251	63	8,704	3,653	134	8,125	3,768	275	7,844	3,936	235	5,831	3,645
	総 数	2,780	5,975	3,652	174	6,435	3,481	354	6,580	3,359	338	7,047	3,436	535	6,863	3,540	651	6,333	3,774	728	4,102	3,199

注) 20歳以上 妊婦・「健康上の理由」により運動習慣の無い者除外