

<p>A single autophosphorylation site on KDR/Flk-1 is essential for VEGF-A-dependent activation of PLC-γ and DNA synthesis in vascular endothelial cells. Takahashi, T., Yamaguchi, S., Chida, K., and <u>Shibuya, M.</u> EMBO J., 20 (11), 2768-2778, 2001.</p>
<p>Amplification of the structurally and functionally altered epidermal growth factor receptor gene (<u>c-erbB</u>) in human brain tumors. Yamazaki, H., Fukui, Y., Ueyama, Y., Tamaoki, N., Kawamoto, T., Taniguchi, S., and <u>Shibuya, M.</u> Mol. Cell. Biol. 8 (4), 1816-1820, 1988.</p>

下遠野 邦忠	慶應義塾大学 医学部 特別研究教授
専門分野 分子生物学、分子ウイルス学	
<p>Miyanari Y, Atsuzawa K, Usuda N, Watashi K, Hishiki T, Zayas M, Bartenschlager R, Wakita T, Hijikata M, Shimotohno K. The lipid droplet is an important organelle for hepatitis C virus production. Nat Cell Biol. 9(9):1089-1097, 2007</p>	
<p>Watashi K, Ishii N, Hijikata M, Inoue D, Murata T, Miyanari Y, Shimotohno K. Cyclophilin B is a functional regulator of hepatitis C virus RNA polymerase. Mol Cell. 19:111-122, 2005.</p>	
<p>「C型肝炎ウイルスの増殖戦略」蛋白質核酸酵素 53巻 666-672 (2008)</p>	

委員：田島和雄	所属・役職：愛知県がんセンター研究所・所長
専門分野：がんの疫学・予防	
<p>Decreased risk of colorectal cancer with the high Natural-Killer (NK)-cell activity NKG2D genotype in Japanese. Furue H, Matsuo K, Kumimoto H, Hiraki A, Suzuki T, Yatabe Y, Komori K, Kanemitsu Y, Hirai T, Kato T, Ueda M, Ishizaki K, <u>Tajima K.</u> Carcinogenesis 29: 316-20, 2008. (他 400 編)</p>	