

「急性血液浄化法」秋澤忠男他・編著、総合医学社、2006
Plasma exchange in hepatic failure. Abe T, Akizawa T, et al. International Society for Apheresis 4 th International Congress, 2003

岡崎 勲	国際医療福祉大学臨床医学研究センター・教授
肝線維化機序、肝硬変、肝癌	
Bone marrow-derived cells express matrix metalloproteinases and contribute to regression of liver fibrosis in mice. Higashiyama R, Inagaki Y, Watanabe T, Okano H, Matsuzaki Y, Shiota G, <u>Okazaki I</u> . Hepatology 2007 Jan; 45(1): 213-222.	
Difference in gene expression for matrix metalloproteinase-1 between early and advanced hepatocellular carcinoma. <u>Okazaki I</u> , Wada N, Nakano M, Saito A, Takasaki K, Doi M, Kameyama K, Otani Y, Kubochi K, Niioka M, Watanabe T, Maruyama K. Hepatology 1997 Mar; 25(3): 580-584.	
“Extracellular Matrix and the Liver -approach to gene therapy” <u>Okazaki I</u> , Ninomiya Y, Friedman SL, Tanikawa K (editors), Academic Press, Amsterdam, Boston, London, New York, 1-467 (2003).	

齋藤英彦	名古屋セントラル病院・院長
血液病学	
Life-threatening human parvovirus B19 infection transmitted by intravenous immune globulin. Hayakawa F, Imada K, Towatari M, <u>Saito H</u> Br J Haematol 118: 1187-1189, 2002.	

Safety and complication of interventional radiology for hepatocellular carcinoma in patients with haemophilia and cirrhosis. Toyoda, H., Fukuda, Y., Yokozaki, S, Hayashi, K., Saito, H., Takamatsu, J. **Brit J Haematol** 112: 1071-1073, 2001

TT virus genotype change frequently in multiply transfused patients with hemophilia but rarely in patients with chronic hepatitis C and healthy subjects. Toyoda, H., Fukuda, Y., Nakano, I., Katano, Y., Yokozaki, S., Hayashi, K., Ito, Y., Suzuki, K., Nakano, H., Saito, H., Takamatsu, J **Transfusion** 41: 1130-1135, 2001

(官職指定)

医政局研究開発振興課長

健康局疾病対策課長

医薬食品局血液対策課長

老健局老人保健課長

国立感染症研究所企画調整主幹

大臣官房厚生科学課長

免疫アレルギー疾患等予防・治療研究事業中間・事後評価委員名簿

(○：委員長)

狩野 庄吾	自治医科大学・名誉教授
内科学（アレルギー・リウマチ学）	
Yoshio, T., Masuyama, J-I., Minota, S., Iwamoto, M., Mimori, A., Takeda, A., Okazaki, H., and Kano, S.: Correlation of serum IgG antibodies to recombinant P0 fusion protein with IgG antibodies to carboxyl-terminal 22 synthetic peptides and carboxyl-terminal 22 amino acid-depleted recombinant P0 fusion protein in patients with systemic lupus erythematosus. <i>Arthritis Rheum.</i> 40: 1364-1365, 1997.	
Masuyama, J., Kaga, S., Kano, S., Minota, S.: A novel costimulation pathway via the 4C8 antigen for the induction of CD4+ regulatory T cells. <i>J Immunol</i> 169:3710-3716, 2002.	

Okazaki, H., Hirata, D., Kamimura, T., Sato, H., Iwamoto, M., Yoshio, T., Masuyama, J., Fujimura, A., Kobayashi, E., Kano, S., Minota, S.: Effects of FTY720 in MRL-lpr/lpr Mice: Therapeutic Potential in Systemic Lupus Erythematosus. *J Rheumatol* 29: 707-716, 2002.

上野川 修一	日本大学生物資源科学部教授
食品免疫学、食品アレルギー学、腸管免疫・細菌学	
H. Nakajima-Adachi, A. Ebihara, A. Kikuchi, T. Ishida, K. Sasaki, K. Hirano, H. Watanabe, K. Asai, Y. Takahashi, Y. Kanamori, N. Shimojo, H. Matsuda, Y. Kohno, S. Hachimura and S. <u>Kaminogawa</u> , Food antigen causes T(H)2-dependent enteropathy followed by tissue repair in T-cell receptor transgenic mice. <i>J. Allergy Clin. Immunol.</i> 117, 1125-1132 (2006)	
K. Takahashi, N. Hayashi, S. <u>Kaminogawa</u> and C. Ra, Molecular mechanisms for transcriptional regulation of human high-affinity IgE receptor beta-chain gene induced by GM-CSF. <i>J. Immunol.</i> 177, 4605-4611 (2006)	
M. Kuraoka, M. Hashiguchi, S. Hachimura and S. <u>Kaminogawa</u> , CD4 ^c -kit ^c CD3 ^c IL-2R ^c Peyer's patch cells are a novel cell subset which secrete IL-5 in response to IL-2: implications for their role in IgA production. <i>Euro. J. Immunol.</i> 34, 1920-1929 (2004)	

近藤 啓文	北里大学北里研究所メディカルセンター病院・病院長
内科学（リウマチ・膠原病）	
Rapid induction of peroxisome proliferator-activated receptor gamma expression in human monocytes by monosodium urate monohydrate crystals. Akahoshi T, Namai R, Murakami Y, Watanabe M, Matsui T, Nishimura A, Kitasato H, Kameya T, Kondo H. <i>Arthritis Rheum</i> , 48 : 231-239, 2003.	

<p>Efficacy and safety of Tacrolimus (FK506) in treatment of rheumatoid arthritis: A randomized, double blind, placebo controlled dose-finding study. Kondo H, Abe T, Hashimoto H, Uchida S, Irimajiri S, Hara M, Sugawara S: <i>J Rheumatol</i> 31:343-351, 2004.</p>
<p>全身性硬化症（強皮症）(SSc). 新臨床内科学 9 版（高久文磨他監修）. 近藤啓文. 医学書院 1446-1450. 2009</p>

<p>笹月 健彦</p>	<p>国立国際医療センター・名誉総長</p>
<p>免疫遺伝学</p>	
<p><u>Sasazuki T</u>, Juji T, Morishima Y, Kinukawa N, Kashiwabara H, Inoko H, Yoshida T, Kimura A, Akaza T, Kamikawaji N, Kodera Y, Takaku F. Effect of matching of class I HLA alleles on clinical outcome after transplantation of hematopoietic stem cells from an unrelated donor. New Engl. J. Med., 339:1177-1185, 1998</p>	
<p>Fukui Y, Hashimoto O, Sanui T, Oono T, Koga H, Abe M, Inayoshi A, Noda M, Oike M, Shirai T, <u>Sasazuki T</u>. Hematopoietic cell-specific CDM family protein DOCK2 is essential for lymphocyte migration. Nature, 412: 826-831, 2001</p>	
<p>Kochi Y, Yamada R, Suzuki A, Harley JB, Shirasawa S, Sawada T, Bae S-C, Tokuhira S, Chang X, Sekine A, Takahashi A, Tsunoda T, Ohnishi Y, Kaufman KM, Kang CP, Kang C, Otsubo S, Yumura W, Mimori A, Koike T, Nakamura Y, <u>Sasazuki T</u>, Yamamoto K. A functional variant in <i>FcRH3</i>, encoding Fc Receptor Homolog 3, is associated with rheumatoid arthritis and several autoimmunities. Nature Genet., 37:478-485, 2005</p>	

<p>戸山 芳昭</p>	<p>慶應義塾大学医学部整形外科学教室・教授</p>
<p>整形外科学、リウマチ学、脊椎・脊髄外科学</p>	
<p>Cell surface colony-stimulating factor 1 can be cleaved by TACE-α converting enzyme or endocytosed in a clathrin-dependent manner : Keisuke Horiuchi, Takeshi Miyamoto, Hironari Takaishi, Akihiro Hakozaiki, Naoto Kosaki, Yoshiteru Miyauchi,</p>	