

医薬品
 医薬部外品 研究報告 調査報告書
 化粧品

識別番号・報告回数		回	報告日 年 月 日	第一報入手日 2006年1月30日	新医薬品等の区分 該当なし	厚生労働省処理欄
一般的名称			研究報告の公表状況	A modified caprylic acid method for manufacturing immunoglobulin G from human plasma with high yield and efficient virus clearance J. Parkkinen, A. Rahola, L. von Bonsdorff, H. Tolo & E. Torma Vox Sanguinis 2006; 90; 97- 104	公表国	
販売名 (企業名)					米国	
研究報告の概要	静注用免疫グロブリン製剤の開発において高収率で IgG が得られる血漿分画法が求められている。本稿では、不活化工程に対して物理化学的に耐性を示すウイルスを効率的に除去しつつ高い収率で IgG を得られる新規の製造法が提示されている。これまでに、分画 II+III から開始してカプリル酸塩沈降法とクロマトグラフィーを組み合わせた精製工程が発表されているが、カプリル酸沈降法は有効なウイルス不活化工程と精製工程を併せ持つ工程として提示されている。本研究では、人血漿から高収率で IgG を精製するための変法カプリル酸法について検討した。濾過条件を最適化し、重合化蛋白を除去することで生成物はウイルス濾過フィルターを容易に通ることができる。この方法により、バルボウイルス B19 などのエンベロプをもたない小さなウイルスを非常に高い確率で除去することが可能になる。本方法では、分画 II+III にカプリル酸塩を添加し、その後 PEG 処理を行った。さらに陰イオン交換クロマトグラフィーを行うことで純度の高い IgG が高収率で得られた。カプリル酸処理によるウイルス除去能は 10^4 であり、全工程におけるウイルス除去能は $10^{13.5}$ と非常に高かった。この製造法が実用化されれば、分画 II+III から安定かつ重合体を含まない IgG が高収率で得られる (すなわちウイルス除去フィルターにより濾過ができる) ことが示された。					使用上の注意記載状況・ その他参考事項等
	報告企業の意見		今後の対応			BYL-2005-0204
弊社の静注用免疫グロブリン製剤の製造工程においてカプリル酸および PEG 処理は行われていないが、本剤のウイルス不活化工程 (分画、透析・限外濾過、S/D 処理、低 pH インキュベーション) によるバルボウイルス B19 のモデルウイルスの除去率は $10^{9.2}$ 以上と非常に高い。しかしながら現時点でもなお感染の可能性を完全には否定できない。本稿において示されたバルボウイルス B19 の不活化工程の実用化については引き続き注視して情報を収集する。			現時点で新たな安全対策上の措置を講じる必要は無いと考える。引き続きバルボウイルス B19 の除去および検出技術に関する関連情報の収集に努める。			

ORIGINAL PAPER

A modified caprylic acid method for manufacturing immunoglobulin G from human plasma with high yield and efficient virus clearance

J. Parkkinen, A. Rahola, L. von Bonsdorff, H. Tölö & E. Törmä

Finnish Red Cross Blood Service, Helsinki, Finland

Vox Sanguinis

Background and Objectives The increasing demand for intravenous immunoglobulin (IVIg) necessitates the development of improved plasma fractionation methods, providing higher immunoglobulin G (IgG) recovery. Here, we describe a new IVIg production process resulting in a high yield of IgG and effective reduction of physico-chemically resistant viruses.

Materials and Methods IgG was purified from Cohn fraction II+III by caprylic acid treatment, polyethylene glycol precipitation, anion-exchange chromatography, nanofiltration and ultrafiltration. Stability of the purified IgG was studied in different formulations. Virus reduction was studied with two viruses: bovine viral diarrhoea virus, assessed by an infectivity assay; and human parvovirus B19, assessed by polymerase chain reaction.

Results The combination of caprylic acid treatment with polyethylene glycol precipitation and a single anion-exchange chromatography yielded polymer-free, pure IgG. The purified IgG could be filtered through a small pore-size virus filter (Millipore V-NFP) with high throughput and excellent yield. The formulated product was stable as a 100 g/l IgG solution. Bovine viral diarrhoea virus was effectively inactivated by the caprylic acid treatment, and parvovirus B19 was effectively removed in the polyethylene glycol precipitation and nanofiltration stages, the total reduction of parvovirus being $\approx 14 \log_{10}$.

Conclusions The new process gives pure and stable IgG solution with an average yield of 4.8 g of IgG per kg of recovered plasma and has a very high capacity to remove even physico-chemically resistant viruses.

Key words: caprylic acid, IgG, intravenous immunoglobulin, nanofiltration, parvovirus, polyethylene glycol.

Received: 23 August 2005,
revised 28 September 2005,
accepted 21 October 2005,
published online 20 December 2005

Introduction

Intravenous immunoglobulins (IVIgs) are concentrated formulations of human immunoglobulin G (IgG) that are prepared by the industrial fractionation of large pools of individual plasma donations. IVIg is widely used for the treatment

of patients with primary and secondary immune deficiencies, as well as for the treatment of various immune-mediated disorders, such as thrombocytopenic purpura and polyneuropathies [1–3].

IgG has traditionally been separated in large scale from human plasma by the cold ethanol fractionation method developed in the 1940s [4,5] and its subsequent modifications. The early IgG preparations could only be administered intramuscularly or subcutaneously because of adverse effects associated with their intravenous infusion [6]. These adverse effects were mainly caused by immunoglobulin aggregates inducing complement activation and occasionally by

Correspondence: Jaakko Parkkinen, Finnish Red Cross Blood Service, Kivihaantie 7, FI-00310 Helsinki, Finland
E-mail: jaakko.parkkinen@bts.redcross.fi

plasma-derived impurities, such as proteases. Therefore, other manufacturing steps were added for the further purification of IgG and removal of aggregates [7]. Until the 1980s, IVIG preparations were thought not to transmit viral infections. However, reports of the transmission of hepatitis C virus (HCV) by a variety of IVIG preparations, which were not subjected to dedicated virus inactivation, led to serious concern about the safety of IVIG with respect to virus transmission [8–10]. This necessitated the addition of specific virus-inactivation steps to the manufacturing process.

The addition of multiple steps to the manufacturing process of IVIG lowers the yield of IgG and raises the manufacturing costs. At the same time, an increasing demand for IVIG has made the yield even more important. Therefore, the emphasis has lately been to develop completely new IVIG manufacturing processes. Recently, Lebing *et al.* [11] described a novel process for IVIG manufacture, which starts from Cohn fraction II+III paste and utilizes caprylic acid treatment and chromatography for purification of IgG. Caprylic acid precipitation serves both as an effective virus inactivation and purification step. This approach resulted in a simplified process with a much improved yield of IgG.

The introduction of sensitive screening assays for viral markers in donated blood and plasma, and the implementation of effective virus-inactivation methods, has greatly improved the safety of current IVIG products. However, a risk of viral transmission may still exist with physico-chemically resistant agents, which are not effectively inactivated by current chemical virus-inactivation methods [3]. Parvovirus B19 is an example of a physico-chemically resistant virus transmitted by plasma products [12]. Parvovirus B19 antibodies present in IVIG are useful in the treatment of severe complications of parvovirus infection [13]. However, the virus itself was detectable, by polymerase chain reaction (PCR), in IVIG preparations and could theoretically pose a threat of infection to recipients [12]. A case of parvovirus B19 infection, transmitted by a heat-treated IVIG preparation, that led to pure red blood cell (RBC) aplasia has recently been reported [14], as well as a possible superinfection with a new strain of parvovirus B19 in an IVIG recipient already infected with B19 [15].

Considering additional reduction steps for physico-chemically resistant viruses, nanofiltration is efficient at removing non-enveloped viruses from solutions of biologically active proteins [16]. However, efficacious nanofiltration of IVIG preparations with filters, which would remove even small viruses, such as parvovirus, has been difficult because of a tendency for the filters to clog. This reduces the filtration capacity, decreases the yield of IgG and increases the filtration costs. In the present study we describe a modified caprylic acid process for the high-yield purification of IgG from human plasma. Owing to the optimization of filtration conditions and lack of polymeric proteins, the product can be efficiently filtered through a small pore-size virus filter.

The described process has a very high capacity to remove non-enveloped viruses.

Materials and methods

Purification of IgG from Cohn fraction II+III

Fraction II+III paste was fractionated by the Cohn method from human plasma (Krijnen's modification). The filter aid free fraction II+III was collected by centrifugation. All experiments were carried out on a laboratory scale using up to 0.5 kg of fraction II+III paste per batch. A flow scheme of the developed process is shown in Fig. 1. The paste was suspended in eight volumes of purified water below 5 °C and the pH was adjusted to pH 4.8 with 0.2 M acetic acid. The solution was brought to room temperature (\approx 22 °C) and caprylic acid was added to a concentration of 50 mM over a 1-h time-period. The suspension was mixed for 2 h and the precipitate was removed by centrifugation. During the development phase, polyethylene glycol (PEG) and caprylic acid were compared as precipitating agents, and different caprylic acid concentrations were tested (10–60 mM). The final conditions were chosen based on IgG recovery and virus-inactivation efficacy of the caprylic acid treatment. The pH of the solution was raised to pH 5.4 with 0.2 M NaOH, PEG 4000 was added and the solution was mixed for 2 h; 2% of diatomaceous earth was then added and the mixture was filtered. Different concentrations of PEG were compared and 3% was chosen for the final process based on clearance of polymers and parvovirus and IgG recovery. The solution conductivity was adjusted to 2.0 mS/cm using sodium acetate buffer. The filtrate was applied to a column (5.0 cm \times 15 cm) of ANX Sepharose FF gel (GE Healthcare, Uppsala, Sweden) equilibrated with 20 mM sodium acetate buffer, pH 5.4. The flow rate was 70 cm/h. The flow-through fraction containing IgG was recovered, and the pH of the solution was adjusted to pH 4.4 with 0.5 M acetic acid. After filtration through a 0.1- μ m prefilter (AcroCap; Pall Life Sciences, Ann Arbor, MI), the solution was filtered through a Millipore V-NFP virus filter (Millipore Corp., Mosheim, France) at 35 °C with a pressure of 3.5 bar. The protein concentration was \approx 8 g/l, and a load of \approx 11 kg of IgG/m² of filter area was used. The filtrate was concentrated by ultrafiltration, diafiltered with water for injection to remove PEG, and finally concentrated. The concentrated solution was formulated to 100 g/l IgG, 0.2 M glycine was added and the pH was adjusted to either 4.4 or 5.3. Alternatively, trehalose was tested as a stabiliser. The formulated solution was sterile filtered and transferred aseptically into containers.

Analytical methods

IgG was determined by immunoturbidimetry using a kit from ThermoClinical Labsystems (Helsinki, Finland). Immunoglobulin




Fig. 1 Flow scheme of the new manufacturing process for intravenous immunoglobulin (IVIg).

A (IgA) was determined by an enzyme immunoassay, as described previously [17], and IgG subclasses were determined using PeliClass enzyme-linked immunosorbent assay (ELISA) kits (Sanquin Reagents, Amsterdam, the Netherlands). Purity was established by zone electrophoresis on agarose, and molecular size distribution was determined by size-exclusion liquid chromatography according to Ph. Eur [18]. Albumin and immunoglobulin M (IgM) were quantified by radial immunodiffusion, using LC Partigen immunoplates (Dade Behring, Marburg, Germany). Prekallikrein activator (PKA) was determined using purified prekallikrein and S-2302 (Chromogenix

AB, Mölndal, Sweden) as the chromogenic substrate, according to Ph. Eur [18]. Direct kallikrein (KAL) activity was measured as hydrolysis of S-2302 without the addition of prekallikrein. Anticomplementary activity (ACA) was determined as consumption of complement and measured by haemolysis of red cells, according to Ph. Eur [18]. Caprylic acid was determined by gas chromatography [18], and PEG as described by Skoog [19].

Virus-reduction studies

Inactivation of bovine viral diarrhoea virus (BVDV, strain NADL; ATCC VR-534, Manassas, VA) was studied at Sanquin Viral Safety Services (Amsterdam, the Netherlands). The virus was propagated and assayed as described by Terpstra *et al.* [20]. The virus inoculum contained $10^{7.7}$ tissue culture infective dose 50% (TCID₅₀)/ml and the spiked starting material contained $10^{6.4}$ TCID₅₀/ml. Parvovirus reduction was studied by spiking the starting solution with high-titre parvovirus B19-positive plasma containing 3.6×10^{12} genome equivalents (geq)/ml of parvovirus DNA (a generous gift from Dr Hideki Abe, Hokkaido Red Cross BTS, Sapporo, Japan). The spiked starting materials contained 10^8 – 10^{10} geq/ml in different experiments. To remove free viral DNA, samples were treated with DNase (RQ1 RNase-Free DNase; Promega, Madison, WI). Nucleic acids were isolated from the starting solution and processed samples were diluted in parvovirus-negative plasma using the MagNA Pure method (Roche, Mannheim, Germany). Parvovirus B19 DNA was determined by real-time PCR using the LightCycler and the Parvovirus B19 Quantification Kit (Roche Diagnostics, Basel, Switzerland).

Results

We compared PEG and caprylic acid precipitation as a first step in the preparation of polymer-free immunoglobulin solution from suspended Cohn fraction II+III. A relatively high concentration of PEG ($\approx 6\%$) was needed for clarification of the crude immunoglobulin solution, which compromised immunoglobulin yield. Using caprylic acid precipitation combined with anion exchange chromatography, pure IgG could be obtained with a good yield. However, some polymeric IgG remained in the product, and nanofiltration downstream in the process was not efficacious owing to a tendency of the filter to clog. We therefore tested different ways to improve the removal of polymeric proteins in the caprylic acid process.

When the supernatant solution, after caprylic acid treatment, was subjected to precipitation with a low concentration of PEG, effective removal of polymers was achieved, while monomeric IgG was recovered with a good yield in the supernatant solution (Fig. 2). Furthermore, the immunoglobulin solution could be effectively filtered through a small pore-size virus filter. Optimal PEG concentration was found to be $\approx 3\%$, which enabled high flux and filtration capacity in nanofiltration




Fig. 2 Influence of polyethylene glycol (PEG) concentration on the removal of immunoglobulin G (IgG) polymers and parvovirus B19 in the PEG precipitation step. B19 removal was measured by polymerase chain reaction (PCR) in spiked process samples before the addition of PEG 4000 and after removal of PEG precipitate by filter aid filtration. Polymers were determined after the anion exchange chromatography step.

(Fig. 3a). The use of higher PEG concentrations decreased the yield of IgG.

Caprylic acid was added as free acid for the inactivation of enveloped viruses and the precipitation of contaminating proteins and lipids. Addition of caprylic acid to a concentration of 50 mM in 1 h resulted in the complete inactivation of BVDV (Fig. 4), which was previously identified as a resistant model virus in caprylic acid inactivation [11]. Immunoglobulin was recovered in the supernatant solution, which was treated with PEG and clarified by filtration in the presence of filter aids.

A combination of PEG precipitation with caprylic acid treatment was found to be beneficial, not only for the removal of polymers but also for the removal of non-enveloped viruses.

When the human parvovirus B19 was used as a model virus, almost 4 log₁₀ of the virus was removed, even with 2% PEG, and with 3% PEG the reduction was ≈ 4.7 log₁₀ (Fig. 2). The effective clearance of parvovirus apparently was a combined effect of residual caprylate remaining after the caprylic acid treatment and the relatively low PEG concentrations, as without caprylate a higher PEG concentration was needed for effective virus clearance (data not shown).

The final purification of IgG was achieved in a single anion-exchange chromatography column using the ANX Sepharose FF gel. Pure IgG was recovered in the flow-through fraction, whereas albumin and other contaminating proteins bound to the column. When the behaviour of parvovirus B19 in the ANX Sepharose column was studied by spiking the starting solution with high-titre parvovirus-positive plasma, the ANX gel bound most of the parvovirus, resulting in an average reduction of 10^{3.1} in the IgG fraction. About 30% of the parvovirus B19 was recovered in the wash fraction eluted with 1 M NaCl. The column was cleaned and sanitized with 0.5 M NaOH at room temperature to destroy and remove potential residual viruses from the column.

During the optimization of the nanofiltration step, it was found that the pH of the IgG solution had a major effect on the flux and filtration capacity. The optimal pH was found to be ≈ 4.4 – 4.8 , whereas the filtration capacity was clearly lower, at pH 4.2, 5.0 and 5.2 (Fig. 3b). At optimal pH, the PEG-treated pure IgG solution could be filtered with high efficacy, yielding ≈ 11 kg of IgG/m² of filter area, with a decrease in flux of $< 50\%$. When different virus filters were compared, the best flux with the process intermediate was achieved by using a composite membrane filter (Millipore V-NVP).

Reduction of parvovirus B19 by the nanofiltration step was studied by PCR. As the PCR assay detects not only DNA in virus particles but also free virus DNA occurring in high-titre plasma, we treated the samples of the spiked starting solution and filtrate with DNase, which destroys free DNA. The reduction factor calculated for the removal of total virus DNA was 3.8 log₁₀, and 4.1 log₁₀ when calculated from the




Fig. 3 Influence of (a) polyethylene glycol (PEG) treatment and (b) pH on the flow rate in nanofiltration. Five-hundred millilitres of the pure immunoglobulin G (IgG) solution (≈ 8 g/l), recovered after the anion-exchange chromatography, was filtered through a 0.1- μ m prefilter and a Millipore V-NVP filter (3.5 cm²) at 35 °C at a constant pressure of 3.5 bar. (a) The starting material treated with the different concentrations of caprylic acid (CA) and PEG indicated was adjusted to pH 4.4. (b) The starting solution that had been treated with 3% PEG was adjusted to the different pH values indicated.




Fig. 4 Reduction of bovine viral diarrhoea virus (BVDV) during caprylic acid precipitation. At 60 min the BVDV was completely inactivated (RF > 5.8 log₁₀). CA, caprylic acid; RF, reduction factor.

Table 1 Reduction of parvovirus B19 DNA in the different process steps

Process step	Log ₁₀ reduction factor
Caprylic acid precipitation	1.7
Polyethylene glycol precipitation	4.6
ANX chromatography	3.1
Nanofiltration	4.1
Total reduction factor	13.5

Table 2 Yield of immunoglobulin G (IgG) in the purification process starting from recovered plasma

Process step	IgG yield	
	Plasma (g/kg)	%
Suspended II+III paste	7.5	100
After caprylic acid treatment	6.8	91
After polyethylene glycol precipitation	5.7	76
After chromatography	5.1	68
After nanofiltration	5.0	67
Final product	4.8	64

DNAse-treated samples. The cumulative reduction of parvovirus in the different process steps was ≈ 14 log₁₀ (Table 1).

The overall yield of IgG from dissolved fraction II+III paste to final product was $\approx 64\%$, corresponding to ≈ 4.8 g/l from recovered plasma. The final product had high purity and did not contain detectable polymers (Table 2). Four batches of pure IgG solution were manufactured at a laboratory scale for the stability studies. The process proved reproducible within

reasonable tolerance limits of the process parameters. The IgG subclass distribution was similar to that of the starting plasma, with somewhat lowered proportions of IgG3 and IgG4. Some IgG3 was lost in the removal of immunoglobulin polymers, and IgG4 was lowered in the anion-exchange chromatography when removal of IgA was optimized (Table 3).

We studied the stability of the final product as a 100 g/l IgG solution at pH 4.4 and 5.3, and compared trehalose, a non-reducing disaccharide, with glycine as a stabiliser. No polymer formation took place in any of the formulations during 12 months at room (25 ± 2 °C) or refrigerator ($2-8$ °C) temperatures. Interestingly, the formation of polymers at an elevated temperature (37 ± 2 °C) was more effectively prevented by trehalose than by glycine. However, an increase in pH from 4.4 to 5.3 was even more effective than replacement of glycine with trehalose in preventing polymer formation at this temperature (Fig. 5). No fragmentation was detected at the refrigerator temperature during 12 months. At room temperature, slight fragmentation could be detected, and fragmentation was clearer at the elevated temperature, but again, to a lower extent, in the formulations at pH 5.3 than at pH 4.4 (Fig. 5).

Discussion

The new method described here makes it possible to purify IgG, at a high yield, from human plasma in a few process steps.

Table 3 Characteristics of intravenous immunoglobulin (IVIg) manufactured by the new process

Analysis	Result
Monomers %	92.9
Dimers %	7.1
Polymers %	0.0
Fragments %	0.0
IgG 1% (in plasma %)	59.0 (55-57)
IgG 2% (in plasma %)	36.0 (32-35)
IgG 3% (in plasma %)	2.4 (3.5-4.1)
IgG 4% (in plasma %)	2.6 (6.0-6.4)
IgG g/l	100
IgA mg/l	14.6
ACA CH50/mg	0.47
IgM mg/l	< 17
PKA IU/ml	< 5
KAL IU/ml	< 5
Albumin mg/l	< 13
PEG g/l	0.16
Caprylate g/l	< 0.17

ACA, anticomplementary activity; IgA, immunoglobulin A; IgG, immunoglobulin G; IgM, immunoglobulin M; KAL, kallikrein; PEG, polyethylene glycol; PKA, prekallikrein activator.




Fig. 5 Polymer formation (a) and fragmentation (b,c) during the storage of immunoglobulin G (IgG) purified by the new process and formulated to 100 g/l solutions at different pH values and with different stabilisers indicated. No polymer formation took place at 2–8 °C or 25 °C, and no fragmentation took place at 2–8 °C.

The main benefit of the new method is its exceptionally high capacity to remove physico-chemically resistant viruses, such as parvovirus. This is based on the efficient precipitation with PEG, which removes both viruses and polymeric proteins and enables efficient nanofiltration downstream in

the process. The pure, essentially polymer-free IgG obtained after a single anion-exchange chromatography can be filtered with high flux and low clogging tendency through a small pore-size virus-removal filter. After concentration and diafiltration, the purified IgG proved stable as a 10% solution.

Caprylic acid was found, by Steinbuch & Audran, to be beneficial in the isolation of IgG [21]. Later, Lundblad & Seng [22] showed that it effectively inactivates enveloped viruses. The caprylic acid concentration used in the current process is similar to the concentration used by Steinbuch & Audran [21] and later shown to be effective in the inactivation of various enveloped viruses by Dichtelmüller *et al.* [23]. In the caprylic acid process developed by Lebing *et al.* [11], sodium caprylate is added in two consecutive steps, and somewhat lower concentrations were found to be effective in virus inactivation at those conditions [24].

PEG precipitation has been used since the 1980s for the removal of polymeric immunoglobulin and so-called ACA of IVIG [25,26]. It has typically been used at concentrations of 4% or higher. In the process described in the present study, excellent removal of polymers was obtained with 2–3% PEG, which evidently was a combined effect of PEG and caprylate. This allowed the recovery of IgG, at a good yield, in the supernatant. Even though no polymers could be detected by size-exclusion liquid chromatography in the purified IgG treated with 2% PEG, the flux and capacity in nanofiltration were clearly increased when the PEG concentration was increased to 3%.

The clogging of small pore-size virus filters with protein aggregates is well known. Hirasaki *et al.* [27] showed that clogging of filter pores with protein aggregates results in decreasing flux and impairs virus removal, probably by shifting the residual flux to larger pores. In the present study we found that in addition to the freedom of polymers, pH had a profound effect on IgG throughput in the nanofiltration. The optimal pH range at 35 °C was surprisingly narrow, at ≈ 4.4 – 4.8 . It has been shown that IgG changes its conformation reversibly at acid pH and elevated temperatures close to 35 °C [28,29]. A conformational change could explain the remarkable improvement in filtrate flux and disappearance of clogging tendency observed in the present study.

Previously, filtration of IVIG products with virus-removal filters, which are capable of effectively removing even small viruses such as parvovirus, has been relatively expensive. This is because of the limited amount of IgG that could be filtered before the filters became clogged. The current method makes it possible to filter even close to 10 kg of IgG with high yield through 1-m² of a virus-removal filter, which lowers manufacturing costs. Effective removal ($\approx 4 \log_{10}$) of parvovirus B19 was observed under the optimized filtration conditions. It is possible that parvovirus antibodies, which are always present in large plasma pools, bound to the viruses (despite the relatively low pH) and contributed to virus removal during

the filtration. The contributing effect of virus antibodies in nanofiltration has been demonstrated previously [30].

The combination of an effective virus-inactivation step with two effective virus-removal steps increases the safety margin of the IVIG purified by the new process. Although the risk of parvovirus transmission with current IVIG products is already very low, based on the presence of protecting antibodies in plasma pools, and limitation of virus load by PCR testing [31], other non-enveloped viruses, with less commonly occurring neutralizing antibodies, may still pose a threat. On the other hand, the original Cohn process has proved effective in removing prions, which are far more resistant to physicochemical agents than non-enveloped viruses. In particular, the precipitation of fraction III effectively removes prions [32,33] and when this step is omitted from IgG manufacturing when aiming at a higher yield, other process steps with corresponding efficacy should be considered for the new process to maintain the same level of safety. Both PEG precipitation and nanofiltration have proved effective in removal of prion infectivity and provide a beneficial combination also in this respect [16,33]. Additionally, caprylic acid precipitation and the subsequent depth filtration have been shown to effectively remove prions [34].

The concentrated pure IgG solution obtained after ultrafiltration proved stable during long-term storage. Interestingly, trehalose prevented, more effectively than glycine, polymer formation of IgG at an elevated temperature. Trehalose is known to be an excellent cryoprotectant but less is known about its ability to protect proteins in solutions [35]. However, an increase of pH to > 5 was even more effective than trehalose in preventing IgG polymerization. This is in accordance with the increase in thermal stability of concentrated IgG solutions when the pH is raised above 5.0, which has been shown by differential scanning calorimetry [36]. The slight fragmentation of IgG found at room temperature was similar to that reported for other liquid IVIG products [37]. Similarly to polymer formation at the elevated temperature, fragmentation was also less pronounced during storage at pH 5.3 than at 4.4.

In conclusion, our results indicate that it is possible to manufacture, with high yield from Cohn fraction II+III, stable, polymer-free IgG that can be filtered with high capacity through a small pore-size virus-removal filter. The polymer removal step also serves as an effective virus-reduction step and, as a whole, the process has very high capacity to remove even physico-chemically stable viruses.

References

- Buckley RH, Schiff RI: The use of intravenous immune globulin in immunodeficiency diseases. *N Engl J Med* 1991; 325:110-117
- Kazatchkine MD, Kaveri SV: Immunomodulation of autoimmune and inflammatory diseases with intravenous immune globulin. *N Engl J Med* 2001; 345:747-755
- Knezevic-Maramica I, Kruskal MS: Intravenous immune globulins: an update for clinicians. *Transfusion* 2003; 43:1460-1480
- Cohn EJ, Strong LE, Hughes WL Jr, Mulford DJ, Ashworth JN, Melin M, Taylor HL: Preparation and properties of serum and plasma proteins III. A system for the separation into fractions of the protein and lipoprotein components of biological tissues and fluids. *J Am Chem Soc* 1946; 68:459-475
- Oncley JL, Melin M, Richert DA, Cameron JW, Gross PM Jr: The separation of the antibodies, isoagglutinins, prothrombin, plasminogen and beta-lipoprotein into subfractions of human plasma. *J Am Chem Soc* 1949; 71:541-550
- Barandun S, Kistler P, Jeunet F, Isliker H: Intravenous administration of human gamma globulin. *Var Sang* 1962; 7:157-174
- Romer J, Spath PJ, Skvaril F, Nydegger UE: Characterization of various immunoglobulin preparations for intravenous application. II. Complement activation and binding to staphylococcus protein A. *Var Sang* 1982; 42:74-80
- Lane RS: Non-A, non-B hepatitis from intravenous immunoglobulin. *Lancet* 1983; ii:974-975
- Lever AM, Webster AD, Brown D, Thomas HC: Non-A, non-B hepatitis occurring in agammaglobulinaemic patients after intravenous immunoglobulin. *Lancet* 1984; ii:1062-1064
- Bjoro K, Froland SS, Yun Z, Samdal HH, Haaland T: Hepatitis C infection in patients with primary hypogammaglobulinemia after treatment with contaminated immune globulin. *N Engl J Med* 1994; 331:1607-1611
- Lebing W, Remington KM, Schreiner C, Paul HI: Properties of a new intravenous immunoglobulin (IGIV-C, 10%) produced by virus inactivation with caprylate and column chromatography. *Var Sang* 2003; 84:193-201
- Prowse C, Ludlam CA, Yap PL: Human parvovirus B19 and blood products. *Var Sang* 1997; 72:1-10
- Young NS: Parvovirus infection and its treatment. *Clin Exp Immunol* 1996; 104:26-30
- Hayakawa F, Imada K, Towatari M, Saito H: Life-threatening human parvovirus B19 infection transmitted by intravenous immune globulin. *Br J Haematol* 2002; 118:1187-1189
- Erdman DD, Anderson BC, Török TJ, Finkel TH, Anderson LJ: Possible transmission of parvovirus B19 from intravenous immune globulin. *J Med Virol* 1997; 53:233-236
- Burnouf Radosevich M: Nanofiltration of plasma-derived biopharmaceutical products. *Haemophilia* 2003; 9:24-37
- Hirvonen M, Koskinen S, Tölö H: A sensitive enzyme immunoassay for the measurement of low concentrations of IgA. *J Immunol Methods* 1993; 163:59-65
- European Pharmacopoeia, 5th edn. Strasbourg, Council of Europe, 2004
- Skoog B: Determination of polyethylene glycols 4000 and 6000 in plasma protein preparations. *Var Sang* 1979; 37:345-349
- Terpstra FG, Parkkinen J, Tölö H, Koenderman AHL, ter Hart HGJ, von Bonsdorff L, Törmä E, van Engelenburg FAC: Viral safety of Nanogam®, a new 15 nm-filtered liquid immunoglobulin product. *Var Sang* 2006; 90:21-32
- Steinbuch M, Audran R: The isolation of IgG from mammalian sera with the aid of caprylic acid. *Arch Biochem Biophys* 1969; 134:279-284
- Lundblad JL, Seng RL: Inactivation of lipid-enveloped viruses in proteins by caprylate. *Var Sang* 1991; 60:75-81

- 23 Dichtelmüller H, Rudnick D, Kloft M: Inactivation of lipid enveloped viruses by octanoic acid treatment of immunoglobulin solution. *Biologicals* 2002; 30:135–142
- 24 Korneyeva M, Hotta J, Lebing W, Rosenthal RS, Franks L, Petteway SR Jr: Enveloped virus inactivation by caprylate: a robust alternative to solvent-detergent treatment in plasma derived intermediates. *Biologicals* 2002; 30:153–162
- 25 Eibl MM, Cairns L, Rosen FS: Safety and efficacy of a monomeric, functionally intact intravenous IgG preparation in patients with primary immunodeficiency syndromes. *Clin Immunol Immunopathol* 1984; 31:151–160
- 26 Uemura Y, Uriyu K, Hirao Y, Takechi K, Ishikawa H, Nakajima T, Kagitani Y, Yokoyama K, Funakoshi S, Nishida M: Inactivation and elimination of viruses during the fractionation of an intravenous immunoglobulin preparation: liquid heat treatment and polyethylene glycol fractionation. *Var Sang* 1989; 56:155–161
- 27 Hirasaki T, Ishikawa G, Nakano H, Manabe S-I, Yamamoto N: Mechanism of virus removal by regenerated cellulose hollow fiber (BMMTM). *Membrane* 1995; 20:135–142
- 28 Zav'yalov V, Troitsky GV, Demchenko AP, Generalov IV: Temperature and pH dependent changes of immunoglobulin G structure. *Biochem Biophys Acta* 1975; 386:155–167
- 29 Calmettes P, Cser L, Rajnavolgyi E: Temperature and pH dependence of immunoglobulin G conformation. *Arch Biochem Biophys* 1991; 291:277–283
- 30 Omar A, Kempf C: Removal of neutralized model parvoviruses and enteroviruses in human IgG solutions by nanofiltration. *Transfusion* 2002; 42:1005–1010
- 31 Schmidt I, Blümel J, Seitz H, Willkommen H, Löwen J: Parvovirus B19 DNA in plasma pools and plasma derivatives. *Var Sang* 2001; 81:228–235
- 32 Reichl HE, Foster PR, Welch AG, Li Q, MacGregor IR, Somerville RA, Fernie K, Steele PJ, Taylor DM: Studies on the removal of a bovine spongiform encephalopathy-derived agent by processes used in the manufacture of human immunoglobulin. *Var Sang* 2002; 83:137–145
- 33 Lee DC, Stenland CJ, Hartwell RC, Ford EK, Cai K, Miller JL, Gilligan KJ, Rubenstein R, Fournel M, Petteway SR Jr: Monitoring plasma processing steps with a sensitive Western blot assay for the detection of the prion protein. *J Virol Methods* 2000; 84:77–89
- 34 Trejo R, Hotta JA, Lebing W, Stenland C, Storms RE, Lee DC, Li H, Petteway S Jr, Remington KM: Evaluation of virus and prion reduction in a new intravenous immunoglobulin manufacturing process. *Var Sang* 2003; 84:176–187
- 35 Xie G, Timasheff SN: The thermodynamic mechanism of protein stabilization by trehalose. *Biophys Chem* 1997; 64:25–43
- 36 Tatford OO, Gomme PT, Bertolini J: Analytical techniques for the evaluation of liquid protein therapeutics. *Biotechnol Appl Biochem* 2004; 40:67–81
- 37 Diemel RV, ter Hart H, Derksen GJA, Koenderman A, Aalberse R: Characterization of immunoglobulin G fragments in liquid intravenous immunoglobulin products. *Transfusion* 2005; 45:1601–1609

医薬品 研究報告 調査報告書

識別番号・報告回数		報告日	第一報入手日	新医薬品等の区分	総合機構処理欄
一般的名称	-	研究報告の 公表状況	http://www.bloodservices.ca/CentreApps/Internet/ UW_V502_MainEngine.nsi/web/ C2364AA876DBB6998525705E0052854F?OpenDocument	公表国	
販売名(企業名)	-			カナダ	
研究報告の概要	<p>英国およびフランスでBSEの流行がピークとなった1980年1月から1996年12月という期間を反映し、1999年9月30日に開始された、変異型クロイツフェルトヤコブ病(vCJD)伝播のリスクから供血システムを保護するための予防措置が変更された。</p> <ol style="list-style-type: none"> 1980年1月1日以降、英国、フランスおよび西ヨーロッパで、輸血または血液製剤の投与を受けたことのあるドナーは、無期限に供血停止とする(これまでは英国に限定)。 1980年1月1日から1996年12月31日の間に、フランスまたは英国に累積合計3ヵ月滞在以上したドナーについては、無期限に供血停止とする(これまでは1997年1月1日以降も対象期間に含まれていた)。 英国またはフランスでの累積3ヵ月間の滞在期間が1980年1月1日から1996年12月31日の間に入らないドナーについては、供血適格者とする。 <p>また、抗体検査法の改良やC型肝炎およびHIVの核酸検査法(NAT)の実施により、輸血伝播性疾患の検査法が著しく進歩していることから、一時的な供血停止措置に対する変更として、刺青、ピアス、針治療、電気治療を受けている人、性的背景が不明のパートナーと性的接触を行っている人、針刺し事故を受けたことのある人、他人の血液と接触したことのある人の供血停止期間を12ヵ月から6ヵ月に短縮する。48時間(これまでは3日間)以上投獄されていた人は、釈放から12ヵ月間は、供血停止とする。</p>				使用上の注意記載状況・ その他参考事項等
	報告企業の意見	今後の対応			
vCJDに対する献血除外基準の緩和、HCV、HIVに対する短期的な供血停止期間変更に関する情報である。vCJDについては日本の献血制限は英国への滞在が1980年～1996年の間に1日以上等、本報告の内容よりも厳しいものとなっている。	今後ともvCJDに関する安全性情報、規制情報等に留意していく。				

Canadian Blood Services Makes Changes to Donor Deferral Criteria
Welcomes back donors who travelled to U.K. and France since 1996

Ottawa, August 15, 2005 – Canadian Blood Services is implementing changes to its donor deferral criteria in order to reflect the most up-to-date scientific research regarding risks to the blood supply. As a result, some donors will be subject to new deferrals while deferral periods for others will be reduced or eliminated.

"Safety and adequacy are two essential components of the blood system," said Dr. Graham Sher, Chief Executive Officer for Canadian Blood Services. "These changes will allow us to keep our commitment to Canadians on both counts."

Changes to Indefinite Deferrals – vCJD

Since September 30, 1999 safeguards have been in place to protect the blood system from the risk of transmission of variant Creutzfeldt Jakob Disease (vCJD). Donors who meet certain criteria under this policy are indefinitely deferred from donating. The following changes directly reflect the most recent information on the safeguards the United Kingdom, France and Western Europe have in place to protect the bovine and human populations:

1. Donors who have received a blood transfusion or received medical treatment with a product made from blood in the United Kingdom, France or Western Europe since January 1, 1980 will now be deferred indefinitely. Previously, this deferral was limited to the United Kingdom.
2. Donors who have spent a cumulative total of three months or more in France or in the United Kingdom between January 1, 1980 and December 31, 1996 will be deferred indefinitely. In the past, donors who had spent a cumulative total of three months or more in France or the United Kingdom since January 1, 1980 were deferred.
3. Donors whose cumulative three month travel period to the UK or France did not occur between January 1, 1980 and December 31, 1996, will once again be eligible to donate.

Since 1992, confirmed BSE cases in the UK and France have been steadily declining. The 1996 cut-off date is reflective of the period between January 1980 and December 1996 when the BSE epidemic was at its peak in the United Kingdom and France. Since that time, cases have continued to decline and BSE monitoring and control mechanisms have been implemented to stop the spread of the disease in the bovine population and thereby decreasing the risk of transmission of vCJD to humans.

For more information on indefinite deferrals (vCJD), [click here](#).

Changes to Temporary Deferrals

Some prospective blood and/or bone marrow donors may be unable to donate for a period of time for reasons of their own health or the safety of the blood supply or marrow product. The following changes are being made to the temporary deferral criteria policies:

1. The following deferral periods will be reduced from 12 months to six months:
 - Persons who have a tattoo, ear or body piercing, or who have undergone acupuncture or electrolysis procedures;
 - Individuals who have had sexual contact with a partner whose sexual background is unknown; and
 - Individuals who have been injured by a needle or who have had contact with

blood from another person.

Reducing the deferral period reflects the latest available medical research on the "window period" – the brief period after the onset of a viral infection during which early signs of a virus cannot be detected by existing tests. Additionally, significant advances in transfusion transmissible disease testing has occurred in recent years, such as improved antibody assays and more recently, the implementation of nucleic acid testing (NAT) for hepatitis C (HCV) and HIV.

2. In order to comply with the Canadian Standards Association standard on Blood and Blood components, persons who have been incarcerated for 48 hours (rather than three days – Canadian Blood Services previous standard) or more will now be deferred from donating blood for 12 months following the date of release from incarceration.

For more information on temporary deferrals, [click here](#).

Message to Donors

If you believe that you may now be eligible to donate or would like more information, please call 1-888-2-DONATE to speak to a Canadian Blood Services representative who will be able to provide more details and/or book an appointment.

About Canadian Blood Services

Canadian Blood Services is a national, not-for-profit charitable organization that manages the blood supply in all provinces and territories outside of Quebec and oversees the country's Unrelated Bone Marrow Donor Registry. Canadian Blood Services operates 41 permanent collection sites and more than 19,000 donor clinics annually. The Provincial and Territorial Ministers of Health provide operational funding to Canadian Blood Services. The federal government, through Health Canada, is responsible for regulating the blood system. For more information, please visit the website at www.bloodservices.ca.

-30-

For further information, please contact:

Derek Mellon
Media Room Relations Manager
Canadian Blood Services
(613) 739-2177
derek.mellon@bloodservices.ca



CANADIAN BLOOD SERVICES SOCIÉTÉ CANADIENNE DU SANG