



トリス(2,3-ジブロモプロピル) ホスフェート<略称 TBPP> (CAS.126-72-7)

①生長曲線



②阻害率曲線



Dose-response curve for EC50 of Algae Growth Test (Logit method)
126727

③毒性値

72hErC50 (実測値に基づく) > 2.7mg/L
72hNOECr (実測値に基づく) = 1.2mg/L

要 約

試験委託者： 環境省

表 題： トリス (2,3-ジブロモプロピル) ホスフェートのオオミジンコ
(*Daphnia magna*) に対する急性遊泳阻害試験

試験番号： A020373-2

試験方法：

- 1) 適用ガイドライン： OECD 化学品テストガイドライン No.202 「ミジンコ類, 急性遊泳阻害試験および繁殖試験」 (1984年)
- 2) 暴露方式： 半止水式 (24時間後に試験液の全量を交換)
水面をテフロンシートで被覆
- 3) 供試生物： オオミジンコ (*Daphnia magna*)
- 4) 暴露期間： 48時間
- 5) 試験濃度： 対照区, 助剤対照区,
(設定値) 0.450, 0.800, 1.42, 2.53, 4.50 mg/L
試験液調製可能最高濃度
公比： 1.8
助剤濃度一定：100 μ L/L (V^oメチルセルロース使用)
- 6) 試験液量： 100 mL/容器
- 7) 連 数： 4容器/試験区
- 8) 供試生物数： 20頭/試験区 (5頭/容器)
- 9) 試験温度： 20 \pm 1 $^{\circ}$ C
- 10) 照 明： 室内光, 16時間明 (800 lux以下) /8時間暗
- 11) 分 析 法： 高速液体クロマトグラフィー質量分析 (LC/MS)

試験結果：

1) 試験液中の被験物質濃度

試験液の分析の結果、測定値の設定値に対する割合は、暴露開始時において97~101%、換水前において88~100%であった。


2) 24時間暴露後の結果

	(mg/L)	95%信頼区間 (mg/L)
半数遊泳阻害濃度 (EiC50)	> 4.50	算出不可
0%阻害最高濃度	2.41	—
100%阻害最低濃度	> 4.50	—

3) 48時間暴露後の結果

	(mg/L)	95%信頼区間 (mg/L)
半数遊泳阻害濃度 (EiC50)	4.16	算出不可
0%阻害最高濃度	2.41	—
100%阻害最低濃度	> 4.50	—

Figure 1 Concentration-Immobility Curve



要 約

試験委託者： 環境省

表 題： トリス (2,3-ジブロモプロピル) ホスフェートのオオミジンコ
(*Daphnia magna*) に対する繁殖阻害試験

試験番号： A020373-3

試験方法：

- 1) 適用ガイドライン： OECD 化学品テストガイドライン No. 211 「オオミジンコ繁殖試験」 (1998年)
- 2) 暴露方式： 半止水式 (毎日試験液の全量を交換)
水面をテフロンシートで被覆
- 3) 供試生物： オオミジンコ (*Daphnia magna*)
- 4) 暴露期間： 21日間
- 5) 試験濃度： 対照区, 助剤対照区, 0.150, 0.350, 0.820, 1.92, 4.50 mg/L
(設定値) (公比: 2.3)
ただし 4.50 mg/Lは試験液調製可能最高濃度
助剤濃度一定: 100 μ L/L (ジメチルホルムアミド使用)
- 6) 試験液量： 80 mL/容器
- 7) 連 数： 10容器/試験区
- 8) 供試生物数： 10頭/試験区 (1頭/容器)
- 9) 試験温度： 20 \pm 1 $^{\circ}$ C
- 10) 照 明： 室内光, 16時間明 (800 lux以下) / 8時間暗
- 11) 分 析 法： 高速液体クロマトグラフィー質量分析 (LC/MS)


試験結果：

1) 試験液中の被験物質濃度

試験液の分析の結果、測定値の設定値に対する割合は、調製時において 93~108%、
換水前において 91~109%であった。


2) 21日間暴露後の結果

	(mg/L)	95%信頼区間 (mg/L)
親ミジンコの半数致死濃度 (LC50)	3.00	1.86~4.84
50%繁殖阻害濃度 (EC50)	1.87	1.67~2.21
最大無作用濃度 (NOEC)	0.832	—
最小作用濃度 (LOEC)	1.86	—

Figure 1 Cumulative Number of Dead Parental *Daphnia*

Values in legend are given in the nominal concentration.

Figure 2 Time Course of $\Sigma F1/P$ for Each Concentration Level



Values in legend are given in the nominal concentration.

—: All parental *Daphnia* were dead during a 21-days testing period.

要 約

試験委託者：環境省

表 題： トリス (2,3-ジブロモプロピル) ホスフェートの
ヒメダカ (*Oryzias latipes*) に対する急性毒性試験

試験番号： A020373-4

試験方法：

- 1) 適用ガイドライン： OECD 化学品テストガイドライン No. 203 「魚類急性毒性試験」
(1992年)
- 2) 暴露方式： 半止水式 (24時間毎に試験液の全量を交換)
水面をテフロンシートで被覆
- 3) 供試生物： ヒメダカ (*Oryzias latipes*)
- 4) 暴露期間： 96時間
- 5) 試験濃度： 対照区, 助剤対照区, 0.450, 0.800, 1.40, 2.50, 4.50 mg/L
(設定値) (試験液調製可能最高濃度)
公比：1.8
助剤濃度一定：100 μ L/L (シキリホウヤミ[®]使用)
- 6) 試験液量： 5.0 L/容器
- 7) 連 数： 1 容器/試験区
- 8) 供試生物数： 10尾/試験区
- 9) 試験温度： 24 \pm 1 $^{\circ}$ C
- 10) 照 明： 室内光, 16時間明 (1000 lux以下) / 8時間暗
- 11) 分 析 法： 高速液体クロマトグラフ質量分析計 (LC/MS)


試験結果：

1) 試験液中の被験物質濃度

試験液の分析の結果、測定値の設定値に対する割合は、暴露開始時において93~109%、24時間後において84~98%であった。

2) 96時間暴露後の半数致死濃度(LC50)： 1.86 mg/L (95%信頼区間： 1.35 ~ 2.57 mg/L)

Figure 1 Concentration-Mortality Curve



要 旨

試験委託者

環境庁

表 題

2,6-ジ-*tert*-ブチル-*p*-クレゾールの藻類 (*Selenastrum capricornutum*) に対する生長阻害試験

試験番号

9 B 4 4 7 G

試験方法

本試験は、OECD 化学品テストガイドライン No. 201「藻類生長阻害試験」(1984年)に準拠して実施した。

- 1) 被験物質: 2,6-ジ-*tert*-ブチル-*p*-クレゾール
- 2) 暴露方式: 止水式(密閉), 振とう培養(100rpm)
- 3) 供試生物: *Selenastrum capricornutum* (ATCC22662)
- 4) 暴露期間: 72時間
- 5) 試験濃度(設定値):
対照区, 助剤対照区, 1.00, 2.15, 4.64, 10.0 mg/L
(分散可能最高濃度)
(公比: 2.2, 助剤濃度一定: 100 mg/L, ジメチルホルムアミド および HCO-40使用)
- 6) 試験液量: 100 mL (OECD培地) / 容器
- 7) 連数: 3 容器 / 濃度区
- 8) 初期細胞濃度: 1×10^4 cells/mL
- 9) 試験温度: 23 ± 2 °C
- 10) 照明: 4000 lux (±20%の変動内, フラスコ液面付近) で連続照明
- 11) 分析法: HPLC法

結 果

1) 試験液中の被験物質濃度

被験物質の測定濃度が開始時において設定値の±20%を超えたものがなかったため、下記の生長阻害濃度の算出には設定値を採用した。

2) 生長曲線下面積の比較による阻害濃度

50%生長阻害濃度 EbC50 (0-72) : >10.0 mg/L (95%信頼区間:算出不可)

最大無作用濃度 NOECb (0-72) : 1.00 mg/L

3) 生長速度の比較による阻害濃度

50%生長阻害濃度 ErC50 (24-48) : >10.0 mg/L (95%信頼区間:算出不可)


最大無作用濃度 NOECr (24-48) : >10.0 mg/L

50%生長阻害濃度 ErC50 (24-72) : >10.0 mg/L (95%信頼区間:算出不可)


最大無作用濃度 NOECr (24-72) : 1.00 mg/L

2,6-ジ-tert-ブチル-p-クレゾール (CAS.128-37-0)

① 生長曲線



②阻害率曲線



③毒性値

48hErC50(実測値に基づく) >7.0mg/L
48hNOECr(実測値に基づく) =1.7mg/L

要 旨

試験委託者

環境庁

表 題

2,6-ジ-*tert*-ブチル-*p*-クレゾールのオオミジンコ (*Daphnia magna*) に対する急性遊泳阻害試験

試験番号

9 B 4 6 9 G

試験方法

本試験は、OECD 化学品テストガイドライン No. 202 「ミジンコ類、急性遊泳阻害試験および繁殖試験」(1984年)に準拠して実施した。

- 1) 被験物質： 2,6-ジ-*tert*-ブチル-*p*-クレゾール
- 2) 暴露方式： 止水式，水面をテフロンシートで被覆
- 3) 供試生物： オオミジンコ (*Daphnia magna*)
- 4) 暴露期間： 48時間
- 5) 試験濃度 (設定値) :
対照区，助剤対照区，0.200, 0.360, 0.630, 1.10, 2.00 mg/L
公比：1.8
助剤濃度一定：40.0 mg/L (HCO-40 および ジメチルホルムアミド 使用)
- 6) 試験液量： 100 mL/容器
- 7) 連数： 4容器/濃度区
- 8) 供試生物数： 20頭/濃度区 (5頭/容器)
- 9) 試験温度： 20±1℃
- 10) 照明： 16時間明/8時間暗
- 11) 分析法： HPLC法

結 果

1) 試験液中の被験物質濃度

被験物質の測定濃度がすべて設定値の±20%以内であったため、各影響濃度の算出には設定値を採用した。

2) 24 時間暴露後の結果

半数遊泳阻害濃度 (EiC50) : >2.00 mg/L (95%信頼限界 : 算出不可)

最大無作用濃度 (NOECi) : 0.630 mg/L

100%阻害最低濃度 : >2.00 mg/L


3) 48 時間暴露後の結果

半数遊泳阻害濃度 (EiC50) : 0.835 mg/L (95%信頼限界 : 0.709~0.985 mg/L)

最大無作用濃度 (NOECi) : 0.360 mg/L

100%阻害最低濃度 : 2.00 mg/L

Figure 1 Concentration-Response (Immobility) Curve



要 旨

試験委託者

環境庁

表 題

2,6-ジ-*tert*-ブチル-*p*-クレゾールのオオミジンコ (*Daphnia magna*) に対する繁殖阻害試験

試験番号

9B491G

試験方法

本試験は、OECD 化学品テストガイドラインNo. 211「オオミジンコ繁殖試験」(1998年)に準拠して実施した。

- 1) 被験物質： 2,6-ジ-*tert*-ブチル-*p*-クレゾール
- 2) 暴露方式： 半止水式 (24時間毎に試験液の全量を交換)
水面をテフロンシートで被覆
- 3) 供試生物： オオミジンコ (*Daphnia magna*)
- 4) 暴露期間： 21日間
- 5) 試験濃度 (設定値)：
対照区, 助剤対照区, 0.008, 0.025, 0.080, 0.250, 0.800 mg/L
公比：3.2
助剤濃度一定：100 mg/L (HCO-60 および ジメチルホルムアミド 使用)
- 6) 試験液量： 80 mL/容器
- 7) 連数： 10容器/濃度区
- 8) 供試生物数：10頭/濃度区 (1頭/容器)
- 9) 試験温度： 20±1℃
- 10) 照明： 16時間明/8時間暗
- 11) 分析法： HPLC法

結 果

1) 試験液中の被験物質濃度

被験物質の測定濃度が設定値の±20%を超えたものがあったため、各影響濃度の算出には測定値（時間加重平均値）を採用した。

2) 21日間暴露の各影響濃度結果を以下に示す。

親ミジンコの半数致死濃度 (LC50) : 0.390 mg/L

(95%信頼限界 : 0.218 ~0.698 mg/L)


50% 繁殖阻害濃度 (EC50) : 0.096 mg/L

(95%信頼限界 : 0.086~0.116 mg/L)

最大無作用濃度 (NOEC) : 0.069 mg/L

最小作用濃度 (LOEC) : 0.218 mg/L

Figure 1 Cumulative Numbers of Dead Parental *Daphnia*




Values in legend are given in the nominal concentration.

Table 4 Mean Cumulative Numbers of Juveniles Produced per Adult Alive for 21 Days ($\Sigma F1/P$)

Nominal Conc.	Days															
	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21
Control	0.0	7.3	7.9	7.9	19.1	28.0	28.0	28.9	56.4	56.4	56.4	63.5	69.9	69.9	69.9	86.2
Solv. cont.	0.0	9.5	9.7	9.7	38.6	40.7	40.7	47.9	76.0	76.0	76.0	93.5	94.9	94.9	99.8	114.3
0.008 mg/L	0.0	7.8	7.8	7.8	38.1	38.4	38.4	52.9	75.8	75.8	75.8	92.2	92.2	92.2	98.5	112.0
0.025 mg/L	0.0	7.2	7.2	7.2	29.8	29.8	29.8	43.7	65.8	65.8	65.8	83.2	83.3	83.3	91.6	105.3
0.080 mg/L	0.0	4.4	4.4	4.4	25.0	25.0	25.0	25.6	57.4	57.4	57.4	74.7	74.7	74.7	88.6	100.9
0.250 mg/L	0.0	0.0	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
0.800 mg/L	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

-: All parental *Daphnia* were dead during a 21-days testing period.

Figure 2 Time Course of $\Sigma F1/P$ for Each Concentration Level



Values in legend are given in the nominal concentration.

要 旨

試験委託者

環境庁

表 題

2,6-ジ-*tert*-ブチル-*p*-クレゾールのヒメダカ (*Oryzias latipes*) に対する急性毒性試験

試験番号

9B513G

試験方法


本試験は、OECD 化学品テストガイドライン No. 203 「魚類毒性試験」 (1992年) に準拠して実施した。

- 1) 被験物質： 2,6-ジ-*tert*-ブチル-*p*-クレゾール
- 2) 暴露方式： 半止水式 (24時間毎に試験液の全量を交換) , 水面をテフロンシートで被覆
- 3) 供試生物： ヒメダカ (*Oryzias latipes*)
- 4) 暴露期間： 96時間
- 5) 試験濃度 (設定値) : 対照区, 助剤対照区, 0.500, 0.900, 1.60, 2.80, 5.00mg/L
公比 ; 1.8, 最大助剤濃度 ; 100 mg/L (メチルセルロース, HCO-40使用)
- 6) 試験液量： 5.0L/容器
- 7) 連数： 1 容器/濃度区
- 8) 供試生物数： 10尾/濃度区
- 9) 試験温度： 24±1℃
- 10) 照明： 室内光, 16時間明/8時間暗
- 11) 分析法： HPLC法

結 果

- 1) 試験液中の被験物質濃度：試験区において設定濃度に対して±20%を越える分析結果があったため、以下の値は測定濃度の幾何平均値を基に示した。
- 2) 96時間の半数致死濃度 (LC50) : 1.10 mg/L (95%信頼区間 : 0.895mg/L~1.36mg/L)

Figure 1 Concentration-Response (Mortality) Curve



SIDS INITIAL ASSESSMENT PROFILE

CAS No.	128-37-0
Chemical Name	2,6-di-tert.-butyl-p-cresol (BHT) Butylated Hydroxytoluene
Structural Formula	
RECOMMENDATIONS	
The chemical is a candidate for further work.	
SUMMARY CONCLUSIONS OF THE SIAR	
Human Health	
<p>BHT is of low acute toxicity. BHT caused acute toxic effects in mammals but there were no specific clinical symptoms. In rats, the oral LD₅₀ was > 2930 mg/kg bw, the LD₅₀ after dermal exposure was > 2000 mg/kg bw. It was slightly irritating to the skin and eyes of rabbits.</p> <p>On chronic oral exposure of rats, liver and thyroid are the main targets. Doses above 25 mg/kg bw/day BHT resulted in thyroid hyperactivity, enlargement of the liver, induction of several liver enzymes. 25 mg/kg bw/day BHT can be considered as NOAEL for chronic exposure. The haemorrhagic effects of high repeated doses of BHT seen in certain strains of mice and rats, but not in other species, may be related to its ability to interact with prothrombin and vitamin K.</p> <p>BHT showed no potential to cause point mutations in several bacterial and mammalian <i>in vitro</i> test systems.</p> <p>Overall, the available studies demonstrate that BHT has no clastogenic activity <i>in vitro</i> or <i>in vivo</i>. Most <i>in vitro</i> chromosome aberration assays were negative as were sister chromatid exchange assays and DNA damage and repair assays. <i>In vivo</i>, micronucleus assays with mice, cytogenetic assays with rats and mice, dominant lethal assays with rats and mice, and the heritable translocation assay with mice were also negative.</p> <p>BHT is not a genotoxic carcinogen. Carcinogenic effects observed in one long-term study with rats probably were caused by the specific study conditions. However, it cannot be completely ruled out that the hepatotoxic effects caused by high and chronic doses of BHT may result in persistent cell proliferation, which is known as a possible mechanism of non-genotoxic carcinogens. In addition, depending on the application regime, BHT may exert either anticarcinogenic or tumour-promoting activity at relatively high doses. For the possible carcinogenic and tumour-promoting effect of BHT, a threshold level of 100 mg/kg bw/day can be assumed. At this dose, no increase in the incidence of liver carcinoma, but a slight increase in liver adenoma were observed after chronic exposure starting <i>in utero</i> as a worst case scenario.</p> <p>The only effects on reproduction were lower numbers of litters of ten or more pups at birth at doses of 100 mg/kg bw/day and above. The NOAEL was 25 mg/kg bw/day.</p> <p>From studies with mice and rats there is no evidence of teratogenic effects of BHT. During pregnancy BHT had maternal effects on mice above oral doses of 240 mg/kg bw/day. The NOEL for developmental toxicity was 800 mg/kg bw day.</p> <p>Despite of being in wide dispersive use as ingredient of various products for many years only very few cases of</p>	

allergic reaction in humans after dermal exposure or oral intake have been described. For the use of BHT as antioxidant in foodstuff an acceptable daily intake (ADI) of 0 - 0.3 mg/kg bw/day has been established.

Environment

BHT has a melting point of ca. 70 °C, a water solubility in the range of 0.6-1.1 mg/l (20-25 °C), a density of 1.03 g/cm³, and a vapor pressure of 1.1 Pa (20 °C). The measured log Kow is determined to be 5.1.

According to a Mackay Level I model calculation, the main target compartment for BHT is air (79-87 %), followed by soil (6.1-10.2 %) and sediment (5.7-9.5 %). Due to the instability of BHT in aqueous solution the estimations reflect a tendency for BHT distribution among environmental compartments. BHT is relatively unstable under environmental conditions. Extent and products of decomposition are dependent on several factors like irradiation, pH, temperature, moisture, presence of soil and soil microorganisms, and oxygen content. In air BHT is indirectly photodegradable by hydroxyl radicals with $t_{1/2} = 7.0$ hours. In aqueous solution BHT is decomposed in natural sunlight with irradiation (ca. 75 %) and without (ca. 40 %), forming different, partly unidentified metabolites. BHT is also not stable in soil. Within one day of incubation 63-82 % of BHT were decomposed in non-sterilized and 25-35 % in sterilized soils. A mineralization up to 30 % was observed under non-sterilized conditions. Depending on the exposure pathways, the compartments air, hydrosphere and soil can be environmental target compartments for this substance and its metabolites. BHT is not readily biodegradable in water according to a modified MITI-I test (4.5 % degradation after 28 days). A wide range of bioconcentration factors (BCF) was found in different experiments. Bioconcentration factors (BCF) in the range of 230-2500 have been determined for fish after 56 days. The BCF values determined after a 28 days exposure period in a model ecosystem with soil were 2-17 for fish, 30 for snails and 38 for algae. It can be assumed that BHT has a moderate to high bioaccumulation potential in aquatic species.

For the toxicity of BHT on aquatic species reliable experimental results from tests with fish, daphnia, and algae are available. Only those effect values are considered for the assessment that did not exceed the low water solubility of BHT (0.6 - 1.1 mg/l) and were based on measured concentrations. The lowest reliable acute toxicity values are:

fish (*Brachydanio rerio*): 96h LC0 ≥ 0.57 mg/l;

invertebrates (*Daphnia magna*): 48h EC0 ≥ 0.17 mg/l;

algae (*Scenedesmus subspicatus*): 72h E_rC₈ = 0.4 mg/l. This value can be used as a NOEC.

In a 21 days reproduction test with *Daphnia magna* a NOEC = 0.07 mg/l was determined. Using an assessment factor of 50, a PNECaqua = 0.0014 mg/l is derived from this long term NOEC.

Exposure

In 2000, the world production capacity of BHT amounts to about 62,000 t/a by more than 20 producers. BHT is a registered antioxidant, licenced for food products, animal feed, cosmetics, and packaging material. It is also used in petroleum products, synthetic rubbers, plastics, elastomers, oils, waxes, soaps, paints, and inks.

Releases into the environment may occur during production of BHT as well as during its use in different applications as stabilizer and during the use of the products that contain the substance. A significant release into the environment is expected from migration of BHT onto the surface of products containing the substance.

NATURE OF FURTHER WORK RECOMMENDED

Environment: The substance is a candidate for further work. Releases into the environment during use of BHT and from products containing the substance have to be assumed but are not quantifiable. In the environment, BHT is rapidly decomposed forming several, partly unidentified, metabolites. BHT is not readily biodegradable, a moderate to high bioaccumulation potential has to be assumed. The NOEC from the long-term toxicity to daphnids was 0.07 mg/l, resulting in a PNEC of 0.0014 mg/l. Therefore, the performance of an environmental risk assessment is recommended. Especially the questions concerning exposure, bioaccumulation as well as toxicity of the metabolites should be clarified.

Human Health: No recommendation for further work, because all SIDS endpoints are adequately covered and because exposure is controlled in occupational settings.

4 HAZARDS TO THE ENVIRONMENT

4.1 Aquatic Effects

Acute Toxicity Test Results

Testing of the ecotoxicological behavior of BHT exhibits difficulties due to its extremely low water solubility of about 0.6 to 1.1 mg/l at room temperature and its tendency to decompose to various degradation products depending on the actual conditions (see chapt. 2.1.2). Based on the instability, low recovery rates of applied BHT from the test solutions are not surprising. Due the aforementioned instability of BHT the data obtained in testing cover not only the toxicity of BHT but also the toxicity of the oxidation/degradation products of BHT.

For the acute toxicity of BHT in aquatic species reliable experimental results from short-term tests with fish, daphnia, algae and microorganisms are available. Only those effect values were considered for the assessment that did not exceed the low water solubility of the compound (0.6 - 1.1 mg/l) and were based on measured concentrations.

A test on the acute toxicity of BHT to fish was conducted according to the European protocol EEC C.1 (equivalent to OECD guideline 203). Semistatic exposure (renewal of test solution every 24 hours) of *Brachydanio rerio* in a limit test with water-saturated concentration of the test substance had no adverse effect within the test period of 96 hours. The BHT concentration measured in the test solution after 24 hours of exposure was 0.57 mg/l (Bayer AG 1994).

In a test according to the European protocol EEC C.2 (equivalent to OECD guideline 202, part 1) the acute toxicity of BHT in invertebrates was determined. *Daphnia magna* were exposed in a limit test with water-saturated concentration of the test substance. No toxic effects were observed during incubation and the 48h EC₀ based on the measured BHT concentration (geometric mean of TS concentrations measured at the start and after 48 h of exposure) was reported to be ≥ 0.17 mg/l (Bayer AG 1994).

A further acute test with *Daphnia pulex* is also available for BHT (Passino, Smith, 1987). This test was performed according to ASTM method. Acetone was used as solvent with a concentration of 0.5 ml/l. A 48h EC₅₀ of 1.44 mg/l related to nominal concentration was found. However the concentration of acetone in this study was higher than proposed by OECD, US-EPA and EU (0.1 ml/l). As there are other valid tests performed without solvent available that are regarded to be more relevant, the test is not used for the effect assessment of BHT.

In a long-term test *Daphnia magna* were exposed to three BHT concentrations (0.1 mg/l; 0.316 mg/l and 1.0 mg/l; nominal) for a test duration of 21 days. The test was performed according to OECD guideline 202 (part 2) under semistatic conditions. The NOEC (endpoint: reproduction) based on measured test substance concentrations (geometric mean of TS concentrations measured at the start and after 48 h and 72 h of exposure at water exchange) was reported to be 0.07 mg/l (Bayer AG 1994).

The acute toxicity of BHT to algae was determined in a test according to the European protocol EEC C.3 (equivalent to OECD guideline 201). The saturated BHT-water concentration was tested in a limit test with the algae *Scenedesmus subspicatus*. Only a slightly lower growth rate was observed after 72 h of incubation (8 % inhibition). Based on measured concentrations (geometric mean of TS concentrations measured at the start and after 72 h of exposure) an E_rC₃ of 0.4 mg/l is derived (Bayer AG 1994). This value can be used as a NOEC.

In a test with activated sludge according to Directive 88/302/EEC, Part C (respiration inhibition test) a 3h $EC_0 = 1000$ mg/l was determined (Bayer AG 2000).

For the protozoan species *Tetrahymena pyriformis* a 24h EC_{50} of 1.7 mg/l was found in a cell multiplication inhibition test (Yoshioka et al., 1985).

The lowest available long-term NOEC of 0.07 mg/l, found in a 21 days reproduction test with *Daphnia magna* is used for the derivation of the $PNEC_{aqua}$. The application of an assessment factor of 50 is justified, as results from two long-term tests (daphnia and algae) are available, resulting in a $PNEC_{aqua}$ of 0.0014 mg/l.

4.2 Terrestrial Effects

No reliable data available.

4.3 Other Environmental Effects

In a feeding study with "White Leghorn" chicken (*Gallus domesticus*) the influence of BHT on aflatoxin toxicity has been studied. 0.1 % BHT in the diet (8x the standard BHT concentration of 0.013 % in feed) had no significant inhibiting effect on weight gain, whereas 0.4% (30 x the standard BHT concentration in feed) resulted in a temporary depression (36 %) of weight gain from day 1 until 3 weeks; from week 3 to week 6 the weight gain was normal (98 % of control) and at the end of BHT treatment (6 weeks) the total weight gain amounted to 78 % of control. Both BHT dosages improved body weight gain and feed efficiency of chicken treated with 3000 ppb aflatoxin in feed (Larsen et al., 1985).

要 約

試験委託者：環境省

表 題：プロピレンテトラマーの藻類 (*Selenastrum capricornutum*) に対する
生長阻害試験

試験番号：A020372-1

試験方法：

- 1) 適用ガイドライン： OECD 化学品テストガイドライン No. 201 「藻類生長阻害試験」(1984年)
- 2) 暴露方式： 止水式 (密閉系), 連続振とう培養 (100rpm)
- 3) 供試生物： *Selenastrum capricornutum* (株名：ATCC22662)
(現在 *Pseudokirchneriella subcapitata*と学名が変更されている。)
- 4) 暴露期間： 72時間
- 5) 試験濃度： 対照区, 助剤対照区, 0.0400 mg/L (試験液調製可能最高濃度での
(設定値) 限度試験)
助剤濃度一定：100 μ L/L (ジメチルホルムアミド使用)
- 6) 試験液量： 100 mL (OECD培地) / 容器
- 7) 連 数： 3 容器 / 試験区
- 8) 初期細胞濃度： 前培養した藻類 1×10^4 cells/mL
- 9) 試験温度： 23 ± 2 $^{\circ}$ C
- 10) 照 明： 4000 lux ($\pm 20\%$ の変動内, フラスコ液面付近) で連続照明
- 11) 分 析 法： ガスクロマトグラフィー質量分析 (GC/MS)

試験結果：

- 1) 試験液および試験培養液中の被験物質濃度

被験物質濃度分析の結果, 測定値の設定値に対する割合は, 暴露開始時の試験液において134%, 暴露終了時の試験培養液において検出限界以下であった。水中からの50%揮散速度は約6時間であることから, 濃度減少の主な原因は揮散と考えられた。阻害濃度の算出には開始時の測定値を用いた。

2) 生長曲線下面積の比較による阻害濃度

50%生長阻害濃度 E_bC_{50} (0-72h) : >0.0534 mg/L (95%信頼区間:算出不可)

最大無作用濃度 $NOEC_b$ (0-72h) : >0.0534 mg/L

3) 生長速度の比較による阻害濃度

50%生長阻害濃度 E_rC_{50} (24-48h) : >0.0534 mg/L (95%信頼区間:算出不可)

最大無作用濃度 $NOEC_r$ (24-48h) : >0.0534 mg/L

50%生長阻害濃度 E_rC_{50} (24-72h) : >0.0534 mg/L (95%信頼区間:算出不可)


最大無作用濃度 $NOEC_r$ (24-72h) : >0.0534 mg/L

4) 藻類の形態観察

暴露終了時の顕微鏡下での細胞形態観察の結果, 0.0400 mg/Lの濃度区では細胞形態の変化(収縮, 膨張, 破裂等)や細胞凝集は認められず, また, 対照区および助剤対照区との相違もなかった。

プロピレンテトラマー (CAS.6842-15-5)

① 生長曲線



② 毒性値

48hErC50 (実測値に基づく) > 0.0032mg/L
48hNOECr (実測値に基づく) = 0.0032mg/L