

Fig. 1 Growth inhibition of CHL/IU cells treated with 2-tert-butylphenol




Fig. 2 Growth inhibition of CHL/IU cells treated with 2-tert-butylphenol

量域にあるものと判断された。

8. 実験群の設定

細胞増殖抑制試験の結果から、染色体異常試験における被験物質の用量は、50%細胞増殖抑制用量の前後が含まれ、かつ3用量以上のデータが得られることを考慮して、S9 mix非存在下では40, 60, 80, 100, 110および120 µg/mL, S9 mix存在下では1.25, 2.5, 5, 7.5, 10および20 µg/mLのそれぞれ6用量を設定した。対照として、溶媒対照群と陽性対照群を設けた。

陽性対照として、短時間処理法S9 mix存在下では3,4-benzo[a]pyrene(B[a]P, Sigma Chemical Co.)を10 µg/mL, S9 mix非存在下では1-methyl-3-nitro-1-nitrosoguanidine(MNNG, Aldrich Chemical Co.)を2.5 µg/mLの用量で用いた。陽性対照物質の溶媒には、いずれもDMSO(和光純薬工業株)を使用した。

9. 染色体標本の作製

培養終了2時間前にコルセミド(Gibco Laboratories)を最終濃度として0.2 µg/mLとなるように添加した。トリプシン処理で細胞を剥離し、遠心分離により細胞を回収した。75 mM塩化カリウム水溶液で低張処理後、用時調製した冷却メタノール・酢酸(3:1)混合液で細胞を固定した。空気乾燥法で染色体標本作製した後、1.4 vol%ギムザ液で約15分間染色した。

10. 染色体の観察

各ディッシュあたり100個、すなわち、1用量当たり2

ディッシュ、200個の分裂中期像を、総合倍率600倍の顕微鏡下で観察した。標本は全てコード化し、盲検法で観察を行った。染色体の分析は、日本環境変異原学会・哺乳動物試験分科会(MMS)による分類法¹⁾に基づいて行い、染色体型あるいは染色体型の切断、交換などの構造異常と倍数性細胞(Polyploid)の有無について観察した。

11. 記録と判定

観察した細胞数、構造異常の種類と数および倍数性細胞の数について集計し、記録した。

染色体構造異常細胞および倍数性細胞の出現頻度について、多試料 χ^2 検定を行い有意差(有意水準5%以下)が認められた場合は、フィッシャーの直接確率法を用いて溶媒対照群と各用量群との間の有意差検定(有意水準は多重性を考慮して、5%または1%を処理群の数で割ったものを用いた)を行った。

その結果、溶媒対照群と比較して、被験物質による染色体異常細胞の出現頻度が2用量以上で有意に増加し、かつ用量依存性あるいは再現性が認められた場合、陽性と判定した。

結果および考察

短時間処理法による結果をTable 1に示す。S9 mix非存在下では、染色体の構造異常および倍数性細胞の誘発作用は認められなかった。一方、S9 mix存在下においては、用量依存的な染色体の構造異常を有する細胞の増

加が認められ、7.5および10 $\mu\text{g}/\text{mL}$ での増加(出現頻度5.5および11.0%)は溶媒対照群と比較して統計学的に有意なものであった。また、10 $\mu\text{g}/\text{mL}$ で倍数性細胞の有意な増加(出現頻度5.0%)も認められた。S9 mix非存在下の110 $\mu\text{g}/\text{mL}$ 以上およびS9 mix存在下の20 $\mu\text{g}/\text{mL}$ においては、被験物質の細胞に対する毒性のため、観察可能な分裂中期像が認められなかった。

以上の成績から、2-tert-ブチルフェノールは、その代謝物に染色体構造異常を誘発する作用があると考えられた。したがって、本実験条件下では、2-tert-ブチルフェノールのCHL/IU細胞に対する染色体異常誘発性は陽性と判定した。陽性結果が得られたため、 D_{20} 値²⁾(分裂中期像の20%に異常を誘発させる被験物質の推定用量)を算出したところ、本被験物質の D_{20} 値は、構造異常に関して0.022 mg/mL、数値異常に関しては、0.043 mg/mLであった。本試験結果は、CHL/IU細胞において、染色体異常を有する細胞の出現頻度が5%以上10%未満を疑陽性、10%以上を陽性とする石館らの判定基準³⁾からみても、陽性と判断されるものであった。

2-tert-ブチルフェノールの類縁化合物については、4-tert-ブチルフェノールでは、CHL/IU細胞を用いた染色体異常試験において連続処理法24時間および48時間処理で倍数性細胞の誘発作用が、また、短時間処理法S9 mix存在下では構造異常および倍数性細胞の誘発作用が報告され⁴⁾、2,4-ジ-tert-ブチルフェノール⁵⁾および6-tert-ブチル-m-クレゾール⁶⁾では同様の試験において、短時間処理法S9 mix存在下で構造異常細胞の誘発作用が報告されている。また、2-sec-ブチルフェノールでもCHL/IU細胞を用いた染色体異常試験において連続処理法48時間処理および短時間処理法S9 mix存在下で構造異常細胞の誘発作用が認められ⁷⁾、さらに、4-sec-ブチルフェノールでは同様の試験において、連続処理法48時間処理および短時間処理法S9 mix存在下での構造異常細胞の誘発に対し、疑陽性の判定が示されている⁸⁾。一方、2,6-tert-ブチルフェノールおよび4-tert-ブチルフェノールでは、ラット肝細胞を用いた染色体異常試験において陰性⁹⁾、また、2,2'-メチレンビス(6-tert-ブチル-p-クレゾール)¹⁰⁾および4,4'-チオビス(6-tert-ブチル-m-クレゾール)¹¹⁾ではCHL/IU細胞を用いた染色体異常試験において陰性と報告されている。

このように、類縁化合物にも染色体異常誘発性を示す物質が多いことから、2-tert-ブチルフェノールの染色体異常誘発性はこれらに共通した化学構造との関連性が考えられる。

文献

- 1) 日本環境変異原学会・哺乳動物試験分科会編, “化学物質による染色体異常アトラス,” 朝倉書店, 東京, 1988, pp.16-37.
- 2) 厚生省生活衛生局企画課生活化学安全対策室 監修, “化審法毒性試験法の解説 改訂版,” 化学工業日報社, 東京, 1992, pp.51-52.

- 3) 石館基監修, “改定増補 染色体異常試験 データ集,” エル・アイ・シー, 東京, 1987, p.19.
- 4) 田中憲穂, 山影康次, 中川ゆずき, 日下部博一, 橋本恵子, 水谷正寛, 古畑紀久子, 化学物質毒性試験報告, 4, 301(1996).
- 5) 野田篤, 昆尚美, 化学物質毒性試験報告, 8, 392(2001).
- 6) 野田篤, 昆尚美, 化学物質毒性試験報告, 7, 200(1999).
- 7) 野田篤, 昆尚美, 化学物質毒性試験報告, 7, 232(1999).
- 8) 田中憲穂, 山影康次, 佐々木澄志, 若栗忍, 日下部博一, 橋本恵子, 化学物質毒性試験報告, 2, 347(1995).
- 9) B. J. Dean, T. M. Brooks, G. Hodson-Walker, D. H. Hutson, *Mutat. Res.*, **153**, 57(1985).
- 10) 田中憲穂, 山影康次, 若栗忍, 日下部博一, 橋本恵子, 長尾哲二, 太田亮, 化学物質毒性試験報告, 4, 427(1996).
- 11) 田中憲穂, 山影康次, 若栗忍, 中川ゆずき, 日下部博一, 橋本恵子, 長尾哲二, 太田亮, 化学物質毒性試験報告, 4, 247(1996).

連絡先

試験責任者: 野田 篤
 試験担当者: 野田 篤, 昆 尚美
 (財)畜産生物科学安全研究所
 〒229-1132 神奈川県相模原市橋本台3-7-11
 Tel 042-762-2775 Fax 042-762-7979

Correspondence

Authors: Atsushi Noda (Study director)
 Naomi Kon
 Research Institute for Animal Science in
 Biochemistry and Toxicology
 3-7-11 Hashimotodai, Sagami-hara-shi, kanagawa,
 229-1132 Japan
 Tel +81-42-762-2775 Fax +81-42-762-7979

Table 1 Chromosome analysis of Chinese hamster cells (CHL/IU) treated with 2-tert-butylphenol with and without S9 mix

Group	Dose ($\mu\text{g}/\text{mL}$)	S9 mix	Time of exposure (hr)	No. of cells analysed	No. of cells with structural aberrations						gap (%)	No. of cells with numerical aberrations			Cell Growth rate(%)
					ctb	cte	csb	cse	oth	total(%) ²		Polyploid	Polyploid ³	total(%) ³	
Solvent ¹	0	-	6-(18)	200	2	1	0	0	0	3(1.5)	0(0)	0	0	0(0)	100.0
2TBP	40	-	6-(18)	200	0	0	0	0	0	0(0)	1(0.5)	0	0	0(0)	105.5
	60	-	6-(18)	200	0	1	0	1	0	2(1.0)	0(0)	0	0	0(0)	97.0
	80	-	6-(18)	200	0	2	0	2	0	4(2.0)	0(0)	1	0	1(0.5)	95.0
	100	-	6-(18)	200	3	6	0	0	0	8(4.0)	0(0)	0	0	0(0)	50.5
	110	-	6-(18)	Toxic	—	—	—	—	—	—	—	—	—	—	13.5
	120	-	6-(18)	Toxic	—	—	—	—	—	—	—	—	—	—	4.0
MNNG	2.5	-	6-(18)	200	64	188	3	1	0	190(95.0)**	3(1.5)	0	0	0(0)	—
Solvent	0	+	6-(18)	200	0	0	0	1	0	1(0.5)	0(0)	1	0	1(0.5)	100.0
2TBP	1.25	+	6-(18)	200	1	1	0	0	0	2(1.0)	0(0)	1	0	1(0.5)	90.0
	2.5	+	6-(18)	200	1	2	0	2	0	4(2.0)	1(0.5)	0	0	0(0)	81.5
	5	+	6-(18)	200	1	2	0	0	0	2(1.0)	0(0)	0	0	0(0)	70.0
	7.5	+	6-(18)	200	2	10	0	0	0	11(5.5)*	1(0.5)	7	0	7(3.5)	43.0
	10	+	6-(18)	200	8	18	0	0	0	22(11.0)**	0(0)	7	3	10(5.0)*	26.5
	20	+	6-(18)	Toxic	—	—	—	—	—	—	—	—	—	—	16.5
BP	10	+	6-(18)	200	13	68	1	0	0	75(37.5)**	0(0)	0	0	0(0)	—

Abbreviations; gap: chromatid gap and chromosome gap, ctb: chromatid break, cte: chromatid exchange, csb: chromosome break, cse: chromosome exchange (dicentric and ring), oth: others, SA: structural aberration, NA: numerical aberration, 2TBP: 2-tert-butylphenol, MNNG: 1-methyl-3-nitro-1-nitrosoguanidine, BP: 3,4-benzo[a]pyrene

1) Dimethyl sulfoxide was used as solvent.

2) Multi-sample χ^2 test was done at $p < 0.05$ and then Fisher's exact test was done at $p < 0.05$ or $p < 0.01$.

3) endoreduplication

*: Significantly different from solvent group data at $p < 0.05$ by Fisher's exact test.

**: Significantly different from solvent group data at $p < 0.01$ by Fisher's exact test.

ORIGINAL ARTICLE

Elevated susceptibility of newborn as compared with young rats to 2-*tert*-butylphenol and 2,4-di-*tert*-butylphenol toxicity

Mutsuko Hirata-Koizumi¹, Masao Hamamura², Hiromi Furukawa², Naemi Fukuda³, Yoshihiko Ito³, Yumi Wako⁴, Kotaro Yamashita⁴, Mika Takahashi¹, Eiichi Kamata¹, Makoto Ema¹, and Ryuichi Hasegawa¹

¹National Institute of Health Sciences, Setagaya-ku, Tokyo; ²Panapharm Laboratories, Uto-shi, Kumamoto; ³Research Institute for Animal Science in Biochemistry and Toxicology, Sagami-hara-shi, Kanagawa; and ⁴Kashima Laboratory, Mitsubishi Chemical Safety Institute Ltd, Ibaraki, Japan

ABSTRACT In order to determine the susceptibility of newborn rats to 2-*tert*-butylphenol (2TBP) and 2,4-di-*tert*-butylphenol (DTBP) toxicity, studies were conducted with oral administration from postnatal days (PND) 4 to 21 and the findings were compared with results for young rats exposed from 5 or 6 weeks of age for 28 days. In the newborn rats, specific effects on physical and sexual development and reflex ontogeny were not observed. While there were no clear differences in toxicological profiles between newborn and young rats, the no-observed-adverse-effect levels (NOAELs) differed markedly. For 2TBP, clinical signs such as ataxic gait, decrease in locomotor activity and effects on liver, such as increase in organ weight, were observed and the NOAELs were concluded to be 20 and 100 mg/kg/day in newborn and young rats, respectively. Based on hepatic and renal toxicity (histopathological changes and increase in organ weight with blood biochemical changes), the respective NOAELs for DTBP were concluded to be 5 and 20 mg/kg/day. Therefore, the susceptibility of newborn rats to 2TBP and DTBP was found to be 4–5 times higher than that of young rats.

Key Words: 2, 4-di-*tert*-butylphenol, 2-*tert*-butylphenol, susceptibility of newborn rats

INTRODUCTION

Protection of humans against disease and injury caused by chemicals in the environment is the ultimate goal of risk assessment and risk management (Landrigan *et al.* 2004). However, the focus has long been solely on adult exposure and toxicity and the fetus via maternal transfer, with little consideration given to early childhood. In the past decade, stimulated especially by the 1993 US National Research Council (NRC) report *Pesticides in the Diets of Infants and Children* (NAS 1993), recognition that special consideration is required for children in risk assessment has grown. The NRC report noted that 'children are not little adults', because of their unique patterns of exposures to environmental hazards and their particular vulnerability.

For the susceptibility of children to environmental chemicals, the early postnatal period (the suckling period) is of particular note. During this period, the infant could be exposed to various chemicals not only through mothers' milk, but also directly, by having

chemical-contaminated baby food, mouthing toys or household materials, and so on; however, current risk assessment gives no consideration to toxic effects resulting from direct exposure to chemicals. An approach that adequately takes into account the susceptibility of infancy is urgently required. However, because there is no standard testing protocol intended for direct exposure of preweaning animals (newborn animals) to chemicals, and toxicity studies using newborn animals are complicated by practical difficulties regarding grouping, direct dosing, and general and functional observation, there is only limited information on susceptibility of the newborn at the present.

We therefore have established a new protocol for repeated dose toxicity studies using newborn rats (newborn rat studies) (Koizumi *et al.* 2001) for systematic application. Results have been compared with those of 28-day repeated dose toxicity studies using young rats (young rat studies) to provide a basis of analyzing susceptibility. Since young rat studies are routinely conducted as one of a battery of minimum toxicity tests and data are stored for many chemicals, comparative analyzes should provide important information for considering effects of direct exposure to chemicals during the suckling period.

We have already reported analytical results for eight chemicals (4-nitrophenol, 2,4-dinitrophenol, 3-aminophenol, 3-methylphenol, 1,3-dibromopropane, 1,1,2,2-tetrabromoethane, 2,4,6-trinitrophenol, and tetrabromobisphenol A) (Koizumi *et al.* 2001, 2002, 2003; Fukuda *et al.* 2004; Takahashi *et al.* 2004; Hirata-Koizumi *et al.* 2005). The susceptibility of newborn rats to the toxicity of the first four agents was four times higher than that of their young counterparts at a maximum. For 1,3-dibromopropane and 1,1,2,2-tetrabromoethane, while the doses causing clear toxicity were lower in newborn rats, doses at which toxic signs began to appear were paradoxically higher in the newborn case. These six chemicals had no impact on development in the newborn period and showed similar toxicity profiles in both age groups. For the other two chemicals, there were marked differences in toxicity profile between the newborn and young rats. Especially, in the case of tetrabromobisphenol A, a specific rather than enhanced renal toxicity was observed in newborn case.

In the present investigation, two *tert*-butylphenols, 2-*tert*-butylphenol (2TBP), and 2,4-di-*tert*-butylphenol (DTBP), were chosen for comparative toxicity analysis. 2TBP has been used in the production of agricultural chemicals, aroma chemicals, and resins (New Chemical Index 2001), and DTBP in the production of antioxidants and ultraviolet absorbers (Chemical Products' Handbook 2004). For either chemical, there is no available toxicity information on human. Regarding toxicity to experimental animals, results from young rat studies of both chemicals are available in

Correspondence: Mutsuko Hirata-Koizumi, BSc, Division of Medicinal Safety Science, National Institute of Health Sciences, 1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan, Email: mkoizumi@nihs.go.jp

Received June 17, 2005; revised and accepted September 2, 2005.

Toxicity Testing Reports of Environmental Chemicals of the Japanese government (MHLW 2001a, 2001b), but no other data have been reported regarding repeated dose toxicity. Since the young rats were only evaluated for toxicity profiles and no-observed-effect levels, we re-evaluated the results for a more practical evaluation index, the no-observed-adverse-effect level (NOAEL), which could serve as the basis for determining tolerable daily intake (TDI) or acceptable daily intake (ADI) for risk assessment, and conducted comparative analyzes with newborn rats.

MATERIALS

2-*tert*-Butylphenol (2TBP, CAS no. 88-18-6, purity: 99.97%) and 2,4-di-*tert*-butylphenol (DTBP, CAS no. 96-76-4, purity: 99.67%), obtained from Dainippon Ink and Chemicals, Incorporated (Tokyo, Japan), were dissolved in olive oil and corn oil, respectively. The test solutions were prepared once a week as stability for eight days had been confirmed. All other reagents used in this study were specific purity grade.

METHODS

All studies were performed under Good Laboratory Practice conditions and in accordance with 'Guidance for Animal Care and Use' of Panapharm Laboratories Co., Ltd, Research Institute for Animal Science in Biochemistry and Toxicology, or Mitsubishi Chemical Safety Institute Ltd.

Animals

In the newborn rat studies of 2TBP and DTBP, pregnant SPF Sprague-Dawley rats [Crj:CD(SD)IGS] were purchased at gestation days 13–15 from Charles River Japan Inc. (Yokohama, Japan), and allowed to deliver spontaneously. All newborn were separated from dams at postnatal day (PND) 3 (the date of birth was defined as PND 0), and pooled according to sex. At the same time, 12 foster mothers were selected among dams, based on the nursing condition. Each foster mother suckled four male and four female newborn, assigned to each of the four dose groups, including the controls, up to weaning on PND 21 (termination of dosing). After weaning, the animals of the recovery-maintenance group (see Study Design) were individually maintained for nine weeks.

In the young rat studies, 4–5 week-old males and females of the same strain were obtained from the same supplier as for the newborn rat studies, and used at ages of 5–6 weeks after acclimation.

All animals were maintained in an environmentally controlled room at 20–26°C with a relative humidity of 40–70%, a ventilation rate of more than ten times per hour, and a 12:12 h light/dark cycle. They were allowed free access to a basal diet (MF: Oriental Yeast Co. Ltd, Tokyo, Japan, or LABO MR Stock: Nihon Nosan Kogyo Inc., Yokohama, Japan) and water (sterile tap water or well water treated with sodium hypochlorite) throughout.

Study design

1. 18-day repeated dose toxicity study in newborn rats (newborn rat study)

Newborn rats (12/sex/dose) were administered the test substances by gastric intubation on PNDs 4–21. On PND 22, six males and six females in each treated group were sacrificed for autopsy (the scheduled-sacrifice group). The remaining animals in all groups (6 rats/sex/dose) were maintained for nine weeks without chemical treatment and then sacrificed at 12 weeks of age (the recovery-maintenance group).

Based on the results of dose-finding studies conducted prior to the main study, the dose, which would show clear toxicity, was selected as the top dose, that without potentially toxic effects as the lowest dose, and the medium dose was set between them. In the dose-finding study for 2TBP (oral administration from PNDs 4–21), some clinical signs and suppressed body weight gain were observed at 200 mg/kg and an increase in relative liver weight at 60 mg/kg and more. For DTBP (oral administration from PNDs 4–17), all of the four males and four females died at 500 mg/kg, and the death of one of the four males, an increase in serum total cholesterol and phospholipid, and increase in relative liver weight were noted in the 100 mg/kg group. Therefore, the doses were set at 0, 20, 60, or 200 mg/kg/day for 2TBP and at 0, 5, 40, or 300 mg/kg/day for DTBP.

During the study, the rats' general condition was observed at least once a day (details of clinical signs noted in this study are described in 'Glossary of terms for toxicity testing' [NIHS 1994]). Body weight and food consumption (only the recovery-maintenance period) was examined once or more a week. As developmental parameters, fur appearance, incisor eruption, pinna detachment and eye opening were assessed for physical development, and testes descent or preputial separation and vaginal opening for sexual development (OECD 2004). In addition, reflex ontogeny, such as visual placing reflex, and surface and mid-air righting reflexes, were also examined (Adams 1986; Jensch & Brent 1988). Urinalysis (color, occult blood, pH, protein, glucose, ketone bodies, bilirubin, urobilinogen, sediment, specific gravity, and volume of the urine) was conducted in the last week of the recovery-maintenance period.

At PNDs 22 and 85, blood was collected from the abdominal aorta under ether anesthesia (for 2TBP) or from the postcaval vein under pentobarbital sodium anesthesia (for DTBP) after overnight starvation for the scheduled-sacrifice and recovery-maintenance groups, respectively. One portion was treated with EDTA-2K and examined for hematological parameters, such as the red blood cell count, hemoglobin, hematocrit, mean corpuscular volume, mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration, white blood cell count, platelet count, reticulocyte count and differential leukocyte count. In the recovery-maintenance group, part of the blood was treated with 3.8% sodium citrate, and blood clotting parameters such as prothrombin time (PT) and activated partial thromboplastin time (APTT) were examined. Serum from the remaining portions of blood for both the scheduled-sacrifice and recovery-maintenance groups were analyzed for blood biochemistry (total protein, albumin, albumin-globulin ratio [A/G ratio], glucose, total cholesterol, triglycerides, phospholipid, total bilirubin, urea nitrogen [BUN], creatinine, glutamate oxaloacetate transaminase, glutamate pyruvate transaminase, alkaline phosphatase, γ -glutamyl transpeptidase [γ -GTP], calcium, inorganic phosphorus, sodium, potassium, and chlorine). Following collection of blood, all animals were sacrificed by exsanguination, and all organs and tissues were macroscopically examined. Then, the brain, pituitary gland, thymus, thyroids, heart, lungs, liver, spleen, kidneys, adrenals, testes, epididymides, and ovaries were removed and weighed. Histopathological examination was conducted for the control and the highest dose groups. The above-listed organs were fixed in 10% buffered formalin-phosphate (following Bouin's fixation for testes and epididymides), and paraffin sections were routinely prepared and stained with Hematoxylin-Eosin for microscopy. For other groups, organs with macroscopically abnormal findings or in which chemical-related effects were evident on microscopic examination for the highest dose group, were similarly investigated.

2. 28-day repeated dose toxicity study in young rats (young rat study)

Five to six week old rats were given the test substances by gastric intubation daily for 28 days and sacrificed following the last treatment (the scheduled-sacrifice group). Recovery groups were maintained for two weeks without chemical treatment and sacrificed at 11 or 12 weeks of age. The number of animals was six for each sex/dose for both scheduled-sacrificed and recovery cases.

The doses were selected in the same way as the newborn rat studies. In the 12-day dose-finding study for 2TBP, ataxic gait was observed at 300 mg/kg and more, and increase in relative liver and kidney weight at 500 mg/kg. For DTBP, with 14-day administration, the death of one of the four females, various changes in some blood biochemical parameters, increase in relative liver weights and light gray macules on kidneys were found at 500 mg/kg. Increase in serum phospholipid and relative liver weights were also demonstrated in the 100 mg/kg group. Based on the results, the doses were determined at 0, 4, 20, 100, or 500 mg/kg/day for 2TBP and at 0, 5, 20, 75, or 300 mg/kg/day for DTBP. Recovery groups were set at 0, 100, 500 mg/kg/day for 2TBP and 0, 300 mg/kg/day for DTBP.

During the study, rats were examined for general condition, body weight, food consumption, urinalysis, hematology and blood biochemistry, necropsy findings, organ weights, and histopathological findings in compliance with the Test Guideline in the Japanese Chemical Control Act (Official Name: Law Concerning the Examination and Regulation of Manufacture, etc. of Chemical Substances).

Statistical analysis

Data for body weights, food consumption, urinalysis findings (except for the results of qualitative analysis), hematological, blood biochemical findings (except for differential leukocyte count), and organ weights were analyzed by the Bartlett's test (Bartlett 1937) for homogeneity of distribution. When homogeneity was recognized, Dunnett's test (Dunnett 1964) was conducted for comparison between control and individual treatment groups ($P < 0.01$ or 0.05). If not homogeneous or for qualitative urinalysis data and differential leukocyte count, the data were analyzed using Steel's multiple comparison tests (Steel 1959), or tests of the Dunnett type (Hollander & Wolfe 1973) ($P < 0.01$ or 0.05). For reflex ontogeny, and physical and sexual development parameters in the newborn rat studies, the χ^2 -test (Fisher 1922) was conducted ($P < 0.01$ or 0.05).

RESULTS

2-tert-butylphenol (2TBP)

Newborn rat study

Various clinical signs such as decrease in locomotor activity, ataxic gait, deep respiration, and muscle weakness were observed throughout the dosing period in the 200 mg/kg group, as shown in Table 1. With 60 mg/kg, transient decrease in locomotor activity was noted on the first dosing day limited to only one of 12 males. Body weights were lowered by 8–17% from dosing day 7 through to the end of the dosing period in males and to recovery-maintenance day 14 in females given 200 mg/kg. At the scheduled sacrifice, there were no hematological changes at any dose, but blood biochemical examination of the 200 mg/kg group showed increases in γ -GTP in both sexes and total protein in males. In addition, significant increase in relative liver weights was noted in 9% of the females in the 60 mg/kg group and in 21–23% of both males and females in the 200 mg/kg group. On histopathological examination, slight hypertrophy of centrilobular hepatocytes was found in one female of the 60 mg/kg group, and in four males and three females from the 200 mg/kg group. During the recovery-maintenance period, no clinical signs were observed and the lowered body weights showed a tendency for recovery. In parameters for physical and sexual development and reflex ontogeny, no definitive changes were detected. At the end of the recovery-maintenance period, no chemical-related changes, also in urinalysis data, were found in any dose group.

The results of the newborn rat study of 2TBP are summarized in Table 2. Since clinical signs and histopathological changes in the liver were observed in the 60 mg/kg group, the NOAEL was concluded to be 20 mg/kg/day.

Young rat study

Ataxic gait were observed sporadically during the dosing period in nine males and 12 females, and decrease in locomotor activity in two females from the 500 mg/kg group. During the dosing period, there were no changes in body weight, food consumption, and urinalysis data. At the scheduled sacrifice, hematological and blood biochemical examination also showed no changes. Eighteen to 19% increases were found in relative liver weights of both sexes receiving 500 mg/kg, but no histopathological changes in liver were observed at any dose. No chemical-related changes were noted during and at the end of the recovery period.

Table 1 Clinical signs observed during the dosing period in the newborn rat study of 2-tert-butylphenol

	Dose (mg/kg/day)			
	0	20	60	200
No. animals (Male/Female)	12/12	12/12	12/12	12/12
No. animals with clinical signs				
Decrease in locomotor activity	0/0	0/0	1†/0	12/12
Ataxic gait	0/0	0/0	0/0	4/6
Deep respiration	0/0	0/0	0/0	12/12
Tremors	0/0	0/0	0/0	2/4
Muscle weakness	0/0	0/0	0/0	12/12
Emaciation	0/0	0/0	0/0	2/2
Pale skin	0/0	0/0	0/0	4/2

†Observed only on the first dosing day.

Table 2 Summary of the results of the newborn and young rat study of 2-*tert*-butylphenol

Newborn rat study				
Dose (mg/kg/day)	20	60	200	
Clinical signs	–	M: Decrease in locomotor activity	Various†	
Body weight changes	–	–	8–17%↓	
Blood biochemical changes	–	–	GTP↑, M: TP↑	
Changes in relative organ weights	–	F: Liver 9%↑	Liver 21–23%↑	
Histopathological findings in liver				
– Slight centrilobular hypertrophy of hepatocytes	–	M: 0/6, F: 1/6	M: 4/6, F: 3/6	
Young rat study				
Dose (mg/kg/day)	4	20	100	500
Clinical signs	–	–	–	Ataxic gait F: Decrease in locomotor activity
Body weight changes	–	–	–	–
Blood biochemical changes	–	–	–	–
Changes in relative organ weights	–	–	–	Liver 18–19%↑
Histopathological findings	n.d.	n.d.	n.d.	–

Statistically significant increases ($P < 0.05$) in body weights, blood biochemical parameters and relative organ weights are shown as ↑, while decreases are shown as ↓. Data on histopathological findings are given as no. of animals with the findings/no. of animals examined, according to sex. Changes observed only in males or females are shown as 'M' or 'F', respectively, while neither 'M' nor 'F' is mentioned in the case of changes noted in both sexes. No chemical-related changes were observed in developmental parameters (conducted only in newborn rat study), urinalysis (only in young rat study), and hematological parameters. †Decrease in locomotor activity, ataxic gait, deep respiration, tremors, muscle weakness, emaciation, and pale skin were observed, as shown in Table 1. GTP, γ -GTP; TP, total protein; –, no change; n.d., not determined.

A summary of the results of the young rat study of 2TBP is given in Table 2. The NOAEL was concluded to be 100 mg/kg/day, at which no changes were observed.

2,4-di-*tert*-butylphenol (DTBP)

Newborn rat study

Two males and one female of the 300 mg/kg group were found dead on dosing days 3, 4, and 7. In this group, decrease in locomotor activity (12 males and 12 females), bradypnea (10 males and 10 females), and hypothermia (one male) were observed from the first dosing day, but then the incidence decreased, with disappearance after dosing day 7. Body weights of the 300 mg/kg group were lowered by 15–25% in males and by 9–20% in females during the dosing period, compared with the control values. There were no definitive changes in parameters for physical development and reflex ontogeny in any dose group. At the scheduled sacrifice, blood biochemical examination showed an increase in total bilirubin and a decrease in the A/G ratio in both sexes, an increase in γ -GTP in males, and an increase in total protein and BUN in females of the 300 mg/kg group. In the 300 mg/kg group, there was a 39–51% increase in relative liver weights, a 37–41% increase in relative kidney weights in both sexes, and a 24% decrease in relative spleen weights in males. In the 40 mg/kg group, 14% increases in relative weight of liver were found in females. On histopathological examination, various changes were observed in livers and kidneys in the 300 mg/kg group, as shown in Table 3. Furthermore, periportal fatty degeneration of hepatocytes was evident in one female given 40 mg/kg, and basophilic tubules in kidneys in one animal of each sex receiving 40 mg/kg and one control group male. Regarding

parameters of sexual development, a slight delay in preputial separation was noted in the 300 mg/kg group (the incidences were 0/5, compared with 2/6 in the control group at PND 42 [recovery-maintenance day 21]; 0/5, 3/6 at PND 43; 2/5, 5/6 at PND 44; 2/5, 6/6 at PND 46; 4/5, 6/6 at PND 47; and 5/5, 6/6 at PND 48). During this observation period, body weights were lowered by approximately 10% in males given 300 mg/kg than control levels, which was not statistically significant. In the last week of the recovery-maintenance period, there were no chemical-related changes on urinalysis in any dose group. At the end of the recovery period, changes noted in the scheduled-sacrifice group were not observed except for histopathological changes in the kidneys, significant in the 300 mg/kg group (Table 3).

A summary of the results of the newborn rat study of DTBP is shown in Table 4. Since fatty degeneration of hepatocytes and increase in liver weight were demonstrated at 40 mg/kg, the NOAEL was concluded to be 5 mg/kg/day.

Young rat study

No chemical-related changes were found in general condition, body weight, and food consumption at any dose. On urinalysis at the fourth week of dosing, an increase in urine volume, and a decrease in specific gravity and osmotic pressure were noted in both sexes of the 300 mg/kg group. At the scheduled sacrifice, hematological examination showed a decrease in hemoglobin and hematocrit, an increase in segmented neutrophils in females, and prolongation of PT and APTT in males at 300 mg/kg. On blood biochemical examination, there was an increase in total bilirubin in males given 300 mg/kg, and an increase in total cholesterol and phospholipid in females given 75 mg/kg and above. For organ weights, there were

Table 3 Histopathological findings for the newborn rat study of 2,4-di-*tert*-butylphenol

Dose (mg/kg/day)	Grade	Scheduled-sacrifice group				Recovery-maintenance group†	
		0	5	40	300	0	300
No. of animals examined (Male/Female)		6/6	6/6	6/6	5/6	6/6	5/5
Liver							
– Fatty degeneration of periportal hepatocytes	+	0/0	0/0	0/1	0/0	0/0	0/0
	++	0/0	0/0	0/0	3/4	0/0	0/0
	+++	0/0	0/0	0/0	2/2	0/0	0/0
Kidneys							
– Basophilic tubules	+	1/0	n.d.	1/1	4/4	0/0	3/0
– Granular casts	+	0/0	n.d.	0/0	4/2	0/0	0/0
– Cystic dilatation of collecting tubules	+	0/0	n.d.	0/0	0/0	0/0	5/4
	++	0/0	n.d.	0/0	3/4	0/0	0/0
	+++	0/0	n.d.	0/0	2/2	0/0	0/0
– Cellular infiltration of neutrophils	+	0/0	n.d.	0/0	2/1	0/0	1/0
	++	0/0	n.d.	0/0	1/1	0/0	1/0
	+++	0/0	n.d.	0/0	1/1	0/0	0/0

†No histopathological examination was conducted at 5 and 40 mg/kg in the recovery-maintenance group. +, mild; ++, moderate; +++, marked; n.d., not determined.

increases in relative liver weights by 40–43% in both sexes given 300 mg/kg, and by 13% in females receiving 75 mg/kg. On histopathological examination, mild to marked changes in livers and kidneys were observed in both sexes from the 300 mg/kg group, as shown in Table 5. At the end of the recovery period, the increase in total cholesterol and phospholipid and renal histopathological changes observed in the scheduled-sacrifice group remained significant in the highest-dose group (Table 5).

The results of the young rat study are summarized in Table 4. Based on increase in the relative liver weights with some changes in blood biochemical parameters in females given 75 mg/kg, the NOAEL was concluded to be 20 mg/kg/day.

DISCUSSION

During development, many rapid and complex biological changes occur, which can have profound consequences on sensitivity to the effects of exogenous chemicals (Scheuplein *et al.* 2002). Although the neonatal body at birth is reasonably well prepared for the abrupt changes associated with parturition, and most functional systems possess a significant portion of their adult capacity (Dourson *et al.* 2002), it is known that the various functions remain immature in early postnatal period and that some organs and tissues, especially in the nervous, immune and reproductive systems, continue to develop after birth (NAS 1993). Therefore, it is important to evaluate toxic effects by exposure to chemicals during the early postnatal period as well as the fetal period for comprehensive risk assessment. However, economic issues and lack of human resources, arising from practical difficulties regarding protocols, have hindered routine implementation of toxicity studies using newborn animals. Our series of comparative analyzes on susceptibility of the newborn are therefore of particular importance for risk assessment.

In the present study on 2TBP and DTBP, there were no clear differences in toxicity profiles between the newborn and young rats in either case. For 2TBP, clinical signs such as a decrease in locomotor activity and ataxic gait, and effects on liver such as an increase in organ weight were observed. In the DTBP case, hepatic and renal toxicity (histopathological changes, increase in organ weight, etc.) were noted. As a characteristic effect of DTBP on male sexual development, slight delay in preputial separation was also observed in the newborn rat study. Preputial separation, an androgen-dependent process which is an early marker of puberty, represents a reliable non-invasive indicator of chemical-induced perturbation of male pubertal development in the rat (Gaytan *et al.* 1988). However, it is known that decreased body weights can result in non-specific delay in puberty (Ashby & Lefevre 2000). Since DTBP lowered body weights in the period of observation of preputial separation and there were no DTBP-related changes in weights or histopathology of the testes and epididymides, well known to be essentially androgen-dependent, no specific effect on male sexual development could be concluded in the present study. As for NOAELs of both chemicals, clear differences were observed between newborn and young rats, with values of 20 and 5 mg/kg/day in newborn rats, and 100 and 20 mg/kg/day in young rats for 2TBP and DTBP, respectively. Therefore, the susceptibility was four- to five-fold higher in newborn than in young rats.

Our previous analysis of 1,3-dibromopropane and 1,1,2,2-tetrabromoethane (Hirata-Koizumi *et al.* 2005) showed dose-response curves to be very different between newborn and young rats. The same was recently reported for the widely used organophosphorus insecticide, chlorpyrifos (Zheng *et al.* 2000), as well as pyrethroid insecticides (Shafer *et al.* 2005). These data showed the importance of estimating unequivocally toxic levels (UETLs), defined for our comparative toxicity analysis as equivalent toxic doses inducing clear toxicity, including death, clinical toxic signs,