

Phase I/II Study of gemtuzumab ozogamicin in combination chemotherapy for CD33+ refractory or relapsed AML: JALSG-AML206

Noriko Usui^{1,2}, Akihiro Takeshita², Chiaki Nakaseko², Nobuaki Dobashi², Hiroyuki Fujita², Hitoshi Kiyoi², Yukio Kobayashi², Yasushi Miyazaki², Shigeki Ohtake², Shuichi Miyawaki², Tomoki Naoe², Ryuzo Ohno², and Kazunori Ohnishi².

¹Dept of Clinical Oncology and Hematology, The Jikei University Daisan Hospital
²Japan Adult Leukemia Study Group,

Background

- Current treatment results in young adults with non-APL-AML are improving (%CR >70%, 5Y-OS 40-50%), however, quite a few patients relapse or refractory to initial therapies.

STUDY	N	CR %	ED %	OS % (3-5 yr)
CALGB	474	72	9	34
GAMLCG	535	74	11	39
HOVON	253	77	7	38
ALFA	345	82	9	38
JALSG	1,064	78	4	51

Tallman M,2007

Study	N	%CR	OS	DFS
AML87	188	79.8	30.2	28.5
AML89	232	78.5	35.1	43.7
AML92	566	77.2	33.5	31.6
AML95	480	80.7	44.3	28
AML97	789	78.7	40.8	35.5
AML201*	1057	78	51	
IDR	532	78.6	53.1	41.8
DNR	525	77.5	49.1	42.2

* Ohtake S, et al. Blood 108:566a #2000,2006

N. Usui, et al. JSMO 2010 001-012

Background

- Patients with relapsed or refractory AML have a poor outcome (%CR<50%, 5Y-OS<30%).
- Gemtuzumab ozogamicin(GO) is a humanized anti-CD33 MoAb conjugated with cytotoxic antitumor antibiotic, calicheamicin, that target CD33 Ag. More than 80% of AML cells have CD33 on their surface.
- GO induces CR+CRp in about 30% of relapsed/refractory AML
- Several studies tested the use of GO in combination chemotherapy, either at diagnosis or at time of relapse and most of them are more promising than monotherapy.

Name	No. of patients	Age (years)	AML status	Phase of GO	Combination	CR (%)	CRp (%)	OS (%)	DFS (%)
MELA	14	28-74	Relapsed AML	II	IDR Ara-C	71	85	20	14
MELA	17	23-78	Relapsed AML	I	IDR Ara-C	59	65	18	4
MELAR	22	18-78	Relapsed AML	I, II	IDR Ara-C GO	64	77	23	4
MELAR	11	16-67	Relapsed AML	I	Ara-C IDR GO	73	82	24	6
MELAR	17	21-68	Relapsed AML	I	Ara-C IDR GO	71	82	24	7
MELAR	44	18-69	Relapsed AML	I, II	Ara-C IDR GO	70	80	24	6

ORR 76%

N. Usui, et al. JSMO 2010 001-012

AIMS and OBJECTIVES

- AIMS**
 - In order to improve outcome of patients with AML, we investigate effective salvage therapies combining new agent (GO) with conventional chemotherapy (JALSG AML201 induction therapy regimen)
- OBJECTIVES**
 - Determine: Maximum tolerated dose (MTD)
 - DLTs:
 - >Grade 4(CTCAE ver 3.0) of FN, Bleeding, Nausea & Vomiting, Infusion reaction, liver toxicity (hyperbilirubinemia, hypertransaminasias)
 - >Grade 3(CTCAE ver 3.0) or more of non hematologic toxicity not related to progression of AML
 - Prolongation of bone marrow suppression (ANC<500 μ L, PLT<20,000 μ L) over 6 weeks not related to AML
 - Toxicity profile (NCI-CTCAE ver3)
 - Response Rate (International Working Group Criteria)

N. Usui, et al. JSMO 2010 001-012

Eligibility

Inclusion Criteria

- CD33+ de novo AML except APL; refractory to the first remission induction therapy or first relapse (>6 months from CR1)
- Age 20 – 64
- ECOG performance status of PS 0 or 1
- After 30 days or more from initial therapy and recovered to baseline from any toxicities of prior chemotherapy
- Adequate hepatic, cardiac, renal, pulmonary function
- Life expectancy \geq 2 months

Exclusion Criteria

- Previous MDS/MP
- Secondary AML
- CNS leukemia
- Received Transplantation
- Women who are pregnant or breastfeeding
- Received therapy with anti-CD33 MoAb
- Concurrent active malignant disease
- Uncontrolled infection
- HBV,HCV or HIV infection
- Treated with investigational drugs
- Previously received cumulative dose of >500 mg/m² of DNR (only for DAG arm)

N. Usui, et al. JSMO 2010 001-012


JALSG AML206-P1: Treatment Schedule

1. IDR arm (IAG)


IDR+Ara-C+GO Combination	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Ara-C 100mg/m ² /day c.i.v.	↓	↓	↓	↓	↓	↓	↓
IDR 12mg/m ² /day d.i.v.	↓	↓	↓	↓	↓	↓	↓
GO 3mg/m ² /day 2hr d.i.v.					↓		

2. DNR arm (DAG)

DNR+Ara-C+GO Combination	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Ara-C 100mg/m ² /day c.i.v.	↓	↓	↓	↓	↓	↓	↓
DNR 50mg/m ² /day d.i.v.	↓	↓	↓	↓	↓	↓	↓
GO 3mg/m ² /day 2hr d.i.v.				↓	↓	↓	↓



N. Usui, et al. JSMO 2010 001-012



Patients Demographics



	Overall	IAG	DAG
N	19	9	10
M:F	9:10	4:5	5:5
Age			
Median	59	61	58
Range	33 - 64	38 - 64	33 - 62
Relapsed	14	7	6
Refractory	5	2	4
AML Subtypes			
M0	1		1
M1	1	2	1
M2	7	3	5
M4	6	3	3
M5	1	1	
Karyotype			
CBF	2	1	1
CN	10	6	4
11q23 t(6;9)	1	1	
Complex	3	1	2
Others	2		2

N. Usui, et al. JSMO 2010 001-012

Patients Demographics-2

	Overall	IAG	DAG
Pre-treatment Status			
WBC (x10 ⁹ /μL)			
Median (Range)	3.0 (0.9-39.2)	3.7 (2.6-39.2)	2.1 (0.9-25.3)
Hb (g/dl)			
Median(Range)	10.9 (7.1-13.6)	12.5 (7.6-13.6)	10.6 (7.1-12.8)
PLT (x10 ⁹ /μL)			
Median (Range)	8.8 (2.6-44.3)	5.9 (2.6-20.7)	13.5 (3.4-44.3)
%Blasts			
Median (Range)	42.8 (7.9-96.8)	56.4 (17.3-88)	29.9 (7.9-96.8)
%CD33 Blasts			
Median (Range)	89.4 (39-100)	92.9 (62.8-100)	80.6 (39-96.9)

N. Usui, et al. JSMO 2010 001-012



IAG: Hematologic Toxicity

Hb	Level-1	Level-2	Level-3
Grade 0	0	0	1
Grade 1	1	1	1
Grade 2	2	1	1
Grade 3/4	0	0	0
Units of RBC-TF	4,4,12	4,6,2	8,16,4

Hb	Level-1	Level-2	Level-3
Days of nadir	Days 26,35,16	Days 18,27	Days 7,23
Days of recovery	Days none,39,35	Days 21,29	Days 30,27

Platelet	Level-1	Level-2	Level-3
Grade 3	3	2	3
Grade 4	0	1	0
Units of PLT-TF	90,130,100	130,130,50	70,220,70

PLT	Level-1	Level-2	Level-3
Days of nadir	Days 11,11,17	Days 13,13,17	Days 14,8,17
Days of recovery	Days 33,none,none	Days none,46,38	Days 28,90,34

N. Usui, et al. JSMO 2010 001-012