

---

1994; 120:272–277.

<sup>46</sup> Morimoto Y, Sugiura T, Tatebayashi S, et al. Reduction in incidence of methicillin-resistant *Staphylococcus aureus* (MRSA) after radical surgery for head and neck cancer. *Spec Care Dentist*. 2006 ;26:209-213.

<sup>47</sup> Nagashima G, Kikuchi T, Tsuyuzaki H, et al. To reduce catheter-related bloodstream infections: is the subclavian route better than the jugular route for central venous catheterization? *J Infect Chemother*. 2006;12:363-365.

<sup>48</sup> Kikuchi T, Nagashima G, Taguchi K, et al. Contaminated oral intubation equipment associated with an outbreak of carbapenem-resistant *Pseudomonas* in an intensive care unit. *J Hosp Infect*. 2007;65:54-57.

<sup>49</sup> Konishi T, Watanabe T, Morikane K, et al. Prospective surveillance effectively reduced rates of surgical site infection associated with elective colorectal surgery at a university hospital in Japan. *Infect Control Hosp Epidemiol*. 2006;27:526-8. Epub 2006 Apr 20.

<sup>50</sup> Yanai M, Uehara Y, Takahashi S. Surveillance of infection control procedures in dialysis units in Japan: a preliminary study. *Ther Apher Dial*. 2006;10:78-86.

<sup>51</sup> Takahashi H, Kramer MH, Yasui Y, et al. Nosocomial *Serratia marcescens* outbreak in Osaka, Japan, from 1999 to 2000. *Infect Control Hosp Epidemiol*. 2004 Feb;25(2):156-61.

<sup>52</sup> Morikane K, Nishioka M, Tanimura H, et al. Using surveillance data to direct infection control efforts to reduce surgical-site infections following clean abdominal operations in Japan. *Infect Control Hosp Epidemiol*. 2002 Jul;23(7):404-406.

<sup>53</sup> Lahaye D, Strauss P, Baleux C, et al. Cost-benefit analysis of hepatitis-B vaccination. *Lancet*. 1987 22:441-3.

<sup>54</sup> Asari S, Deguchi M, Tahara K, et al. Seroprevalence survey of measles, rubella, varicella, and mumps antibodies in health care workers and evaluation of a vaccination program in a tertiary care hospital in Japan. *Am J Infect Control*. 2003;31:157-162.