

- 2.5.1 カテーテルロックを実施する場合は、作り置きしたヘパリン生食は使用しない。
448, 449(ⅢA)
- 2.6 静脈炎発生時の対応
 - 2.6.1 静脈炎の徴候(発赤、腫脹、疼痛)がある場合は、カテーテルを抜去する。⁴⁵⁰
(ⅢA)
- 2.7 静脈炎予防薬剤の使用
 - 2.7.1 静脈炎予防のためのステロイド剤、ヘパリン、血管拡張剤は、使用しない方が
良い。^{451, 452}(ⅡB)
- 2.8 刺入部のドレッシング管理法
 - 2.8.1 カテーテル刺入部は滅菌のドレッシングで被覆し、カテーテル入れ換え時に交
換した方がよい。^{453, 454, 455}(ⅢB)
- 2.9 末梢のアミノ酸製剤
 - 2.9.1 アミノ酸加糖電解質製剤を投与する場合は側注を避けるなどの厳密な衛生管
理下においてのみ使用する。⁴⁵⁶(ⅢA)

²⁰⁸ Gallagher-Allred CR, Voss AC, Finn SC, et al: Malnutrition and clinical outcomes: the case for medical nutritional therapy. *J Am Diet Assoc* 1996;96:361-366.

²⁰⁹ Bistrain BR, Blackburn GL, Vitale J, et al: Prevalence of malnutrition in general medical patients. *JAMA* 1976;23:1567-1570.

²¹⁰ Kaminski MV, Jr: The case for nutrition support: Eliminating hospital-acquired malnutrition improves outcome and reduces costs. *Health Progress* 73:69-78,1992.

²¹¹ Messner RL, Stephens N, Wheeler WE, et al: Effect of admission nutritional status on length of hospital stay. *Gastroenterol Nurs* 13:202-205:1991.

²¹² Reilly JJ Jr, Hull SF, Albert N, et al: Economic impact of malnutrition: a model system for hospitalized patients. *JPEN* 12:371-376,1988.

²¹³ Martin CM, Doig GS, Heyland DK, et al: Multicentre, cluster-randomized clinical trial of algorithms for critical-care enteral and parenteral therapy (ACCEPT). *CMAJ* 2004;170:197-20.

²¹⁴ Moore EE, Jones TN: Benefits of immediate jejunostomy feeding after major abdominal trauma-a prospective, randomized study. *J Trauma* 1986;26:784-881.

²¹⁵ Windsor AC, Kanwar S, Li AG, et al: Compared with parenteral nutrition, enteral feeding attenuates the acute phase response and improves disease severity in acute pancreatitis. *Gut* 1998;42:431-435.

²¹⁶ Kudsk KA, Croce MA, Fabian TC, et al: Enteral versus parenteral feeding. Effects on septic morbidity after blunt and penetrating abdominal trauma. *Ann Surg* 1992;215:503-511.

²¹⁷ Lipman TO: Bacterial translocation and enteral nutrition in humans: an outsider looks in. *JPEN* 1995;19:156-165.

²¹⁸ Reynolds JV, Kanwar S, Welsh FK, et al: Does the route of feeding modify gut barrier function and clinical outcome in patients after major upper gastrointestinal surgery? *JPEN* 1997;21:196-201.

²¹⁹ Braunschweig CL, Levy P, Sheean PM, et al: Enteral compared with parenteral nutrition: a meta-analysis. *Am J Clin Nutr* 2001;74:534-542.

²²⁰ Lim ST, Choa RG, Lam KH, et al: Total parenteral nutrition versus gastrostomy in the preoperative preparation of patients with carcinoma of the oesophagus. *Br J Surg* 1981;68:69-72.