

表9-2-4 解熱剤と怒り出す

		発現数 (%)								①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)		
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^②		既使用 ^③				総計	未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用
第1病日	朝	18	0.8%	2	2.4%	2	2.4%	—	20	0.8%	0.188	3.14	2,310	83	0	2,393	0.8%	2.4%
	昼	15	0.7%	7	2.5%	5	2.5%	2	22	0.9%	0.045	3.54	2,092	200	81	2,373	1.5%	4.8%
	夜	37	2.3%	24	3.3%	15	3.3%	9	61	2.6%	0.353	1.48	1,628	449	274	2,351	3.7%	8.0%
第2病日	朝	6	0.4%	5	0.6%	4	5.3%	1	11	0.5%	0.001	1.64	1,516	75	699	2,290	4.1%	8.6%
	昼	17	1.2%	12	1.3%	2	1.6%	10	29	1.3%	0.935	1.10	1,386	124	769	2,279	5.3%	9.8%
	夜	23	1.8%	17	1.7%	6	5.5%	11	40	1.8%	0.029	0.94	1,259	110	881	2,250	7.0%	11.4%
第3病日	朝	4	0.3%	3	0.3%	0	0.0%	3	7	0.3%	0.982	0.94	1,231	5	974	2,210	7.3%	11.6%
	昼	3	0.2%	3	0.3%	0	0.0%	3	6	0.3%	0.940	1.24	1,218	9	976	2,203	7.5%	11.9%
	夜	3	0.2%	3	0.3%	0	0.0%	3	6	0.3%	0.950	1.22	1,208	7	982	2,197	7.8%	12.2%
第4病日	朝	2	0.2%	1	0.1%	0	0.0%	1	3	0.1%	0.917	0.61	1,204	1	986	2,191	7.9%	12.3%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%			1,199	3	986	2,188	7.9%	12.3%
	夜	0	0.0%	1	0.1%	0	0.0%	1	1	0.0%	0.452		1,195	4	989	2,188	7.9%	12.4%
第5病日	朝	0	0.0%	0	0.0%	—	—	0	0	0.0%			1,195	0	992	2,187	7.9%	12.4%
	昼	0	0.0%	1	0.1%	0	0.0%	1	1	0.0%	0.454		1,193	2	992	2,187	7.9%	12.4%
	夜	0	0.0%	1	0.1%	—	—	1	1	0.0%	0.209		1,193	0	993	2,186	7.9%	12.5%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0	0.0%			1,193	0	992	2,185	7.9%	12.5%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%			1,192	1	992	2,185	7.9%	12.5%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%			1,191	1	993	2,185	7.9%	12.5%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0	0.0%			1,191	0	994	2,185	7.9%	12.5%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%			1,190	1	994	2,185	7.9%	12.5%
	夜	0	0.0%	0	0.0%	—	—	0	0	0.0%			1,190	0	995	2,185	7.9%	12.5%
累積発現率 (参考)		128		80		34		46	208	8.7%							7.9%	12.5%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.39 (1.04 , 1.87)	0.029
第1～第2病日	1.40 (1.02 , 1.92)	0.038
第1～第4病日	1.36 (1.01 , 1.83)	0.042

表9-1-5 タミフルとけいれん

		発現数 (%)							①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)				
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^② 既使用 ^③		総計			未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用			
第1病日	朝	4	0.2%	1	0.3%	1	0.3%	—	5	0.2%	0.635	1.76	2,134	304	0	2,438	0.2%	0.3%	
	昼	11	0.6%	5	0.8%	2	0.7%	3	1.0%	16	0.7%	0.758	1.42	1,841	289	303	2,433	0.8%	1.2%
	夜	17	1.2%	11	1.1%	8	2.1%	3	0.5%	28	1.2%	0.078	0.97	1,450	380	587	2,417	1.9%	2.3%
第2病日	朝	4	0.4%	6	0.4%	4	0.8%	2	0.2%	10	0.4%	0.272	0.96	932	501	956	2,389	2.4%	2.7%
	昼	4	0.8%	8	0.4%	4	1.0%	4	0.3%	12	0.5%	0.143	0.56	523	405	1,451	2,379	3.1%	3.1%
	夜	1	0.3%	5	0.2%	2	1.2%	3	0.2%	6	0.3%	0.154	0.87	350	169	1,848	2,367	3.4%	3.4%
第3病日	朝	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0.008		269	80	2,012	2,361	3.4%	3.4%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%		213	56	2,092	2,361	3.4%	3.4%	
	夜	0	0.0%	1	0.0%	1	5.0%	0	0.0%	1	0.0%		193	20	2,148	2,361	3.4%	3.4%	
第4病日	朝	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			186	7	2,167	2,360	3.4%	3.4%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	180	6	2,174	2,360	3.4%	3.4%		
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	178	2	2,180	2,360	3.4%	3.4%		
第5病日	朝	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			177	1	2,182	2,360	3.4%	3.4%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	177	0	2,183	2,360	3.4%	3.4%		
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	177	0	2,183	2,360	3.4%	3.4%		
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			177	0	2,183	2,360	3.4%	3.4%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	177	0	2,183	2,360	3.4%	3.4%		
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	176	1	2,183	2,360	3.4%	3.4%		
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			176	0	2,184	2,360	3.4%	3.4%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	176	0	2,184	2,360	3.4%	3.4%		
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	176	0	2,184	2,360	3.4%	3.4%		
累積発生率 (参考)		41		37		22		15		78	3.2%						3.4%	3.4%	

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	0.99 (0.60 , 1.62)	0.957
第1～第2病日	0.98 (0.60 , 1.62)	0.941
第1～第4病日	0.99 (0.60 , 1.62)	0.957

表9-2-5 解熱剤とけいれん

		発現数 (%)								①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)		
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^②		既使用 ^③				総計	未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用
第1病日	朝	2	0.1%	3	3.6%	3	3.6%	—	5	0.2%	0.000	43.06	2,327	84	0	2,411	0.1%	3.6%
	昼	13	0.6%	3	1.1%	3	1.5%	0	0.0%	0.260	1.74	2,123	202	81	2,406	0.7%	4.6%	
	夜	11	0.7%	17	2.3%	12	2.7%	5	1.8%	0.003	3.56	1,658	452	280	2,390	1.4%	6.8%	
第2病日	朝	3	0.2%	7	0.9%	2	2.5%	5	0.7%	0.024	4.64	1,568	79	715	2,362	1.5%	7.6%	
	昼	8	0.6%	4	0.4%	2	1.5%	2	0.3%	0.216	0.78	1,435	130	787	2,352	2.1%	8.0%	
	夜	1	0.1%	5	0.5%	1	0.9%	4	0.4%	0.115	6.42	1,313	114	913	2,340	2.2%	8.5%	
第3病日	朝	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0.444		1,306	6	1,022	2,334	2.2%	8.5%	
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%		1,297	9	1,028	2,334	2.2%	8.5%		
	夜	0	0.0%	1	0.1%	0	0.0%	1	0.1%		1,289	8	1,037	2,334	2.2%	8.6%		
第4病日	朝	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,288	1	1,044	2,333	2.2%	8.6%	
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,285	3	1,045	2,333	2.2%	8.6%	
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,280	5	1,048	2,333	2.2%	8.6%	
第5病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%			1,280	0	1,053	2,333	2.2%	8.6%	
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,277	3	1,053	2,333	2.2%	8.6%	
	夜	0	0.0%	0	0.0%	—	—	0	0.0%			1,277	0	1,056	2,333	2.2%	8.6%	
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%			1,277	0	1,056	2,333	2.2%	8.6%	
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,276	1	1,056	2,333	2.2%	8.6%	
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,275	1	1,057	2,333	2.2%	8.6%	
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%			1,275	0	1,058	2,333	2.2%	8.6%	
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,274	1	1,058	2,333	2.2%	8.6%	
	夜	0	0.0%	0	0.0%	—	—	0	0.0%			1,274	0	1,059	2,333	2.2%	8.6%	
累積発現率 (参考)		38		40		23		17		78	3.2%					2.2%	8.6%	

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	3.04 (1.89 , 4.88)	<.0001
第1～第2病日	2.99 (1.86 , 4.81)	<.0001
第1～第4病日	3.04 (1.89 , 4.88)	<.0001

表9-1-6 タミフルと意識消失

		発現数 (%)								①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)			
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^②		既使用 ^③				総計	未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用	
																			1
第1病日	朝	1	0.0%	0	0.0%	0	0.0%	—	1	0.0%	0.606	0.00	2,126	303	0	2,429	0.0%	0.0%	
	昼	5	0.3%	1	0.2%	1	0.3%	0	0.0%	6	0.2%	0.439	0.62	1,836	289	303	2,428	0.3%	0.2%
	夜	7	0.5%	4	0.4%	3	0.8%	1	0.2%	11	0.5%	0.339	0.85	1,446	385	591	2,422	0.8%	0.6%
第2病日	朝	3	0.3%	2	0.1%	2	0.4%	0	0.0%	5	0.2%	0.074	0.42	931	508	972	2,411	1.1%	0.7%
	昼	0	0.0%	3	0.2%	3	0.7%	0	0.0%	3	0.1%	0.005		523	405	1,478	2,406	1.1%	0.9%
	夜	1	0.3%	4	0.2%	2	1.2%	2	0.1%	5	0.2%	0.088	0.69	352	171	1,880	2,403	1.4%	1.1%
第3病日	朝	0	0.0%	1	0.0%	1	1.3%	0	0.0%	1	0.0%	0.033		271	80	2,047	2,398	1.4%	1.1%
	昼	0	0.0%	1	0.0%	0	0.0%	1	0.0%	1	0.0%	0.887		212	59	2,126	2,397	1.4%	1.2%
	夜	1	0.5%	1	0.0%	1	4.8%	0	0.0%	2	0.1%	0.003	0.09	191	21	2,184	2,396	1.9%	1.2%
第4病日	朝	0	0.0%	1	0.0%	0	0.0%	1	0.0%	1	0.0%	0.921		183	7	2,204	2,394	1.9%	1.2%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			177	6	2,210	2,393	1.9%	1.2%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			175	2	2,216	2,393	1.9%	1.2%
第5病日	朝	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			174	1	2,218	2,393	1.9%	1.2%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			174	0	2,219	2,393	1.9%	1.2%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			174	0	2,219	2,393	1.9%	1.2%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			174	0	2,219	2,393	1.9%	1.2%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			174	0	2,219	2,393	1.9%	1.2%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			173	1	2,219	2,393	1.9%	1.2%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			173	0	2,220	2,393	1.9%	1.2%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			173	0	2,220	2,393	1.9%	1.2%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			173	0	2,220	2,393	1.9%	1.2%
累積発生率 (参考)		18		18		13		5		36	1.5%					1.9%	1.2%		

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	0.71 (0.33 , 1.52)	0.377
第1～第2病日	0.76 (0.34 , 1.68)	0.493
第1～第4病日	0.71 (0.33 , 1.52)	0.377

表9-2-6 解熱剤と意識消失

		発現数 (%)								①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)	
		薬剤 未使用 ^①		薬剤使用		(内訳)		総計				未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用
				計	使用開始 ^②	既使用 ^③											
第1病日	朝	0	0.0%	1	1.2%	1	1.2%	—	1	0.0%	0.010	2,317	84	0	2,401	0.0%	1.2%
	昼	5	0.2%	1	0.4%	1	0.5%	0	0.0%	0.666	1.49	2,116	201	83	2,400	0.2%	1.5%
	夜	3	0.2%	8	1.1%	6	1.3%	2	0.7%	0.012	6.03	1,655	456	283	2,394	0.4%	2.6%
第2病日	朝	1	0.1%	4	0.5%	1	1.3%	3	0.4%	0.072	7.82	1,574	78	731	2,383	0.5%	3.1%
	昼	1	0.1%	2	0.2%	0	0.0%	2	0.2%	0.470	3.10	1,445	128	805	2,378	0.5%	3.3%
	夜	0	0.0%	5	0.5%	1	0.9%	4	0.4%	0.014		1,328	116	931	2,375	0.5%	3.8%
第3病日	朝	1	0.1%	0	0.0%	0	0.0%	0	0.0%	0.558	0.00	1,322	6	1,042	2,370	0.6%	3.8%
	昼	1	0.1%	0	0.0%	0	0.0%	0	0.0%	0.554	0.00	1,312	9	1,048	2,369	0.7%	3.8%
	夜	1	0.1%	1	0.1%	0	0.0%	1	0.1%	0.983	1.22	1,303	8	1,057	2,368	0.8%	3.8%
第4病日	朝	0	0.0%	1	0.1%	0	0.0%	1	0.1%	0.450		1,301	1	1,064	2,366	0.8%	3.9%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,298	3	1,064	2,365	0.8%	3.9%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,293	5	1,067	2,365	0.8%	3.9%
第5病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%			1,293	0	1,072	2,365	0.8%	3.9%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,290	3	1,072	2,365	0.8%	3.9%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%			1,290	0	1,075	2,365	0.8%	3.9%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%			1,290	0	1,075	2,365	0.8%	3.9%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,289	1	1,075	2,365	0.8%	3.9%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,288	1	1,076	2,365	0.8%	3.9%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%			1,288	0	1,077	2,365	0.8%	3.9%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,287	1	1,077	2,365	0.8%	3.9%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%			1,287	0	1,078	2,365	0.8%	3.9%
累積発現率 (参考)	13		23		10		13		36	1.5%						0.8%	3.9%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	4.33 (2.11 , 8.92)	<.0001
第1～第2病日	5.86 (2.63 , 13.05)	<.0001
第1～第4病日	4.33 (2.11 , 8.92)	<.0001

表9-1-7 タミフルとひどい咳

		発現数 (%)								①②③間の		期間ごとの オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)	
		薬剤 未使用 ^①		薬剤使用		(内訳)		総計	χ ² 検定 p値	未使用	使用開始		既使用	総計	薬剤未使用	薬剤使用		
				計	使用開始 ^②	既使用 ^③												
第1病日	朝	323	15.1%	74	24.6%	74	24.6%	—	397	16.2%	<.0001	1.84	2,146	301	0	2,447	15.1%	24.6%
	昼	73	4.6%	20	4.3%	11	4.6%	9	93	4.5%	0.905	0.93	1,584	239	227	2,050	19.0%	27.8%
	夜	151	12.6%	49	6.5%	24	7.7%	25	200	10.2%	<.0001	0.48	1,198	313	446	1,957	29.2%	32.5%
第2病日	朝	34	5.0%	28	2.6%	16	4.3%	12	62	3.5%	0.001	0.50	674	373	710	1,757	32.8%	34.2%
	昼	8	2.3%	19	1.4%	9	3.1%	10	27	1.6%	0.023	0.61	351	289	1,055	1,695	34.3%	35.2%
	夜	13	5.8%	35	2.4%	6	5.0%	29	48	2.9%	0.009	0.40	223	120	1,325	1,668	38.1%	36.7%
第3病日	朝	16	9.6%	26	1.8%	2	4.5%	24	42	2.6%	<.0001	0.17	166	44	1,410	1,620	44.1%	37.9%
	昼	2	1.7%	14	1.0%	2	5.7%	12	16	1.0%	0.097	0.55	115	35	1,428	1,578	45.1%	38.5%
	夜	4	3.9%	34	2.3%	0	0.0%	34	38	2.4%	0.498	0.58	102	11	1,449	1,562	47.2%	39.9%
第4病日	朝	6	6.3%	20	1.4%	1	33.3%	19	26	1.7%	0.002	0.21	95	3	1,426	1,524	50.5%	40.7%
	昼	3	3.4%	4	0.3%	0	0.0%	4	7	0.5%	0.018	0.08	87	2	1,409	1,498	52.2%	40.9%
	夜	1	1.2%	25	1.8%	1	100.0%	24	26	1.7%	0.016	1.48	83	1	1,407	1,491	52.8%	41.9%
第5病日	朝	2	2.4%	9	0.7%	—	—	9	11	0.8%	0.141	0.26	82	0	1,383	1,465	54.0%	42.3%
	昼	2	2.5%	5	0.4%	—	—	5	7	0.5%	0.051	0.14	80	0	1,374	1,454	55.1%	42.5%
	夜	4	5.1%	12	0.9%	—	—	12	16	1.1%	0.009	0.16	78	0	1,369	1,447	57.4%	43.0%
第6病日	朝	0	0.0%	4	0.3%	—	—	4	4	0.3%	0.514		74	0	1,357	1,431	57.4%	43.2%
	昼	0	0.0%	1	0.1%	—	—	1	1	0.1%	0.744		74	0	1,353	1,427	57.4%	43.2%
	夜	0	0.0%	6	0.4%	—	—	6	6	0.4%	0.423		74	0	1,352	1,426	57.4%	43.5%
第7病日	朝	0	0.0%	2	0.1%	—	—	2	2	0.1%	0.644		74	0	1,346	1,420	57.4%	43.6%
	昼	0	0.0%	2	0.1%	—	—	2	2	0.1%	0.643		74	0	1,344	1,418	57.4%	43.7%
	夜	1	1.4%	0	0.0%	—	—	0	1	0.1%	0.015	0.00	74	0	1,342	1,416	58.0%	43.7%
累積発現率 (参考)	643		389		146		243	1032	42.2%								58.0%	43.7%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	0.69 (0.58 , 0.82)	<.0001
第1～第2病日	0.83 (0.70 , 0.99)	0.040
第1～第4病日	0.72 (0.61 , 0.86)	0.000

表9-2-7 解熱剤とひどい咳

		発現数 (%)								①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)			
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^②		既使用 ^③				総計	未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用	
第1病日	朝	362	15.5%	32	38.1%	32	38.1%	—	—	394	16.3%	<.0001	3.35	2,334	84	0	2,418	15.5%	38.1%
	夜	81	4.5%	12	5.5%	10	6.0%	2	3.8%	93	4.6%	0.682	1.23	1,804	168	52	2,024	19.3%	41.5%
第2病日	朝	139	10.3%	59	10.1%	42	11.2%	17	8.2%	198	10.3%	0.491	0.98	1,349	374	208	1,931	27.6%	47.4%
	夜	41	3.6%	21	3.6%	4	7.0%	17	3.3%	62	3.6%	0.426	1.02	1,153	57	523	1,733	30.2%	49.3%
第3病日	朝	12	1.2%	14	2.2%	2	2.2%	12	2.1%	26	1.6%	0.296	1.86	1,022	90	559	1,671	31.0%	50.4%
	夜	22	2.4%	26	3.7%	5	6.5%	21	3.3%	48	2.9%	0.134	1.57	933	77	635	1,645	32.6%	52.2%
第4病日	朝	24	2.6%	16	2.3%	0	0.0%	16	2.3%	40	2.5%	0.835	0.87	907	4	686	1,597	34.4%	53.3%
	夜	3	0.3%	13	1.9%	1	16.7%	12	1.8%	16	1.0%	0.002	5.68	877	6	674	1,557	34.6%	54.2%
第5病日	朝	20	2.3%	17	2.5%	0	0.0%	17	2.5%	37	2.4%	0.863	1.10	870	4	667	1,541	36.1%	55.4%
	夜	13	1.5%	13	2.0%	0	0.0%	13	2.0%	26	1.7%	0.785	1.30	849	1	654	1,504	37.1%	56.3%
第6病日	朝	4	0.5%	2	0.3%	0	0.0%	2	0.3%	6	0.4%	0.871	0.65	834	2	642	1,478	37.4%	56.4%
	夜	12	1.5%	14	2.2%	0	0.0%	14	2.2%	26	1.8%	0.540	1.50	826	4	642	1,472	38.3%	57.3%
第7病日	朝	3	0.4%	8	1.3%	—	—	8	1.3%	11	0.8%	0.050	3.47	814	0	632	1,446	38.6%	57.9%
	夜	3	0.4%	4	0.6%	0	0.0%	4	0.6%	7	0.5%	0.762	1.73	809	2	624	1,435	38.8%	58.2%
第8病日	朝	7	0.9%	9	1.4%	—	—	9	1.4%	16	1.1%	0.306	1.68	806	0	622	1,428	39.3%	58.8%
	夜	2	0.3%	2	0.3%	—	—	2	0.3%	4	0.3%	0.791	1.30	799	0	613	1,412	39.5%	58.9%
第9病日	朝	1	0.1%	0	0.0%	0	0.0%	0	0.0%	1	0.1%	0.565	0.00	796	1	611	1,408	39.6%	58.9%
	夜	0	0.0%	6	1.0%	0	0.0%	6	1.0%	6	0.4%	0.007	—	794	1	612	1,407	39.6%	59.3%
第10病日	朝	1	0.1%	1	0.2%	—	—	1	0.2%	2	0.1%	0.850	1.31	794	0	607	1,401	39.6%	59.4%
	夜	1	0.1%	1	0.2%	—	—	1	0.2%	2	0.1%	0.849	1.31	793	0	606	1,399	39.7%	59.4%
第11病日	朝	0	0.0%	1	0.2%	—	—	1	0.2%	1	0.1%	0.196	—	792	0	605	1,397	39.7%	59.5%
	夜	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
累積発現率 (参考)		751		271		96		175		1022	42.3%							39.7%	59.5%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.40 (1.19 , 1.63)	<.0001
第1～第2病日	1.36 (1.12 , 1.64)	0.002
第1～第4病日	1.34 (1.14 , 1.58)	0.001

表9-1-8 タミフルと嘔吐・下痢

		発現数 (%)								①②③間の		期間ごとの オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)		
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^②		既使用 ^③		総計			χ ² 検定 p値	未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用
第1病日	朝	79	3.7%	46	15.2%	46	15.2%	—	—	125	5.1%	<.0001	4.69	2,148	303	0	2,451	3.7%	15.2%
	昼	54	3.0%	53	9.9%	35	12.5%	18	7.0%	107	4.6%	<.0001	3.53	1,790	279	257	2,326	6.6%	23.6%
	夜	110	7.9%	77	9.3%	46	13.2%	31	6.4%	187	8.4%	0.002	1.18	1,387	349	483	2,219	14.0%	30.6%
第2病日	朝	38	4.6%	63	5.2%	30	6.7%	33	4.4%	101	5.0%	0.193	1.14	826	451	755	2,032	17.9%	34.3%
	昼	15	3.4%	85	5.7%	32	9.4%	53	4.6%	100	5.2%	0.001	1.74	446	342	1,143	1,931	20.7%	38.0%
	夜	11	3.9%	73	4.7%	11	7.4%	62	4.4%	84	4.6%	0.264	1.22	282	149	1,400	1,831	23.8%	40.9%
第3病日	朝	8	3.9%	48	3.1%	6	9.4%	42	2.8%	56	3.2%	0.046	0.80	207	64	1,476	1,747	26.7%	42.8%
	昼	5	3.3%	30	1.9%	0	0.0%	30	2.0%	35	2.1%	0.222	0.58	151	48	1,492	1,691	29.2%	43.9%
	夜	4	3.1%	21	1.4%	1	6.7%	20	1.3%	25	1.5%	0.178	0.44	131	15	1,510	1,656	31.3%	44.7%
第4病日	朝	0	0.0%	19	1.3%	1	20.0%	18	1.2%	19	1.2%	0.032	—	122	5	1,504	1,631	31.3%	45.4%
	昼	1	0.8%	8	0.5%	0	0.0%	8	0.5%	9	0.6%	0.908	0.64	119	3	1,490	1,612	31.9%	45.7%
	夜	1	0.9%	12	0.8%	0	0.0%	12	0.8%	13	0.8%	0.982	0.94	116	2	1,485	1,603	32.5%	46.1%
第5病日	朝	0	0.0%	4	0.3%	—	—	4	0.3%	4	0.3%	0.438	—	115	0	1,475	1,590	32.5%	46.2%
	昼	1	0.9%	2	0.1%	—	—	2	0.1%	3	0.2%	0.188	0.16	115	0	1,471	1,586	33.1%	46.3%
	夜	0	0.0%	8	0.5%	—	—	8	0.5%	8	0.5%	0.274	—	114	0	1,469	1,583	33.1%	46.6%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	114	0	1,461	1,575	33.1%	46.6%
	昼	1	0.9%	2	0.1%	—	—	2	0.1%	3	0.2%	0.187	0.15	114	0	1,461	1,575	33.7%	46.7%
	夜	1	0.9%	2	0.1%	0	0.0%	2	0.1%	3	0.2%	0.413	0.15	112	1	1,459	1,572	34.3%	46.8%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	111	0	1,458	1,569	34.3%	46.8%
	昼	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.1%	0.702	—	111	0	1,458	1,569	34.3%	46.8%
	夜	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.1%	0.702	—	111	0	1,457	1,568	34.3%	46.8%
累積発生率 (参考)		329		555		208		347		884	36.1%							34.3%	46.8%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.70 (1.44 , 2.01)	<.0001
第1～第2病日	1.87 (1.57 , 2.22)	<.0001
第1～第4病日	1.73 (1.46 , 2.04)	<.0001

表9-2-8 解熱剤と嘔吐・下痢

		発現数 (%)									①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)		
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^②		既使用 ^③		総計			未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用	
第1病日	朝	109	4.7%	14	16.3%	14	16.3%	—	—	123	5.1%	<.0001	3.97	2,337	86	0	2,423	4.7%	16.3%
	昼	82	4.0%	23	8.9%	15	8.1%	8	11.1%	105	4.6%	0.004	2.35	2,043	185	72	2,300	8.5%	23.8%
	夜	121	7.9%	65	9.9%	46	10.8%	19	8.1%	186	8.5%	0.163	1.28	1,537	424	234	2,195	15.7%	31.3%
第2病日	朝	58	4.3%	40	6.0%	5	7.1%	35	5.9%	98	4.9%	0.233	1.43	1,346	70	593	2,009	19.3%	35.4%
	昼	59	5.0%	40	5.5%	9	8.3%	31	5.0%	99	5.2%	0.369	1.10	1,180	108	623	1,911	23.4%	39.0%
	夜	34	3.3%	50	6.4%	5	5.4%	45	6.5%	84	4.6%	0.008	2.00	1,029	92	691	1,812	25.9%	42.9%
第3病日	朝	31	3.1%	25	3.4%	1	16.7%	24	3.3%	56	3.2%	0.403	1.08	989	6	733	1,728	28.2%	44.8%
	昼	13	1.4%	22	3.1%	0	0.0%	22	3.1%	35	2.1%	0.049	2.28	952	6	714	1,672	29.2%	46.5%
	夜	13	1.4%	12	1.7%	0	0.0%	12	1.7%	25	1.5%	0.782	1.22	932	7	698	1,637	30.2%	47.4%
第4病日	朝	7	0.8%	12	1.7%	0	0.0%	12	1.7%	19	1.2%	0.205	2.29	918	1	693	1,612	30.7%	48.3%
	昼	7	0.8%	2	0.3%	0	0.0%	2	0.3%	9	0.6%	0.423	0.38	909	2	682	1,593	31.2%	48.5%
	夜	7	0.8%	6	0.9%	0	0.0%	6	0.9%	13	0.8%	0.952	1.13	899	3	682	1,584	31.8%	48.9%
第5病日	朝	3	0.3%	1	0.1%	—	—	1	0.1%	4	0.3%	0.448	0.44	892	0	679	1,571	32.0%	49.0%
	昼	2	0.2%	1	0.1%	—	—	1	0.1%	3	0.2%	0.725	0.66	889	0	678	1,567	32.2%	49.1%
	夜	3	0.3%	4	0.6%	—	—	4	0.6%	7	0.4%	0.461	1.75	887	0	677	1,564	32.4%	49.4%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	884	0	673	1,557	32.4%	49.4%
	昼	2	0.2%	1	0.1%	—	—	1	0.1%	3	0.2%	0.726	0.66	884	0	673	1,557	32.6%	49.4%
	夜	2	0.2%	1	0.1%	0	0.0%	1	0.1%	3	0.2%	0.938	0.65	881	1	672	1,554	32.7%	49.5%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	879	0	672	1,551	32.7%	49.5%
	昼	0	0.0%	1	0.1%	0	0.0%	1	0.1%	1	0.1%	0.433	—	878	1	672	1,551	32.7%	49.6%
	夜	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.1%	0.196	—	878	0	672	1,550	32.7%	49.7%
累積発生率 (参考)	553		321		95		226		874	36.1%								32.7%	49.7%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.48 (1.28 , 1.72)	<.0001
第1～第2病日	1.54 (1.30 , 1.83)	<.0001
第1～第4病日	1.50 (1.29 , 1.74)	<.0001

表9-1-9 タミフルと筋肉痛

		発現数 (%)							①②③間の x2検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)				
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^② 既使用 ^③		総計			未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用			
		人数	%	人数	%	人数	%										人数	%	
第1病日	朝	97	5.1%	23	8.1%	23	8.1%	—	120	5.5%	0.050	1.64	1,903	284	0	2,187	5.1%	8.1%	
	昼	29	1.8%	13	2.7%	6	2.6%	7	2.7%	42	2.0%	0.547	1.46	1,579	227	261	2,067	6.8%	10.5%
	夜	75	6.1%	26	3.3%	16	5.0%	10	2.1%	101	5.0%	0.001	0.52	1,232	318	475	2,025	12.5%	13.5%
第2病日	朝	14	1.9%	22	1.9%	10	2.5%	12	1.6%	36	1.9%	0.571	1.01	751	406	767	1,924	14.1%	15.1%
	昼	5	1.2%	13	0.9%	4	1.2%	9	0.8%	18	1.0%	0.639	0.71	406	331	1,151	1,888	15.2%	15.8%
	夜	5	1.9%	13	0.8%	2	1.5%	11	0.7%	18	1.0%	0.234	0.43	269	132	1,469	1,870	16.8%	16.5%
第3病日	朝	1	0.5%	8	0.5%	0	0.0%	8	0.5%	9	0.5%	0.746	1.00	205	59	1,588	1,852	17.2%	16.9%
	昼	0	0.0%	1	0.1%	0	0.0%	1	0.1%	1	0.1%	0.889	0.24	152	52	1,639	1,843	17.2%	17.0%
	夜	1	0.7%	3	0.2%	0	0.0%	3	0.2%	4	0.2%	0.536	0.24	135	17	1,690	1,842	17.8%	17.1%
第4病日	朝	0	0.0%	1	0.1%	1	16.7%	0	0.0%	1	0.1%	0.003	—	128	6	1,704	1,838	17.8%	17.2%
	昼	0	0.0%	1	0.1%	0	0.0%	1	0.1%	1	0.1%	0.930	—	123	5	1,709	1,837	17.8%	17.2%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	—	—	121	2	1,713	1,836	17.8%	17.2%
第5病日	朝	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.1%	0.712	—	121	0	1,715	1,836	17.8%	17.3%
	昼	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.1%	0.712	—	121	0	1,714	1,835	17.8%	17.3%
	夜	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.1%	0.712	—	121	0	1,713	1,834	17.8%	17.4%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	121	0	1,712	1,833	17.8%	17.4%
	昼	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.1%	0.712	—	121	0	1,712	1,833	17.8%	17.4%
	夜	1	0.8%	0	0.0%	0	0.0%	0	0.0%	1	0.1%	0.065	0.00	120	1	1,711	1,832	18.5%	17.4%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	119	0	1,712	1,831	18.5%	17.4%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	119	0	1,712	1,831	18.5%	17.4%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	119	0	1,712	1,831	18.5%	17.4%
累積発現率 (参考)		228		128		62		66		356	16.3%						18.5%	17.4%	

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	0.88 (0.68 , 1.13)	0.315
第1～第2病日	0.89 (0.69 , 1.16)	0.386
第1～第4病日	0.89 (0.69 , 1.15)	0.357

表9-2-9 解熱剤と筋肉痛

		発現数 (%)								①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)		
		薬剤 未使用 ^①		薬剤使用		(内訳)		総計				未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用	
				計	使用開始 ^②													既使用 ^③
第1病日	朝	107	5.1%	11	14.9%	11	14.9%	—	118	5.5%	0.002	3.23	2,088	74	0	2,162	5.1%	14.9%
	昼	37	2.0%	4	1.7%	3	1.8%	1	41	2.0%	0.947	0.85	1,814	167	63	2,044	7.1%	16.3%
	夜	67	4.8%	33	5.4%	25	6.6%	8	100	5.0%	0.215	1.14	1,397	380	226	2,003	11.5%	20.9%
第2病日	朝	24	1.9%	12	1.9%	5	7.7%	7	36	1.9%	0.016	0.99	1,265	65	573	1,903	13.2%	22.4%
	昼	12	1.1%	6	0.8%	2	2.0%	4	18	1.0%	0.420	0.78	1,141	100	626	1,867	14.1%	23.0%
	夜	8	0.8%	10	1.2%	1	1.1%	9	18	1.0%	0.604	1.60	1,037	92	720	1,849	14.8%	24.0%
第3病日	朝	1	0.1%	8	1.0%	0	0.0%	8	9	0.5%	0.017	10.27	1,025	4	802	1,831	14.9%	24.7%
	昼	1	0.1%	0	0.0%	0	0.0%	0	1	0.1%	0.558	0.00	1,017	7	798	1,822	14.9%	24.7%
	夜	1	0.1%	3	0.4%	0	0.0%	3	4	0.2%	0.454	3.76	1,012	4	805	1,821	15.0%	25.0%
第4病日	朝	1	0.1%	0	0.0%	—	—	0	1	0.1%	0.279	0.00	1,011	0	806	1,817	15.1%	25.0%
	昼	0	0.0%	1	0.1%	0	0.0%	1	1	0.1%	0.444	—	1,009	1	806	1,816	15.1%	25.1%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%	—	—	1,006	3	806	1,815	15.1%	25.1%
第5病日	朝	1	0.1%	0	0.0%	—	—	0	1	0.1%	0.277	0.00	1,006	0	809	1,815	15.2%	25.1%
	昼	0	0.0%	1	0.1%	0	0.0%	1	1	0.1%	0.446	—	1,002	3	809	1,814	15.2%	25.2%
	夜	1	0.1%	0	0.0%	—	—	0	1	0.1%	0.276	0.00	1,002	0	811	1,813	15.3%	25.2%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0	0.0%	—	—	1,001	0	811	1,812	15.3%	25.2%
	昼	1	0.1%	0	0.0%	0	0.0%	1	1	0.1%	0.552	0.00	1,000	1	811	1,812	15.4%	25.2%
	夜	0	0.0%	1	0.1%	—	—	1	1	0.1%	0.205	—	999	0	812	1,811	15.4%	25.3%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0	0.0%	—	—	999	0	811	1,810	15.4%	25.3%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%	—	—	998	1	811	1,810	15.4%	25.3%
	夜	0	0.0%	0	0.0%	—	—	0	0	0.0%	—	—	998	0	812	1,810	15.4%	25.3%
累積発現率 (参考)	262		90		47		43	352	16.3%								15.4%	25.3%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.33 (1.02 , 1.72)	0.035
(症候出現後に使用とした場合)	1.24 (0.94 , 1.64)	0.125
第1～第4病日	1.34 (1.03 , 1.75)	0.030

表9-1-10 タミフルと関節痛

		発現数 (%)									①②③間の χ ² 検定 p値	期間ごとの オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)	
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^②		既使用 ^③		総計			未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用
第1病日	朝	131	6.9%	42	15.1%	42	15.1%	—	173	8.0%	<.0001	2.40	1,896	278	0	2,174	6.9%	15.1%
	昼	46	3.0%	14	3.1%	7	3.2%	7	60	3.0%	0.982	1.04	1,547	218	236	2,001	9.7%	17.7%
	夜	123	10.3%	35	4.7%	20	6.6%	15	158	8.1%	<.0001	0.43	1,199	302	440	1,941	18.9%	21.6%
第2病日	朝	27	3.8%	22	2.1%	16	4.4%	6	49	2.7%	<.0001	0.53	711	365	707	1,783	22.0%	23.2%
	昼	4	1.0%	20	1.5%	8	2.7%	12	24	1.4%	0.153	1.44	386	298	1,050	1,734	22.8%	24.4%
	夜	3	1.2%	11	0.8%	4	3.2%	7	14	0.8%	0.030	0.65	258	124	1,328	1,710	23.7%	24.9%
第3病日	朝	0	0.0%	4	0.3%	0	0.0%	4	4	0.2%	0.521	—	197	58	1,441	1,696	23.7%	25.1%
	昼	1	0.7%	2	0.1%	1	2.0%	1	3	0.2%	0.060	0.19	148	49	1,495	1,692	24.2%	25.2%
	夜	0	0.0%	4	0.3%	0	0.0%	4	4	0.2%	0.694	—	130	17	1,542	1,689	24.2%	25.4%
第4病日	朝	0	0.0%	1	0.1%	1	16.7%	0	1	0.1%	0.003	—	124	6	1,555	1,685	24.2%	25.5%
	昼	0	0.0%	2	0.1%	0	0.0%	2	2	0.1%	0.858	—	120	4	1,560	1,684	24.2%	25.6%
	夜	0	0.0%	1	0.1%	0	0.0%	1	1	0.1%	0.929	—	118	2	1,562	1,682	24.2%	25.6%
第5病日	朝	0	0.0%	1	0.1%	0	0.0%	1	1	0.1%	0.930	—	117	1	1,563	1,681	24.2%	25.7%
	昼	0	0.0%	1	0.1%	—	—	1	1	0.1%	0.704	—	117	0	1,563	1,680	24.2%	25.7%
	夜	1	0.9%	1	0.1%	—	—	1	2	0.1%	0.100	0.07	117	0	1,562	1,679	24.9%	25.8%
第6病日	朝	0	0.0%	1	0.1%	—	—	1	1	0.1%	0.705	—	116	0	1,561	1,677	24.9%	25.8%
	昼	1	0.9%	0	0.0%	—	—	0	1	0.1%	0.021	0.00	116	0	1,560	1,676	25.5%	25.8%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%	—	—	114	1	1,560	1,675	25.5%	25.8%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0	0.0%	—	—	114	0	1,561	1,675	25.5%	25.8%
	昼	0	0.0%	0	0.0%	—	—	0	0	0.0%	—	—	114	0	1,561	1,675	25.5%	25.8%
	夜	0	0.0%	0	0.0%	—	—	0	0	0.0%	—	—	114	0	1,561	1,675	25.5%	25.8%
累積発現率 (参考)		337		162		99		63	499	23.0%							25.5%	25.8%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	0.85 (0.68 , 1.05)	0.138
第1～第2病日	0.86 (0.69 , 1.07)	0.174
第1～第4病日	0.87 (0.70 , 1.07)	0.189

表9-2-10 解熱剤と関節痛

		発現数 (%)								①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)			
		薬剤 未使用 ^①		薬剤使用		(内訳)		総計				未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用		
				計	使用開始 ^②	既使用 ^③													
第1病日	朝	160	7.7%	12	16.4%	12	16.4%	—	172	8.0%	0.016	2.35	2,075	73	0	2,148	7.7%	16.4%	
	昼	51	2.9%	7	3.2%	5	3.2%	2	3.3%	58	2.9%	0.962	1.12	1,760	155	61	1,976	10.4%	19.1%
	夜	111	8.2%	47	8.3%	38	10.7%	9	4.3%	158	8.2%	0.021	1.02	1,354	355	209	1,918	17.7%	25.9%
第2病日	朝	38	3.2%	12	2.1%	5	8.8%	7	1.4%	50	2.8%	0.006	0.65	1,186	57	517	1,760	20.4%	27.4%
	昼	16	1.5%	8	1.2%	2	2.2%	6	1.1%	24	1.4%	0.615	0.81	1,059	89	562	1,710	21.6%	28.3%
	夜	4	0.4%	10	1.4%	1	1.2%	9	1.4%	14	0.8%	0.098	3.33	959	84	643	1,686	21.9%	29.3%
第3病日	朝	1	0.1%	3	0.4%	0	0.0%	3	0.4%	4	0.2%	0.424	3.97	951	4	717	1,672	22.0%	29.6%
	昼	1	0.1%	2	0.3%	0	0.0%	2	0.3%	3	0.2%	0.707	2.61	943	7	718	1,668	22.1%	29.8%
	夜	1	0.1%	3	0.4%	0	0.0%	3	0.4%	4	0.2%	0.436	3.87	937	5	723	1,665	22.1%	30.1%
第4病日	朝	1	0.1%	0	0.0%	—	—	0	0.0%	1	0.1%	0.284	0.00	936	0	725	1,661	22.2%	30.1%
	昼	1	0.1%	1	0.1%	0	0.0%	1	0.1%	2	0.1%	0.983	1.29	934	1	725	1,660	22.3%	30.2%
	夜	0	0.0%	1	0.1%	0	0.0%	1	0.1%	1	0.1%	0.437	—	929	4	725	1,658	22.3%	30.3%
第5病日	朝	1	0.1%	0	0.0%	—	—	0	0.0%	1	0.1%	0.282	0.00	929	0	728	1,657	22.4%	30.3%
	昼	1	0.1%	0	0.0%	0	0.0%	0	0.0%	1	0.1%	0.560	0.00	927	1	728	1,656	22.5%	30.3%
	夜	0	0.0%	2	0.3%	—	—	2	0.3%	2	0.1%	0.070	—	926	0	729	1,655	22.5%	30.5%
第6病日	朝	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.1%	0.200	—	926	0	727	1,653	22.5%	30.6%
	昼	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.1%	0.200	—	926	0	726	1,652	22.5%	30.7%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	926	0	725	1,651	22.5%	30.7%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	926	0	725	1,651	22.5%	30.7%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	—	—	925	1	725	1,651	22.5%	30.7%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	925	0	726	1,651	22.5%	30.7%
累積発生率 (参考)	387		110		63		47		497	23.1%							22.5%	30.7%	

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.17 (0.93 , 1.47)	0.188
第1～第2病日	1.10 (0.86 , 1.40)	0.445
第1～第4病日	1.15 (0.91 , 1.45)	0.244

表9-1-11 タミフルとノドの痛み

		発現数 (%)								①②③間の χ ² 検定 p値	期間ごとの オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)			
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^②		既使用 ^③				総計	未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用	
		人数	%	人数	%	人数	%	人数	%										
第1病日	朝	390	20.1%	92	32.4%	92	32.4%	—	—	482	21.7%	<.0001	1.91	1,942	284	0	2,226	20.1%	32.4%
	昼	70	5.1%	21	5.8%	14	8.1%	7	3.6%	91	5.2%	0.159	1.15	1,380	172	192	1,744	24.1%	36.3%
	夜	191	18.3%	50	8.2%	35	13.2%	15	4.4%	241	14.6%	<.0001	0.40	1,045	265	343	1,653	38.0%	41.5%
第2病日	朝	40	7.0%	46	5.5%	23	8.2%	23	4.1%	86	6.1%	0.030	0.78	574	280	558	1,412	42.3%	44.7%
	昼	10	3.4%	14	1.4%	7	2.9%	7	0.9%	24	1.8%	0.009	0.39	295	239	792	1,326	44.3%	45.5%
	夜	7	3.6%	22	2.0%	1	1.1%	21	2.1%	29	2.2%	0.348	0.55	196	89	1,017	1,302	46.3%	46.6%
第3病日	朝	6	4.1%	8	0.7%	1	2.4%	7	0.6%	14	1.1%	0.007	0.17	147	42	1,084	1,273	48.5%	47.0%
	昼	3	2.9%	6	0.5%	1	2.6%	5	0.4%	9	0.7%	0.040	0.17	103	38	1,118	1,259	50.0%	47.2%
	夜	2	2.2%	10	0.9%	0	0.0%	10	0.9%	12	1.0%	0.487	0.38	89	11	1,150	1,250	51.1%	47.7%
第4病日	朝	4	4.8%	7	0.6%	2	50.0%	5	0.4%	11	0.9%	<.0001	0.12	83	4	1,151	1,238	53.4%	48.0%
	昼	0	0.0%	4	0.3%	1	20.0%	3	0.3%	4	0.3%	0.029		74	5	1,148	1,227	53.4%	48.2%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			73	1	1,149	1,223	53.4%	48.2%
第5病日	朝	0	0.0%	1	0.1%	0	0.0%	1	0.1%	1	0.1%	0.940		72	1	1,150	1,223	53.4%	48.2%
	昼	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.1%	0.727		72	0	1,150	1,222	53.4%	48.3%
	夜	1	1.4%	2	0.2%	—	—	2	0.2%	3	0.2%	0.148	0.12	72	0	1,149	1,221	54.1%	48.4%
第6病日	朝	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.1%	0.729		71	0	1,147	1,218	54.1%	48.4%
	昼	1	1.4%	0	0.0%	—	—	0	0.0%	1	0.1%	0.017	0.00	71	0	1,146	1,217	54.7%	48.4%
	夜	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.1%	0.731		70	0	1,146	1,216	54.7%	48.5%
第7病日	朝	0	0.0%	2	0.2%	—	—	2	0.2%	2	0.2%	0.626		70	0	1,145	1,215	54.7%	48.5%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			70	0	1,143	1,213	54.7%	48.5%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			70	0	1,143	1,213	54.7%	48.5%
累積発生率 (参考)	725		288		177		111		1013	45.5%								54.7%	48.5%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	0.82 (0.70 , 0.96)	0.015
第1～第2病日	0.89 (0.75 , 1.04)	0.145
第1～第4病日	0.83 (0.70 , 0.97)	0.021

表9-2-11 解熱剤とノドの痛み

	薬剤 未使用 ^①	発現数 (%)						①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)	
		薬剤使用 計		(内訳)		総計				未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用
		未使用 ^①	計	使用開始 ^②	既使用 ^③	計	総計								
第1病日	朝	447 21.0%	30 40.0%	30 40.0%	—	477 21.7%	0.000	2.50	2,125	75	0	2,200	21.0%	40.0%	
	昼	79 5.1%	11 6.0%	9 6.5%	2 4.4%	90 5.2%	0.770	1.18	1,540	138	45	1,723	25.1%	43.6%	
	夜	170 14.8%	65 13.4%	51 16.3%	14 8.1%	235 14.4%	0.025	0.89	1,148	313	172	1,633	36.2%	51.2%	
第2病日	朝	63 6.7%	23 5.1%	5 14.7%	18 4.3%	86 6.2%	0.040	0.75	944	34	420	1,398	40.4%	53.6%	
	昼	10 1.2%	13 2.6%	6 9.2%	7 1.6%	23 1.8%	0.002	2.17	816	65	431	1,312	41.2%	54.9%	
	夜	18 2.4%	11 2.0%	4 7.1%	7 1.4%	29 2.2%	0.061	0.85	750	56	483	1,289	42.6%	55.8%	
第3病日	朝	9 1.2%	5 0.9%	1 33.3%	4 0.8%	14 1.1%	0.050	0.76	729	3	528	1,260	43.3%	56.2%	
	昼	5 0.7%	4 0.8%	1 16.7%	3 0.6%	9 0.7%	0.095	1.07	714	6	526	1,246	43.7%	56.5%	
	夜	4 0.6%	8 1.5%	—	8 1.5%	12 1.0%	0.093	2.71	709	0	528	1,237	44.0%	57.2%	
第4病日	朝	4 0.6%	6 1.2%	—	6 1.2%	10 0.8%	0.263	2.05	705	0	520	1,225	44.3%	57.7%	
	昼	3 0.4%	1 0.2%	0 0.0%	1 0.2%	4 0.3%	0.766	0.45	700	1	514	1,215	44.6%	57.8%	
	夜	0 0.0%	0 0.0%	0 0.0%	0 0.0%	0 0.0%	—	—	695	2	514	1,211	44.6%	57.8%	
第5病日	朝	1 0.1%	0 0.0%	—	0 0.0%	1 0.1%	0.292	0.00	695	0	516	1,211	44.6%	57.8%	
	昼	0 0.0%	1 0.2%	0 0.0%	1 0.2%	1 0.1%	0.426	—	692	2	516	1,210	44.6%	57.8%	
	夜	1 0.1%	2 0.4%	—	2 0.4%	3 0.2%	0.404	2.68	692	0	517	1,209	44.7%	58.0%	
第6病日	朝	0 0.0%	1 0.2%	—	1 0.2%	1 0.1%	0.192	—	691	0	515	1,206	44.7%	58.1%	
	昼	0 0.0%	1 0.2%	0 0.0%	1 0.2%	1 0.1%	0.426	—	690	1	514	1,205	44.7%	58.2%	
	夜	1 0.1%	0 0.0%	—	0 0.0%	1 0.1%	0.291	0.00	690	0	514	1,204	44.8%	58.2%	
第7病日	朝	1 0.1%	1 0.2%	—	1 0.2%	2 0.2%	0.836	1.34	689	0	514	1,203	44.9%	58.2%	
	昼	0 0.0%	0 0.0%	0 0.0%	0 0.0%	0 0.0%	—	—	687	1	513	1,201	44.9%	58.2%	
	夜	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%	—	—	687	0	514	1,201	44.9%	58.2%	
累積発現率 (参考)	816	183	107	76	999	45.4%	—	—	—	—	—	—	44.9%	58.2%	

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.13 (0.95 , 1.35)	0.171
第1～第2病日	1.10 (0.91 , 1.33)	0.329
第1～第4病日	1.12 (0.94 , 1.34)	0.221