

参考文献

- 1) Moise AR, Noy N, Palczewski K, et al. Delivery of retinoid-based therapies to target tissues. *Biochemistry* 2008; 46: 4449–58.
- 2) Debier C, Larondelle Y. Vitamins A and E: metabolism, roles and transfer to offspring. *Br J Nutr* 2005; 93: 153–74.
- 3) During A, Harrison EH. Intestinal absorption and metabolism of carotenoids: insights from cell culture. *Arch Biochem Biophys* 2004; 430: 77–88.
- 4) van het Hof KH, West CE, Weststrate JA, et al. Dietary factors that affect the bioavailability of carotenoids. *J Nutr* 2000; 130: 503–6.
- 5) van het Hof KH, Brouwer IA, West CE, et al. Bioavailability of lutein from vegetables is five times higher than that of beta-carotene. *Am J Clin Nutr* 1999; 70: 261–8.
- 6) Food and Nutrition Board, Institute of Medicine. Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. 2nd ed. National Academy Press, Washington D.C., 2002.
- 7) Sigmundsdottir H, Butcher EC. Environmental cues, dendritic cells and the programming of tissue-selective lymphocyte trafficking. *Nature Immunol* 2008; 9: 981–7.
- 8) Sauberlich HE, Hodges RE, Wallace DL, et al. Vitamin A metabolism and requirements in the human studied with the use of labeled retinol. *Vitam Horm* 1974; 32: 251–75.
- 9) Ahmad SM, Haskell MJ, Raqib R, et al. Men with low vitamin A stores respond adequately to primary yellow fever and secondary tetanus toxoid vaccination. *J Nutr* 2008; 138: 2276–83.
- 10) Olson JA. Recommended dietary intakes (RDI) of vitamin A in humans. *Am J Clin Nutr* 1987; 45: 704–16.
- 11) Cifelli CJ, Green JB, Wang Z, et al. Kinetic analysis shows that vitamin A disposal rate in humans is positively correlated with vitamin A stores. *J Nutr* 2008; 138: 971–7.
- 12) Cifelli CJ, Green JB, Green MH. Use of model-based compartmental analysis to study vitamin A kinetics and metabolism. *Vitam Horm* 2007; 75: 161–95.
- 13) Furr HC, Green MH, Haskell M, et al. Stable isotope dilution techniques for assessing vitamin A status and bioefficiency of provitamin A carotenoids in humans. *Public Health Nutr* 2005; 8: 596–607.
- 14) 島田馨. 内科学書：改訂第6版. 中山書店, 東京, 2002.
- 15) Raica N, Scott J, Lowry L, et al. Vitamin A concentration in human tissues collected from five areas in the United States. *Am J Clin Nutr* 1972; 25: 291–6.
- 16) Joint FAO/WHO Expert Group. Human vitamin and mineral requirements, 2nd edition. Chapter 2. Vitamin A. WHO/FAO, 2004: 17–44.
- 17) Canfield LM, Clandinin MT, Davies DP, et al. Multinational study of major breast milk carotenoids of healthy mothers. *Eur J Nutr* 2003; 42: 133–41.
- 18) Sakurai T, Furukawa M, Asoh M, et al. Fat-soluble and water-soluble vitamin contents of breast milk from Japanese women. *J Nutr Sci Vitaminol* 2005; 51: 239–47.
- 19) Kamao M, Tsugawa N, Suhara Y, et al. Quantification of fat-soluble vitamins in human

- breast milk by liquid chromatography-tandem mass spectrometry. *J Chromatogr B* 2007; 859: 192-200.
- 20) 鈴木久美子, 佐々木晶子, 新澤佳代, 他. 離乳前乳児の哺乳量に関する研究. *栄養学雑誌* 2004; 62: 369-72.
- 21) 廣瀬潤子, 遠藤美佳, 柴田克己, 他. 日本人母乳栄養児（0～5ヵ月）の哺乳量. *日本母乳哺育学会雑誌* 2008; 2: 23-8.
- 22) Montreewasuwat N, Olson JA. Serum and liver concentrations of vitamin A in Thai fetuses as a function of gestational age. *Am J Clin Nutr* 1979; 32: 601-6.
- 23) Strobel M, Tinz J, Biesalski HK. The importance of beta-carotene as a source of vitamin A with special regard to pregnant and breastfeeding women. *Eur J Nutr* 2007; 46(Suppl 1): S1-20.
- 24) Penniston KL, Tanumihardjo SA. The acute and chronic toxic effects of vitamin A. *Am J Clin Nutr* 2006; 83: 191-201.
- 25) Azais-Braesco V, Pascal G. Vitamin A in pregnancy: requirements and safety limits. *Am J Clin Nutr* 2000; 71 (5 Suppl): S1325-33.
- 26) Rothman KJ, Moore LL, Singer MR, et al. Teratogenicity of high vitamin A intake. *N Engl J Med* 1995; 333: 1369-73.
- 27) Minuk GY, Kelly JK, Hwang WS. Vitamin A hepatotoxicity in multiple family members. *Hepatology* 1988; 8: 272-5.
- 28) Persson B, Tunell R, Ekengren K. Chronic vitamin A intoxication during the first year of life. *Acta Paediatr Scand* 1965; 54: 49-60.
- 29) Michaelsson K, Lithell H, Vessby B, et al. Serum retinol levels and the risk of fracture. *N Engl J Med* 2003; 348: 287-94.
- 30) Ribaya-Mercado JD, Blumberg JB. Vitamin A: is it a risk factor for osteoporosis and bone fracture? *Nutr Rev* 2007; 65: 425-38.
- 31) Mannisto S, Smith-Warner SA, Spiegelman D, et al. Dietary carotenoids and risk of lung cancer in a pooled analysis of seven cohort studies. *Cancer Epidemiol Biomarkers Prev* 2004; 13: 40-8.
- 32) The Alpha-tocopherol, Beta-carotene Cancer Prevention Study Group. The effect of vitamin E and beta-carotene on the incidence of lung cancer and other cancers in male smokers. *N Engl J Med* 1994; 330: 1029-35.
- 33) Albanes D, Heinonen OP, Taylor PR, et al. Alpha-tocopherol and beta-carotene supplements and lung cancer incidence in the alpha-tocopherol, beta-carotene cancer prevention study: effects of base-line characteristics and study compliance. *J Natl Cancer Inst* 1996; 88: 1560-70.
- 34) Omenn GS, Goodman GE, Thornquist MD, et al. Effects of a combination of beta-carotene and vitamin A on lung cancer and cardiovascular disease. *N Eng J Med* 1996; 334: 1150-5.
- 35) Hennekens CH, Buring JE, Manson JE, et al. Lack of effect of long-term supplementation with beta-carotene on the incidence of malignant neoplasms and cardiovascular disease. *N Eng J Med* 1996; 334: 1145-9.
- 36) Kavanagh CJ, Trumbo PR, Ellwood KC. The US. Food and Drug Administration's

- evidence-based review for qualified health claims: tomatoes, lycopene, and cancer. *J Nat Cancer Inst* 2007; 99: 1074–85.
- 37) Van Patten CL, deBoer JG, Tomlinson Guns ES. Diet and dietary supplement intervention trials for the prevention of prostate cancer recurrence: a review of the randomized controlled trial evidence. *J Urol* 2008; 180: 2314–21.
 - 38) Chong EW, Wong TY, Kreis AJ, et al. Dietary antioxidants and primary prevention of age related macular degeneration: systematic review and meta-analysis. *BMJ* 2007; 335: 755.
 - 39) Leung IY-F. Macular pigment: new clinical methods of detection and the role of carotenoids in age-related macular degeneration. *Optometry* 2008; 79: 266–72.
 - 40) Stahl W, Sies H. Carotenoids and flavonoids contribute to nutritional protection against skin damage from sunlight. *Mol Biotechnol* 2007; 37: 26–30.
 - 41) Brustad M, Alsaker E, Engelsen O, et al. Vitamin D status of middle-aged women at 65–71 degrees N in relation to dietary intake and exposure to ultraviolet radiation. *Public Health Nutr* 2004; 7: 327–35.
 - 42) Holick MF. Vitamin D. In: Holick MF, Dawson-Hughes B, eds. *Nutrition and bone health*. Humana Press Inc, Totowa, 2004: 403–40.
 - 43) Malabanan A, Veronikis IE, Holick MF. Redefining vitamin D insufficiency. *Lancet* 1998; 351: 805–6.
 - 44) Jesudason D, Need AG, Horowitz M, et al. Relationship between serum 25-hydroxyvitamin D and bone resorption markers in vitamin D insufficiency. *Bone* 2002; 31: 626–30.
 - 45) Dawson-Hughes B, Heaney RP, Holick MF, et al. Estimates of optimal vitamin D status. *Osteoporosis Int* 2005; 24: 395–400.
 - 46) Nakamura K, Tsugawa N, Saito T, et al. Vitamin D status, bone mass, and bone metabolism in home-dwelling postmenopausal Japanese women: Yokogishi Study. *Bone* 2008; 42: 271–7.
 - 47) Krall EA, Sahyoun N, Tannenbaum S, et al. Effect of vitamin D intake on seasonal variations in parathyroid hormone secretion in postmenopausal women. *N Engl J Med* 1989; 32: 1777–83.
 - 48) Nakamura K, Nashimoto M, Tsuchiya Y, et al. Vitamin D insufficiency in Japanese female college students: a preliminary report. *Int J Vitam Nutr Res* 2001; 7: 302–5.
 - 49) Nakamura K, Nashimoto M, Matsuyama S, et al. Low serum concentrations of 25-hydroxyvitamin D in young adult Japanese women: a cross sectional study. *Nutrition* 2001; 17: 921–5.
 - 50) Nakamura K, Nashimoto M, Hori Y, et al. Serum 25-hydroxyvitamin D levels in active women of middle and advanced age in a rural community in Japan. *Nutrition* 1999; 15: 870–3.
 - 51) Nakamura K, Nashimoto M, Yamamoto M. Summer/winter differences in the serum 25-hydroxyvitamin D₃ and parathyroid hormone levels of Japanese women. *Int J Biometeorol* 2000; 44: 186–9.
 - 52) Nakamura K, Nashimoto M, Hori Y, et al. Serum 25-hydroxyvitamin D concentrations and related dietary factors in peri-and postmenopausal Japanese women. *Am J Clin Nutr*

- 2000; 71: 1161–5.
- 53) Nakamura K, Nashimoto M, Yamamoto M. Are the serum 25-hydroxyvitamin D concentrations in winter associated with forearm bone mineral density in healthy elderly Japanese women? *Int J Vitam Nutr Res* 2001; 71: 25–9.
- 54) 平成 15 年度厚生労働科学研究費補助金・老人保健健康増進等事業報告書：運動が骨密度に及ぼす影響に関する個人差の解明事業. 2004: 23–75.
- 55) 厚生労働省. 平成 17 年国民健康・栄養調査報告, 東京. 2007.
- 56) 厚生労働省. 平成 18 年国民健康・栄養調査報告, 東京. 2009.
- 57) 平成 18 年度厚生労働科学研究費補助金・循環器疾患等生活習慣病対策総合研究事業報告書：日本人の食事摂取基準（栄養所要量）の策定に関する研究—平成 18 年度総括分担研究報告書. 2007: 246–91.
- 58) Koenig J, Elmada I. Status of calcium and vitamin D of different population groups in Austria. *Int J Vitam Nutr Res* 2000; 70: 214–20.
- 59) Lehtonen-Veromaa MK, Mottonen TT, Nuotio IO, et al. Vitamin D and attainment of peak bone mass among peripubertal Finnish girls: a 3-y prospective study. *Am J Clin Nutr* 2002; 76: 1446–53.
- 60) Taha W, Chin D, Silverberg AI, et al. Reduced spinal bone mineral density in adolescents of an Ultra-Orthodox Jewish community in Brooklyn. *Pediatrics* 2001; 107: E79.
- 61) Yorifuji J, Yorifuki T, Tachibana K, et al. Craniotabes in normal newborns : the earliest sign of subclinical vitamin D deficiency. *J Clin Endocrinol Metab* 2008; 93: 1784–8.
- 62) Nakao H. Nutritional significance of human milk vitamin D in neonatal period. *Kobe J Med Sci* 1988; 34: 121–8.
- 63) Ziegler EE, Hollis BW, Nelson SE, et al. Vitamin D deficiency in breastfed infants in Iowa. *Pediatrics* 2006; 118: 603–10.
- 64) 文部科学省科学技術・学術審議会資源調査分科会報告. 五訂増補日本食品標準成分表. 国立印刷局. 東京. 2005.
- 65) Specker BL, Valanis B, Hertzberg V, et al. Sunshine exposure and serum 25-hydroxyvitamin D concentrations in exclusively breast-fed infants. *J Pediatrics* 1985; 107: 372–6.
- 66) Specker BL, Ho ML, Oestreich A, et al. Prospective study of vitamin D supplementation and rickets in China. *J Pediatr* 1992; 120: 733–9.
- 67) Leung SS, Liu S, Swaminathan R. Vitamin D status of Hong Kong Chinese infants. *Acta Paediatr Scand* 1989; 78: 303–6.
- 68) MacLennan WJ, Hamilton JC, Darmady JM. The effects of season and stage of pregnancy on plasma 25-hydroxyvitamin D concentrations in pregnant women. *Postgrad Med J* 1980; 56: 75–9.
- 69) Henriksen C, Brunvand L, Stoltenberg C, et al. Diet and vitamin D status among pregnant Pakistani women in Oslo. *Eur J Clin Nutr* 1995; 49: 211–8.
- 70) Narang NK, Gupta RC, Jain MK. Role of vitamin D in pulmonary tuberculosis. *J Assoc Physicians India* 1984; 32: 185–8.
- 71) Honkanen R, Alhava E, Parviainen M, et al. The necessity and safety of calcium and vitamin D in the elderly. *J Am Geriatr Soc* 1990; 38: 862–6.

- 72) Ala-Houhala M, Koskinen T, Terho A, et al. Maternal compared with infant vitamin D supplementation. *Arch Dis Child* 1986; 61: 1159–63.
- 73) Fomon SJ, Younoszai MK, Thomas LN. Influence of vitamin D on linear growth of normal full-term infants. *J Nutr* 1966; 88: 345–50.
- 74) Food and Nutrition Board, Institute of Medicine. Vitamin D. In: Institute of Medicine, ed. Dietary reference intakes for calcium, phosphorus, magnesium, vitamin D, and fluoride. National Academy Press, Washington D.C., 1998; 250–87.
- 75) Keller J, Losoasky MS. The absorption of alpha-tocopherol in man. *Br J Nutr* 1970; 24: 1033–47.
- 76) Blomstrand R, Forsgren L. Labeled tocopherols in man, intestinal absorption and thoracic-duct lymph transport of dl-alpha-tocopheryl-3,4-¹⁴C₂ acetate, dl-alpha-tocopheramine-3,4-¹⁴C₂, dl-alpha-tocopherol-(5-methyl-³H) and N-(methyl-³H)-dl-gamma-tocopheramine. *Z Vitaminforsch* 1968; 38: 328–44.
- 77) Traber MG, Arai H. Molecular mechanisms of vitamin E transport. *Ann Rev Nutr* 1999; 19: 343–55.
- 78) Horwitt MK, Century B, Zeman AA. Erythrocyte survival time and reticulocyte levels after tocopherol depletion in man. *Am J Clin Nutr* 1963; 12: 99–106.
- 79) Farrell PM, Bieri JG, Fratanton JF, et al. The occurrence and effects of human vitamin E deficiency. A study in patients with cystic fibrosis. *J Clin Invest* 1977; 60: 233–41.
- 80) Horwitt MK. Vitamin E and lipid metabolism in man. *Am J Clin Nutr* 1960; 8: 451–61.
- 81) Sasaki S, Ushio F, Amano K, et al. Serum biomarker-based validation of a self-administered diet history questionnaire for Japanese subjects. *J Nutr Sci Vitaminol* 2000; 46: 285–96.
- 82) Hiraoka M. Nutritional status of vitamin A, E, C, B₁, B₂, B₆, nicotinic acid, B₁₂, folate, and beta-carotene in young women. *J Nutr Sci Vitaminol* 2001; 47: 20–7.
- 83) Maruyama C, Imamura K, Oshima S, et al. Effects of tomato juice consumption on plasma and lipoprotein carotenoid concentrations and the susceptibility of low density lipoprotein to oxidative modification. *J Nutr Sci Vitaminol* 2001; 47: 213–21.
- 84) Jansson L, Akesson B, Holmberg L. Vitamin E and fatty acid composition of human milk. *Am J Clin Nutr* 1981; 34: 8–13.
- 85) Lammi-Keefe CJ, Jensen RG, Clark RM, et al. Alpha tocopherol, total lipid and linoleic acid contents of human milk at 2, 6, 12 and 16 weeks. In: Schaub J, eds. Composition and physiological properties of human milk. Elsevier Science, New York, 1985: 241–5.
- 86) Herrera E, Ortega H, Alvino G, et al. Relationship between plasma fatty acid profile and antioxidant vitamins during normal pregnancy. *Eur J Clin Nutr* 2004; 58: 1231–8.
- 87) Morinobu T, Ban R, Yoshikawa S, et al. The safety of high-dose vitamin E supplementation in healthy Japanese male adults. *J Nutr Sci Vitaminol* 2002; 48: 6–9.
- 88) Miller ER 3rd, Pastor-Barriuso R, Dalal D, et al. Meta-analysis: high-dosage vitamin E supplementation may increase all-cause mortality. *Ann Intern Med* 2005; 142: 37–46.
- 89) Bjelakovic G, Nikolova D, Gluud LL, et al. Mortality in randomized trials of antioxidant supplementations for primary and secondary prevention: systematic review and meta-

- analysis. *JAMA* 2005; 297: 842–57.
- 90) Asleh R, Blum S, Kalet-Litman S, et al. Correction of HDL dysfunction in individuals with diabetes and the haptoglobin 2–2 genotype. *Diabetes* 2008; 57: 2794–800.
- 91) Milman U, Blum S, Shapira C, et al. Vitamin E supplementation reduces cardiovascular events in a subgroup of middle-aged individuals with both type 2 diabetes mellitus and the haptoglobin 2–2 genotype: a prospective double-blinded clinical trial. *Arterioscler Thromb Vasc Biol* 2008; 28: 341–7.
- 92) Shearer MJ, Bach A, Kohlmeier M. Chemistry, nutritional sources, tissue distribution and metabolism of vitamin K with special reference to bone health. *J Nutr* 1996; 126: 1181S–6S.
- 93) Schurgers LJ, Vermeer C. Differential lipoprotein transport pathways of K-vitamins in healthy subjects. *Biochim Biophys Acta* 2002; 1570: 27–32.
- 94) Shearer MJ. Vitamin K. *Lancet* 1995; 345: 229–34.
- 95) Okano T, Shimomura Y, Yamane M, et al. Conversion of phylloquinone (vitamin K₁) into menaquinone-4 (vitamin K₂) in mice: two possible routes for menaquinone-4 accumulation in cerebra of mice. *J Biol Chem* 2008; 283: 11270–9.
- 96) Suttie JW, Mumma-Schendel LL, Sha DV, et al. Vitamin K deficiency from dietary vitamin K restriction in humans. *Am J Clin Nutr* 1988; 47: 475–80.
- 97) Frick PG, Riedler G, Brogli HJ. Dose response and minimal daily requirement for vitamin K in man. *Appl Physiol* 1967; 23: 387–9.
- 98) Suttie JW. The importance of menaquinones in human nutrition. *Annu Rev Nutr* 1995; 15: 399–417.
- 99) Merli GJ, Fink J. Vitamin K and thrombosis. *Vitam Horm* 2008; 78: 265–79.
- 100) Bügel S. Vitamin K and bone health in adult humans. *Vitam Horm* 2008; 78: 393–416.
- 101) Cranenburg EC, Schurgers LJ, Vermeer C. Vitamin K: the coagulation vitamin that became omnipotent. *Thromb Haemost* 2007; 98: 120–5.
- 102) Feskanich D, Weber P, Willett WC, et al. Vitamin K intake and hip fractures in women: a prospective study. *Am J Clin Nutr* 1999; 69: 74–9.
- 103) Booth SL, Tucker KL, Chen H, et al. Dietary vitamin K intakes are associated with hip fracture but not with bone mineral density in elderly men and women. *Am J Clin Nutr* 2000; 71: 1201–8.
- 104) Binkley NC, Krueger DC, Kawahara TN, et al. A high phylloquinone intake is required to achieve maximal osteocalcin gamma-carboxylation. *Am J Clin Nutr* 2002; 76: 1055–60.
- 105) Bügel S, Sørensen AD, Hels O, et al. Effect of phylloquinone supplementation on biochemical markers of vitamin K status and bone turnover in postmenopausal women. *Br J Nutr* 2007; 97: 373–80.
- 106) Cockayne S, Adamson J, Lanham-New S, et al. Vitamin K and the prevention of fractures: systematic review and meta-analysis of randomized controlled trials. *Arch Intern Med* 2006; 166: 1256–61.
- 107) Tsugawa N, Shiraki M, Suhara Y, et al. Vitamin K status of healthy Japanese women: age-related vitamin K requirement for gamma-carboxylation of osteocalcin. *Am J Clin Nutr*

- 2006; 83: 380–6.
- 108) Kojima T, Asoh M, Yamawaki N, et al. Vitamin K concentrations in the maternal milk of Japanese women. *Acta Paediatr* 2004; 93: 457–63.
 - 109) Shearer MJ, Rahim S, Barkhan P, et al. Plasma vitamin K₁ in mothers and their newborn babies. *Lancet* 1982; 2: 460–3.
 - 110) Greer FR, Mummah-Schendel LL, Marshall S, et al. Vitamin K₁ (phylloquinone) and vitamin K₂ (menaquinone) status in newborns during the first week of life. *Pediatrics* 1988; 81: 137–40.
 - 111) Puckett RM, Offringa M. Prophylactic vitamin K for vitamin K deficiency bleeding in neonates. *Cochrane Database Syst Rev* 2000: CD002776.
 - 112) 骨粗鬆症の予防と治療ガイドライン作成委員会. 骨粗鬆症の予防と治療ガイドライン（2006年版）. ライフサイエンス出版, 2006.