

◆第28表の1◆ ヘモグロビンA_{1c}の分布（男、年齢階級別）

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~69歳	(再掲) 70~74歳	(再掲) 75~79歳	(再掲) 80~84歳	(再掲) 85歳以上
	総 数 (実 数)	100.0 (2,100)	100.0 (161)	100.0 (263)	100.0 (268)	100.0 (402)	100.0 (510)	100.0 (496)	100.0 (262)	100.0 (238)	100.0 (144)	100.0 (77)	100.0 (37)
男	3.9%未満	0.2	0.6	0.4	0.0	0.2	0.4	0.0	0.4	0.0	0.0	0.0	0.0
	4.0~4.1	0.1	0.6	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	4.2~4.3	1.1	4.3	1.1	3.0	0.2	0.6	0.4	0.8	0.8	0.0	0.0	0.0
	4.4~4.5	2.2	7.5	6.5	2.6	0.7	0.6	1.0	0.4	0.4	1.4	0.0	5.4
	4.6~4.7	7.3	22.4	15.2	10.4	6.2	2.7	2.2	2.3	2.9	0.7	2.6	2.7
	4.8~4.9	13.5	24.2	19.4	16.0	13.4	10.8	8.5	11.1	5.9	15.3	5.2	5.4
	5.0~5.1	18.2	21.1	25.1	20.5	18.9	15.3	14.7	15.3	11.8	12.5	27.3	16.2
	5.2~5.3	19.0	15.5	17.1	20.1	23.4	19.2	16.9	20.2	18.9	17.4	15.6	5.4
	5.4~5.5	14.1	1.2	10.3	14.6	12.2	16.7	19.2	16.0	20.2	14.6	19.5	29.7
	5.6~5.7	6.7	1.2	1.9	4.5	6.5	8.8	10.1	6.9	8.8	11.8	11.7	8.1
	5.8~5.9	4.0	0.0	0.8	2.2	3.7	6.3	5.8	6.9	7.6	5.6	3.9	0.0
	6.0	1.8	0.0	0.4	0.4	2.2	1.2	4.2	0.8	4.2	4.2	2.6	8.1
	6.1	0.9	0.0	0.0	0.7	1.7	0.4	1.6	0.0	1.7	1.4	2.6	0.0
	6.2	1.0	0.0	0.0	0.0	1.0	1.6	1.8	1.5	1.3	2.1	1.3	5.4
	6.3	0.6	0.6	0.0	0.0	0.2	1.2	0.8	1.1	0.8	1.4	0.0	0.0
	6.4	0.9	0.0	0.0	0.0	0.2	2.0	1.6	2.7	2.5	0.0	2.6	0.0
	6.5	0.7	0.0	0.0	0.7	0.5	0.6	1.6	0.8	1.7	2.1	0.0	2.7
	6.6	0.6	0.0	0.0	0.7	0.2	0.6	1.2	1.1	0.8	2.1	0.0	2.7
	6.7	0.6	0.0	0.0	0.4	0.5	0.8	1.0	1.5	1.7	0.7	0.0	0.0
	6.8	0.2	0.0	0.0	0.0	0.5	0.2	0.2	0.0	0.4	0.0	0.0	0.0
	6.9	0.5	0.0	0.0	0.4	1.0	0.2	1.0	0.4	0.8	1.4	1.3	0.0
	7.0~7.1	0.7	0.0	0.0	0.0	0.5	1.4	1.2	1.1	1.7	0.7	0.0	2.7
	7.2~7.3	0.2	0.0	0.4	0.0	0.0	0.6	0.0	0.8	0.0	0.0	0.0	0.0
	7.4~7.5	0.6	0.0	0.0	0.0	0.2	1.4	1.0	1.5	1.7	0.7	0.0	0.0
	7.6~7.7	0.3	0.0	0.0	0.0	0.2	0.6	0.6	0.4	1.3	0.0	0.0	0.0
	7.8~7.9	0.5	0.0	0.0	0.0	1.0	0.6	0.6	0.8	0.0	1.4	0.0	2.7
	8.0~8.1	0.6	0.6	0.0	0.0	0.5	0.8	1.0	0.4	0.4	0.7	2.6	2.7
	8.2~8.3	0.4	0.0	0.0	0.4	0.5	0.8	0.2	1.1	0.0	0.7	0.0	0.0
	8.4~8.5	0.1	0.0	0.0	0.4	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0
	8.6~8.7	0.3	0.0	0.0	0.0	0.7	0.2	0.6	0.4	0.4	0.7	1.3	0.0
	8.8~8.9	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	9.0%以上	1.9	0.0	1.1	1.9	2.2	3.5	0.8	3.4	1.3	0.7	0.0	0.0
	6.1%以上(再掲)	11.6	1.2	1.5	5.6	12.2	17.5	16.9	19.1	18.5	16.7	11.7	18.9
平均 値 %		5.43	4.89	5.07	5.24	5.48	5.66	5.60	5.69	5.64	5.58	5.50	5.61
標準 偏 差		0.92	0.41	0.75	0.81	0.96	1.10	0.78	1.14	0.80	0.80	0.70	0.80

◆第28表の1◆ ヘモグロビンA_{1c}の分布（女、年齢階級別）

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~69歳	(再掲) 70~74歳	(再掲) 75~79歳	(再掲) 80~84歳	(再掲) 85歳以上
	総 数 (実 数)	100.0 (3,185)	100.0 (243)	100.0 (461)	100.0 (474)	100.0 (663)	100.0 (686)	100.0 (658)	100.0 (355)	100.0 (283)	100.0 (198)	100.0 (113)	100.0 (64)
女	3.9%未満	0.1	0.0	0.2	0.0	0.2	0.1	0.0	0.0	0.0	0.0	0.0	0.0
	4.0~4.1	0.2	0.0	0.4	0.0	0.3	0.0	0.2	0.0	0.0	0.5	0.0	0.0
	4.2~4.3	0.7	1.2	1.7	0.6	0.5	0.1	0.5	0.3	0.0	1.0	0.9	0.0
	4.4~4.5	2.6	9.5	4.3	3.6	1.8	0.7	0.8	0.3	0.4	1.0	1.8	0.0
	4.6~4.7	7.2	19.3	11.7	11.0	5.4	3.5	2.3	3.7	2.1	2.5	1.8	3.1
	4.8~4.9	12.7	28.0	18.0	15.6	11.3	6.9	8.7	6.2	10.2	7.1	8.8	6.3
	5.0~5.1	20.4	23.5	30.6	24.5	17.8	17.1	15.2	16.1	14.1	14.1	16.8	20.3
	5.2~5.3	19.6	10.7	16.9	19.0	23.7	19.5	21.3	19.7	20.5	17.7	24.8	29.7
	5.4~5.5	15.2	6.2	8.9	13.5	17.9	19.2	17.2	18.0	17.3	16.7	19.5	14.1
	5.6~5.7	8.2	1.2	4.1	5.1	9.0	11.5	11.6	13.5	12.0	10.1	11.5	14.1
	5.8~5.9	4.1	0.0	0.7	2.7	4.1	6.9	5.9	7.9	5.7	6.6	5.3	6.3
	6.0	1.6	0.0	0.2	1.3	2.1	2.0	2.4	1.7	1.4	5.6	0.9	0.0
	6.1	1.2	0.0	0.9	0.4	1.1	1.3	2.3	2.0	3.2	2.5	0.0	1.6
	6.2	0.6	0.0	0.2	0.6	0.3	1.3	0.8	2.3	0.7	1.5	0.0	0.0
	6.3	0.5	0.4	0.0	0.2	0.2	0.6	1.5	0.3	2.1	1.5	0.9	0.0
	6.4	0.4	0.0	0.0	0.0	0.3	0.9	0.6	0.6	0.7	1.0	0.0	0.0
	6.5	0.3	0.0	0.2	0.0	0.0	0.7	0.8	0.8	0.0	2.0	0.9	0.0
	6.6	0.4	0.0	0.0	0.0	0.6	0.7	0.5	0.3	0.7	0.5	0.0	0.0
	6.7	0.2	0.0	0.0	0.4	0.0	0.6	0.2	0.6	0.0	0.5	0.0	0.0
	6.8	0.3	0.0	0.0	0.0	0.2	0.4	1.1	0.6	0.7	1.5	0.9	1.6
	6.9	0.3	0.0	0.0	0.2	0.2	0.4	0.5	0.6	0.7	0.5	0.0	0.0
	7.0~7.1	0.6	0.0	0.2	0.0	0.3	1.2	1.1	1.4	0.7	1.5	1.8	0.0
	7.2~7.3	0.3	0.0	0.2	0.0	0.2	0.6	0.6	0.3	0.7	1.0	0.0	0.0
	7.4~7.5	0.3	0.0	0.0	0.0	0.3	0.4	0.5	0.3	1.1	0.0	0.0	0.0
	7.6~7.7	0.3	0.0	0.2	0.2	0.0	0.3	0.6	0.6	1.4	0.0	0.0	0.0
	7.8~7.9	0.3	0.0	0.0	0.2	0.2	0.7	0.5	0.3	0.7	0.0	0.0	1.6
	8.0~8.1	0.2	0.0	0.0	0.0	0.0	0.4	0.6	0.3	0.4	1.0	0.9	0.0
	8.2~8.3	0.2	0.0	0.0	0.0	0.2	0.3	0.5	0.3	0.0	1.0	0.9	0.0
	8.4~8.5	0.2	0.0	0.2	0.0	0.5	0.0	0.2	0.0	0.4	0.0	0.0	0.0
	8.6~8.7	0.2	0.0	0.0	0.0	0.2	0.4	0.2	0.3	0.0	0.5	0.0	0.0
	8.8~8.9	0.1	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0
	9.0%以上	0.9	0.0	0.0	0.8	1.5	0.7	1.5	1.1	2.1	0.5	1.8	1.6
	6.1%以上(再掲)	7.6	0.4	2.2	3.2	5.9	12.4	14.1	12.7	16.3	17.2	8.0	6.3
平均 値 %		5.34	4.91	5.05	5.19	5.38	5.53	5.57	5.54	5.62	5.58	5.51	5.43
標準 偏 差		0.78	0.29	0.42	0.66	0.89	0.80	0.91	0.81	0.97	0.76	1.07	0.69

◆第28表の2◆ ヘモグロビンA_{1c}の分布（全国、年齢階級別）

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~69歳	(再掲) 70~74歳	(再掲) 75~79歳	(再掲) 80~84歳	(再掲) 85歳以上
全 国	総 数 (実 数)	100.0 (5,070)	100.0 (404)	100.0 (719)	100.0 (730)	100.0 (1,029)	100.0 (1,126)	100.0 (1,062)	100.0 (575)	100.0 (478)	100.0 (309)	100.0 (179)	100.0 (96)
	3.9%未満	0.2	0.2	0.3	0.0	0.2	0.3	0.0	0.2	0.0	0.0	0.0	0.0
4.0~4.1	0.1	0.2	0.4	0.0	0.2	0.0	0.1	0.0	0.0	0.0	0.3	0.0	0.0
4.2~4.3	0.9	2.5	1.5	1.5	0.4	0.4	0.4	0.5	0.2	0.6	0.6	0.6	0.0
4.4~4.5	2.5	8.7	5.1	3.3	1.5	0.7	0.9	0.3	0.4	1.3	1.1	2.1	
4.6~4.7	7.5	20.5	13.1	10.8	5.9	3.4	2.4	3.3	2.7	1.9	2.2	3.1	
4.8~4.9	13.5	26.5	18.6	16.0	12.4	9.0	9.0	8.7	8.8	11.0	7.8	6.3	
5.0~5.1	20.2	22.5	28.7	23.3	18.7	17.3	16.0	16.9	14.0	14.9	21.2	19.8	
5.2~5.3	20.1	12.6	17.1	19.7	24.2	20.6	20.9	21.4	21.3	19.1	22.3	21.9	
5.4~5.5	15.3	4.2	9.3	14.0	16.3	19.2	19.4	18.3	20.3	17.2	20.1	20.8	
5.6~5.7	7.8	1.2	3.2	4.9	8.3	10.8	11.6	11.3	11.3	11.7	12.3	11.5	
5.8~5.9	4.1	0.0	0.7	2.3	4.0	6.9	6.3	7.8	6.9	6.8	5.0	4.2	
6.0	1.7	0.0	0.3	1.0	2.1	1.8	3.3	1.4	2.9	4.9	1.7	3.1	
6.1	1.1	0.0	0.6	0.5	1.4	1.0	2.1	1.2	2.5	2.3	1.1	1.0	
6.2	0.7	0.0	0.1	0.3	0.6	1.3	1.0	1.7	0.8	1.6	0.6	1.0	
6.3	0.4	0.5	0.0	0.1	0.2	0.7	0.8	0.3	1.0	1.0	0.0	0.0	
6.4	0.4	0.0	0.0	0.0	0.3	1.0	0.8	1.0	0.8	0.6	1.1	0.0	
6.5	0.4	0.0	0.1	0.3	0.2	0.7	0.8	0.9	0.6	1.6	0.6	0.0	
6.6	0.4	0.0	0.0	0.1	0.5	0.5	0.6	0.5	0.6	0.6	0.0	1.0	
6.7	0.2	0.0	0.0	0.3	0.0	0.4	0.4	0.7	0.6	0.3	0.0	0.0	
6.8	0.1	0.0	0.0	0.0	0.1	0.2	0.2	0.2	0.2	0.0	0.0	1.0	
6.9	0.3	0.0	0.0	0.3	0.5	0.4	0.2	0.5	0.2	0.3	0.0	0.0	
7.0~7.1	0.4	0.0	0.1	0.0	0.3	0.7	0.7	0.9	0.4	1.0	1.1	0.0	
7.2~7.3	0.1	0.0	0.3	0.0	0.0	0.4	0.1	0.2	0.2	0.0	0.0	0.0	
7.4~7.5	0.2	0.0	0.0	0.0	0.2	0.4	0.2	0.3	0.4	0.0	0.0	0.0	
7.6~7.7	0.1	0.0	0.0	0.0	0.0	0.2	0.4	0.3	0.8	0.0	0.0	0.0	
7.8~7.9	0.1	0.0	0.0	0.1	0.2	0.2	0.2	0.0	0.2	0.0	0.0	1.0	
8.0~8.1	0.2	0.2	0.0	0.0	0.1	0.4	0.4	0.2	0.2	0.3	0.6	1.0	
8.2~8.3	0.1	0.0	0.0	0.1	0.1	0.2	0.1	0.2	0.0	0.3	0.0	0.0	
8.4~8.5	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0	
8.6~8.7	0.1	0.0	0.0	0.0	0.1	0.0	0.2	0.0	0.2	0.3	0.0	0.0	
8.8~8.9	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	
9.0%以上	0.7	0.0	0.4	0.8	1.2	0.8	0.7	0.7	1.0	0.0	0.6	1.0	
6.1%以上(再掲)	6.1	0.7	1.7	3.2	5.8	9.7	9.6	9.9	11.1	10.4	5.6	7.3	
平均 値 %	5.29	4.90	5.05	5.17	5.34	5.45	5.46	5.45	5.51	5.44	5.39	5.44	
標準 偏 差	0.68	0.35	0.54	0.63	0.77	0.71	0.68	0.68	0.73	0.57	0.69	0.68	

注) インスリン注射又は血糖を下げる薬服用者除外。

◆第28表の2◆ ヘモグロビンA_{1c}の分布（男、年齢階級別）

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~69歳	(再掲) 70~74歳	(再掲) 75~79歳	(再掲) 80~84歳	(再掲) 85歳以上
	総 数 (実 数)	100.0 (1,986)	100.0 (161)	100.0 (262)	100.0 (262)	100.0 (384)	100.0 (469)	100.0 (448)	100.0 (236)	100.0 (216)	100.0 (128)	100.0 (71)	100.0 (33)
男	3.9%未満	0.3	0.6	0.4	0.0	0.3	0.4	0.0	0.4	0.0	0.0	0.0	0.0
	4.0~4.1	0.1	0.6	0.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	4.2~4.3	1.2	4.3	1.1	3.1	0.3	0.6	0.2	0.8	0.5	0.0	0.0	0.0
	4.4~4.5	2.4	7.5	6.5	2.7	0.8	0.6	1.1	0.4	0.5	1.6	0.0	6.1
	4.6~4.7	7.8	22.4	15.3	10.7	6.5	3.0	2.5	2.5	3.2	0.8	2.8	3.0
	4.8~4.9	14.1	24.2	19.5	16.4	13.8	11.5	9.2	11.9	6.5	16.4	5.6	6.1
	5.0~5.1	19.0	21.1	24.8	20.6	19.5	16.6	15.8	16.9	12.5	14.1	28.2	18.2
	5.2~5.3	20.0	15.5	17.2	20.6	24.5	20.9	18.3	22.5	20.4	18.8	16.9	6.1
	5.4~5.5	14.8	1.2	10.3	14.5	12.8	17.9	21.0	17.4	22.2	16.4	19.7	33.3
	5.6~5.7	6.9	1.2	1.9	4.6	6.8	9.2	10.9	7.2	9.3	13.3	12.7	9.1
	5.8~5.9	4.2	0.0	0.8	2.3	3.9	6.8	6.5	7.6	8.3	6.3	4.2	0.0
	6.0	1.9	0.0	0.4	0.4	2.3	1.3	4.5	0.8	4.6	3.9	2.8	9.1
	6.1	1.0	0.0	0.0	0.8	1.8	0.4	1.8	0.0	1.9	1.6	2.8	0.0
	6.2	1.0	0.0	0.0	0.0	1.0	1.7	1.6	1.7	1.4	1.6	1.4	3.0
	6.3	0.5	0.6	0.0	0.0	0.3	0.9	0.7	0.4	0.9	0.8	0.0	0.0
	6.4	0.6	0.0	0.0	0.0	0.3	1.3	1.1	1.7	1.4	0.0	2.8	0.0
	6.5	0.6	0.0	0.0	0.8	0.5	0.6	0.9	0.8	1.4	0.8	0.0	0.0
	6.6	0.4	0.0	0.0	0.4	0.3	0.6	0.7	1.3	0.5	0.8	0.0	3.0
	6.7	0.3	0.0	0.0	0.0	0.0	0.4	0.7	0.8	1.4	0.0	0.0	0.0
	6.8	—	—	—	—	—	—	—	—	—	—	—	—
	6.9	0.4	0.0	0.0	0.4	1.0	0.2	0.2	0.4	0.0	0.8	0.0	0.0
	7.0~7.1	0.4	0.0	0.0	0.0	0.5	0.9	0.4	0.8	0.5	0.8	0.0	0.0
	7.2~7.3	0.2	0.0	0.4	0.0	0.0	0.4	0.0	0.4	0.0	0.0	0.0	0.0
	7.4~7.5	0.3	0.0	0.0	0.0	0.3	0.6	0.2	0.8	0.5	0.0	0.0	0.0
	7.6~7.7	0.2	0.0	0.0	0.0	0.0	0.2	0.4	0.4	0.9	0.0	0.0	0.0
	7.8~7.9	0.1	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	8.0~8.1	0.3	0.6	0.0	0.0	0.3	0.6	0.2	0.4	0.0	0.0	0.0	3.0
	8.2~8.3	0.3	0.0	0.0	0.4	0.3	0.4	0.2	0.4	0.0	0.8	0.0	0.0
	8.4~8.5	0.1	0.0	0.0	0.4	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0
	8.6~8.7	0.2	0.0	0.0	0.0	0.3	0.0	0.4	0.0	0.5	0.8	0.0	0.0
	8.8~8.9	—	—	—	—	—	—	—	—	—	—	—	—
	9.0%以上	1.1	0.0	1.1	1.1	1.6	1.5	0.4	0.8	0.9	0.0	0.0	0.0
	6.1%以上(再掲)	7.4	1.2	1.5	4.2	8.6	11.1	10.0	11.4	12.0	8.6	7.0	9.1
平均 値 %		5.32	4.89	5.07	5.19	5.38	5.48	5.47	5.46	5.53	5.43	5.37	5.45
標準 偏 差		0.74	0.41	0.75	0.66	0.79	0.82	0.62	0.75	0.68	0.60	0.39	0.66

注) インスリン注射又は血糖を下げる薬服用者除外。

◆第28表の2◆ ヘモグロビンA_{1c}の分布（女、年齢階級別）

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~69歳	(再掲) 70~74歳	(再掲) 75~79歳	(再掲) 80~84歳	(再掲) 85歳以上
	総 数 (実 数)	100.0 (3,084)	100.0 (243)	100.0 (457)	100.0 (468)	100.0 (645)	100.0 (657)	100.0 (614)	100.0 (339)	100.0 (262)	100.0 (181)	100.0 (108)	100.0 (63)
女	3.9%未満	0.1	0.0	0.2	0.0	0.2	0.2	0.0	0.0	0.0	0.0	0.0	0.0
	4.0~4.1	0.2	0.0	0.4	0.0	0.3	0.0	0.2	0.0	0.0	0.6	0.0	0.0
	4.2~4.3	0.7	1.2	1.8	0.6	0.5	0.2	0.5	0.3	0.0	1.1	0.9	0.0
	4.4~4.5	2.7	9.5	4.4	3.6	1.9	0.8	0.8	0.3	0.4	1.1	1.9	0.0
	4.6~4.7	7.4	19.3	11.8	10.9	5.6	3.7	2.4	3.8	2.3	2.8	1.9	3.2
	4.8~4.9	13.0	28.0	18.2	15.8	11.6	7.2	9.0	6.5	10.7	7.2	9.3	6.3
	5.0~5.1	21.0	23.5	30.9	24.8	18.1	17.8	16.1	16.8	15.3	15.5	16.7	20.6
	5.2~5.3	20.2	10.7	17.1	19.2	24.0	20.4	22.8	20.6	22.1	19.3	25.9	30.2
	5.4~5.5	15.6	6.2	8.8	13.7	18.4	20.1	18.2	18.9	18.7	17.7	20.4	14.3
	5.6~5.7	8.3	1.2	3.9	5.1	9.1	12.0	12.1	14.2	13.0	10.5	12.0	12.7
	5.8~5.9	4.0	0.0	0.7	2.4	4.0	7.0	6.2	8.0	5.7	7.2	5.6	6.3
	6.0	1.6	0.0	0.2	1.3	2.0	2.1	2.4	1.8	1.5	5.5	0.9	0.0
	6.1	1.2	0.0	0.9	0.4	1.1	1.4	2.3	2.1	3.1	2.8	0.0	1.6
	6.2	0.5	0.0	0.2	0.4	0.3	1.1	0.7	1.8	0.4	1.7	0.0	0.0
	6.3	0.4	0.4	0.0	0.2	0.2	0.6	0.8	0.3	1.1	1.1	0.0	0.0
	6.4	0.3	0.0	0.0	0.0	0.3	0.8	0.5	0.6	0.4	1.1	0.0	0.0
	6.5	0.4	0.0	0.2	0.0	0.0	0.8	0.8	0.9	0.0	2.2	0.9	0.0
	6.6	0.3	0.0	0.0	0.0	0.6	0.5	0.5	0.0	0.8	0.6	0.0	0.0
	6.7	0.2	0.0	0.0	0.4	0.0	0.5	0.2	0.6	0.0	0.6	0.0	0.0
	6.8	0.2	0.0	0.0	0.0	0.2	0.3	0.3	0.3	0.4	0.0	0.0	1.6
	6.9	0.2	0.0	0.0	0.2	0.2	0.5	0.2	0.6	0.4	0.0	0.0	0.0
	7.0~7.1	0.4	0.0	0.2	0.0	0.2	0.6	0.8	0.9	0.4	1.1	1.9	0.0
	7.2~7.3	0.1	0.0	0.2	0.0	0.0	0.3	0.2	0.0	0.4	0.0	0.0	0.0
	7.4~7.5	0.1	0.0	0.0	0.0	0.2	0.2	0.2	0.0	0.4	0.0	0.0	0.0
	7.6~7.7	0.1	0.0	0.0	0.0	0.0	0.2	0.3	0.3	0.8	0.0	0.0	0.0
	7.8~7.9	0.2	0.0	0.0	0.2	0.2	0.3	0.3	0.0	0.4	0.0	0.0	1.6
	8.0~8.1	0.2	0.0	0.0	0.0	0.0	0.3	0.5	0.0	0.4	0.6	0.9	0.0
	8.2~8.3	—	—	—	—	—	—	—	—	—	—	—	—
	8.4~8.5	—	—	—	—	—	—	—	—	—	—	—	—
	8.6~8.7	—	—	—	—	—	—	—	—	—	—	—	—
	8.8~8.9	0.1	0.0	0.0	0.0	0.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	9.0%以上	0.5	0.0	0.0	0.6	0.9	0.3	0.8	0.6	1.1	0.0	0.9	1.6
	6.1%以上(再掲)	5.3	0.4	1.8	2.6	4.2	8.7	9.3	8.8	10.3	11.6	4.6	6.3
平均 値 %		5.27	4.91	5.04	5.17	5.32	5.44	5.46	5.45	5.49	5.45	5.41	5.43
標準 偏 差		0.65	0.29	0.37	0.60	0.75	0.62	0.72	0.62	0.78	0.54	0.83	0.69

注) インスリン注射又は血糖を下げる薬服用者除外。

◆第29表の1◆ 血糖値の分布（食後3時間以上、性・年齢階級別）

(%)

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~69歳	(再掲) 70~74歳	(再掲) 75~79歳	(再掲) 80~84歳	(再掲) 85歳以上
性別	総 数 (実 数)	100.0 (1,616)	100.0 (125)	100.0 (214)	100.0 (213)	100.0 (307)	100.0 (393)	100.0 (364)	100.0 (198)	100.0 (181)	100.0 (103)	100.0 (53)	100.0 (27)
	40 mg/dl 未満	—	—	—	—	—	—	—	—	—	—	—	—
男	40~ 59	0.1	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.6	0.0	0.0	0.0
	60~ 79	5.0	10.4	4.7	2.8	6.5	4.3	3.8	6.6	2.2	5.8	0.0	14.8
	80~ 99	50.5	70.4	75.2	59.2	50.8	38.9	36.3	35.9	37.0	44.7	28.3	14.8
	100~ 119	29.3	16.0	16.4	28.2	29.0	34.9	36.3	36.4	38.1	30.1	47.2	25.9
	120~ 139	7.4	2.4	1.9	4.7	7.8	10.2	10.7	9.1	9.9	7.8	15.1	18.5
	140~ 159	2.5	0.8	0.0	2.8	1.3	2.8	5.2	1.5	3.9	5.8	3.8	14.8
	160~ 179	1.2	0.0	0.0	0.5	1.3	2.5	1.1	4.5	0.6	1.9	0.0	3.7
	180~ 199	1.2	0.0	0.0	0.5	1.3	1.3	2.5	1.0	3.9	0.0	3.8	0.0
	200~ 219	1.1	0.0	0.0	0.9	0.7	2.0	1.4	2.5	1.7	1.9	0.0	0.0
	220~ 239	0.2	0.0	0.0	0.5	0.0	0.3	0.3	0.5	0.6	0.0	0.0	0.0
	240~ 259	0.3	0.0	0.0	0.0	1.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0
	260 mg/dl 以上	1.4	0.0	1.9	0.0	0.3	2.3	2.2	2.0	1.7	1.9	1.9	7.4
	110 以上(再掲)	24.1	9.6	6.5	16.9	19.9	32.1	38.7	30.8	37.6	33.0	43.4	59.3
	平均値 mg/dl	106.6	92.7	97.2	101.4	104.0	113.6	114.5	112.5	114.3	110.0	116.2	129.9
	標準偏差	36.7	12.3	30.0	21.8	29.6	46.3	42.5	41.0	41.1	37.2	41.2	66.3
		総 数	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
性別	総 数 (実 数)	100.0 (2,280)	100.0 (177)	100.0 (316)	100.0 (335)	100.0 (480)	100.0 (500)	100.0 (472)	100.0 (249)	100.0 (200)	100.0 (142)	100.0 (85)	100.0 (45)
	40 mg/dl 未満	—	—	—	—	—	—	—	—	—	—	—	—
女	40~ 59	—	—	—	—	—	—	—	—	—	—	—	—
	60~ 79	4.2	6.8	10.1	4.5	4.4	1.6	1.7	1.6	2.5	0.7	0.0	4.4
	80~ 99	51.0	80.8	68.7	65.7	52.7	37.0	30.5	35.3	34.5	33.1	20.0	24.4
	100~ 119	32.1	10.2	18.4	25.1	33.1	42.4	42.6	43.8	43.0	39.4	43.5	48.9
	120~ 139	7.2	2.3	2.5	2.7	6.3	9.4	14.0	9.6	10.0	17.6	20.0	8.9
	140~ 159	2.8	0.0	0.3	1.2	1.5	5.2	5.5	5.6	5.5	4.2	8.2	4.4
	160~ 179	1.0	0.0	0.0	0.6	0.6	1.2	2.5	1.2	2.5	1.4	3.5	4.4
	180~ 199	0.5	0.0	0.0	0.0	0.0	1.0	1.3	0.8	0.5	2.1	1.2	2.2
	200~ 219	0.4	0.0	0.0	0.0	0.8	0.4	0.6	0.4	0.0	0.7	1.2	2.2
	220~ 239	0.1	0.0	0.0	0.3	0.2	0.2	0.0	0.4	0.0	0.0	0.0	0.0
	240~ 259	0.1	0.0	0.0	0.0	0.0	0.2	0.2	0.0	0.5	0.0	0.0	0.0
	260 mg/dl 以上	0.6	0.0	0.0	0.0	0.4	1.4	1.1	1.2	1.0	0.7	2.4	0.0
	110 以上(再掲)	23.3	5.1	7.3	12.8	19.4	34.8	40.3	37.8	34.5	41.5	52.9	37.8
	平均値 mg/dl	103.5	91.9	92.8	97.2	101.8	110.7	113.6	110.5	110.7	112.1	123.5	112.1
	標準偏差	26.8	10.6	11.5	14.9	25.3	31.1	34.4	28.8	32.3	26.4	50.1	25.9

◆第29表の2 ◆ 血糖値の分布（食後3時間以上、性・年齢階級別）

(%)

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~69歳	(再掲) 70~74歳	(再掲) 75~79歳	(再掲) 80~84歳	(再掲) 85歳以上
男	総 数 (実 数)	100.0 (1,528)	100.0 (125)	100.0 (213)	100.0 (208)	100.0 (294)	100.0 (364)	100.0 (324)	100.0 (180)	100.0 (163)	100.0 (89)	100.0 (48)	100.0 (24)
	40 mg/dl 未満	—	—	—	—	—	—	—	—	—	—	—	—
男	40~ 59	0.1	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.6	0.0	0.0	0.0
	60~ 79	4.9	10.4	4.7	2.4	6.8	4.4	3.4	6.7	2.5	3.4	0.0	16.7
	80~ 99	52.4	70.4	75.1	59.6	52.4	41.2	38.6	37.8	38.7	49.4	29.2	16.7
	100~ 119	30.2	16.0	16.4	28.8	29.3	36.8	38.9	38.9	39.9	33.7	50.0	29.2
	120~ 139	7.0	2.4	1.9	4.8	7.1	10.2	9.9	9.4	9.2	6.7	16.7	12.5
	140~ 159	2.4	0.8	0.0	2.4	1.4	2.2	5.6	1.1	4.3	5.6	4.2	16.7
	160~ 179	0.9	0.0	0.0	0.5	1.0	1.6	0.9	3.3	0.6	1.1	0.0	4.2
	180~ 199	0.7	0.0	0.0	0.5	0.7	1.4	0.9	1.1	1.8	0.0	0.0	0.0
	200~ 219	0.6	0.0	0.0	1.0	0.3	1.1	0.6	1.1	1.2	0.0	0.0	0.0
	220~ 239	—	—	—	—	—	—	—	—	—	—	—	—
	240~ 259	0.2	0.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	260 mg/dl 以上	0.7	0.0	1.9	0.0	0.0	1.1	0.9	0.6	1.2	0.0	0.0	4.2
	110 以上(再掲)	21.5	9.6	6.6	16.3	17.7	28.6	34.9	27.2	33.7	29.2	39.6	54.2
平均値 mg/dl		103.2	92.7	97.2	100.7	101.7	108.0	109.0	106.5	110.5	103.8	108.4	118.7
標準偏差		29.0	12.3	30.1	19.5	23.6	36.1	31.5	27.8	37.8	17.4	16.5	43.6
女	総 数 (実 数)	100.0 (2,213)	100.0 (177)	100.0 (315)	100.0 (331)	100.0 (468)	100.0 (477)	100.0 (445)	100.0 (237)	100.0 (188)	100.0 (132)	100.0 (81)	100.0 (44)
	40 mg/dl 未満	—	—	—	—	—	—	—	—	—	—	—	—
女	40~ 59	—	—	—	—	—	—	—	—	—	—	—	—
	60~ 79	4.3	6.8	10.2	4.5	4.5	1.7	1.6	1.7	2.7	0.0	0.0	4.5
	80~ 99	52.2	80.8	68.9	66.5	53.6	38.6	31.5	36.7	35.1	34.8	21.0	25.0
	100~ 119	32.4	10.2	18.1	24.5	33.8	43.6	43.8	45.6	44.7	41.7	43.2	47.7
	120~ 139	7.1	2.3	2.5	2.4	6.2	9.6	14.2	10.1	9.6	18.2	21.0	9.1
	140~ 159	2.4	0.0	0.3	1.2	1.3	4.4	4.9	4.2	5.3	3.0	7.4	4.5
	160~ 179	0.8	0.0	0.0	0.6	0.2	0.8	2.2	0.8	2.1	0.8	3.7	4.5
	180~ 199	0.2	0.0	0.0	0.0	0.0	0.4	0.4	0.4	0.0	0.0	1.2	2.3
	200~ 219	0.2	0.0	0.0	0.0	0.2	0.0	0.7	0.0	0.0	0.8	1.2	2.3
	220~ 239	0.1	0.0	0.0	0.3	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	240~ 259	0.1	0.0	0.0	0.0	0.0	0.2	0.2	0.0	0.5	0.0	0.0	0.0
	260 mg/dl 以上	0.2	0.0	0.0	0.0	0.0	0.6	0.4	0.4	0.0	0.8	1.2	0.0
	110 以上(再掲)	21.8	5.1	7.3	12.4	17.9	32.7	38.2	35.4	32.4	38.6	50.6	38.6
平均値 mg/dl		101.5	91.9	92.7	96.9	99.6	107.4	110.8	107.0	107.6	109.7	119.2	112.2
標準偏差		20.3	10.6	11.5	14.9	16.0	23.6	25.3	21.3	21.5	23.1	33.7	26.2

注) インスリン注射又は血糖を下げる薬服用者除外。

◆第30表の1◆ 総コレステロール値の分布（性・年齢階級別）

(%)

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~69歳	(再掲) 70~74歳	(再掲) 75~79歳	(再掲) 80~84歳	(再掲) 85歳以上
性	総 数 (実 数)	100.0 (2,108)	100.0 (162)	100.0 (263)	100.0 (270)	100.0 (401)	100.0 (513)	100.0 (499)	100.0 (264)	100.0 (239)	100.0 (144)	100.0 (79)	100.0 (37)
	120 mg/dl 未満	0.7	1.9	0.0	0.4	0.0	1.4	0.6	0.8	0.4	1.4	0.0	0.0
男	120~139	2.8	8.0	1.5	2.6	1.2	2.5	3.4	1.9	2.9	3.5	3.8	5.4
	140~159	9.3	17.3	8.4	3.7	7.2	7.8	13.4	8.3	9.2	14.6	21.5	18.9
	160~179	18.2	23.5	21.7	19.3	12.5	12.1	24.8	12.5	25.9	23.6	22.8	27.0
	180~199	23.1	23.5	21.7	19.6	21.7	26.7	22.8	26.9	23.8	22.2	24.1	16.2
	200~219	20.8	17.9	19.4	23.3	21.7	23.6	17.6	22.0	18.0	18.8	12.7	21.6
	220~239	13.7	5.6	13.7	14.8	18.2	14.2	11.4	14.8	12.1	11.8	8.9	10.8
	240~259	6.5	1.2	6.8	9.3	11.0	6.2	3.4	8.0	3.8	2.8	5.1	0.0
	260~279	3.3	1.2	4.2	5.6	4.0	3.7	1.4	3.0	2.5	0.7	0.0	0.0
	280~299	1.0	0.0	1.9	0.7	1.2	1.0	1.0	1.1	1.3	0.7	1.3	0.0
	300 以上	0.6	0.0	0.8	0.7	1.2	0.8	0.0	0.8	0.0	0.0	0.0	0.0
	220 以上(再掲)	25.2	8.0	27.4	31.1	35.7	25.9	17.2	27.7	19.7	16.0	15.2	10.8
	240 以上(再掲)	11.5	2.5	13.7	16.3	17.5	11.7	5.8	12.9	7.5	4.2	6.3	0.0
平均値 mg/dl		198.0	178.8	201.4	204.8	207.8	200.5	188.3	201.6	192.2	186.7	183.6	180.1
標準偏差		35.0	30.3	35.8	35.4	34.9	34.2	32.1	33.6	32.7	31.7	31.7	27.7
女	総 数 (実 数)	100.0 (3,190)	100.0 (243)	100.0 (463)	100.0 (476)	100.0 (662)	100.0 (690)	100.0 (656)	100.0 (358)	100.0 (283)	100.0 (196)	100.0 (113)	100.0 (64)
	120 mg/dl 未満	0.2	0.8	0.2	0.2	0.0	0.1	0.3	0.0	0.0	0.5	0.0	1.6
	120~139	1.4	8.6	1.7	1.3	0.3	0.6	0.6	0.8	0.7	0.5	0.9	0.0
	140~159	6.6	15.2	15.8	9.7	2.1	2.8	3.5	3.4	2.8	2.0	5.3	7.8
	160~179	14.8	27.6	24.0	17.0	10.4	8.7	13.0	8.7	9.5	14.3	18.6	14.1
	180~199	21.3	22.2	27.0	25.6	16.5	18.1	22.0	19.6	21.6	21.9	23.0	21.9
	200~219	22.5	14.4	17.5	21.2	24.2	23.5	27.3	23.5	27.9	26.0	27.4	28.1
	220~239	16.2	7.8	7.6	14.1	17.5	21.4	20.3	23.2	20.5	22.4	19.5	14.1
	240~259	9.4	2.1	4.8	5.0	16.9	12.6	7.6	11.7	9.2	7.7	4.4	6.3
	260~279	4.8	0.0	0.9	4.4	7.9	8.0	3.4	6.7	5.3	3.1	0.9	0.0
	280~299	1.8	0.4	0.6	1.1	2.7	2.9	1.4	1.7	0.7	1.5	0.0	6.3
	300 以上	0.9	0.8	0.0	0.4	1.5	1.3	0.8	0.8	1.8	0.0	0.0	0.0
220 以上(再掲)		33.1	11.1	13.8	25.0	46.5	46.2	33.4	44.1	37.5	34.7	24.8	26.6
240 以上(再掲)		16.9	3.3	6.3	10.9	29.0	24.8	13.1	20.9	17.0	12.2	5.3	12.5
平均値 mg/dl		206.3	180.6	187.9	199.7	219.5	218.2	207.8	215.6	212.6	207.0	199.7	202.9
標準偏差		35.5	31.8	29.7	34.7	34.3	34.6	31.3	32.2	32.9	28.9	26.2	36.0

◆第30表の2 ◆ 総コレステロール値の分布（性・年齢階級別）

(%)

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~69歳	(再掲) 70~74歳	(再掲) 75~79歳	(再掲) 80~84歳	(再掲) 85歳以上
	総 数 (実 数)	100.0 (1,954)	100.0 (161)	100.0 (260)	100.0 (264)	100.0 (378)	100.0 (461)	100.0 (430)	100.0 (230)	100.0 (198)	100.0 (129)	100.0 (69)	100.0 (34)
男	120 mg/dl 未満	0.7	1.9	0.0	0.4	0.0	1.5	0.7	0.9	0.5	1.6	0.0	0.0
	120~139	2.9	8.1	1.5	2.7	1.3	2.6	3.7	2.2	3.0	3.9	4.3	5.9
	140~159	9.4	17.4	8.1	3.4	7.1	8.2	14.0	8.7	9.1	15.5	21.7	20.6
	160~179	18.3	23.6	21.9	19.3	12.7	12.1	24.9	12.6	26.8	23.3	20.3	29.4
	180~199	22.9	23.0	21.9	19.3	21.7	25.4	24.0	24.8	24.7	24.0	24.6	17.6
	200~219	21.1	18.0	18.8	23.9	22.5	24.3	17.2	22.6	18.2	17.8	14.5	14.7
	220~239	13.5	5.6	13.8	15.2	18.0	13.9	10.9	14.8	11.6	11.6	7.2	11.8
	240~259	6.7	1.2	6.9	9.1	11.1	6.5	3.3	8.3	4.0	1.6	5.8	0.0
	260~279	3.1	1.2	4.2	5.3	3.4	3.7	0.7	3.0	1.5	0.0	0.0	0.0
	280~299	0.9	0.0	1.9	0.8	0.8	1.1	0.7	1.3	0.5	0.8	1.4	0.0
	300 以上	0.6	0.0	0.8	0.8	1.3	0.7	0.0	0.9	0.0	0.0	0.0	0.0
	220 以上(再掲)	24.8	8.1	27.7	31.1	34.7	25.8	15.6	28.3	17.7	14.0	14.5	11.8
	240 以上(再掲)	11.3	2.5	13.8	15.9	16.7	11.9	4.7	13.5	6.1	2.3	7.2	0.0
平均値 mg/dl		197.6	178.7	201.6	204.9	207.2	200.1	186.7	201.7	190.5	184.5	183.9	178.1
標準偏差		34.80	30.4	35.8	35.2	34.4	34.3	30.9	34.4	30.7	30.6	32.3	28.1
女	総 数 (実 数)	100.0 (2,842)	100.0 (242)	100.0 (458)	100.0 (467)	100.0 (603)	100.0 (559)	100.0 (513)	100.0 (288)	100.0 (215)	100.0 (145)	100.0 (95)	100.0 (58)
	120 mg/dl 未満	0.2	0.8	0.2	0.2	0.0	0.2	0.4	0.0	0.0	0.7	0.0	1.7
	120~139	1.6	8.7	1.7	1.3	0.3	0.7	0.8	1.0	0.9	0.7	1.1	0.0
	140~159	6.9	15.3	15.9	9.6	1.5	2.3	3.5	3.1	2.8	2.1	6.3	5.2
	160~179	15.4	27.7	24.2	17.3	10.3	8.1	14.2	8.0	11.2	15.9	17.9	15.5
	180~199	21.0	22.3	26.9	25.5	16.7	16.8	20.7	18.1	19.5	19.3	24.2	22.4
	200~219	21.9	14.5	17.7	21.6	24.2	22.9	25.5	22.9	25.6	22.1	28.4	29.3
	220~239	15.9	7.4	7.2	14.1	17.7	22.5	19.9	24.3	20.9	23.4	15.8	13.8
	240~259	9.6	2.1	4.6	4.7	16.9	13.2	9.4	12.5	12.1	9.7	5.3	5.2
	260~279	4.8	0.0	0.9	4.3	8.0	8.6	3.3	7.3	4.7	4.1	1.1	0.0
	280~299	1.8	0.4	0.7	0.9	2.7	3.2	1.6	2.1	0.5	2.1	0.0	6.9
	300 以上	0.9	0.8	0.0	0.4	1.7	1.4	0.8	0.7	1.9	0.0	0.0	0.0
	220 以上(再掲)	33.0	10.7	13.3	24.4	46.9	49.0	34.9	46.9	40.0	39.3	22.1	25.9
	240 以上(再掲)	17.1	3.3	6.1	10.3	29.2	26.5	15.0	22.6	19.1	15.9	6.3	12.1
平均値 mg/dl		205.9	180.4	187.5	199.2	220.1	220.1	208.0	217.1	213.2	208.4	198.5	203.4
標準偏差		36.1	31.7	29.6	34.3	34.2	35.2	32.8	32.7	34.3	31.6	26.6	36.1

注) コレステロールを下げる薬服用者除外。

◆第31表の1◆ トリグリセリド値の分布（性・年齢階級別）

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~69歳	(再掲) 70~74歳	(再掲) 75~79歳	(再掲) 80~84歳	(再掲) 85歳以上
男	総 数 (実 数)	100.0 (2,108)	100.0 (162)	100.0 (263)	100.0 (270)	100.0 (401)	100.0 (513)	100.0 (499)	100.0 (264)	100.0 (239)	100.0 (144)	100.0 (79)	100.0 (37)
	20 mg/dl 未満	—	—	—	—	—	—	—	—	—	—	—	—
男	20~49	3.2	10.5	2.7	3.0	1.5	1.9	3.8	1.9	0.8	6.9	5.1	8.1
	50~79	17.3	31.5	20.5	13.0	12.0	12.7	22.2	13.3	19.7	27.1	24.1	16.2
	80~109	20.7	22.2	16.7	20.7	20.0	18.1	25.7	17.8	25.9	19.4	29.1	40.5
	110~139	16.8	11.7	18.3	11.1	16.2	22.0	15.8	20.1	16.3	18.1	13.9	8.1
	140~169	12.0	8.6	12.9	10.7	13.7	11.9	11.8	12.1	10.9	11.1	16.5	10.8
	170~199	8.0	4.9	6.8	10.4	9.7	9.4	5.6	11.0	5.9	4.9	3.8	10.8
	200 以上	22.1	10.5	22.1	31.1	26.9	24.0	15.0	23.9	20.5	12.5	7.6	5.4
	150 以上(再掲)	37.5	22.2	36.5	48.1	45.9	40.9	26.9	42.8	31.8	22.9	21.5	21.6
	平均値 mg/dl	158.0	117.9	162.7	186.2	176.2	164.6	131.9	164.8	147.2	120.7	116.6	110.1
	標準偏差	118.2	90.7	138.2	139.7	130.9	113.1	83.7	115.3	97.5	68.2	66.3	53.6
女	総 数 (実 数)	100.0 (3,190)	100.0 (243)	100.0 (463)	100.0 (476)	100.0 (662)	100.0 (690)	100.0 (656)	100.0 (358)	100.0 (283)	100.0 (196)	100.0 (113)	100.0 (64)
	20 mg/dl 未満	—	—	—	—	—	—	—	—	—	—	—	—
女	20~49	6.5	19.8	12.7	8.8	4.1	2.2	2.3	2.2	2.1	2.0	1.8	4.7
	50~79	22.7	39.1	35.0	30.3	16.9	11.7	20.0	11.7	16.6	20.4	23.9	26.6
	80~109	23.6	20.6	26.8	22.1	26.9	23.5	20.3	24.0	19.4	16.8	30.1	17.2
	110~139	17.1	10.3	11.4	15.5	19.0	19.1	20.4	19.3	20.8	20.9	16.8	23.4
	140~169	10.9	3.7	5.6	6.9	14.5	15.1	12.3	15.4	14.5	13.3	8.0	7.8
	170~199	6.0	2.5	2.8	3.4	5.4	8.4	9.6	8.1	8.5	11.2	9.7	9.4
	200 以上	13.2	4.1	5.6	13.0	13.1	20.0	15.1	19.3	18.0	15.3	9.7	10.9
	150 以上(再掲)	26.1	8.6	11.7	20.0	27.2	38.4	33.1	38.0	35.3	35.2	27.4	26.6
	平均値 mg/dl	126.9	85.4	97.4	123.7	132.2	149.1	136.5	148.6	143.1	138.2	122.2	127.2
	標準偏差	90.4	50.2	67.8	133.1	88.6	82.3	76.2	81.6	79.4	75.7	66.3	76.7

◆第31表の2◆ トリグリセリド値の分布（性・年齢階級別）

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~69歳	(再掲) 70~74歳	(再掲) 75~79歳	(再掲) 80~84歳	(再掲) 85歳以上
男	総 数 (実 数)	100.0 (1,954)	100.0 (161)	100.0 (260)	100.0 (264)	100.0 (378)	100.0 (461)	100.0 (430)	100.0 (230)	100.0 (198)	100.0 (129)	100.0 (69)	100.0 (34)
	20 mg/dl 未満	—	—	—	—	—	—	—	—	—	—	—	—
男	20~49	3.3	10.6	2.7	3.0	1.3	2.0	4.4	1.7	1.0	7.8	5.8	8.8
	50~79	18.0	31.7	20.8	12.9	12.7	13.9	23.3	14.8	20.7	27.1	26.1	17.6
	80~109	20.2	22.4	16.2	20.1	19.8	18.4	24.2	17.8	23.7	19.4	27.5	38.2
	110~139	17.0	11.8	18.5	11.4	16.1	21.9	17.2	19.6	18.7	18.6	14.5	8.8
	140~169	11.9	8.1	13.1	11.0	12.7	11.5	13.0	11.7	12.6	12.4	15.9	11.8
	170~199	8.1	5.0	6.5	10.6	9.5	9.5	6.0	11.7	6.1	5.4	4.3	11.8
	200 以上	21.4	10.6	22.3	31.1	27.8	22.8	11.9	22.6	17.2	9.3	5.8	2.9
	150 以上(再掲)	36.9	21.7	36.5	48.5	46.0	39.5	24.9	41.3	29.3	20.9	21.7	20.6
	平均値 mg/dl	155.6	117.6	163.2	186.1	176.2	159.9	123.6	161.2	136.0	114.9	112.9	106.7
	標準偏差	116.0	90.9	138.8	139.0	131.3	107.0	68.1	112.2	75.6	59.9	62.9	47.8
女	総 数 (実 数)	100.0 (2,842)	100.0 (242)	100.0 (458)	100.0 (467)	100.0 (603)	100.0 (559)	100.0 (513)	100.0 (288)	100.0 (215)	100.0 (145)	100.0 (95)	100.0 (58)
	20 mg/dl 未満	—	—	—	—	—	—	—	—	—	—	—	—
女	20~49	7.1	19.8	12.7	9.0	4.3	2.1	2.9	2.1	2.8	2.8	2.1	5.2
	50~79	23.9	38.8	35.2	30.4	17.2	12.2	21.4	12.2	17.7	22.1	24.2	29.3
	80~109	23.5	20.7	26.9	21.8	26.2	23.3	20.3	23.3	17.2	18.6	30.5	19.0
	110~139	16.6	10.3	11.6	15.8	19.1	18.6	19.5	19.1	20.5	18.6	17.9	20.7
	140~169	10.8	3.7	5.7	6.6	14.8	15.2	13.1	14.9	15.3	13.8	9.5	8.6
	170~199	5.6	2.5	2.8	3.4	5.6	8.2	8.8	7.3	8.8	9.7	6.3	10.3
	200 以上	12.6	4.1	5.2	12.8	12.8	20.4	14.0	21.2	17.7	14.5	9.5	6.9
	150 以上(再掲)	24.9	8.7	11.4	19.7	27.2	39.4	31.2	39.6	34.9	32.4	25.3	24.1
	平均値 mg/dl	123.9	85.5	95.8	122.5	131.7	148.9	132.1	151.3	141.2	133.9	117.3	118.4
	標準偏差	89.4	50.3	60.7	132.1	88.8	81.2	73.1	84.9	78.5	74.8	55.2	69.3

注) コレステロールを下げる薬服用者除外。

◆第32表の1◆ HDL-コレステロール値の分布 (性・年齢階級別)

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~69歳	(再掲) 70~74歳	(再掲) 75~79歳	(再掲) 80~84歳	(再掲) 85歳以上
男	総 数 (実 数)	100.0 (2,108)	100.0 (162)	100.0 (263)	100.0 (270)	100.0 (401)	100.0 (513)	100.0 (499)	100.0 (264)	100.0 (239)	100.0 (144)	100.0 (79)	100.0 (37)
	30 mg/dl 未満	1.4	0.0	1.1	1.5	0.5	1.9	2.0	2.7	2.9	0.7	2.5	0.0
	30~39	11.8	4.9	8.4	11.5	12.7	12.9	14.2	13.6	16.7	13.2	7.6	16.2
	40~49	23.7	17.3	21.7	22.6	23.7	25.7	25.5	26.5	25.1	27.8	24.1	21.6
	50~59	26.9	31.5	30.8	27.8	24.7	25.5	26.1	27.3	23.0	28.5	26.6	35.1
	60~69	18.1	26.5	17.5	17.8	18.2	18.1	15.8	17.8	15.9	12.5	21.5	16.2
	70~79	11.1	13.0	11.4	7.8	13.0	9.9	11.6	7.6	10.0	11.8	16.5	10.8
	80~89	4.2	3.7	5.3	4.8	5.2	3.9	3.0	2.3	4.2	2.8	1.3	0.0
	90~99	2.0	2.5	3.4	4.8	1.0	1.4	1.2	1.9	2.1	0.7	0.0	0.0
	100 以上	0.8	0.6	0.4	1.5	1.0	0.6	0.6	0.4	0.0	2.1	0.0	0.0
平均値 mg/dl	86.8	95.1	90.5	87.0	86.8	85.2	83.8	83.7	80.3	86.1	89.9	83.8	
	標準偏差	15.2	13.4	15.2	16.7	15.1	15.1	14.9	14.5	15.0	16.4	12.9	11.8
女	総 数 (実 数)	100.0 (3,190)	100.0 (243)	100.0 (463)	100.0 (476)	100.0 (662)	100.0 (690)	100.0 (656)	100.0 (358)	100.0 (283)	100.0 (196)	100.0 (113)	100.0 (64)
	30 mg/dl 未満	0.2	0.0	0.0	0.0	0.2	0.6	0.2	0.6	0.0	0.5	0.0	0.0
	30~39	3.4	0.8	1.1	2.5	2.3	4.2	6.7	5.6	6.4	7.7	6.2	6.3
	40~49	13.2	7.4	7.3	9.5	10.9	19.3	18.3	20.9	17.7	16.8	21.2	20.3
	50~59	22.8	18.5	17.5	21.4	22.1	26.4	26.1	25.1	25.4	26.0	28.3	25.0
	60~69	25.8	28.4	28.5	26.9	28.1	23.3	22.4	26.3	23.7	20.9	19.5	26.6
	70~79	17.9	24.3	23.1	21.2	15.9	13.5	16.2	10.9	17.3	14.8	17.7	12.5
	80~89	10.5	11.9	13.0	11.1	14.0	8.3	6.4	6.4	4.6	8.7	5.3	9.4
	90~99	4.2	5.8	5.0	5.5	4.2	3.2	3.0	3.4	3.5	4.1	1.8	0.0
	100 以上	2.1	2.9	4.5	1.9	2.4	1.3	0.8	0.8	1.4	0.5	0.0	0.0
平均値 mg/dl	96.5	99.2	98.9	97.5	97.6	95.2	93.1	93.9	93.6	91.8	93.8	93.8	
	標準偏差	15.6	14.2	15.4	15.0	15.5	15.5	14.8	14.9	15.2	15.4	13.6	13.5

◆第32表の2◆ HDL-コレステロール値の分布 (性・年齢階級別)

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~69歳	(再掲) 70~74歳	(再掲) 75~79歳	(再掲) 80~84歳	(再掲) 85歳以上
男	総 数 (実 数)	100.0 (1,954)	100.0 (161)	100.0 (260)	100.0 (264)	100.0 (378)	100.0 (461)	100.0 (430)	100.0 (230)	100.0 (198)	100.0 (129)	100.0 (69)	100.0 (34)
	30 mg/dl 未満	1.3	0.0	1.2	1.5	0.3	2.2	1.6	3.0	2.5	0.8	1.4	0.0
	30~39	11.4	5.0	8.5	11.4	12.7	12.6	13.3	13.5	15.2	13.2	7.2	14.7
	40~49	23.2	17.4	21.5	22.7	22.8	24.5	25.6	25.2	25.3	27.1	24.6	23.5
	50~59	27.1	31.7	30.8	27.7	25.7	24.9	26.5	26.1	23.2	29.5	27.5	32.4
	60~69	18.4	26.1	17.7	18.2	18.5	18.7	15.6	18.7	16.7	10.9	20.3	17.6
	70~79	11.5	13.0	11.5	7.6	12.4	11.1	12.8	8.7	11.6	12.4	17.4	11.8
	80~89	4.5	3.7	5.4	4.9	5.6	4.3	3.3	2.6	4.5	3.1	1.4	0.0
	90~99	1.9	2.5	3.1	4.5	1.1	1.3	0.7	2.2	1.0	0.8	0.0	0.0
	100 以上	0.8	0.6	0.4	1.5	1.1	0.4	0.7	0.0	0.0	2.3	0.0	0.0
平均値 mg/dl	87.3	95.0	90.4	87.1	87.0	85.2	85.1	83.5	82.3	86.0	91.3	85.3	
	標準偏差	15.2	13.5	15.1	16.7	15.1	15.3	14.7	14.7	14.4	16.8	12.6	12.1
女	総 数 (実 数)	100.0 (2,842)	100.0 (242)	100.0 (458)	100.0 (467)	100.0 (603)	100.0 (559)	100.0 (513)	100.0 (288)	100.0 (215)	100.0 (145)	100.0 (95)	100.0 (58)
	30 mg/dl 未満	0.1	0.0	0.0	0.0	0.0	0.5	0.2	0.7	0.0	0.7	0.0	0.0
	30~39	3.2	0.8	0.9	2.6	2.3	4.1	7.0	5.9	6.0	8.3	7.4	6.9
	40~49	13.0	7.4	7.4	9.0	10.9	20.2	18.9	21.9	20.0	14.5	22.1	20.7
	50~59	22.6	18.2	17.2	21.4	21.4	27.0	27.1	26.0	25.6	29.7	27.4	25.9
	60~69	25.7	28.5	28.8	27.4	28.2	22.4	20.9	26.7	21.4	17.9	21.1	25.9
	70~79	18.4	24.4	23.4	21.2	16.6	13.1	16.6	9.0	17.7	17.2	15.8	12.1
	80~89	10.6	12.0	12.9	11.1	13.6	8.4	6.0	5.9	4.7	7.6	5.3	8.6
	90~99	4.2	5.8	4.8	5.4	4.5	3.0	2.5	2.8	3.3	3.4	1.1	0.0
	100 以上	2.2	2.9	4.6	1.9	2.5	1.3	0.8	1.0	1.4	0.7	0.0	0.0
平均値 mg/dl	96.7	99.2	99.1	97.4	97.7	95.3	92.8	93.4	94.0	91.0	92.6	93.1	
	標準偏差	15.6	14.2	15.3	14.9	15.5	15.4	14.8	14.8	15.3	15.4	13.5	13.4

注) コレステロールを下げる薬服用者除外。

◆第33表◆ 血中コチニン濃度の分布（性・年齢階級別）

(%)

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上
全 国	総 数 (実数)	100.0 (1,207)	100.0 (90)	100.0 (166)	100.0 (179)	100.0 (248)	100.0 (268)	100.0 (256)
	1.0 ng/ml 未満	51.3	33.3	41.6	35.8	43.1	62.3	71.1
	1.0~2.0未満	17.3	20.0	23.5	23.5	17.7	11.9	13.3
	2.0~3.0	3.1	5.6	1.8	2.8	4.4	3.7	1.6
	3.0~4.0	1.2	1.1	0.6	1.1	2.4	1.1	0.4
	4.0~5.0	0.6	1.1	0.6	0.6	0.4	1.1	0.0
	5.0~20	1.2	1.1	0.0	1.1	2.0	1.1	1.2
	20~50	2.0	4.4	3.6	2.8	2.0	0.4	1.2
	50~100	2.9	2.2	1.8	2.8	3.6	3.4	2.7
	100~200	6.9	8.9	8.4	9.5	5.6	7.1	4.3
	200~300	7.3	13.3	9.6	10.1	9.7	4.1	2.7
	300~400	3.6	5.6	3.6	5.6	4.4	3.0	1.2
	400~500	1.4	2.2	1.8	2.8	2.8	0.0	0.0
	500ng/ml以上	1.3	1.1	3.0	1.7	1.6	0.7	0.4
	平均 値	58.1	86.9	79.3	84.8	72.0	39.0	22.5
	標準 偏 差	121.1	135.2	146.5	145.3	133.2	94.9	69.9
男	総 数 (実数)	100.0 (478)	100.0 (34)	100.0 (62)	100.0 (67)	100.0 (87)	100.0 (118)	100.0 (110)
	1.0 ng/ml 未満	38.1	14.7	16.1	19.4	23.0	54.2	63.6
	1.0~2.0未満	13.6	23.5	27.4	14.9	9.2	10.2	9.1
	2.0~3.0	1.5	2.9	0.0	3.0	1.1	1.7	0.9
	3.0~4.0	0.8	2.9	0.0	0.0	3.4	0.0	0.0
	4.0~5.0	0.4	2.9	1.6	0.0	0.0	0.0	0.0
	5.0~20	1.9	0.0	0.0	1.5	4.6	1.7	1.8
	20~50	2.3	2.9	3.2	3.0	3.4	0.8	1.8
	50~100	4.0	0.0	1.6	3.0	3.4	5.9	5.5
	100~200	12.3	17.6	16.1	13.4	9.2	12.7	10.0
	200~300	12.1	14.7	16.1	16.4	20.7	6.8	5.5
	300~400	6.7	11.8	6.5	13.4	9.2	5.1	0.9
	400~500	3.3	5.9	3.2	7.5	8.0	0.0	0.0
	500ng/ml以上	2.9	0.0	8.1	4.5	4.6	0.8	0.9
	平均 値	108.2	132.9	153.2	172.9	160.2	64.5	41.5
	標準 偏 差	155.2	150.6	189.1	186.6	174.2	110.5	88.0
女	総 数 (実数)	100.0 (729)	100.0 (56)	100.0 (104)	100.0 (112)	100.0 (161)	100.0 (150)	100.0 (146)
	1.0 ng/ml 未満	59.9	44.6	56.7	45.5	54.0	68.7	76.7
	1.0~2.0未満	19.8	17.9	21.2	28.6	22.4	13.3	16.4
	2.0~3.0	4.3	7.1	2.9	2.7	6.2	5.3	2.1
	3.0~4.0	1.4	0.0	1.0	1.8	1.9	2.0	0.7
	4.0~5.0	0.7	0.0	0.0	0.9	0.6	2.0	0.0
	5.0~20	0.7	1.8	0.0	0.9	0.6	0.7	0.7
	20~50	1.8	5.4	3.8	2.7	1.2	0.0	0.7
	50~100	2.2	3.6	1.9	2.7	3.7	1.3	0.7
	100~200	3.3	3.6	3.8	7.1	3.7	2.7	0.0
	200~300	4.1	12.5	5.8	6.3	3.7	2.0	0.7
	300~400	1.5	1.8	1.9	0.9	1.9	1.3	1.4
	400~500	0.1	0.0	1.0	0.0	0.0	0.0	0.0
	500ng/ml以上	0.3	1.8	0.0	0.0	0.0	0.7	0.0
	平均 値	25.3	59.0	34.8	32.0	24.3	18.9	8.2
	標準 偏 差	76.1	117.8	88.4	75.3	67.3	75.1	47.8

◆第34表◆ 運動習慣の状況（性・年齢階級別）

		総 数	20～29歳	30～39歳	40～49歳	50～59歳	60～69歳	70歳以上	(再掲) 65～74歳	(再掲) 75歳以上
全 国	総 数 (実 数)	100.0 (6,040)	100.0 (534)	100.0 (862)	100.0 (852)	100.0 (1,200)	100.0 (1,280)	100.0 (1,312)	100.0 (1,235)	100.0 (738)
	習慣無(健康上の理由)	6.7	1.5	1.7	2.5	3.6	6.9	17.5	10.8	20.5
	習慣無(その他)	67.1	80.3	81.9	79.5	70.7	56.2	51.1	55.5	48.5
	運動の習慣有	26.3	18.2	16.4	18.1	25.8	37.0	31.4	33.8	31.0
男	総 数 (実 数)	100.0 (2,474)	100.0 (226)	100.0 (328)	100.0 (336)	100.0 (472)	100.0 (553)	100.0 (559)	100.0 (540)	100.0 (302)
	習慣無(健康上の理由)	5.1	0	1.8	1.5	3.0	4.3	14.0	7.0	17.2
	習慣無(その他)	65.6	77.9	76.5	78.3	73.9	56.1	48.8	54.4	45.4
	運動の習慣有	29.3	22.1	21.6	20.2	23.1	39.6	37.2	38.5	37.4
女	総 数 (実 数)	100.0 (3,566)	100.0 (308)	100.0 (534)	100.0 (516)	100.0 (728)	100.0 (727)	100.0 (753)	100.0 (695)	100.0 (436)
	習慣無(健康上の理由)	7.8	2.6	1.7	3.1	4.0	8.8	20.1	13.7	22.7
	習慣無(その他)	68.1	82.1	85.2	80.2	68.5	56.3	52.9	56.3	50.7
	運動の習慣有	24.1	15.3	13.1	16.7	27.5	34.9	27.1	30.1	26.6

◆第35表◆ 1週間の運動日数（性・年齢階級別）

		総 数	20～29歳	30～39歳	40～49歳	50～59歳	60～69歳	70歳以上	(再掲) 65～74歳	(再掲) 75歳以上
全 国	総 数 (実 数)	100.0 (6,040)	100.0 (534)	100.0 (862)	100.0 (852)	100.0 (1,200)	100.0 (1,280)	100.0 (1,312)	100.0 (1,235)	100.0 (738)
	運動無し	73.7	81.8	83.6	81.9	74.3	63.0	68.6	66.2	69.0
	2日／週	5.2	5.2	4.3	5.8	5.2	6.2	4.6	4.6	5.0
	3	4.1	3.4	3.2	3.1	4.4	5.4	4.0	4.6	3.7
男	4	2.3	1.3	1.3	2.2	2.3	3.4	2.2	3.2	1.9
	5	2.8	1.3	1.7	2.0	2.9	4.2	3.0	3.1	3.7
	6	1.3	1.1	0.2	0.7	1.3	2.1	1.8	2.0	1.9
	7日	9.0	2.4	2.9	3.5	8.3	13.9	15.0	15.1	14.2
女	運動有・頻度不明	1.6	3.4	2.7	0.8	1.3	1.7	0.7	1.1	0.7
	総 数 (実 数)	100.0 (2,474)	100.0 (226)	100.0 (328)	100.0 (336)	100.0 (472)	100.0 (553)	100.0 (559)	100.0 (540)	100.0 (302)
	運動無し	70.7	77.9	78.4	79.8	76.9	60.4	62.8	61.5	62.6
	2日／週	6.2	7.1	5.5	7.4	5.1	7.6	5.2	5.7	5.0
男	3	4.4	4.0	4.9	2.7	3.8	5.6	4.7	5.4	4.3
	4	2.1	0.9	1.8	2.1	1.7	3.1	2.3	3.5	2.0
	5	3.1	0.9	3.0	1.5	3.0	4.0	4.1	3.3	4.6
	6	1.1	0.9	0.0	0.9	1.1	1.6	1.6	1.9	2.0
女	7日	10.7	3.5	2.7	4.5	7.4	16.3	19.1	18.0	19.5
	運動有・頻度不明	1.7	4.9	3.7	1.2	1.1	1.4	0.2	0.7	0.0
	総 数 (実 数)	100.0 (3,566)	100.0 (308)	100.0 (534)	100.0 (516)	100.0 (728)	100.0 (727)	100.0 (753)	100.0 (695)	100.0 (436)
	運動無し	75.9	84.7	86.9	83.3	72.5	65.1	72.9	69.9	73.4
女	2日／週	4.5	3.9	3.6	4.7	5.2	5.1	4.2	3.7	5.0
	3	3.8	2.9	2.2	3.3	4.8	5.2	3.5	4.0	3.2
	4	2.4	1.6	0.9	2.3	2.6	3.7	2.1	2.9	1.8
	5	2.6	1.6	0.9	2.3	2.9	4.4	2.3	2.9	3.0
男	6	1.5	1.3	0.4	0.6	1.5	2.5	2.0	2.2	1.8
	7日	7.8	1.6	3.0	2.9	8.9	12.1	12.0	12.9	10.6
	運動有・頻度不明	1.5	2.3	2.1	0.6	1.5	1.9	1.1	1.4	1.1

◆第36表◆ 運動を行う日の平均運動時間（性・年齢階級別）

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~74歳	(再掲) 75歳以上
全 国	総 数 (実 数)	100.0 (6,040)	100.0 (534)	100.0 (862)	100.0 (852)	100.0 (1,200)	100.0 (1,280)	100.0 (1,312)	100.0 (1,235)	100.0 (738)
	運動無し	73.7	81.8	83.6	81.9	74.3	63.0	68.6	66.2	69.0
	30分以上60分未満	10.8	6.0	5.2	6.1	11.8	14.6	14.7	14.7	14.6
	60 120	9.8	5.6	6.3	7.6	9.9	14.4	10.7	11.7	11.1
	120 180	3.8	4.5	3.8	3.2	2.8	5.3	3.5	4.8	2.7
	180分以上	1.8	2.1	1.0	1.1	1.2	2.7	2.4	2.5	2.6
	運動有・時間不明	0.0	0.0	0.0	0.1	0.1	0.0	0.1	0.1	0.0
男	総 数 (実 数)	100.0 (2,474)	100.0 (226)	100.0 (328)	100.0 (336)	100.0 (472)	100.0 (553)	100.0 (559)	100.0 (540)	100.0 (302)
	運動無し	70.7	77.9	78.4	79.8	76.9	60.4	62.8	61.5	62.6
	30分以上60分未満	11.0	6.2	6.4	7.1	9.7	14.5	15.4	13.9	17.5
	60 120	10.7	5.3	8.5	7.7	8.1	15.2	13.8	14.1	13.6
	120 180	4.6	6.6	5.2	3.9	3.4	6.1	3.6	6.1	2.3
	180分以上	2.9	4.0	1.5	1.2	1.7	3.8	4.3	4.3	4.0
	運動有・時間不明	0.1	0.0	0.0	0.3	0.2	0.0	0.2	0.2	0.0
女	総 数 (実 数)	100.0 (3,566)	100.0 (308)	100.0 (534)	100.0 (516)	100.0 (728)	100.0 (727)	100.0 (753)	100.0 (695)	100.0 (436)
	運動無し	75.9	84.7	86.9	83.3	72.5	65.1	72.9	69.9	73.4
	30分以上60分未満	10.7	5.8	4.5	5.4	13.2	14.7	14.2	15.4	12.6
	60 120	9.2	5.8	4.9	7.6	11.1	13.8	8.5	9.8	9.4
	120 180	3.3	2.9	3.0	2.7	2.3	4.7	3.5	3.7	3.0
	180分以上	1.0	0.6	0.7	1.0	0.8	1.8	0.9	1.2	1.6
	運動有・時間不明	—	—	—	—	—	—	—	—	—

◆第37表◆ 運動強度の状況（性・年齢階級別）

		総 数	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~74歳	(再掲) 75歳以上
全 国	総 数 (実 数)	100.0 (6,040)	100.0 (534)	100.0 (862)	100.0 (852)	100.0 (1,200)	100.0 (1,280)	100.0 (1,312)	100.0 (1,235)	100.0 (738)
	運動無し	73.7	81.8	83.6	81.9	74.3	63.0	68.6	66.2	69.0
	高強度	2.1	4.9	3.7	2.1	1.8	1.8	0.3	1.1	0.3
	中強度	8.3	7.7	7.2	8.2	9.3	11.1	5.9	8.3	4.7
	低強度	15.7	5.6	5.5	7.7	14.3	23.9	25.2	24.3	26.0
	運動有・強度不明	0.1	0.0	0.0	0.0	0.3	0.2	0.1	0.2	0.0
	総 数 (実 数)	100.0 (2,474)	100.0 (226)	100.0 (328)	100.0 (336)	100.0 (472)	100.0 (553)	100.0 (559)	100.0 (540)	100.0 (302)
男	運動無し	70.7	77.9	78.4	79.8	76.9	60.4	62.8	61.5	62.6
	高強度	2.9	8.4	5.5	3.0	1.9	2.4	0.5	1.3	0.7
	中強度	10.0	8.0	11.6	8.6	9.1	13.0	8.4	10.6	7.3
	低強度	16.2	5.8	4.6	8.6	11.4	23.9	28.1	26.3	29.5
	運動有・強度不明	0.2	0.0	0.0	0.0	0.6	0.4	0.2	0.4	0.0
	総 数 (実 数)	100.0 (3,566)	100.0 (308)	100.0 (534)	100.0 (516)	100.0 (728)	100.0 (727)	100.0 (753)	100.0 (695)	100.0 (436)
	運動無し	75.9	84.7	86.9	83.3	72.5	65.1	72.9	69.9	73.4
女	高強度	1.5	2.3	2.6	1.6	1.8	1.4	0.1	0.9	0.0
	中強度	7.2	7.5	4.5	7.9	9.3	9.6	4.0	6.5	3.0
	低強度	15.5	5.5	6.0	7.2	16.2	23.9	23.0	22.7	23.6
	運動有・強度不明	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0

◆第38表◆ 運動習慣別栄養素等摂取量

栄養素等別	全 国		男		女	
	習慣有	習慣無	習慣有	習慣無	習慣有	習慣無
調査人數	1,543	3,922	710	1,575	833	2,347
エネルギー kcal	1,987	1,961	2,220	2,239	1,789	1,775
たんぱく質 g	75.6	72.9	82.2	81.0	69.9	67.4
うち動物性 g	39.3	38.2	43.2	43.2	35.9	34.9
脂質 g	52.8	52.9	56.9	56.8	49.2	50.2
うち動物性 g	25.9	26.0	28.6	28.6	23.7	24.2
炭水化物 g	283.8	278.6	312.2	314.8	259.6	254.3
食塩(ナトリウム×2.54/1,000) g	12.3	11.8	13.3	13.0	11.4	11.1
カリウム mg	2,713	2,435	2,792	2,544	2,646	2,362
カルシウム mg	602	538	606	546	599	532
マグネシウム mg	290	265	305	286	277	251
リン mg	1,090	1,033	1,167	1,130	1,026	969
鉄 mg	9.1	8.8	9.4	9.1	8.9	8.6
亜鉛 mg	8.6	8.5	9.4	9.5	8.0	7.8
銅 mg	1.33	1.26	1.42	1.38	1.26	1.17
ビタミンA μgRE	1,081	923	1,087	931	1,076	919
ビタミンD μg	9.5	8.8	10.2	9.9	8.9	8.1
ビタミンE $\text{mg}\alpha\text{-TE}$	13.8	10.7	12.8	9.9	14.7	11.2
ビタミンK μg	337	278	337	282	337	276
ビタミンB ₁ mg	1.64	1.66	1.61	1.75	1.66	1.60
ビタミンB ₂ mg	2.43	2.14	3.16	1.61	1.82	2.49
ナイアシン mgNE	16.4	15.7	18.0	17.8	15.0	14.2
ビタミンB ₆ mg	1.98	2.16	1.93	2.11	2.02	2.19
ビタミンB ₁₂ μg	8.4	7.8	9.2	8.9	7.7	7.0
葉酸 μg	368	328	372	338	364	322
パントテン酸 mg	6.00	5.57	6.35	6.01	5.71	5.28
ビタミンC mg	156	130	145	116	166	139
コレステロール mg	320	331	347	365	297	308
食物繊維 g	16.8	14.9	16.8	15.2	16.7	14.8
うち水溶性 g	3.9	3.5	3.9	3.5	3.8	3.5
うち不溶性 g	12.9	11.5	12.9	11.7	12.8	11.3
脂肪エネルギー比率 %	23.6	24.1	22.7	22.6	24.3	25.1
炭水化物エネルギー比率 %	61.1	60.9	62.4	62.8	59.9	59.7
動物性たんぱく質比率 %	50.3	50.5	50.9	51.5	49.8	49.8

注) 妊婦、「健康上の理由」により運動習慣の無い者除外。

◆第39表◆ 歩行数の平均値及び標準偏差（性・年齢階級別）

		人 数	平均歩行数	標準偏差
全 国	総 数	8,867	7,168	4,248
	15～19歳	500	8,260	4,295
	20～29	984	8,007	4,398
	30～39	1,296	7,925	3,979
	40～49	1,257	7,961	4,146
	50～59	1,653	7,909	4,169
	60～69	1,568	7,016	4,024
	70歳以上	1,609	4,474	3,540
	(再掲)65～74	1,489	6,084	3,823
男	(再掲)75歳以上	912	3,687	3,265
	総 数	4,089	7,575	4,580
	15～19歳	240	8,718	4,600
	20～29	465	8,925	4,878
	30～39	607	8,543	4,459
	40～49	592	8,044	4,477
	50～59	766	8,028	4,552
	60～69	726	7,201	4,290
	70歳以上	693	4,915	3,714
女	(再掲)65～74	689	6,279	3,953
	(再掲)75歳以上	363	4,208	3,653
	総 数	4,778	6,821	3,909
	15～19歳	260	7,837	3,955
	20～29	519	7,185	3,735
	30～39	689	7,381	3,413
	40～49	665	7,888	3,829
	50～59	887	7,807	3,806
	60～69	842	6,857	3,775
	70歳以上	916	4,142	3,367
	(再掲)65～74	800	5,915	3,702
	(再掲)75歳以上	549	3,343	2,935

◆第40表◆ 歩行数の分布（性・年齢階級別）

		総 数	15~19歳	20~29歳	30~39歳	40~49歳	50~59歳	60~69歳	70歳以上	(再掲) 65~74歳	(再掲) 75歳以上
全 国	総 数 (実 数)	100.0 (8,867)	100.0 (500)	100.0 (984)	100.0 (1,296)	100.0 (1,257)	100.0 (1,653)	100.0 (1,568)	100.0 (1,609)	100.0 (1,489)	100.0 (912)
	2,000歩未満	8.6	2.6	4.8	3.2	3.8	3.4	7.1	27.8	12.6	36.0
	2,000~	15.8	12.2	13.1	11.2	12.0	12.9	17.5	26.7	21.5	29.5
	4,000~	19.1	17.6	19.9	20.1	18.7	19.7	21.0	16.3	21.1	13.3
	6,000~	18.6	19.4	18.1	21.9	20.2	21.1	17.9	13.1	16.7	10.2
	8,000~	15.1	18.4	14.4	17.6	18.7	16.0	16.1	8.0	12.8	6.5
	10,000歩以上	22.6	29.8	29.7	26.0	26.6	26.9	20.3	8.0	15.4	4.6
男	総 数 (実 数)	100.0 (4,089)	100.0 (240)	100.0 (465)	100.0 (607)	100.0 (592)	100.0 (766)	100.0 (726)	100.0 (693)	100.0 (689)	100.0 (363)
	2,000歩未満	7.7	2.9	5.2	2.6	4.2	3.7	7.4	23.4	12.0	31.1
	2,000~	14.9	9.6	10.1	9.9	11.5	14.2	16.5	26.3	20.5	28.4
	4,000~	18.7	16.3	15.1	19.3	20.6	20.6	20.2	16.0	19.4	14.0
	6,000~	17.9	20.4	16.1	20.6	19.3	17.8	16.7	15.9	18.4	12.1
	8,000~	15.3	17.5	15.9	16.6	17.6	15.7	17.5	8.1	12.8	7.2
	10,000歩以上	25.6	33.3	37.6	31.0	26.9	28.1	21.6	10.4	16.8	7.2
女	総 数 (実 数)	100.0 (4,778)	100.0 (260)	100.0 (519)	100.0 (689)	100.0 (665)	100.0 (887)	100.0 (842)	100.0 (916)	100.0 (800)	100.0 (549)
	2,000歩未満	9.4	2.3	4.4	3.6	3.5	3.3	6.8	31.1	13.0	39.2
	2,000~	16.7	14.6	15.8	12.3	12.5	11.8	18.4	27.1	22.4	30.2
	4,000~	19.5	18.8	24.3	20.9	17.0	18.9	21.6	16.6	22.5	12.8
	6,000~	19.3	18.5	19.8	23.1	21.1	23.9	19.0	11.0	15.3	8.9
	8,000~	15.0	19.2	13.1	18.4	19.7	16.2	15.0	8.0	12.8	6.0
	10,000歩以上	20.1	26.5	22.5	21.6	26.3	25.8	19.2	6.2	14.1	2.9

