

A randomized, double-blind study comparing the safety and efficacy of clopidogrel versus ticlopidine in Japanese patients with noncardioembolic cerebral infarction.
Fukuuchi Y, Tohgi H, Okudera T, Ikeda Y, Miyanaga Y, Uchiyama S, Hirano M, Shinohara Y, Matsumoto M, Yamaguchi T, Cerebrovasc Dis. 2008;25(1-2):40-9.

水澤 英洋	東京医科歯科大学大学院脳神経病態学分野・教授
神経内科学	
Nishina K, Unno T, Uno Y, Kubodera T, Kanouchi T, <u>Mizusawa H</u> , Yokota T. Efficient in vivo delivery of siRNA to the liver by Conjugation of α -Tocopherol. <i>Mol Ther</i> 2008; 16:734-740.	
<u>Mizusawa H</u> , Clark HB, Koppen AH: Spinocerebellar Ataxia. (Olsson Y, Dickson D, Berberon C, Hardy JA, Hauw JJ, Jellinger K, Lantos P, Mizuno Y, <u>Mizusawa H</u> , eds), Neurodegeneration: The Molecular Pathology of Dementia and Movement Disorders, International Society of Neuropathology Press, Woerthsee, p242-256, 2003	
<u>Mizusawa H</u> . Special Lecture. Treatment of focal brain ischemia using stem cells. 26th Annual Meeting of Korean Neurological Association, Seoul, 2007.10.27.	

○ 矢崎 義雄	独立行政法人国立病院機構
循環器学	
A novel potent vasoconstrictor peptide produced by vascular endothelial cells. <i>Nature</i> 332(6163):411-415, 1988. M. Yanagisawa, et. al.	
Tyrosine phosphorylation of the EGE receptor by the kinase JAK2 is induced by growth hormone. <i>Nature</i> 390(6655):91-96, 1997. T. Yamauchi, et al.	