

第2部 身体状況調査の結果

第11表 身長・体重の平均値及び標準偏差（性・年齢階級別）

年齢(歳)	男 性						女 性					
	身 長(cm)			体 重(kg)			身 長(cm)			体 重(kg)		
	人 数	平均値	標準偏差	人 数	平均値	標準偏差	人 数	平均値	標準偏差	人 数	平均値	標準偏差
1	20	81.0	4.5	20	11.2	1.6	27	80.6	7.7	27	10.2	1.0
2	37	89.8	8.0	37	12.7	1.3	21	86.5	6.0	21	11.9	1.4
3	16	96.6	4.2	16	14.5	1.9	27	94.0	4.0	26	14.0	1.7
4	34	104.1	5.3	34	16.6	2.3	27	103.2	5.1	27	16.0	1.6
5	19	108.7	4.5	19	17.8	1.5	22	110.9	4.9	22	19.6	3.0
6	30	116.2	4.5	30	20.9	2.7	24	115.7	4.8	24	20.2	2.1
7	25	121.0	5.4	25	23.6	4.2	39	120.5	4.5	39	22.6	2.7
8	31	127.8	5.5	31	26.9	4.9	40	127.7	5.9	40	26.1	5.9
9	43	132.0	6.2	43	28.4	5.0	38	133.2	7.0	38	29.4	6.5
10	42	139.2	6.4	42	34.5	7.1	26	140.5	6.5	26	35.9	8.3
11	32	145.3	5.5	32	38.2	7.2	42	143.5	7.4	42	35.0	5.9
12	30	150.6	8.2	30	42.0	8.6	36	152.7	8.1	36	42.4	8.4
13	28	159.4	6.3	28	48.2	6.6	28	153.4	6.2	28	44.5	6.5
14	24	164.5	6.0	24	52.8	11.7	22	154.1	6.1	22	44.3	6.1
15	31	170.7	5.0	31	58.8	8.9	27	157.4	5.2	27	50.1	7.0
16	37	169.5	5.9	37	59.7	9.2	32	157.3	4.8	32	53.0	9.0
17	28	170.5	6.0	28	61.9	10.9	34	157.4	5.3	32	50.4	5.7
18	20	171.1	7.4	18	66.3	15.6	33	157.1	4.8	33	51.3	6.2
19	19	170.7	4.8	19	63.9	14.8	16	157.0	4.6	16	51.2	5.5
20	26	168.7	5.2	26	64.6	13.5	23	157.7	5.5	23	52.4	8.4
21	18	170.2	6.1	18	62.6	11.3	22	157.5	8.6	21	55.2	14.5
22	20	170.4	5.2	20	66.2	14.6	24	159.0	5.4	24	51.4	5.3
23	20	170.2	5.8	20	65.5	9.9	23	156.0	4.7	23	50.2	10.0
24	20	170.4	5.1	20	62.6	8.1	31	158.5	5.0	31	49.9	5.8
25	15	169.7	5.2	15	67.5	16.6	19	159.1	5.8	18	53.0	11.5
26-29	86	172.6	6.6	84	68.9	13.2	122	157.4	5.0	116	51.9	8.9
30-39	374	171.5	5.8	371	70.0	11.9	446	158.1	5.3	426	53.9	8.9
40-49	368	169.7	6.1	368	69.6	11.3	451	158.0	5.1	448	55.9	9.5
50-59	431	168.8	6.0	432	68.5	10.9	486	155.4	5.2	485	54.9	9.3
60-69	542	165.7	5.7	542	65.2	9.3	668	152.9	5.2	668	53.4	8.6
70歳以上	668	161.4	6.7	669	60.4	9.3	846	147.4	6.5	849	49.7	8.5
(再掲)												
20歳以上	2,588	166.9	7.2	2,585	65.9	11.2	3,161	153.7	7.0	3,132	52.9	9.2
20-29	205	171.0	6.1	203	66.5	12.9	264	157.7	5.5	256	51.9	9.2
60-64	300	166.5	5.5	300	65.8	9.5	368	153.8	5.1	368	53.5	8.1
65-69	242	164.6	5.6	242	64.4	9.2	300	151.8	5.1	300	53.2	9.2
70-74	257	163.2	6.5	257	62.5	8.4	305	149.6	5.5	305	51.8	8.0
75-79	212	160.9	6.1	212	60.8	9.3	247	148.1	6.1	248	51.1	8.2
80歳以上	199	159.4	6.9	200	57.2	9.5	294	144.4	6.7	296	46.3	8.2
総 数	3,134	161.7	17.9	3,722	60.8	17.1	3,129	150.5	13.9	3,690	50.0	12.5

注) 体重は妊婦除外

(23年)

第12表 BMIの平均値及び標準偏差（15歳以上，性・年齢階級別）

年齢(歳)	男 性			女 性		
	人 数	平均値	標準偏差	人 数	平均値	標準偏差
15-19歳	133	21.09	3.31	140	20.72	2.62
20-29歳	203	22.66	3.94	256	20.80	3.39
30-39歳	371	23.81	3.89	426	21.54	3.37
40-49歳	368	24.16	3.67	448	22.37	3.48
50-59歳	431	23.99	3.29	485	22.75	3.65
60-69歳	542	23.74	3.11	668	22.86	3.62
70歳以上	668	23.15	3.07	846	22.86	3.54
(再掲)						
20歳以上	2,583	23.61	3.43	3,129	22.43	3.59
20-69歳	1,915	23.78	3.53	2,283	22.26	3.60
40-69歳	1,341	23.94	3.33	1,601	22.69	3.59
65-69歳	242	23.76	3.12	300	23.09	3.81
70-74歳	257	23.43	2.74	305	23.13	3.37
75-79歳	212	23.44	3.10	247	23.31	3.54
80-84歳	139	22.75	3.13	170	22.27	3.37
80歳以上	199	22.48	3.34	294	22.21	3.63
85歳以上	60	21.86	3.72	124	22.13	3.97
総 数	2,716	23.49	3.46	3,269	22.35	3.57

注) 妊婦除外

(23年)

第13表 BMIの分布（15歳以上，性別）

	総 数		男 性		女 性	
	人 数	%	人 数	%	人 数	%
15未満	10	0.2	2	0.1	8	0.2
15-16未満	28	0.5	6	0.2	22	0.7
16-17未満	76	1.3	26	1.0	50	1.5
17-18未満	201	3.4	55	2.0	146	4.5
18-19未満	392	6.5	112	4.1	280	8.6
19-20未満	553	9.2	174	6.4	379	11.6
20-21未満	648	10.8	246	9.1	402	12.3
21-22未満	722	12.1	315	11.6	407	12.5
22-23未満	722	12.1	349	12.8	373	11.4
23-24未満	594	9.9	310	11.4	284	8.7
24-25未満	560	9.4	321	11.8	239	7.3
25-26未満	430	7.2	238	8.8	192	5.9
26-27未満	313	5.2	174	6.4	139	4.3
27-28未満	231	3.9	113	4.2	118	3.6
28-29未満	151	2.5	90	3.3	61	1.9
29-30未満	119	2.0	65	2.4	54	1.7
30-31未満	78	1.3	39	1.4	39	1.2
31-32未満	50	0.8	25	0.9	25	0.8
32-33未満	33	0.6	20	0.7	13	0.4
33-34未満	30	0.5	13	0.5	17	0.5
34-35未満	12	0.2	6	0.2	6	0.2
35以上	32	0.5	17	0.6	15	0.5
総 数	5,985	100.0	2,716	100.0	3,269	100.0

注) 妊婦除外

(23年)

第14表 BMIの状況（15歳以上、性・年齢階級別）

		総 数		や せ		普 通		肥 満		(再掲) 20以下		(再掲) 25以上30未満		(再掲) 30以上	
				18.5未満		18.5以上25未満		25以上							
		人 数	%	人 数	%	人 数	%	人 数	%	人 数	%	人 数	%	人 数	%
総数	15-19歳	273	100.0	47	17.2	203	74.4	23	8.4	123	45.1	20	7.3	3	1.1
	20-29歳	459	100.0	73	15.9	317	69.1	69	15.0	182	39.7	49	10.7	20	4.4
	30-39歳	797	100.0	76	9.5	544	68.3	177	22.2	220	27.6	134	16.8	43	5.4
	40-49歳	816	100.0	43	5.3	551	67.5	222	27.2	157	19.2	180	22.1	42	5.1
	50-59歳	916	100.0	47	5.1	613	66.9	256	27.9	155	16.9	211	23.0	45	4.9
	60-69歳	1,210	100.0	79	6.5	797	65.9	334	27.6	200	16.5	288	23.8	46	3.8
	70歳以上	1,514	100.0	125	8.3	991	65.5	398	26.3	287	19.0	362	23.9	36	2.4
	(再掲)														
	20歳以上	5,712	100.0	443	7.8	3,813	66.8	1,456	25.5	1,201	21.0	1,224	21.4	232	4.1
	20-69歳	4,198	100.0	318	7.6	2,822	67.2	1,058	25.2	914	21.8	862	20.5	196	4.7
	40-69歳	2,942	100.0	169	5.7	1,961	66.7	812	27.6	512	17.4	679	23.1	133	4.5
	65-69歳	542	100.0	34	6.3	352	64.9	156	28.8	87	16.1	131	24.2	25	4.6
	70-74歳	562	100.0	35	6.2	373	66.4	154	27.4	78	13.9	142	25.3	12	2.1
	75-79歳	459	100.0	31	6.8	287	62.5	141	30.7	71	15.5	126	27.5	15	3.3
	80-84歳	309	100.0	33	10.7	212	68.6	64	20.7	78	25.2	59	19.1	5	1.6
	80歳以上	493	100.0	59	12.0	331	67.1	103	20.9	138	28.0	94	19.1	9	1.8
	85歳以上	184	100.0	26	14.1	119	64.7	39	21.2	60	32.6	35	19.0	4	2.2
	総 数	5,985	100.0	490	8.2	4,016	67.1	1,479	24.7	1,324	22.1	1,244	20.8	235	3.9
男性	15-19歳	133	100.0	23	17.3	93	69.9	17	12.8	57	42.9	15	11.3	2	1.5
	20-29歳	203	100.0	17	8.4	143	70.4	43	21.2	46	22.7	31	15.3	12	5.9
	30-39歳	371	100.0	19	5.1	230	62.0	122	32.9	59	15.9	95	25.6	27	7.3
	40-49歳	368	100.0	9	2.4	231	62.8	128	34.8	39	10.6	100	27.2	28	7.6
	50-59歳	431	100.0	10	2.3	277	64.3	144	33.4	40	9.3	124	28.8	20	4.6
	60-69歳	542	100.0	21	3.9	350	64.6	171	31.5	56	10.3	152	28.0	19	3.5
	70歳以上	668	100.0	43	6.4	450	67.4	175	26.2	97	14.5	163	24.4	12	1.8
	(再掲)														
	20歳以上	2,583	100.0	119	4.6	1,681	65.1	783	30.3	337	13.0	665	25.7	118	4.6
	20-69歳	1,915	100.0	76	4.0	1,231	64.3	608	31.7	240	12.5	502	26.2	106	5.5
	40-69歳	1,341	100.0	40	3.0	858	64.0	443	33.0	135	10.1	376	28.0	67	5.0
	65-69歳	242	100.0	9	3.7	160	66.1	73	30.2	23	9.5	65	26.9	8	3.3
	70-74歳	257	100.0	9	3.5	175	68.1	73	28.4	26	10.1	70	27.2	3	1.2
	75-79歳	212	100.0	14	6.6	133	62.7	65	30.7	25	11.8	61	28.8	4	1.9
	80-84歳	139	100.0	10	7.2	101	72.7	28	20.1	27	19.4	25	18.0	3	2.2
	80歳以上	199	100.0	20	10.1	142	71.4	37	18.6	46	23.1	32	16.1	5	2.5
	85歳以上	60	100.0	10	16.7	41	68.3	9	15.0	19	31.7	7	11.7	2	3.3
	総 数	2,716	100.0	142	5.2	1,774	65.3	800	29.5	394	14.5	680	25.0	120	4.4
女性	15-19歳	140	100.0	24	17.1	110	78.6	6	4.3	66	47.1	5	3.6	1	0.7
	20-29歳	256	100.0	56	21.9	174	68.0	26	10.2	136	53.1	18	7.0	8	3.1
	30-39歳	426	100.0	57	13.4	314	73.7	55	12.9	161	37.8	39	9.2	16	3.8
	40-49歳	448	100.0	34	7.6	320	71.4	94	21.0	118	26.3	80	17.9	14	3.1
	50-59歳	485	100.0	37	7.6	336	69.3	112	23.1	115	23.7	87	17.9	25	5.2
	60-69歳	668	100.0	58	8.7	447	66.9	163	24.4	144	21.6	136	20.4	27	4.0
	70歳以上	846	100.0	82	9.7	541	63.9	223	26.4	190	22.5	199	23.5	24	2.8
	(再掲)														
	20歳以上	3,129	100.0	324	10.4	2,132	68.1	673	21.5	864	27.6	559	17.9	114	3.6
	20-69歳	2,283	100.0	242	10.6	1,591	69.7	450	19.7	674	29.5	360	15.8	90	3.9
	40-69歳	1,601	100.0	129	8.1	1,103	68.9	369	23.0	377	23.5	303	18.9	66	4.1
	65-69歳	300	100.0	25	8.3	192	64.0	83	27.7	64	21.3	66	22.0	17	5.7
	70-74歳	305	100.0	26	8.5	198	64.9	81	26.6	52	17.0	72	23.6	9	3.0
	75-79歳	247	100.0	17	6.9	154	62.3	76	30.8	46	18.6	65	26.3	11	4.5
	80-84歳	170	100.0	23	13.5	111	65.3	36	21.2	51	30.0	34	20.0	2	1.2
	80歳以上	294	100.0	39	13.3	189	64.3	66	22.4	92	31.3	62	21.1	4	1.4
	85歳以上	124	100.0	16	12.9	78	62.9	30	24.2	41	33.1	28	22.6	2	1.6
	総 数	3,269	100.0	348	10.6	2,242	68.6	679	20.8	930	28.4	564	17.3	115	3.5

注) 妊婦除外。BMIの判定はp.17参照

(23年)

第15表 学校保健統計調査方式による肥満区分（6－14歳，性・年齢階級別）

		総 数		6－8歳		9－11歳		12－14歳	
		人 数	%	人 数	%	人 数	%	人 数	%
総数	－30%以下	9	1.6	8	4.2	1	0.4	0	0.0
	－30%超～－20%以下	41	7.1	17	9.0	13	5.8	11	6.5
	－20%超～－10%以下	128	22.1	35	18.5	47	21.1	46	27.4
	－10%超～ 0%未満	193	33.3	68	36.0	76	34.1	49	29.2
	0%以上～+10%未満	120	20.7	43	22.8	42	18.8	35	20.8
	+10%以上～+20%未満	54	9.3	13	6.9	23	10.3	18	10.7
	+20%以上～+30%未満	18	3.1	2	1.1	11	4.9	5	3.0
	+30%以上～+40%未満	10	1.7	2	1.1	5	2.2	3	1.8
	+40%以上～+50%未満	2	0.3	0	0.0	1	0.4	1	0.6
	+50%以上	5	0.9	1	0.5	4	1.8	0	0.0
	総 数	580	100.0	189	100.0	223	100.0	168	100.0
男性	－30%以下	8	2.8	8	9.3	0	0.0	0	0.0
	－30%超～－20%以下	25	8.8	16	18.6	6	5.1	3	3.7
	－20%超～－10%以下	54	18.9	10	11.6	26	22.2	18	22.0
	－10%超～ 0%未満	97	34.0	30	34.9	41	35.0	26	31.7
	0%以上～+10%未満	53	18.6	15	17.4	19	16.2	19	23.2
	+10%以上～+20%未満	30	10.5	4	4.7	17	14.5	9	11.0
	+20%以上～+30%未満	7	2.5	1	1.2	2	1.7	4	4.9
	+30%以上～+40%未満	7	2.5	2	2.3	3	2.6	2	2.4
	+40%以上～+50%未満	2	0.7	0	0.0	1	0.9	1	1.2
	+50%以上	2	0.7	0	0.0	2	1.7	0	0.0
	総 数	285	100.0	86	100.0	117	100.0	82	100.0
女性	－30%以下	1	0.3	0	0.0	1	0.9	0	0.0
	－30%超～－20%以下	16	5.4	1	1.0	7	6.6	8	9.3
	－20%超～－10%以下	74	25.1	25	24.3	21	19.8	28	32.6
	－10%超～ 0%未満	96	32.5	38	36.9	35	33.0	23	26.7
	0%以上～+10%未満	67	22.7	28	27.2	23	21.7	16	18.6
	+10%以上～+20%未満	24	8.1	9	8.7	6	5.7	9	10.5
	+20%以上～+30%未満	11	3.7	1	1.0	9	8.5	1	1.2
	+30%以上～+40%未満	3	1.0	0	0.0	2	1.9	1	1.2
	+40%以上～+50%未満	0	0.0	0	0.0	0	0.0	0	0.0
	+50%以上	3	1.0	1	1.0	2	1.9	0	0.0
	総 数	295	100.0	103	100.0	106	100.0	86	100.0

注) 学校保健統計調査方式による区分は，p.17 参照

(23年)

第16表 腹囲の分布（6歳以上、性・年齢階級別）

	総 数		6-8歳		9-11歳		12-14歳		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 20歳以上		(再掲) 40-74歳		
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
総 数	50cm未満	21	0.3	16	8.8	3	1.5	1	0.6	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.0	0	0.0
	50－55cm未満	100	1.6	73	40.1	25	12.4	1	0.6	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1	0	0.0	1	0.0	1	0.0
	55－60	186	3.0	67	36.8	72	35.6	21	13.1	9	3.7	8	2.0	1	0.1	3	0.4	0	0.0	2	0.2	3	0.2	17	0.3	7	0.2
	60－65	259	4.2	13	7.1	56	27.7	42	26.3	29	11.9	27	6.7	31	4.2	16	2.1	9	1.0	13	1.1	23	1.6	119	2.2	44	1.3
	65－70	483	7.8	6	3.3	25	12.4	47	29.4	52	21.4	76	18.8	75	10.2	67	8.8	49	5.6	38	3.3	48	3.3	353	6.6	165	5.0
	70－75	779	12.7	5	2.7	10	5.0	28	17.5	70	28.8	90	22.2	129	17.6	118	15.5	94	10.8	108	9.4	127	8.8	666	12.4	360	10.8
	75－80	960	15.6	1	0.5	3	1.5	10	6.3	43	17.7	91	22.5	163	22.2	127	16.7	143	16.4	177	15.4	202	14.0	903	16.8	522	15.7
	80－85	1085	17.6	0	0.0	6	3.0	6	3.8	14	5.8	51	12.6	127	17.3	151	19.8	205	23.5	239	20.8	286	19.8	1059	19.7	709	21.3
	85－90	1001	16.3	1	0.5	1	0.5	2	1.3	13	5.3	23	5.7	93	12.7	133	17.5	166	19.0	248	21.6	321	22.2	984	18.3	680	20.4
	90－95	635	10.3	0	0.0	0	0.0	1	0.6	6	2.5	14	3.5	54	7.4	72	9.4	88	10.1	179	15.6	221	15.3	628	11.7	425	12.8
	95－100	365	5.9	0	0.0	1	0.5	0	0.0	2	0.8	9	2.2	33	4.5	42	5.5	72	8.2	88	7.7	118	8.2	362	6.7	251	7.5
	100－105	160	2.6	0	0.0	0	0.0	1	0.6	2	0.8	7	1.7	13	1.8	19	2.5	24	2.7	35	3.0	59	4.1	157	2.9	97	2.9
	105－110	72	1.2	0	0.0	0	0.0	0	0.0	2	0.8	5	1.2	10	1.4	5	0.7	14	1.6	10	0.9	26	1.8	70	1.3	37	1.1
	110－115	24	0.4	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	3	0.4	3	0.4	3	0.3	8	0.7	6	0.4	24	0.4	14	0.4
	115－120	16	0.3	0	0.0	0	0.0	0	0.0	1	0.4	1	0.2	0	0.0	4	0.5	5	0.6	3	0.3	2	0.1	15	0.3	13	0.4
	120cm以上	8	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	2	0.3	2	0.3	1	0.1	1	0.1	1	0.1	8	0.1	5	0.2
総 数	6,154	100.0	182	100.0	202	100.0	160	100.0	243	100.0	405	100.0	734	100.0	762	100.0	873	100.0	1,150	100.0	1,443	100.0	5,367	100.0	3,330	100.0	
男 性	50cm未満	8	0.3	8	9.6	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	50－55cm未満	46	1.6	31	37.3	15	14.2	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	55－60	90	3.2	32	38.6	44	41.5	10	12.8	0	0.0	4	2.2	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	4	0.2	0	0.0
	60－65	52	1.9	4	4.8	21	19.8	17	21.8	4	3.4	1	0.5	0	0.0	0	0.0	0	0.0	0	0.0	5	0.8	6	0.2	0	0.0
	65－70	138	4.9	4	4.8	12	11.3	27	34.6	31	26.7	16	8.7	14	4.1	7	2.1	6	1.5	6	1.2	15	2.3	64	2.6	23	1.5
	70－75	253	9.0	3	3.6	8	7.5	11	14.1	35	30.2	36	19.6	35	10.3	28	8.4	22	5.4	31	6.1	44	6.8	196	8.1	93	6.2
	75－80	379	13.5	1	1.2	0	0.0	5	6.4	19	16.4	51	27.7	71	20.8	47	14.1	53	13.1	64	12.5	68	10.6	354	14.6	187	12.5
	80－85	553	19.7	0	0.0	5	4.7	5	6.4	5	4.3	33	17.9	79	23.2	78	23.4	103	25.4	103	20.1	134	22.1	538	22.2	346	23.1
	85－90	535	19.1	0	0.0	1	0.9	1	1.3	10	8.6	19	10.3	52	15.2	77	23.1	86	21.2	130	25.4	159	24.7	523	21.6	361	24.1
	90－95	373	13.3	0	0.0	0	0.0	1	1.3	6	5.2	8	4.3	40	11.7	45	13.5	59	14.6	103	20.1	111	17.3	366	15.1	248	16.6
	95－100	208	7.4	0	0.0	0	0.0	0	0.0	2	1.7	7	3.8	27	7.9	26	7.8	42	10.4	45	8.8	59	9.2	206	8.5	137	9.1
	100－105	97	3.5	0	0.0	0	0.0	1	1.3	1	0.9	4	2.2	9	2.6	15	4.5	19	4.7	20	3.9	28	4.4	95	3.9	65	4.3
	105－110	40	1.4	0	0.0	0	0.0	0	0.0	2	1.7	2	1.1	9	2.6	3	0.9	10	2.5	5	1.0	9	1.4	38	1.6	21	1.4
	110－115	16	0.6	0	0.0	0	0.0	0	0.0	0	0.0	1	0.5	3	0.9	3	0.9	2	0.5	4	0.8	3	0.5	16	0.7	9	0.6
	115－120	9	0.3	0	0.0	0	0.0	0	0.0	1	0.9	1	0.5	0	0.0	3	0.9	3	0.7	1	0.2	0	0.0	8	0.3	7	0.5
	120cm以上	4	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.5	2	0.6	1	0.3	0	0.0	0	0.0	0	0.0	4	0.2	1	0.1
総 数	2,801	100.0	83	100.0	106	100.0	78	100.0	116	100.0	184	100.0	341	100.0	333	100.0	405	100.0	512	100.0	643	100.0	2,418	100.0	1,498	100.0	
女 性	50cm未満	13	0.4	8	8.1	3	3.1	1	1.2	0	0.0	1	0.5	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.0	0	0.0
	50－55cm未満	54	1.6	42	42.4	10	10.4	1	1.2	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.0	1	0.1
	55－60	96	2.9	35	35.4	28	29.2	11	13.4	9	7.1	4	1.8	1	0.3	3	0.7	0	0.0	2	0.3	3	0.4	13	0.4	7	0.4
	60－65	207	6.2	9	9.1	35	36.5	25	30.5	25	19.7	26	11.8	31	7.9	16	3.7	9	1.9	13	2.0	18	2.3	113	3.8	44	2.4
	65－70	345	10.3	2	2.0	13	13.5	20	24.4	21	16.5	60	27.1	61	15.5	60	14.0	43	9.2	32	5.0	33	4.1	289	9.8	142	7.8
	70－75	526	15.7	2	2.0	2	2.1	17	20.7	35	27.6	54	24.4	94	23.9	90	21.0	72	15.4	77	12.1	83	10.4	470	15.9	267	14.6
	75－80	581	17.3	0	0.0	3	3.1	5	6.1	24	18.9	40	18.1	92	23.4	80	18.6	90	19.2	113	17.7	134	16.8	549	18.6	335	18.3
	80－85	532	15.9	0	0.0	1	1.0	1	1.2	9	7.1	18	8.1	48	12.2	73	17.0	102	21.8	136	21.3	144	18.0	521	17.7	363	19.8
	85－90	466	13.9	1	1.0	0	0.0	1	1.2	3	2.4	4	1.8	41	10.4	56	13.1	80	17.1	118	18.5	162	20.3	461	15.6	319	17.4
	90－95	262	7.8	0	0.0	0	0.0	0	0.0	0	0.0	6	2.7	14	3.6	27	6.3	29	6.2	76	11.9	110	13.8	262	8.9	177	9.7
	95－100	157	4.7	0	0.0	1	1.0	0	0.0	0	0.0	2	0.9	6	1.5	16	3.7	30	6.4	43	6.7	59	7.4	156	5.3	114	6.2
	100－105	63	1.9	0	0.0	0	0.0	0	0.0	1	0.8	3	1.4	4	1.0	4	0.9	5	1.1	15	2.4	31	3.9	62	2.1	32	1.7
	105－110	32	1.0	0	0.0	0	0.0	0	0.0	0	0.0	3	1.4	1	0.3	2	0.5	4	0.9	5	0.8	17	2.1	32	1.1	16	0.9
	110－115	8	0.2	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	4	0.6	3	0.4	8	0.3	5	0.3
	115－120	7	0.2	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	2	0.4	2	0.3	2	0.3	7	0.2	6	0.3
	120cm以上	4	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	1	0.2	1	0.1	4	0.1	4	0.2
総 数	3,353	100.0	99	100.0	96	100.0	82	100.0	127	100.0	221	100.0	393	100.0	429	100.0	468	100.0	638	100.0	800	100.0	2,949	100.0	1,832	100.0	

注) 妊婦除外

(23年)

第 17 表 腹囲の区分（20 歳以上，性・年齢階級別）

	総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	<85cm(男)/<90cm(女)		3,580	66.7	348	85.9	567	77.2	538	70.6	696	60.5	851	59.0	2,127	63.9
	≥85cm(男)/≥90cm(女)		1,787	33.3	57	14.1	167	22.8	224	29.4	454	39.5	592	41.0	1,203	36.1
	総 数		5,367	100.0	405	100.0	734	100.0	762	100.0	1,150	100.0	1,443	100.0	3,330	100.0
男性	<85cm		1,162	48.1	141	76.6	199	58.4	160	48.0	204	39.8	274	42.6	649	43.3
	≥85cm		1,256	51.9	43	23.4	142	41.6	173	52.0	308	60.2	369	57.4	849	56.7
	総 数		2,418	100.0	184	100.0	341	100.0	333	100.0	512	100.0	643	100.0	1,498	100.0
女性	<90cm		2,418	82.0	207	93.7	368	93.6	378	88.1	492	77.1	577	72.1	1,478	80.7
	≥90cm		531	18.0	14	6.3	25	6.4	51	11.9	146	22.9	223	27.9	354	19.3
	総 数		2,949	100.0	221	100.0	393	100.0	429	100.0	638	100.0	800	100.0	1,832	100.0

注) 妊婦除外

(23年)

第 18 表 BMI・腹囲の区分（20 歳以上，性・年齢階級別）

	総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	BMI<25腹囲<85cm(男)/90cm(女)		3,254	61.1	324	81.2	525	71.9	479	63.4	632	55.3	770	53.6	1,920	58.1
	BMIのみ超(BMI≥25)		298	5.6	19	4.8	38	5.2	53	7.2	58	5.1	76	5.3	191	5.8
	腹囲のみ超(腹囲≥85cm(男)/90cm(女))		705	13.2	16	4.0	43	5.9	102	11.8	191	16.7	283	19.7	473	14.3
	BMI≥25腹囲≥85cm(男)/90cm(女)		1,073	20.1	40	10.0	124	17.0	187	20.1	262	22.9	308	21.4	723	21.9
	総 数		5,330	100.0	399	100.0	730	100.0	755	100.0	1,143	100.0	1,437	100.0	3,307	100.0
男性	BMI<25腹囲<85cm		1,099	45.8	131	72.4	186	54.9	174	44.5	197	38.8	264	41.06	614	41.3
	BMIのみ超(BMI≥25)		53	2.2	8	4.4	11	3.2	8	3.6	4	0.8	10	1.56	29	2.0
	腹囲のみ超(腹囲≥85cm)		563	23.4	13	7.2	39	11.5	89	19.7	149	29.3	208	32.35	384	25.8
	BMI≥25腹囲≥85cm		686	28.6	29	16.0	103	30.4	129	32.1	158	31.1	161	25.04	459	30.9
	総 数		2,401	100.0	181	100.0	339	100.0	330	100.0	508	100.0	643	100.0	1,486	100.0
女性	BMI<25腹囲<90cm		2,155	73.6	193	88.5	339	86.7	332	78.1	435	68.5	506	63.7	1,306	71.7
	BMIのみ超(BMI≥25)		245	8.4	11	5.0	27	6.9	45	9.9	54	8.5	66	8.3	162	8.9
	腹囲のみ超(腹囲≥90cm)		142	4.8	3	1.4	4	1.0	13	2.8	42	6.6	75	9.4	89	4.9
	BMI≥25腹囲≥90cm		387	13.2	11	5.0	21	5.4	58	10.8	104	16.4	147	18.5	264	14.5
	総 数		2,929	100.0	218	100.0	391	100.0	466	100.0	635	100.0	794	100.0	1,821	100.0

注) 妊婦除外

(23年)

【参考】 内臓脂肪型肥満の診断基準：

- BMI 25 以上で，男性のウエスト周囲径 85cm 以上，女性のウエスト周囲径 90cm 以上を上半身肥満の疑いとする。
- 上半身肥満の疑いと判定され，腹部CT法による内臓脂肪面積 100cm²以上（男女とも）を内臓脂肪型肥満と診断する。（日本肥満学会肥満症診断基準検討委員会，2000 年）

※国民健康・栄養調査の「腹囲」は，「立位のへその高さ」で計測したが，計測位置は「ウエスト周囲径」と同じである。

第19表 薬の服用状況（性・年齢階級別）

		総数		20－29歳		30－39歳		40－49歳		50－59歳		60－69歳		70歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	使用なし	2,685	64.1	273	99.6	506	96.7	465	90.1	487	73.7	506	52.9	448	35.7
	使用あり(一つ以上)	1,502	35.9	1	0.4	17	3.3	51	9.9	174	26.3	451	47.1	808	64.3
	血圧を下げる薬	1,150	27.5	0	0.0	7	1.3	17	3.3	131	19.8	341	35.6	654	52.1
	脈の乱れを治す薬	165	3.9	0	0.0	1	0.2	0	0.0	10	1.5	34	3.6	120	9.6
	インスリン注射又は血糖を下げる薬	249	5.9	1	0.4	2	0.4	8	1.6	28	4.2	83	8.7	127	10.1
	コレステロールを下げる薬	586	14.0	0	0.0	3	0.6	21	4.1	58	8.8	202	21.1	302	24.0
	中性脂肪(トリグリセライド)を下げる薬	176	4.2	0	0.0	1	0.2	6	1.2	19	2.9	54	5.6	96	7.6
	貧血治療のための薬(鉄剤)	68	1.6	0	0.0	5	1.0	13	2.5	7	1.1	12	1.3	31	2.5
	総 数	4,187	100.0	274	100.0	523	100.0	516	100.0	661	100.0	957	100.0	1,256	100.0
男性	使用なし	1,099	61.9	112	100.0	209	96.3	175	88.4	196	70.8	213	51.6	194	34.7
	使用あり(一つ以上)	677	38.1	0	0.0	8	3.7	23	11.6	81	29.2	200	48.4	365	65.3
	血圧を下げる薬	535	30.1	0	0.0	6	2.8	10	5.1	59	21.3	151	36.6	309	55.3
	脈の乱れを治す薬	88	5.0	0	0.0	0	0.0	0	0.0	4	1.4	21	5.1	63	11.3
	インスリン注射又は血糖を下げる薬	153	8.6	0	0.0	1	0.5	4	2.0	17	6.1	53	12.8	78	14.0
	コレステロールを下げる薬	227	12.8	0	0.0	2	0.9	12	6.1	22	7.9	74	17.9	117	20.9
	中性脂肪(トリグリセライド)を下げる薬	95	5.3	0	0.0	1	0.5	5	2.5	13	4.7	27	6.5	49	8.8
	貧血治療のための薬(鉄剤)	19	1.1	0	0.0	0	0.0	0	0.0	1	0.4	5	1.2	13	2.3
	総 数	1,776	100.0	112	100.0	217	100.0	198	100.0	277	100.0	413	100.0	559	100.0
女性	使用なし	1,602	66.0	166	98.8	308	96.9	290	91.2	291	75.8	293	53.9	254	36.4
	使用あり(一つ以上)	827	34.0	2	1.2	10	3.1	28	8.8	93	24.2	251	46.1	443	63.6
	血圧を下げる薬	615	25.3	0	0.0	1	0.3	7	2.2	72	18.8	190	34.9	345	49.5
	脈の乱れを治す薬	77	3.2	0	0.0	1	0.3	0	0.0	6	1.6	13	2.4	57	8.2
	インスリン注射又は血糖を下げる薬	96	4.0	1	0.6	1	0.3	4	1.3	11	2.9	30	5.5	49	7.0
	コレステロールを下げる薬	359	14.8	0	0.0	1	0.3	9	2.8	36	9.4	128	23.5	185	26.5
	中性脂肪(トリグリセライド)を下げる薬	81	3.3	0	0.0	0	0.0	1	0.3	6	1.6	27	5.0	47	6.7
	貧血治療のための薬(鉄剤)	51	2.1	1	0.6	6	1.9	13	4.1	6	1.6	7	1.3	18	2.6
	総 数	2,429	100.0	168	100.0	318	100.0	318	100.0	384	100.0	544	100.0	697	100.0

* 複数回答

(23年)

第20表の1 血圧の状況（性・年齢階級別）

		総数		至適血圧		正常血圧		正常高値血圧		Ⅰ度高血圧		Ⅱ度高血圧		Ⅲ度高血圧		(再掲) 収縮期高血圧	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%*
総数	15-19歳	115	100.0	95	82.6	16	13.9	2	1.7	2	1.7	0	0.0	0	0.0	1	0.9
	20-29歳	219	100.0	168	76.7	26	11.9	14	6.4	10	4.6	0	0.0	1	0.5	3	1.4
	30-39歳	483	100.0	282	58.4	96	19.9	54	11.2	41	8.5	6	1.2	4	0.8	6	1.2
	40-49歳	457	100.0	209	45.7	93	20.4	71	15.5	61	13.3	15	3.3	8	1.8	20	4.4
	50-59歳	593	100.0	131	22.1	137	23.1	121	20.4	146	24.6	46	7.8	12	2.0	63	10.6
	60-69歳	908	100.0	104	11.5	164	18.1	242	26.7	297	32.7	81	8.9	20	2.2	202	22.2
	70歳以上	1,184	100.0	102	8.6	155	13.1	275	23.2	458	38.7	159	13.4	35	3.0	483	40.8
	(再掲) 65-74歳	890	100.0	91	10.2	138	15.5	248	27.9	293	32.9	100	11.2	20	2.2	272	30.6
	(再掲) 75歳以上	720	100.0	52	7.2	88	12.2	159	22.1	301	41.8	97	13.5	23	3.2	327	45.4
	総 数	3,959	100.0	1,091	27.6	687	17.4	779	19.7	1,015	25.6	307	7.8	80	2.0	778	19.7
男性	15-19歳	55	100.0	38	69.1	13	23.6	2	3.6	2	3.6	0	0.0	0	0.0	1	1.8
	20-29歳	79	100.0	41	51.9	20	25.3	10	12.7	7	8.9	0	0.0	1	1.3	3	3.8
	30-39歳	188	100.0	64	34.0	52	27.7	34	18.1	31	16.5	3	1.6	4	2.1	3	1.6
	40-49歳	164	100.0	53	32.3	34	20.7	34	20.7	30	18.3	7	4.3	6	3.7	9	5.5
	50-59歳	236	100.0	36	15.3	42	17.8	55	23.3	71	30.1	25	10.6	7	3.0	25	10.6
	60-69歳	388	100.0	32	8.2	55	14.2	91	23.5	156	40.2	45	11.6	9	2.3	97	25.0
	70歳以上	528	100.0	38	7.2	73	13.8	119	22.5	201	38.1	77	14.6	20	3.8	205	38.8
	(再掲) 65-74歳	388	100.0	28	7.2	62	16.0	96	24.7	142	36.6	50	12.9	10	2.6	121	31.2
	(再掲) 75歳以上	321	100.0	21	6.5	39	12.1	71	22.1	129	40.2	48	15.0	13	4.0	142	44.2
	総 数	1,638	100.0	302	18.4	289	17.6	345	21.1	498	30.4	157	9.6	47	2.9	343	20.9
女性	15-19歳	60	100.0	57	95.0	3	5.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	20-29歳	140	100.0	127	90.7	6	4.3	4	2.9	3	2.1	0	0.0	0	0.0	0	0.0
	30-39歳	295	100.0	218	73.9	44	14.9	20	6.8	10	3.4	3	1.0	0	0.0	3	1.0
	40-49歳	293	100.0	156	53.2	59	20.1	37	12.6	31	10.6	8	2.7	2	0.7	11	3.8
	50-59歳	357	100.0	95	26.6	95	26.6	66	18.5	75	21.0	21	5.9	5	1.4	38	10.6
	60-69歳	520	100.0	72	13.8	109	21.0	151	29.0	141	27.1	36	6.9	11	2.1	105	20.2
	70歳以上	656	100.0	64	9.8	82	12.5	156	23.8	257	39.2	82	12.5	15	2.3	278	42.4
	(再掲) 65-74歳	502	100.0	63	12.5	76	15.1	152	30.3	151	30.1	50	10.0	10	2.0	151	30.1
	(再掲) 75歳以上	399	100.0	31	7.8	49	12.3	88	22.1	172	43.1	49	12.3	10	2.5	185	46.4
	総 数	2,321	100.0	789	34.0	398	17.1	434	18.7	517	22.3	150	6.5	33	1.4	435	18.7

注) 日本高血圧学会による血圧の分類は p.16 を参照。

(23年)

数値は2回の測定値の平均値を用いた。なお、1回しか測定できなかった者については、その値を採用した。

* 総数に対する割合

第 20 表の 2 血圧の状況（性・年齢階級別）〔服薬者除外〕

		総数		至適血圧		正常血圧		正常高値血圧		Ⅰ度高血圧		Ⅱ度高血圧		Ⅲ度高血圧		(再掲) 収縮期高血圧	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%*
総数	15-19歳	115	100.0	95	82.6	16	13.9	2	1.7	2	1.7	0	0.0	0	0.0	1	0.9
	20-29歳	218	100.0	167	76.6	26	11.9	14	6.4	10	4.6	0	0.0	1	0.5	3	1.4
	30-39歳	475	100.0	281	59.2	95	20	52	10.9	37	7.8	6	1.3	4	0.8	6	1.3
	40-49歳	440	100.0	209	47.5	91	20.7	66	15.0	57	13.0	11	2.5	6	1.4	19	4.3
	50-59歳	470	100.0	120	25.5	115	24.5	88	18.7	101	21.5	36	7.7	10	2.1	43	9.1
	60-69歳	587	100.0	93	15.8	115	19.6	156	26.6	170	29.0	38	6.5	15	2.6	103	17.5
	70歳以上	566	100.0	73	12.9	80	14.1	136	24.0	195	34.5	67	11.8	15	2.7	201	35.5
	(再掲) 65-74歳	522	100.0	74	14.2	83	15.9	149	28.5	163	31.2	42	8.0	11	2.1	142	27.2
	(再掲) 75歳以上	305	100.0	33	10.8	44	14.4	69	22.6	110	36.1	39	12.8	10	3.3	119	39.0
	総 数	2,871	100.0	1,038	36.2	538	18.7	514	17.9	572	19.9	158	5.5	51	1.8	376	13.1
男性	15-19歳	55	100.0	38	69.1	13	23.6	2	3.6	2	3.6	0	0.0	0	0.0	1	1.8
	20-29歳	79	100.0	41	51.9	20	25.3	10	12.7	7	8.9	0	0.0	1	1.3	3	3.8
	30-39歳	182	100.0	64	35.2	51	28.0	32	17.6	28	15.4	3	1.6	4	2.2	3	1.6
	40-49歳	155	100.0	53	34.2	32	20.6	33	21.3	28	18.1	5	3.2	4	2.6	8	5.2
	50-59歳	179	100.0	32	17.9	34	19.0	36	20.1	49	27.4	22	12.3	6	3.4	18	10.1
	60-69歳	247	100.0	29	11.7	38	15.4	60	24.3	91	36.8	22	8.9	7	2.8	48	19.4
	70歳以上	234	100.0	24	10.3	39	16.7	56	23.9	77	32.9	28	12.0	10	4.3	79	33.8
	(再掲) 65-74歳	207	100.0	22	10.6	35	16.9	56	27.1	71	34.3	18	8.7	5	2.4	58	28.0
	(再掲) 75歳以上	130	100.0	12	9.2	23	17.7	27	20.8	45	34.6	16	12.3	7	5.4	49	37.7
	総 数	1,131	100.0	281	24.8	227	20.1	229	20.2	282	24.9	80	7.1	32	2.8	160	14.1
女性	15-19歳	60	100.0	57	95.0	3	5.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	20-29歳	139	100.0	126	90.6	6	4.3	4	2.9	3	2.2	0	0.0	0	0.0	0	0.0
	30-39歳	293	100.0	217	74.1	44	15.0	20	6.8	9	3.1	3	1.0	0	0.0	3	1.0
	40-49歳	285	100.0	156	54.7	59	20.7	33	11.6	29	10.2	6	2.1	2	0.7	11	3.9
	50-59歳	291	100.0	88	30.2	81	27.8	52	17.9	52	17.9	14	4.8	4	1.4	25	8.6
	60-69歳	340	100.0	64	18.8	77	22.6	96	28.2	79	23.2	16	4.7	8	2.4	55	16.2
	70歳以上	332	100.0	49	14.8	41	12.3	80	24.1	118	35.5	39	11.7	5	1.5	122	36.7
	(再掲) 65-74歳	315	100.0	52	16.5	48	15.2	93	29.5	92	29.2	24	7.6	6	1.9	84	26.7
	(再掲) 75歳以上	175	100.0	21	12.0	21	12.0	42	24.0	65	37.1	23	13.1	3	1.7	70	40.0
	総 数	1,740	100.0	757	43.5	311	17.9	285	16.4	290	16.7	78	4.5	19	1.1	216	12.4

注) 血圧を下げる薬服用者除外。日本高血圧学会による血圧の分類は p.16 を参照。

(23年)

数値は 2 回の測定値の平均値を用いた。なお、1 回しか測定できなかった者については、その値を採用した。

* 総数に対する割合

第21表の1 収縮期（最高）・拡張期（最低）血圧の分布（性・年齢階級別）

			総 数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-89歳	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	最高血圧	90mmHg未満	2	0.1	2	3.6	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		90-99	20	1.2	4	7.3	3	3.8	3	1.6	3	1.8	2	0.8	2	0.5	3	0.6	10	0.8
		100-109	92	5.6	14	25.5	12	15.2	21	11.2	22	13.4	9	3.8	6	1.5	8	1.5	45	3.4
		110-119	241	14.7	20	36.4	30	38.0	52	27.7	40	24.4	35	14.8	32	8.2	32	6.1	139	10.6
		120-129	313	19.1	11	20.0	20	25.3	55	29.3	36	22.0	49	20.8	62	16.0	80	15.2	227	17.4
		130-139	366	22.3	2	3.6	8	10.1	36	19.1	35	21.3	64	27.1	102	26.3	119	22.5	319	24.4
		140-149	270	16.5	2	3.6	4	5.1	16	8.5	14	8.5	30	12.7	84	21.6	120	22.7	245	18.7
		150-159	168	10.3	0	0.0	1	1.3	2	1.1	8	4.9	28	11.9	53	13.7	76	14.4	163	12.5
		160-169	78	4.8	0	0.0	1	1.3	0	0.0	1	0.6	7	3.0	28	7.2	41	7.8	77	5.9
		170-179	56	3.4	0	0.0	0	0.0	2	1.1	0	0.0	9	3.8	12	3.1	33	6.3	53	4.1
		180mmHg以上	32	2.0	0	0.0	0	0.0	1	0.5	5	3.0	3	1.3	7	1.8	16	3.0	30	2.3
		総 数	1,638	100.0	55	100.0	79	100.0	188	100.0	164	100.0	236	100.0	388	100.0	528	100.0	1,308	100.0
		平均値(mmHg)	135.0		113.5		119.6		124.5		126.4		134.8		139.6		142.6		138.2	
		標準偏差	19.1		12.3		12.2		14.2		18.8		17.3		16.6		18.9		18.7	
女性	最低血圧	40mmHg未満	1	0.1	1	1.8	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		40-49	3	0.2	2	3.6	1	1.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		50-59	29	1.8	5	9.1	2	2.5	3	1.6	0	0.0	1	0.4	2	0.5	16	3.0	19	1.5
		60-69	197	12.0	15	27.3	26	32.9	23	12.2	20	12.2	16	6.8	28	7.2	69	13.1	133	10.2
		70-79	534	32.6	28	50.9	30	38.0	76	40.4	54	32.9	52	22.0	110	28.4	184	34.8	397	30.4
		80-89	515	31.4	3	5.5	15	19.0	51	27.1	56	34.1	89	37.7	135	34.8	166	31.4	442	33.8
		90-99	269	16.4	1	1.8	4	5.1	29	15.4	21	12.8	55	23.3	85	21.9	74	14.0	235	18.0
		100-109	65	4.0	0	0.0	0	0.0	2	1.1	8	4.9	18	7.6	24	6.2	13	2.5	62	4.7
		110-119	19	1.2	0	0.0	1	1.3	2	1.1	3	1.8	4	1.7	4	1.0	5	0.9	16	1.2
		120-129	3	0.2	0	0.0	0	0.0	1	0.5	1	0.6	1	0.4	0	0.0	0	0.0	2	0.2
		130-139	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		140mmHg以上	3	0.2	0	0.0	0	0.0	1	0.5	1	0.6	0	0.0	0	0.0	1	0.2	2	0.2
		総 数	1,638	100.0	55	100.0	79	100.0	188	100.0	164	100.0	236	100.0	388	100.0	528	100.0	1,308	100.0
		平均値(mmHg)	81.2		69.3		74.3		80.2		82.4		85.8		83.6		79.4		82.2	
		標準偏差	11.7		10.1		10.1		11.6		12.4		10.9		10.6		11.5		11.5	
女性	最高血圧	90mmHg未満	11	0.5	3	5.0	5	3.6	0	0.0	1	0.3	1	0.3	1	0.2	0	0.0	3	0.2
		90-99	111	4.8	14	23.3	32	22.9	37	12.5	15	5.1	12	3.4	1	0.2	0	0.0	28	1.5
		100-109	328	14.1	27	45.0	58	41.4	106	35.9	68	23.2	34	9.5	23	4.4	12	1.8	136	7.5
		110-119	405	17.4	15	25.0	36	25.7	87	29.5	85	29.0	67	18.8	61	11.7	54	8.2	267	14.7
		120-129	399	17.2	1	1.7	5	3.6	43	14.6	58	19.8	87	24.4	114	21.9	91	13.9	348	19.2
		130-139	432	18.6	0	0.0	4	2.9	16	5.4	34	11.6	70	19.6	153	29.4	155	23.6	411	22.7
		140-149	306	13.2	0	0.0	0	0.0	4	1.4	18	6.1	36	10.1	83	16.0	165	25.2	299	16.5
		150-159	180	7.8	0	0.0	0	0.0	2	0.7	9	3.1	31	8.7	51	9.8	87	13.3	174	9.6
		160-169	92	4.0	0	0.0	0	0.0	0	0.0	3	1.0	11	3.1	21	4.0	57	8.7	91	5.0
		170-179	32	1.4	0	0.0	0	0.0	0	0.0	1	0.3	5	1.4	5	1.0	21	3.2	30	1.7
		180mmHg以上	25	1.1	0	0.0	0	0.0	0	0.0	1	0.3	3	0.8	7	1.3	14	2.1	25	1.4
		総 数	2,321	100.0	60	100.0	140	100.0	295	100.0	293	100.0	357	100.0	520	100.0	656	100.0	1,812	100.0
		平均値(mmHg)	128.0		104.1		106.0		111.8		119.3		128.7		134.1		140.8		133.0	
		標準偏差	19.6		8.5		9.8		11.4		15.8		18.1		15.8		17.1		18.3	
女性	最低血圧	40mmHg未満	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.3	2	0.1
		40-49	8	0.3	1	1.7	6	4.3	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	1	0.1
		50-59	106	4.6	20	33.3	25	17.9	29	9.8	6	2.0	5	1.4	3	0.6	18	2.7	31	1.7
		60-69	499	21.5	28	46.7	63	45.0	96	32.5	75	25.6	45	12.6	73	14.0	119	18.1	310	17.1
		70-79	810	34.9	8	13.3	37	26.4	120	40.7	118	40.3	117	32.8	179	34.4	231	35.2	639	35.3
		80-89	631	27.2	3	5.0	6	4.3	40	13.6	63	21.5	127	35.6	182	35.0	210	32.0	580	32.0
		90-99	201	8.7	0	0.0	3	2.1	7	2.4	21	7.2	49	13.7	60	11.5	61	9.3	189	10.4
		100-109	53	2.3	0	0.0	0	0.0	3	1.0	8	2.7	12	3.4	17	3.3	13	2.0	50	2.8
		110-119	9	0.4	0	0.0	0	0.0	0	0.0	1	0.3	2	0.6	5	1.0	1	0.2	8	0.4
		120-129	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.1
		130-139	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.1
		140mmHg以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		総 数	2,321	100.0	60	100.0	140	100.0	295	100.0	293	100.0	357	100.0	520	100.0	656	100.0	1,812	100.0
		平均値(mmHg)	76.6		63.2		66.1		71.2		75.9		80.4		80.3		77.8		78.7	
		標準偏差	11.2		7.7		9.7		9.4		10.3		10.4		10.2		10.8		10.5	

注) 数値は2回の測定値の平均値を用いた。なお、1回しか測定できなかった者については、その値を採用した。

(23年)

第 21 表の 2 収縮期（最高）・拡張期（最低）血圧の分布（性・年齢階級別）〔服薬者除外〕

			総 数		15－19歳		20－29歳		30－39歳		40－49歳		50－59歳		60－69歳		70歳以上		(再掲) 40－89歳	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	最高血圧	90mmHg未満	2	0.2	2	3.6	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		90－99	18	1.6	4	7.3	3	3.8	3	1.6	3	1.9	2	1.1	2	0.8	1	0.4	8	1.0
		100－109	89	7.9	14	25.5	12	15.2	21	11.5	22	14.2	9	5.0	6	2.4	5	2.1	42	5.2
		110－119	221	19.5	20	36.4	30	38.0	52	28.6	39	25.2	29	16.2	28	11.3	23	9.8	119	14.7
		120－129	238	21.0	11	20.0	20	25.3	54	29.7	34	21.9	38	21.2	41	16.6	40	17.1	153	18.9
		130－139	242	21.4	2	3.6	8	10.1	32	17.6	33	21.3	44	24.6	68	27.5	55	23.5	199	24.5
		140－149	146	12.9	2	3.6	4	5.1	15	8.2	13	8.4	23	12.8	46	18.6	43	18.4	125	15.4
		150－159	90	8.0	0	0.0	1	1.3	2	1.1	6	3.9	18	10.1	31	12.6	32	13.7	85	10.5
		160－169	35	3.1	0	0.0	1	1.3	0	0.0	1	0.6	6	3.4	14	5.7	13	5.6	34	4.2
		170－179	30	2.7	0	0.0	0	0.0	2	1.1	0	0.0	8	4.5	6	2.4	14	6.0	28	3.5
		180mmHg以上	20	1.8	0	0.0	0	0.0	1	0.5	4	2.6	2	1.1	5	2.0	8	3.4	18	2.2
		総 数	1,131	100.0	55	100.0	79	100.0	182	100.0	155	100.0	179	100.0	247	100.0	234	100.0	811	100.0
		平均値(mmHg)	131.3		113.5		119.6		124.2		125.5		134.0		137.5		140.3		135.2	
		標準偏差	18.9		12.3		12.2		14.3		18.4		18.0		17.1		20.0		19.1	
	最低血圧	40mmHg未満	1	0.1	1	1.8	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		40－49	3	0.3	2	3.6	1	1.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		50－59	16	1.4	5	9.1	2	2.5	3	1.6	0	0.0	1	0.6	2	0.8	3	1.3	6	0.7
		60－69	143	12.6	15	27.3	26	32.9	23	12.6	20	12.9	15	8.4	18	7.3	26	11.1	79	9.7
		70－79	385	34.0	28	50.9	30	38.0	75	41.2	53	34.2	41	22.9	74	30.0	84	35.9	251	30.9
		80－89	349	30.9	3	5.5	15	19.0	49	26.9	53	34.2	63	35.2	81	32.8	85	36.3	280	34.5
		90－99	177	15.6	1	1.8	4	5.1	26	14.3	20	12.9	39	21.8	57	23.1	30	12.8	146	18.0
		100－109	37	3.3	0	0.0	0	0.0	2	1.1	6	3.9	15	8.4	11	4.5	3	1.3	34	4.2
		110－119	14	1.2	0	0.0	1	1.3	2	1.1	1	0.6	4	2.2	4	1.6	2	0.9	11	1.4
		120－129	3	0.3	0	0.0	0	0.0	1	0.5	1	0.6	1	0.6	0	0.0	0	0.0	2	0.2
女性	最高血圧	130－139	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		140mmHg以上	3	0.3	0	0.0	0	0.0	1	0.5	1	0.6	0	0.0	0	0.0	1	0.4	2	0.2
		総 数	1,131	100.0	55	100.0	79	100.0	182	100.0	155	100.0	179	100.0	247	100.0	234	100.0	811	100.0
		平均値(mmHg)	80.9		69.3		74.3		80.0		81.6		85.4		83.4		80.1		82.5	
		標準偏差	11.8		10.1		10.1		11.6		12.0		11.6		10.8		11.0		11.4	
	最低血圧	40mmHg未満	11	0.6	3	5.0	5	3.6	0	0.0	1	0.4	1	0.3	1	0.3	0	0.0	3	0.2
		90－99	110	6.3	14	23.3	32	23.0	36	12.3	15	5.3	12	4.1	1	0.3	0	0.0	28	2.3
		100－109	322	18.5	27	45.0	58	41.7	106	36.2	68	23.9	33	11.3	20	5.9	10	3.0	130	10.5
		110－119	372	21.4	15	25.0	35	25.2	87	29.7	85	29.8	58	19.9	52	15.3	40	12.0	235	18.9
		120－129	315	18.1	1	1.7	5	3.6	43	14.7	58	20.4	75	25.8	84	24.7	49	14.8	265	21.4
		130－139	269	15.5	0	0.0	4	2.9	15	5.1	29	10.2	54	18.6	91	26.8	76	22.9	250	20.1
		140－149	174	10.0	0	0.0	0	0.0	4	1.4	18	6.3	28	9.6	50	14.7	74	22.3	168	13.5
		150－159	94	5.4	0	0.0	0	0.0	2	0.7	7	2.5	18	6.2	26	7.6	41	12.3	92	7.4
		160－169	47	2.7	0	0.0	0	0.0	0	0.0	2	0.7	8	2.7	8	2.4	29	8.7	46	3.7
		170－179	13	0.7	0	0.0	0	0.0	0	0.0	1	0.4	1	0.3	2	0.6	9	2.7	11	0.9
		180mmHg以上	13	0.7	0	0.0	0	0.0	0	0.0	1	0.4	3	1.0	5	1.5	4	1.2	13	1.0
		総 数	1,740	100.0	60	100.0	139	100.0	293	100.0	285	100.0	291	100.0	340	100.0	332	100.0	1,241	100.0
		平均値(mmHg)	123.5		104.1		106.0		111.8		118.6		126.3		131.5		138.3		129.0	
		標準偏差	18.6		8.5		9.8		11.3		15.3		17.6		15.7		17.1		17.8	
	最低血圧	40mmHg未満	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	1	0.1
		40－49	8	0.5	1	1.7	6	4.3	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0	1	0.1
		50－59	94	5.4	20	33.3	25	18.0	29	9.9	6	2.1	5	1.7	1	0.3	8	2.4	19	1.5
		60－69	428	24.6	28	46.7	63	45.3	95	32.4	75	26.3	41	14.1	58	17.1	68	20.5	241	19.4
		70－79	612	35.2	8	13.3	36	25.9	120	41.0	118	41.4	102	35.1	119	35.0	109	32.8	446	35.9
		80－89	426	24.5	3	5.0	6	4.3	40	13.7	59	20.7	98	33.7	114	33.5	106	31.9	376	30.3
		90－99	129	7.4	0	0.0	3	2.2	6	2.0	19	6.7	33	11.3	35	10.3	33	9.9	119	9.6
		100－109	34	2.0	0	0.0	0	0.0	3	1.0	6	2.1	11	3.8	9	2.6	5	1.5	31	2.5
		110－119	7	0.4	0	0.0	0	0.0	0	0.0	1	0.4	1	0.3	4	1.2	1	0.3	6	0.5
		120－129	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		130－139	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	1	0.1
		140mmHg以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		総 数	1,740	100.0	60	100.0	139	100.0	293	100.0	285	100.0	291	100.0	340	100.0	332	100.0	1,241	100.0
		平均値(mmHg)	75.5		63.2		66.0		71.1		75.4		79.4		79.5		77.9		78.1	
		標準偏差	11.1		7.7		9.6		9.4		10.0		10.5		10.0		11.0		10.5	

注) 血圧を下げる薬服用者除外。

(23年)

数値は 2 回の測定値の平均値を用いた。なお、1 回しか測定できなかった者については、その値を採用した。

第22表の1 血色素量の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	100g/dL未満	11	0.8	0	0.0	0	0.0	0	0.0	0	0.0	3	0.8	8	1.7	5	1.4
	100-109	14	1.0	0	0.0	0	0.0	0	0.0	2	0.9	1	0.3	11	2.3	3	0.8
	110-119	49	3.4	0	0.0	1	0.6	0	0.0	1	0.5	9	2.5	38	7.9	11	3.1
	120-129	93	6.4	0	0.0	1	0.6	2	1.3	10	4.7	19	5.3	61	12.7	27	7.5
	130-139	233	16.0	2	2.7	16	8.8	12	7.8	31	14.4	59	16.6	113	23.5	60	16.7
	140-149	436	29.9	9	12.3	50	27.6	51	33.3	64	29.8	136	38.2	126	26.2	131	36.5
	150-159	416	28.5	34	46.6	78	43.1	52	34.0	71	33.0	94	26.4	87	18.1	84	23.4
	160-169	165	11.3	22	30.1	26	14.4	33	21.6	27	12.6	28	7.9	29	6.0	28	7.8
	170-179	35	2.4	5	6.8	9	5.0	3	2.0	8	3.7	4	1.1	6	1.2	6	1.7
	180g/dL以上	7	0.5	1	1.4	0	0.0	0	0.0	1	0.5	3	0.8	2	0.4	4	1.1
	総 数	1,459	100.0	73	100.0	181	100.0	153	100.0	215	100.0	356	100.0	481	100.0	359	100.0
女性	100g/dL未満	50	2.4	1	0.8	10	3.7	20	7.2	2	0.6	0	0.0	17	2.9	4	0.9
	100-109	68	3.3	6	4.8	12	4.4	12	4.3	5	1.5	6	1.2	27	4.6	9	1.9
	110-119	229	11.0	11	8.8	33	12.1	29	10.4	36	8.3	36	7.3	93	15.8	50	10.8
	120-129	635	30.5	39	31.2	98	36.0	73	26.3	90	27.5	144	29.3	191	32.4	131	28.4
	130-139	696	33.4	48	38.4	87	32.0	92	33.1	128	39.1	170	34.6	171	29.0	166	35.9
	140-149	344	16.5	18	14.4	29	10.7	46	16.5	60	18.3	121	24.6	70	11.9	82	17.7
	150-159	56	2.7	2	1.6	3	1.1	6	2.2	14	4.3	14	2.8	17	2.9	17	3.7
	160-169	5	0.2	0	0.0	0	0.0	0	0.0	1	0.3	1	0.2	3	0.5	3	0.6
	170-179	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	180g/dL以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	総 数	2,083	100.0	125	100.0	272	100.0	278	100.0	327	100.0	492	100.0	589	100.0	462	100.0
	平均値(g/dL)	12.9		12.9		12.7		12.7		13.2		13.3		12.8		13.1	
	標準偏差	1.2		1.0		1.2		1.5		1.1		1.0		1.3		1.1	

(23年)

第22表の2 血色素量の分布（性・年齢階級別）〔服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	100g/dL未満	10	0.7	0	0.0	0	0.0	0	0.0	0	0.0	2	0.6	8	1.7	4	1.1
	100-109	13	0.9	0	0.0	0	0.0	0	0.0	2	0.9	1	0.3	10	2.1	3	0.8
	110-119	46	3.2	0	0.0	1	0.6	0	0.0	1	0.5	8	2.3	36	7.7	10	2.8
	120-129	90	6.3	0	0.0	1	0.6	2	1.3	10	4.7	18	5.1	59	12.6	27	7.6
	130-139	226	15.7	2	2.7	16	8.8	12	7.8	31	14.5	58	16.5	107	22.9	57	16.1
	140-149	433	30.1	9	12.3	50	27.6	51	33.3	63	29.4	136	38.7	124	26.6	131	37.0
	150-159	414	28.8	34	46.6	78	43.1	52	34.0	71	33.2	93	26.5	86	18.4	84	23.7
	160-169	165	11.5	22	30.1	26	14.4	33	21.6	27	12.6	28	8.0	29	6.2	28	7.9
	170-179	35	2.4	5	6.8	9	5.0	3	2.0	8	3.7	4	1.1	6	1.3	6	1.7
	180g/dL以上	7	0.5	1	1.4	0	0.0	0	0.0	1	0.5	3	0.9	2	0.4	4	1.1
	総 数	1,439	100.0	73	100.0	181	100.0	153	100.0	214	100.0	351	100.0	467	100.0	354	100.0
女性	100g/dL未満	47	2.3	1	0.8	10	3.8	18	6.7	2	0.6	0	0.0	16	2.8	4	0.9
	100-109	63	3.1	6	4.8	10	3.8	11	4.1	5	1.5	6	1.2	25	4.4	9	2.0
	110-119	221	10.8	11	8.8	32	12.1	27	10.1	27	8.3	35	7.2	89	15.6	47	10.4
	120-129	619	30.4	39	31.2	95	36.0	70	26.2	87	26.9	141	29.0	187	32.7	128	28.3
	130-139	691	33.9	48	38.4	86	32.6	91	34.1	128	39.5	170	35.0	168	29.4	165	36.4
	140-149	338	16.6	18	14.4	28	10.6	44	16.5	60	18.5	120	24.7	68	11.9	80	17.7
	150-159	55	2.7	2	1.6	3	1.1	6	2.2	14	4.3	13	2.7	17	3.0	17	3.8
	160-169	4	0.2	0	0.0	0	0.0	0	0.0	1	0.3	1	0.2	2	0.3	3	0.7
	170-179	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	180g/dL以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	総 数	2,038	100.0	125	100.0	264	100.0	267	100.0	324	100.0	486	100.0	572	100.0	453	100.0
	平均値(g/dL)	13.0		12.9		12.7		12.8		13.2		13.3		12.8		13.1	
	標準偏差	1.2		1.0		1.2		1.5		1.1		1.0		1.2		1.1	

注) 貧血治療のための薬服用者除外。

(23年)

第 23 表 赤血球数の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	400万個/mm ³ 未満	124	8.5	0	0.0	0	0.0	0	0.0	9	4.2	23	6.5	92	19.1	29	8.1	71	24.7
	400-449	328	22.5	0	0.0	16	8.8	9	5.9	43	20.0	102	28.7	158	32.8	105	29.2	103	35.8
	450-499	619	42.4	21	28.8	76	42.0	76	49.7	99	46.0	176	49.4	171	35.6	162	45.1	87	30.2
	500-449	351	24.1	45	61.6	83	45.9	62	40.5	58	27.0	50	14.0	53	11.0	59	16.4	21	7.3
	550-599	36	2.5	7	9.6	6	3.3	6	3.9	6	2.8	4	1.1	7	1.5	3	0.8	6	2.1
	600-649	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	1	0.3	0	0.0
	650万個/mm ³ 以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	総 数	1,459	100.0	73	100.0	181	100.0	153	100.0	215	100.0	356	100.0	481	100.0	359	100.0	288	100.0
	平均値(万個/mm ³) 標準偏差	469.3 48.4		517.5 28.0		499.0 32.5		495.3 30.9		476.1 43.5		463.5 41.3		443.8 50.8		459.7 44.9		436.1 52.0	
女性	400万個/mm ³ 未満	333	16.0	11	8.8	27	9.9	31	11.2	29	8.9	62	12.6	173	29.4	82	17.7	124	34.9
	400-449	1,048	50.3	68	54.4	160	58.8	144	51.8	174	53.2	229	46.5	273	46.3	227	49.1	157	44.2
	450-499	623	29.9	43	34.4	70	25.7	93	33.5	103	31.5	186	37.8	128	21.7	141	30.5	66	18.6
	500-449	76	3.6	3	2.4	14	5.1	10	3.6	19	5.8	15	3.0	15	2.5	12	2.6	8	2.3
	550-599	3	0.1	0	0.0	1	0.4	0	0.0	2	0.6	0	0.0	0	0.0	0	0.0	0	0.0
	600-649	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	650万個/mm ³ 以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	総 数	2,083	100.0	125	100.0	272	100.0	278	100.0	327	100.0	492	100.0	589	100.0	462	100.0	355	100.0
	平均値(万個/mm ³) 標準偏差	434.4 37.5		440.8 31.8		438.5 33.7		439.9 34.2		442.1 35.9		439.6 34.3		419.8 41.2		433.4 36.3		413.6 42.4	

(23年)

第 24 表 ヘマトクリット値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	25.0%未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	25.0-29.9	6	0.4	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	5	1.0	2	0.6	3	1.0
	30.0-34.9	28	1.9	0	0.0	0	0.0	0	0.0	1	0.5	3	0.8	24	5.0	6	1.7	19	6.6
	35.0-39.9	130	8.9	0	0.0	2	1.1	2	1.3	12	5.6	31	8.7	83	17.3	37	10.3	64	22.2
	40.0-44.9	580	39.8	12	16.4	64	35.4	54	35.3	89	41.4	158	44.4	203	42.2	155	43.2	123	42.7
	45.0-49.9	626	42.9	51	69.9	93	51.4	89	58.2	101	47.0	148	41.6	144	29.9	141	39.3	64	22.2
	50.0-54.9	84	5.8	9	12.3	21	11.6	8	5.2	12	5.6	15	4.2	19	4.0	18	5.0	12	4.2
	55.0%以上	5	0.3	1	1.4	1	0.6	0	0.0	0	0.0	0	0.0	3	0.6	0	0.0	3	1.0
	(再掲)39.8未満	119	8.2	0	0.0	1	0.6	1	0.7	8	3.7	22	6.2	87	18.1	32	8.9	66	22.9
	(再掲)51.9以上	25	1.7	2	2.7	6	3.3	0	0.0	4	1.9	5	1.4	8	1.7	6	1.7	7	2.4
	総 数	1,459	100.0	73	100.0	181	100.0	153	100.0	215	100.0	356	100.0	481	100.0	359	100.0	288	100.0
	平均値(%) 標準偏差	44.5 4.0		47.3 2.5		46.3 3.1		45.9 2.8		45.1 3.3		44.4 3.6		42.9 4.7		44.2 4.0		42.2 4.8	
女性	25.0%未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	25.0-29.9	17	0.8	0	0.0	3	1.1	6	2.2	1	0.3	0	0.0	7	1.2	1	0.2	6	1.7
	30.0-34.9	101	4.8	5	4.0	17	6.3	23	8.3	6	1.8	7	1.4	43	7.3	12	2.6	34	9.6
	35.0-39.9	746	35.8	54	43.2	112	41.2	107	38.5	96	29.4	147	29.9	230	39.0	148	32.0	153	43.1
	40.0-44.9	1,051	50.5	63	50.4	127	46.7	123	44.2	191	58.4	283	57.5	264	44.8	253	54.8	140	39.4
	45.0-49.9	163	7.8	3	2.4	13	4.8	19	6.8	32	9.8	55	11.2	41	7.0	44	9.5	22	6.2
	50.0-54.9	5	0.2	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	4	0.7	4	0.9	0	0.0
	55.0%以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	(再掲)33.4未満	58	2.8	4	3.2	10	3.7	18	6.5	3	0.9	0	0.0	23	3.9	3	0.6	20	5.6
	(再掲)45.0以上	168	8.1	3	2.4	13	4.8	19	6.8	33	10.1	55	11.2	45	7.6	48	10.4	22	6.2
	総 数	2,083	100.0	125	100.0	272	100.0	278	100.0	327	100.0	492	100.0	589	100.0	462	100.0	355	100.0
	平均値(%) 標準偏差	40.5 3.4		39.9 2.8		39.7 3.3		39.8 3.8		41.2 3.0		41.5 2.9		40.0 3.8		41.1 3.3		39.5 3.8	

(23年)

第25表 平均赤血球容積（MCV）の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	70f未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	70-75未満	3	0.2	0	0.0	0	0.0	0	0.0	1	0.5	1	0.3	1	0.2	1	0.3	1	0.3
	75-80未満	4	0.3	0	0.0	0	0.0	0	0.0	0	0.0	2	0.6	2	0.4	0	0.0	2	0.7
	80-85未満	26	1.8	2	2.7	2	1.1	4	2.6	6	2.8	6	1.7	6	1.2	6	1.7	3	1.0
	85-90未満	193	13.2	25	34.2	44	24.3	30	19.6	24	11.2	31	8.7	39	8.1	32	8.9	23	8.0
	90-95未満	509	34.9	33	45.2	81	44.8	78	51.0	79	36.7	108	30.3	130	27.0	106	29.5	77	26.7
	95-100未満	455	31.2	13	17.8	46	25.4	32	20.9	70	32.6	126	35.4	168	34.9	125	34.8	99	34.4
	100-105未満	211	14.5	0	0.0	7	3.9	9	5.9	29	13.5	64	18.0	102	21.2	71	19.8	58	20.1
	105-110未満	42	2.9	0	0.0	1	0.6	0	0.0	2	0.9	15	4.2	24	5.0	16	4.5	17	5.9
	110-115未満	13	0.9	0	0.0	0	0.0	0	0.0	3	1.4	2	0.6	8	1.7	0	0.0	8	2.8
	115-120未満	2	0.1	0	0.0	0	0.0	0	0.0	1	0.5	1	0.3	0	0.0	1	0.3	0	0.0
	120f以上	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.3	0	0.0
総 数		1,459	100.0	73	100.0	181	100.0	153	100.0	215	100.0	356	100.0	481	100.0	359	100.0	288	100.0
平均値(f)		95.2		91.4		92.8		92.8		95.0		96.1		96.8		96.3		97.0	
標準偏差		5.5		3.5		4.1		4.1		5.7		5.6		5.7		5.5		5.9	
女性	70f未満	11	0.5	0	0.0	5	1.8	5	1.8	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0
	70-75未満	14	0.7	1	0.8	6	2.2	7	2.5	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	75-80未満	29	1.4	3	2.4	9	3.3	14	5.0	1	0.3	0	0.0	2	0.3	1	0.2	1	0.3
	80-85未満	81	3.9	7	5.6	16	5.9	19	6.8	15	4.6	6	1.2	18	3.1	11	2.4	11	3.1
	85-90未満	342	16.4	36	28.8	66	24.3	65	23.4	59	18.0	62	12.6	54	9.2	45	9.7	32	9.0
	90-95未満	806	38.7	56	44.8	104	38.2	98	35.3	129	39.4	222	45.1	197	33.4	183	39.6	112	31.5
	95-100未満	553	26.5	20	16.0	49	18.0	54	19.4	87	26.6	138	28.0	205	34.8	151	32.7	122	34.4
	100-105未満	201	9.6	1	0.8	15	5.5	13	4.7	31	9.5	51	10.4	90	15.3	58	12.6	62	17.5
	105-110未満	40	1.9	1	0.8	2	0.7	2	0.7	4	1.2	12	2.4	19	3.2	10	2.2	13	3.7
	110-115未満	5	0.2	0	0.0	0	0.0	1	0.4	0	0.0	1	0.2	3	0.5	3	0.6	1	0.3
	115-120未満	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.3
	120f以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
総 数		2,083	100.0	125	100.0	272	100.0	278	100.0	327	100.0	492	100.0	589	100.0	462	100.0	355	100.0
平均値(f)		93.4		90.7		90.8		90.6		93.4		94.5		95.4		95.0		95.7	
標準偏差		6.0		5.0		6.8		7.1		5.3		4.6		5.4		5.0		5.5	

(23年)

第26表 平均赤血球血色素量（MCH）の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	20pg未満	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.3
	20-25未満	8	0.5	0	0.0	0	0.0	0	0.0	2	0.9	3	0.8	3	0.6	2	0.6	2	0.7
	25-27未満	20	1.4	0	0.0	0	0.0	3	2.0	1	0.5	6	1.7	10	2.1	9	2.5	6	2.1
	27-29未満	99	6.8	7	9.6	21	11.6	9	5.9	15	7.0	15	4.2	32	6.7	18	5.0	20	6.9
	29-31未満	526	36.1	43	58.9	97	53.6	83	54.2	71	33.0	102	28.7	130	27.0	102	28.4	79	27.4
	31-33未満	621	42.6	21	28.8	57	31.5	51	33.3	94	43.7	178	50.0	220	45.7	176	49.0	126	43.8
	33-35未満	161	11.0	2	2.7	6	3.3	7	4.6	28	13.0	40	11.2	78	16.2	42	11.7	52	18.1
	35-40未満	22	1.5	0	0.0	0	0.0	0	0.0	4	1.9	12	3.4	6	1.2	9	2.5	2	0.7
	40pg以上	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.3	0	0.0
	総 数	1,459	100.0	73	100.0	181	100.0	153	100.0	215	100.0	356	100.0	481	100.0	359	100.0	288	100.0
女性	20pg未満	10	0.5	1	0.8	4	1.5	4	1.4	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0
	20-25未満	66	3.2	2	1.6	18	6.6	26	9.4	7	2.1	2	0.4	11	1.9	6	1.3	6	1.7
	25-27未満	79	3.8	8	6.4	18	6.6	20	7.2	13	4.0	5	1.0	15	2.5	7	1.5	11	3.1
	27-29未満	299	14.4	32	25.6	60	22.1	42	15.1	54	16.5	61	12.4	50	8.5	42	9.1	33	9.3
	29-31未満	1,050	50.4	68	54.4	127	46.7	128	46.0	164	50.2	283	57.5	280	47.5	254	55.0	156	43.9
	31-33未満	516	24.8	13	10.4	40	14.7	50	18.0	81	24.8	123	25.0	209	35.5	136	29.4	131	36.9
	33-35未満	58	2.8	1	0.8	5	1.8	7	2.5	6	1.8	16	3.3	23	3.9	16	3.5	17	4.8
	35-40未満	5	0.2	0	0.0	0	0.0	1	0.4	1	0.3	2	0.4	1	0.2	1	0.2	1	0.3
	40pg以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	総 数	2,083	100.0	125	100.0	272	100.0	278	100.0	327	100.0	492	100.0	589	100.0	462	100.0	355	100.0
	平均値(pg)	29.9		29.3		29.0		29.0		29.9		30.3		30.4		30.3		30.5	
	標準偏差	2.2		1.9		2.7		3.0		1.8		1.5		1.8		1.6		1.8	

(23年)

第 27 表 平均赤血球血色素濃度（MCHC）の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	22%未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	22-24未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	24-26未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	26-28未満	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.3
	28-30未満	24	1.6	1	1.4	0	0.0	0	0.0	4	1.9	3	0.8	16	3.3	3	0.8	15	5.2
	30-32未満	327	22.4	6	8.2	44	24.3	22	14.4	35	16.3	83	23.3	137	28.5	93	25.9	84	29.2
	32-34未満	900	61.7	43	58.9	111	61.3	103	67.3	143	66.5	222	62.4	278	57.8	217	60.4	160	55.6
	34%以上	207	14.2	23	31.5	26	14.4	28	18.3	33	15.3	48	13.5	49	10.2	46	12.8	28	9.7
	総 数	1,459	100.0	73	100.0	181	100.0	153	100.0	215	100.0	356	100.0	481	100.0	359	100.0	288	100.0
	平均値(%) 標準偏差	32.7 1.2		33.3 1.1		32.8 1.0		33.0 1.0		32.9 1.1		32.7 1.1		32.4 1.3		32.6 1.1		32.3 1.3	
女性	22%未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	22-24未満	1	0.0	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	24-26未満	5	0.2	0	0.0	2	0.7	2	0.7	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0
	26-28未満	12	0.6	1	0.8	2	0.7	6	2.2	0	0.0	0.0	0.0	3	0.5	2	0.4	1	0.3
	28-30未満	118	5.7	5	4.0	24	8.8	20	7.2	18	5.5	20	4.1	31	5.3	22	4.8	21	5.9
	30-32未満	771	37.0	28	22.4	94	34.6	85	30.6	120	36.7	196	39.8	248	42.1	183	39.6	160	45.1
	32-34未満	1,116	53.6	87	69.6	137	50.4	154	55.4	177	54.1	270	54.9	291	49.4	247	53.5	164	46.2
	34%以上	60	2.9	4	3.2	12	4.4	11	4.0	11	3.4	6	1.2	16	2.7	8	1.7	9	2.5
	総 数	2,083	100.0	125	100.0	272	100.0	278	100.0	327	100.0	492	100.0	589	100.0	462	100.0	355	100.0
	平均値(%) 標準偏差	32.0 1.3		32.4 1.2		31.9 1.6		31.9 1.6		32.1 1.2		32.0 1.1		31.9 1.2		32.0 1.2		31.9 1.2	

(23年)

第 28 表 血清総たんぱく質値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75-79歳		(再掲) 80歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	6.0g/dL未満	3	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	2	0.4	1	0.3	1	0.7	0	0.0
	6.0-6.4	17	1.2	0	0.0	1	0.6	0	0.0	2	0.9	4	1.1	10	2.1	4	1.1	5	3.3	3	2.2
	6.5-6.9	148	10.1	2	2.7	8	4.4	17	11.0	17	7.9	39	11.0	65	13.4	48	13.3	19	12.6	18	12.9
	7.0-7.4	640	43.7	18	24.7	71	39.2	68	44.2	97	45.1	169	47.5	217	44.8	158	43.9	68	45.0	66	47.5
	7.5-7.9	517	35.3	45	61.6	82	45.3	59	38.3	78	36.3	113	31.7	140	28.9	112	31.1	40	26.5	38	27.3
	8.0-8.4	129	8.8	8	11.0	18	9.9	9	5.8	19	8.8	28	7.9	47	9.7	34	9.4	17	11.3	14	10.1
	8.5g/dL以上	9	0.6	0	0.0	1	0.6	1	0.6	2	0.9	2	0.6	3	0.6	3	0.8	1	0.7	0	0.0
	(再掲)6.5未満	20	1.4	0	0.0	1	0.6	0	0.0	2	0.9	5	1.4	12	2.5	5	1.4	6	4.0	3	2.2
	総 数	1,467	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	360	100.0	151	100.0	139	100.0
	平均値(g/dL) 標準偏差	7.4 0.4		7.6 0.3		7.5 0.4		7.4 0.4		7.4 0.4		7.4 0.4		7.4 0.4		7.4 0.4		7.3 0.5		7.4 0.4	
女性	6.0g/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	6.0-6.4	18	0.9	2	1.6	6	2.2	2	0.7	1	0.3	1	0.2	6	1.0	1	0.2	3	1.7	3	1.7
	6.5-6.9	210	10.0	7	5.5	21	7.7	21	7.6	25	7.6	54	10.9	82	13.8	51	11.0	25	13.8	32	17.9
	7.0-7.4	875	41.8	46	36.2	121	44.3	125	45.1	138	42.1	193	39.1	252	42.4	197	42.5	76	42.0	69	38.5
	7.5-7.9	782	37.3	59	46.5	104	38.1	106	38.3	128	39.0	179	36.2	206	34.6	164	35.3	63	34.8	63	35.2
	8.0-8.4	190	9.1	13	10.2	20	7.3	22	7.9	31	9.5	59	11.9	45	7.6	47	10.1	13	7.2	10	5.6
	8.5g/dL以上	19	0.9	0	0.0	1	0.4	1	0.4	5	1.5	8	1.6	4	0.7	4	0.9	1	0.6	2	1.1
	(再掲)6.5未満	18	0.9	2	1.6	6	2.2	2	0.7	1	0.3	1	0.2	6	1.0	1	0.2	3	1.7	3	1.7
	総 数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	181	100.0	179	100.0
	平均値(g/dL) 標準偏差	7.4 0.4		7.5 0.4		7.4 0.4		7.4 0.4		7.5 0.4		7.5 0.4		7.4 0.4		7.4 0.4		7.4 0.4		7.4 0.4	

(23年)

第29表 血清アルブミン値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75-79歳		(再掲) 80歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	3.0g/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	3.0-3.1	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	2	0.6	0	0.0	0	0.0
	3.2-3.3	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.3	0	0.0	0	0.0
	3.4-3.5	5	0.3	0	0.0	0	0.0	0	0.0	1	0.5	0	0.0	4	0.8	1	0.3	2	1.3	1	0.7
	3.6-3.7	9	0.6	0	0.0	0	0.0	0	0.0	0	0.0	4	1.1	5	1.0	2	0.6	3	2.0	2	1.4
	3.8-3.9	35	2.4	0	0.0	1	0.6	0	0.0	3	1.4	4	1.1	27	5.6	5	1.4	11	7.3	12	8.6
	4.0-4.1	87	5.9	1	1.4	0	0.0	2	1.3	6	2.8	18	5.1	60	12.4	31	8.6	18	11.9	22	15.8
	4.2-4.3	221	15.1	1	1.4	3	1.7	5	3.2	22	10.2	69	19.4	121	25.0	84	23.3	40	26.5	38	27.3
	4.4-4.5	375	25.6	2	2.7	19	10.5	37	24.0	61	28.4	114	32.0	142	29.3	103	28.6	43	28.5	41	29.5
	4.6-4.7	370	25.3	10	13.7	60	33.1	57	37.0	63	29.3	100	28.1	80	16.5	85	23.6	18	11.9	17	12.2
	4.8-4.9	236	16.1	25	34.2	60	33.1	37	24.0	41	19.1	37	10.4	36	7.4	40	11.1	14	9.3	5	3.6
	5.0-5.1	99	6.8	21	28.8	32	17.7	14	9.1	17	7.9	10	2.8	5	1.0	6	1.7	1	0.7	1	0.7
	5.2g/dL以上 (再掲)3.5以下	23	1.6	13	17.8	6	3.3	2	1.3	1	0.5	0	0.0	1	0.2	0	0.0	1	0.7	0	0.0
総 数		1,463	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	360	100.0	151	100.0	139	100.0
平均値(g/dL) 標準偏差		4.5 0.3		4.9 0.3		4.8 0.2		4.7 0.2		4.6 0.3		4.5 0.2		4.4 0.3		4.4 0.3		4.4 0.3		4.3 0.3	
女性	3.0g/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	3.0-3.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	3.2-3.3	1	0.0	1	0.8	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	3.4-3.5	7	0.3	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0	6	1.0	0	0.0	4	2.2	2	1.1
	3.6-3.7	7	0.3	0	0.0	3	1.1	0	0.0	2	0.6	0	0.0	2	0.3	0	0.0	1	0.6	1	0.6
	3.8-3.9	44	2.1	3	2.4	3	1.1	3	1.1	4	1.2	8	1.6	23	3.9	11	2.4	7	3.9	11	6.1
	4.0-4.1	113	5.4	0	0.0	11	4.0	7	2.5	8	2.4	22	4.5	65	10.9	32	6.9	21	11.6	24	13.4
	4.2-4.3	360	17.2	11	8.7	34	12.5	37	13.4	36	11.0	82	16.6	160	26.9	96	20.7	57	31.5	50	27.9
	4.4-4.5	659	31.5	33	26.0	77	28.2	99	35.7	103	31.4	167	33.8	180	30.3	151	32.5	53	29.3	54	30.2
	4.6-4.7	549	26.2	31	24.4	82	30.0	88	31.8	105	32.0	127	25.7	116	19.5	119	25.6	32	17.7	25	14.0
	4.8-4.9	274	13.1	36	28.3	49	17.9	37	13.4	51	15.5	66	13.4	35	5.9	42	9.1	6	3.3	11	6.1
	5.0-5.1	71	3.4	11	8.7	13	4.8	4	1.4	17	5.2	20	4.0	6	1.0	11	2.4	0	0.0	1	0.6
	5.2g/dL以上 (再掲)3.5以下	9	0.4	1	0.8	0	0.0	2	0.7	2	0.6	2	0.4	2	0.3	2	0.4	0	0.0	0	0.0
総 数		2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	181	100.0	179	100.0
平均値(g/dL) 標準偏差		4.5 0.3		4.6 0.3		4.5 0.3		4.5 0.2		4.6 0.2		4.5 0.2		4.4 0.3		4.5 0.2		4.3 0.3		4.3 0.3	

(23年)

第 30 表の 1 ヘモグロビン A1c の分布 (性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	3.9%以下	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1	1	0.1	0	0.0
	4.0-4.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	4.2-4.3	17	0.5	4	2.0	2	0.4	4	0.9	2	0.4	2	0.2	3	0.3	0	0.0	3	0.5
	4.4-4.5	75	2.1	20	10.1	21	4.6	11	2.6	7	1.3	8	0.9	8	0.7	6	0.7	6	0.9
	4.6-4.7	254	7.2	47	23.6	75	16.6	49	11.4	26	4.8	30	3.5	27	2.5	24	2.9	14	2.2
	4.8-4.9	554	15.6	66	33.2	138	30.5	93	21.6	73	13.5	82	9.7	102	9.5	77	9.4	56	8.7
	5.0-5.1	731	20.6	38	19.1	115	25.4	117	27.1	127	23.4	160	18.9	174	16.2	140	17.0	106	16.4
	5.2	396	11.2	10	5.0	37	8.2	46	10.7	72	13.3	102	12.0	129	12.0	93	11.3	77	11.9
	5.3	310	8.7	5	2.5	22	4.9	44	10.2	49	9.0	85	10.0	105	9.8	84	10.2	62	9.6
	5.4	258	7.3	5	2.5	14	3.1	23	5.3	49	9.0	75	8.8	92	8.6	70	8.5	55	8.5
	5.5	215	6.1	1	0.5	10	2.2	9	2.1	44	8.1	65	7.7	86	8.0	77	9.4	53	8.2
	5.6	126	3.6	2	1.0	5	1.1	11	2.6	16	3.0	35	4.1	57	5.3	43	5.2	33	5.1
	5.7	104	2.9	0	0.0	2	0.4	3	0.7	17	3.1	31	3.7	51	4.8	28	3.4	35	5.4
	5.8	89	2.5	0	0.0	1	0.2	9	2.1	9	1.7	26	3.1	44	4.1	27	3.3	28	4.3
	5.9	61	1.7	0	0.0	3	0.7	0	0.0	5	0.9	23	2.7	30	2.8	27	3.3	15	2.3
	6.0	55	1.6	0	0.0	2	0.4	1	0.2	9	1.7	15	1.8	28	2.6	21	2.6	15	2.3
	6.1	37	1.0	0	0.0	1	0.2	1	0.2	5	0.9	14	1.7	16	1.5	9	1.1	13	2.0
	6.2-6.3	58	1.6	0	0.0	2	0.4	4	0.9	4	0.7	21	2.5	27	2.5	20	2.4	18	2.8
	6.4-6.5	45	1.3	0	0.0	1	0.2	0	0.0	7	1.3	19	2.2	18	1.7	17	2.1	10	1.6
	6.6-6.7	32	0.9	0	0.0	1	0.2	0	0.0	6	1.1	11	1.3	14	1.3	13	1.6	7	1.1
	6.8-6.9	30	0.8	0	0.0	1	0.2	0	0.0	2	0.4	4	0.5	23	2.1	12	1.5	13	2.0
	7.0-7.1	22	0.6	0	0.0	0	0.0	0	0.0	1	0.2	10	1.2	11	1.0	9	1.1	8	1.2
	7.2-7.3	9	0.3	0	0.0	0	0.0	0	0.0	2	0.4	4	0.5	3	0.3	3	0.4	2	0.3
	7.4-7.5	13	0.4	0	0.0	0	0.0	1	0.2	4	0.7	4	0.5	4	0.4	3	0.4	3	0.5
	7.6-7.7	13	0.4	0	0.0	0	0.0	0	0.0	2	0.4	7	0.8	4	0.4	4	0.5	3	0.5
	7.8-7.9	10	0.3	0	0.0	0	0.0	1	0.2	0	0.0	3	0.4	6	0.6	5	0.6	4	0.6
	8.0-8.1	3	0.1	1	0.5	0	0.0	0	0.0	0	0.0	0	0.0	2	0.2	2	0.2	0	0.0
	8.2-8.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.4-8.5	5	0.1	0	0.0	0	0.0	0	0.0	2	0.4	2	0.2	1	0.1	2	0.2	1	0.2
	8.6-8.7	4	0.1	0	0.0	0	0.0	0	0.0	0	0.0	2	0.2	2	0.2	0	0.0	2	0.3
	8.8-8.9	4	0.1	0	0.0	0	0.0	1	0.2	0	0.0	1	0.1	2	0.2	1	0.1	1	0.2
	9.0%以上	15	0.4	0	0.0	0	0.0	3	0.7	2	0.4	7	0.8	3	0.3	4	0.5	2	0.3
	(再掲)6.1以上	300	8.5	1	0.5	6	1.3	11	2.6	37	6.8	109	12.9	136	12.7	104	12.7	87	13.5
	総 数	3,546	100.0	199	100.0	453	100.0	431	100.0	542	100.0	848	100.0	1,073	100.0	822	100.0	645	100.0
	平均値 (%) 標準偏差	5.3 0.7		4.9 0.3		5.0 0.3		5.1 0.6		5.3 0.6		5.5 0.8		5.5 0.7		5.5 0.7		5.5 0.7	
男 性	3.9%以下	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.3	0	0.0
	4.0-4.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	4.2-4.3	7	0.5	0	0.0	1	0.6	3	2.0	1	0.5	1	0.3	1	0.2	0	0.0	1	0.3
	4.4-4.5	37	2.5	10	13.7	9	5.0	4	2.6	2	0.9	6	1.7	6	1.2	3	0.8	5	1.7
	4.6-4.7	99	6.8	23	31.5	28	15.5	12	7.8	7	3.3	16	4.5	13	2.7	10	2.8	7	2.4
	4.8-4.9	211	14.5	25	34.2	53	29.3	30	19.6	33	15.3	34	9.6	36	7.5	29	8.1	18	6.2
	5.0-5.1	264	18.1	9	12.3	43	23.8	41	26.8	43	20.0	60	16.9	68	14.1	62	17.3	35	12.1
	5.2	167	11.4	4	5.5	19	10.5	22	14.4	30	14.0	45	12.6	47	9.8	37	10.3	30	10.4
	5.3	110	7.5	1	1.4	7	3.9	13	8.5	15	7.0	33	9.3	41	8.5	31	8.6	27	9.3
	5.4	89	6.1	0	0.0	5	2.8	10	6.5	20	9.3	20	5.6	34	7.1	24	6.7	20	6.9
	5.5	93	6.4	0	0.0	6	3.3	1	0.7	19	8.8	29	8.1	38	7.9	28	7.8	26	9.0
	5.6	47	3.2	1	1.4	2	1.1	3	2.0	6	2.8	10	2.8	25	5.2	18	5.0	14	4.8
	5.7	46	3.2	0	0.0	1	0.6	1	0.7	7	3.3	9	2.5	28	5.8	11	3.1	19	6.6
	5.8	53	3.6	0	0.0	0	0.0	6	3.9	4	1.9	16	4.5	27	5.6	15	4.2	18	6.2
	5.9	33	2.3	0	0.0	3	1.7	0	0.0	3	1.4	10	2.8	17	3.5	18	5.0	6	2.1
	6.0	23	1.6	0	0.0	1	0.6	0	0.0	2	0.9	7	2.0	13	2.7	10	2.8	8	2.8
	6.1	21	1.4	0	0.0	0	0.0	1	0.7	4	1.9	8	2.2	8	1.7	6	1.7	6	2.1
	6.2-6.3	38	2.6	0	0.0	1	0.6	4	2.6	1	0.5	12	3.4	20	4.1	15	4.2	12	4.2
	6.4-6.5	24	1.6	0	0.0	1	0.6	0	0.0	4	1.9	7	2.0	12	2.5	7	1.9	7	2.4
	6.6-6.7	20	1.4	0	0.0	1	0.6	0	0.0	4	1.9	7	2.0	8	1.7	7	1.9	5	1.7
	6.8-6.9	15	1.0	0	0.0	0	0.0	0	0.0	0	0.0	3	0.8	12	2.5	6	1.7	7	2.4
	7.0-7.1	13	0.9	0	0.0	0	0.0	0	0.0	1	0.5	6	1.7	6	1.2	8	2.2	3	1.0
	7.2-7.3	7	0.5	0	0.0	0	0.0	0	0.0	1	0.5	3	0.8	3	0.6	2	0.6	2	0.7
	7.4-7.5	9	0.6	0	0.0	0	0.0	1	0.7	4	1.9	2	0.6	2	0.4	1	0.3	2	0.7
	7.6-7.7	8	0.5	0	0.0	0	0.0	0	0.0	2	0.9	3	0.8	3	0.6	2	0.6	2	0.7
	7.8-7.9	9	0.6	0	0.0	0	0.0	0	0.0	0	0.0	3	0.8	6	1.2	5	1.4	4	1.4
	8.0-8.1	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	2	0.6	0	0.0
	8.2-8.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.4-8.5	4	0.3	0	0.0	0	0.0	0	0.0	2	0.9	1	0.3	1	0.2	1	0.3	1	0.3
	8.6-8.7	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	1	0.2	0	0.0	1	0.3
	8.8-8.9	3	0.2	0	0.0	0	0.0	1	0.7	0	0.0	1	0.3	1	0.2	0	0.0	1	0.3
	9.0%以上	5	0.3	0	0.0	0	0.0	0	0.0	0	0.0	3	0.8	2	0.4	0	0.0	2	0.7
	(再掲)6.1以上	180	12.3	0	0.0	3	1.7	7	4.6	23	10.7	60	16.9	87	18.0	62	17.3	55	19.0
	総 数	1,460	100.0	73	100.0	181	100.0	153	100.0	215	100.0	356	100.0	482	100.0	359	100.0	289	100.0
	平均値 (%) 標準偏差	5.4 0.7		4.8 0.2		5.0 0.4		5.1 0.5		5.4 0.6		5.5 0.8		5.6 0.8		5.6 0.7		5.6 0.8	

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
女 性	3.9%以下	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	4.0-4.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	4.2-4.3	10	0.5	4	3.2	1	0.4	1	0.4	1	0.3	1	0.2	2	0.3	0	0.0	2	0.6
	4.4-4.5	38	1.8	10	7.9	12	4.4	7	2.5	5	1.5	2	0.4	2	0.3	3	0.6	1	0.3
	4.6-4.7	155	7.4	24	19.0	47	17.3	37	13.3	19	5.8	14	2.8	14	2.4	14	3.0	7	2.0
	4.8-4.9	343	16.4	41	32.5	85	31.3	63	22.7	40	12.2	48	9.8	66	11.2	48	10.4	38	10.7
	5.0-5.1	467	22.4	29	23.0	72	26.5	76	27.3	84	25.7	100	20.3	106	17.9	78	16.8	71	19.9
	5.2	229	11.0	6	4.8	18	6.6	24	8.6	42	12.8	57	11.6	82	13.9	56	12.1	47	13.2
	5.3	200	9.6	4	3.2	15	5.5	31	11.2	34	10.4	52	10.6	64	10.8	53	11.4	35	9.8
	5.4	169	8.1	5	4.0	9	3.3	13	4.7	29	8.9	55	11.2	58	9.8	46	9.9	35	9.8
	5.5	122	5.8	1	0.8	4	1.5	8	2.9	25	7.6	36	7.3	48	8.1	49	10.6	27	7.6
	5.6	79	3.8	1	0.8	3	1.1	8	2.9	10	3.1	25	5.1	32	5.4	25	5.4	19	5.3
	5.7	58	2.8	0	0.0	1	0.4	2	0.7	10	3.1	22	4.5	23	3.9	17	3.7	16	4.5
	5.8	36	1.7	0	0.0	1	0.4	3	1.1	5	1.5	10	2.0	17	2.9	12	2.6	10	2.8
	5.9	28	1.3	0	0.0	0	0.0	0	0.0	2	0.6	13	2.6	13	2.2	9	1.9	9	2.5
	6.0	32	1.5	0	0.0	1	0.4	1	0.4	7	2.1	8	1.6	15	2.5	11	2.4	7	2.0
	6.1	16	0.8	0	0.0	1	0.4	0	0.0	1	0.3	6	1.2	8	1.4	3	0.6	7	2.0
	6.2-6.3	20	1.0	0	0.0	1	0.4	0	0.0	3	0.9	9	1.8	7	1.2	5	1.1	6	1.7
	6.4-6.5	21	1.0	0	0.0	0	0.0	0	0.0	3	0.9	12	2.4	6	1.0	10	2.2	3	0.8
	6.6-6.7	12	0.6	0	0.0	0	0.0	0	0.0	2	0.6	4	0.8	6	1.0	6	1.3	2	0.6
	6.8-6.9	15	0.7	0	0.0	1	0.4	0	0.0	2	0.6	1	0.2	11	1.9	6	1.3	6	1.7
	7.0-7.1	9	0.4	0	0.0	0	0.0	0	0.0	0	0.0	4	0.8	5	0.8	1	0.2	5	1.4
	7.2-7.3	2	0.1	0	0.0	0	0.0	0	0.0	1	0.3	1	0.2	0	0.0	1	0.2	0	0.0
	7.4-7.5	4	0.2	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	2	0.3	2	0.4	1	0.3
	7.6-7.7	5	0.2	0	0.0	0	0.0	0	0.0	0	0.0	4	0.8	1	0.2	2	0.4	1	0.3
	7.8-7.9	1	0.0	0	0.0	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.0-8.1	1	0.0	1	0.8	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.2-8.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.4-8.5	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.2	0	0.0
	8.6-8.7	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	0	0.0	1	0.3
	8.8-8.9	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	0	0.0
	9.0%以上	10	0.5	0	0.0	0	0.0	3	1.1	2	0.6	4	0.8	1	0.2	4	0.9	0	0.0
	(再掲)6.1以上	120	5.8	1	0.8	3	1.1	4	1.4	14	4.3	49	10.0	49	8.3	42	9.1	32	9.0
総 数		2,086	100.0	126	100.0	272	100.0	278	100.0	327	100.0	492	100.0	591	100.0	463	100.0	356	100.0
平均値 (%)		5.3		4.9		5.0		5.1		5.3		5.4		5.4		5.4		5.4	
標準偏差		0.6		0.4		0.3		0.6		0.6		0.7		0.6		0.8		0.5	

(23年)

第 30 表の 2 ヘモグロビン A1c の分布（性・年齢階級別）〔インスリン注射または服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	3.9%以下	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1	1	0.1	0	0.0
	4.0-4.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	4.2-4.3	17	0.5	4	2.0	2	0.4	4	0.9	2	0.4	2	0.3	3	0.3	0	0.0	3	0.5
	4.4-4.5	75	2.3	20	10.1	21	4.7	11	2.6	7	1.3	8	1.0	8	0.8	6	0.8	6	1.0
	4.6-4.7	252	7.6	47	23.7	75	16.7	49	11.6	26	5.0	29	3.8	26	2.7	23	3.2	13	2.3
	4.8-4.9	552	16.6	66	33.3	138	30.7	93	22.0	72	13.8	82	10.6	101	10.5	77	10.6	55	9.6
	5.0-5.1	727	21.9	38	19.2	114	25.4	117	27.7	127	24.4	159	20.6	172	17.9	138	18.9	105	18.3
	5.2	394	11.8	10	5.1	36	8.0	46	10.9	72	13.8	102	13.2	128	13.3	93	12.8	76	13.2
	5.3	308	9.3	5	2.5	21	4.7	44	10.4	49	9.4	84	10.9	105	10.9	83	11.4	62	10.8
	5.4	250	7.5	5	2.5	14	3.1	23	5.4	49	9.4	75	9.7	84	8.7	67	9.2	50	8.7
	5.5	210	6.3	1	0.5	10	2.2	9	2.1	43	8.3	62	8.0	85	8.8	75	10.3	52	9.1
	5.6	124	3.7	2	1.0	5	1.1	11	2.6	16	3.1	35	4.5	55	5.7	41	5.6	33	5.7
	5.7	99	3.0	0	0.0	2	0.4	3	0.7	17	3.3	29	3.8	48	5.0	25	3.4	34	5.9
	5.8	82	2.5	0	0.0	1	0.2	8	1.9	9	1.7	24	3.1	40	4.2	26	3.6	24	4.2
	5.9	50	1.5	0	0.0	3	0.7	0	0.0	5	1.0	19	2.5	23	2.4	21	2.9	11	1.9
	6.0	46	1.4	0	0.0	2	0.4	1	0.2	8	1.5	11	1.4	24	2.5	16	2.2	12	2.1
	6.1	23	0.7	0	0.0	1	0.2	1	0.2	1	0.2	9	1.2	11	1.1	5	0.7	9	1.6
	6.2-6.3	36	1.1	0	0.0	2	0.4	2	0.5	3	0.6	12	1.6	17	1.8	9	1.2	12	2.1
	6.4-6.5	26	0.8	0	0.0	1	0.2	0	0.0	5	1.0	13	1.7	7	0.7	6	0.8	4	0.7
	6.6-6.7	11	0.3	0	0.0	1	0.2	0	0.0	2	0.4	4	0.5	4	0.4	4	0.5	2	0.3
	6.8-6.9	12	0.4	0	0.0	0	0.0	0	0.0	2	0.4	1	0.1	9	0.9	4	0.5	5	0.9
	7.0-7.1	7	0.2	0	0.0	0	0.0	0	0.0	1	0.2	4	0.5	2	0.2	3	0.4	0	0.0
	7.2-7.3	3	0.1	0	0.0	0	0.0	0	0.0	2	0.4	0	0.0	1	0.1	0	0.0	1	0.2
	7.4-7.5	6	0.2	0	0.0	0	0.0	1	0.2	2	0.4	1	0.1	2	0.2	1	0.1	2	0.3
	7.6-7.7	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	7.8-7.9	3	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	3	0.3	1	0.1	2	0.3
	8.0-8.1	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1	1	0.1	0	0.0
	8.2-8.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.4-8.5	3	0.1	0	0.0	0	0.0	0	0.0	1	0.2	1	0.1	1	0.1	1	0.1	1	0.2
	8.6-8.7	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
	8.8-8.9	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1	0	0.0	0	0.0	0	0.0
	9.0%以上	6	0.2	0	0.0	0	0.0	0	0.0	0	0.0	5	0.6	1	0.1	2	0.3	0	0.0
	(再掲)6.1以上	139	4.2	0	0.0	5	1.1	4	0.9	19	3.6	52	6.7	59	6.1	37	5.1	38	6.6
	総 数	3,326	100.0	198	100.0	449	100.0	423	100.0	521	100.0	773	100.0	962	100.0	729	100.0	574	100.0
	平均値 (%) 標準偏差	5.2 0.5		4.8 0.3		5.0 0.3		5.1 0.3		5.2 0.4		5.4 0.6		5.4 0.5		5.4 0.6		5.4 0.5	
男 性	3.9%以下	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.3	0	0.0
	4.0-4.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	4.2-4.3	7	0.5	0	0.0	1	0.6	3	2.0	1	0.5	1	0.3	1	0.2	0	0.0	1	0.4
	4.4-4.5	37	2.8	10	13.7	9	5.0	4	2.7	2	1.0	6	1.9	6	1.4	3	1.0	5	2.0
	4.6-4.7	98	7.4	23	31.5	28	15.6	12	8.1	7	3.5	15	4.9	13	3.1	9	3.0	7	2.9
	4.8-4.9	209	15.8	25	34.2	53	29.4	30	20.1	32	15.8	34	11.0	35	8.5	29	9.6	17	7.0
	5.0-5.1	262	19.8	9	12.3	43	23.9	41	27.5	43	21.3	59	19.2	67	16.2	60	19.9	35	14.3
	5.2	165	12.4	4	5.5	18	10.0	22	14.8	30	14.9	45	14.6	46	11.1	37	12.3	29	11.9
	5.3	110	8.3	1	1.4	7	3.9	13	8.7	15	7.4	33	10.7	41	9.9	31	10.3	27	11.1
	5.4	86	6.5	0	0.0	5	2.8	10	6.7	20	9.9	20	6.5	31	7.5	23	7.6	18	7.4
	5.5	89	6.7	0	0.0	6	3.3	1	0.7	19	9.4	26	8.4	37	8.9	26	8.6	25	10.2
	5.6	46	3.5	1	1.4	2	1.1	3	2.0	6	3.0	10	3.2	24	5.8	17	5.6	14	5.7
	5.7	43	3.2	0	0.0	1	0.6	1	0.7	7	3.5	7	2.3	27	6.5	10	3.3	18	7.4
	5.8	48	3.6	0	0.0	0	0.0	5	3.4	4	2.0	15	4.9	24	5.8	14	4.7	15	6.1
	5.9	27	2.0	0	0.0	3	1.7	0	0.0	3	1.5	8	2.6	13	3.1	14	4.7	4	1.6
	6.0	18	1.4	0	0.0	1	0.6	0	0.0	2	1.0	4	1.3	11	2.7	7	2.3	6	2.5
	6.1	11	0.8	0	0.0	0	0.0	1	0.7	1	0.5	4	1.3	5	1.2	3	1.0	4	1.6
	6.2-6.3	24	1.8	0	0.0	1	0.6	2	1.3	1	0.5	7	2.3	13	3.1	7	2.3	9	3.7
	6.4-6.5	12	0.9	0	0.0	1	0.6	0	0.0	3	1.5	5	1.6	3	0.7	2	0.7	1	0.4
	6.6-6.7	6	0.5	0	0.0	1	0.6	0	0.0	1	0.5	2	0.6	2	0.5	2	0.7	1	0.4
	6.8-6.9	6	0.5	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	5	1.2	2	0.7	3	1.2
	7.0-7.1	4	0.3	0	0.0	0	0.0	0	0.0	1	0.5	1	0.3	2	0.5	2	0.7	0	0.0
	7.2-7.3	2	0.2	0	0.0	0	0.0	0	0.0	1	0.5	0	0.0	1	0.2	0	0.0	1	0.4
	7.4-7.5	4	0.3	0	0.0	0	0.0	1	0.7	2	1.0	0	0.0	1	0.2	0	0.0	1	0.4
	7.6-7.7	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	7.8-7.9	3	0.2	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	3	0.7	1	0.3	2	0.8
	8.0-8.1	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.3	0	0.0
	8.2-8.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.4-8.5	2	0.2	0	0.0	0	0.0	0	0.0	1	0.5	0	0.0	1	0.2	0	0.0	1	0.4
	8.6-8.7	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0
	8.8-8.9	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0
	9.0%以上	3	0.2	0	0.0	0	0.0	0	0.0	0	0.0	3	1.0	0	0.0	0	0.0	0	0.0
	(再掲)6.1以上	80	6.0	0	0.0	3	1.7	4	2.7	11	5.4	25	8.1	37	8.9	20	6.6	23	9.4
	総 数	1,326	100.0	73	100.0	180	100.0	149	100.0	202	100.0	308	100.0	414	100.0	301	100.0	244	100.0
	平均値 (%) 標準偏差	5.3 0.6		4.8 0.2		5.0 0.4		5.1 0.4		5.3 0.5		5.4 0.7		5.4 0.6		5.4 0.5		5.4 0.6	

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
女 性	3.9%以下	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	4.0-4.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	4.2-4.3	10	0.5	4	3.2	1	0.4	1	0.4	1	0.3	1	0.2	2	0.4	0	0.0	2	0.6
	4.4-4.5	38	1.9	10	8.0	12	4.5	7	2.6	5	1.6	2	0.4	2	0.4	3	0.7	1	0.3
	4.6-4.7	154	7.7	24	19.2	47	17.5	37	13.5	19	6.0	14	3.0	13	2.4	14	3.3	6	1.8
	4.8-4.9	343	17.2	41	32.8	85	31.6	63	23.0	40	12.5	48	10.3	66	12.0	48	11.2	38	11.5
	5.0-5.1	465	23.3	29	23.2	71	26.4	76	27.7	84	26.3	100	21.5	105	19.2	78	18.2	70	21.2
	5.2	229	11.5	6	4.8	18	6.7	24	8.8	42	13.2	57	12.3	82	15.0	56	13.1	47	14.2
	5.3	198	9.9	4	3.2	14	5.2	31	11.3	34	10.7	51	11.0	64	11.7	52	12.1	35	10.6
	5.4	164	8.2	5	4.0	9	3.3	13	4.7	29	9.1	55	11.8	53	9.7	44	10.3	32	9.7
	5.5	121	6.1	1	0.8	4	1.5	8	2.9	24	7.5	36	7.7	48	8.8	49	11.4	27	8.2
	5.6	78	3.9	1	0.8	3	1.1	8	2.9	10	3.1	25	5.4	31	5.7	24	5.6	19	5.8
	5.7	56	2.8	0	0.0	1	0.4	2	0.7	10	3.1	22	4.7	21	3.8	15	3.5	16	4.8
	5.8	34	1.7	0	0.0	1	0.4	3	1.1	5	1.6	9	1.9	16	2.9	12	2.8	9	2.7
	5.9	23	1.2	0	0.0	0	0.0	0	0.0	2	0.6	11	2.4	10	1.8	7	1.6	7	2.1
	6.0	28	1.4	0	0.0	1	0.4	1	0.4	6	1.9	7	1.5	13	2.4	9	2.1	6	1.8
	6.1	12	0.6	0	0.0	1	0.4	0	0.0	0	0.0	5	1.1	6	1.1	2	0.5	5	1.5
	6.2-6.3	12	0.6	0	0.0	1	0.4	0	0.0	2	0.6	5	1.1	4	0.7	2	0.5	3	0.9
	6.4-6.5	14	0.7	0	0.0	0	0.0	0	0.0	2	0.6	8	1.7	4	0.7	4	0.9	3	0.9
	6.6-6.7	5	0.3	0	0.0	0	0.0	0	0.0	1	0.3	2	0.4	2	0.4	2	0.5	1	0.3
	6.8-6.9	6	0.3	0	0.0	0	0.0	0	0.0	2	0.6	0	0.0	4	0.7	2	0.5	2	0.6
	7.0-7.1	3	0.2	0	0.0	0	0.0	0	0.0	0	0.0	3	0.6	0	0.0	1	0.2	0	0.0
	7.2-7.3	1	0.1	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0
	7.4-7.5	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	1	0.2	1	0.3
	7.6-7.7	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	7.8-7.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.0-8.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.2-8.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.4-8.5	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.2	0	0.0
	8.6-8.7	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.8-8.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	9.0%以上	3	0.2	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	1	0.2	2	0.5	0	0.0
	(再掲)6.1以上	59	3.0	0	0.0	2	0.7	0	0.0	8	2.5	27	5.8	22	4.0	17	4.0	15	4.5
	総 数	2,000	100.0	125	100.0	269	100.0	274	100.0	319	100.0	465	100.0	548	100.0	428	100.0	330	100.0
	平均値 (%)	5.2		4.9		5.0		5.0		5.2		5.3		5.3		5.3		5.3	
	標準偏差	0.5		0.3		0.3		0.3		0.4		0.6		0.5		0.6		0.4	

注) インスリン注射または血糖を下げる薬服用者除外

(23年)

第 31 表の 1 血糖値の分布（食後 3 時間以上，性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	40mg/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	40-59	5	0.4	0	0.0	0	0.0	1	0.8	1	0.6	0	0.0	3	0.8	0	0.0	3	1.4
	60-79	48	4.2	7	11.5	9	6.2	6	4.9	7	4.1	9	3.2	10	2.7	10	3.5	5	2.3
	80-99	626	54.4	44	72.1	114	78.1	85	69.7	109	64.5	137	48.2	137	37.2	125	44.3	66	30.8
	100-119	281	24.4	10	16.4	21	14.4	19	15.6	36	21.3	80	28.2	115	31.3	78	27.7	71	33.2
	120-140	96	8.3	0	0.0	1	0.7	6	4.9	8	4.7	34	12.0	47	12.8	37	13.1	30	14.0
	140-160	37	3.2	0	0.0	1	0.7	3	2.5	2	1.2	12	4.2	19	5.2	10	3.5	17	7.9
	160-180	20	1.7	0	0.0	0	0.0	0	0.0	3	1.8	3	1.1	14	3.8	8	2.8	9	4.2
	180-200	16	1.4	0	0.0	0	0.0	2	1.6	1	0.6	3	1.1	10	2.7	4	1.4	8	3.7
	200-220	6	0.5	0	0.0	0	0.0	0	0.0	2	1.2	2	0.7	2	0.5	3	1.1	0	0.0
	220-240	3	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	3	0.8	1	0.4	2	0.9
	240-260	3	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	3	0.8	2	0.7	1	0.5
	260mg/dL以上	9	0.8	0	0.0	0	0.0	0	0.0	0	0.0	4	1.4	5	1.4	4	1.4	2	0.9
	(再掲)110以上	287	25.0	5	8.2	6	4.1	14	11.5	27	16.0	85	29.9	150	40.8	100	35.5	97	45.3
	総 数	1,150	100.0	61	100.0	146	100.0	122	100.0	169	100.0	284	100.0	368	100.0	282	100.0	214	100.0
	平均値(mg/dL) 標準偏差	104.8 30.4		92.1 10.4		92.4 10.0		96.2 18.3		99.1 21.6		107.9 34.1		115.0 37.0		111.1 34.5		117.1 35.4	
女 性	40mg/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	40-59	1	0.1	1	1.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	60-79	62	3.9	7	6.7	17	8.4	8	3.9	12	4.7	12	3.2	6	1.3	5	1.4	5	1.8
	80-99	920	57.3	82	78.8	157	77.7	163	79.1	164	63.8	187	50.0	167	36.1	143	41.3	101	35.6
	100-119	431	26.8	12	11.5	22	10.9	28	13.6	59	23.0	119	31.8	191	41.3	132	38.2	117	41.2
	120-140	121	7.5	1	1.0	6	3.0	2	1.0	15	5.8	35	9.4	62	13.4	47	13.6	34	12.0
	140-160	33	2.1	0	0.0	0	0.0	0	0.0	3	1.2	11	2.9	19	4.1	8	2.3	13	4.6
	160-180	15	0.9	0	0.0	0	0.0	0	0.0	2	0.8	2	0.5	11	2.4	3	0.9	10	3.5
	180-200	8	0.5	1	1.0	0	0.0	0	0.0	1	0.4	4	1.1	2	0.4	4	1.2	1	0.4
	200-220	4	0.2	0	0.0	0	0.0	2	1.0	0	0.0	0	0.0	2	0.4	1	0.3	1	0.4
	220-240	2	0.1	0	0.0	0	0.0	1	0.5	0	0.0	1	0.3	0	0.0	1	0.3	0	0.0
	240-260	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	0	0.0	2	0.7
	260mg/dL以上	7	0.4	0	0.0	0	0.0	2	1.0	1	0.4	3	0.8	1	0.2	2	0.6	0	0.0
	(再掲)110以上	338	21.0	6	5.8	12	5.9	17	8.3	36	14.0	94	25.1	173	37.4	109	31.5	111	39.1
	総 数	1,606	100.0	104	100.0	202	100.0	206	100.0	257	100.0	374	100.0	463	100.0	346	100.0	284	100.0
	平均値(mg/dL) 標準偏差	101.6 24.6		91.2 14.2		91.4 10.2		96.3 28.6		98.8 20.6		104.8 28.2		109.8 24.8		108.0 28.8		110.2 23.7	

(23年)

第31表の2 血糖値の分布（食後3時間以上、性・年齢階級別）〔インスリン注射または服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	40mg/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	40-59	4	0.4	0	0.0	0	0.0	1	0.8	1	0.6	0	0.0	2	0.6	0	0.0	2	1.1
	60-79	45	4.3	7	11.5	8	5.5	6	5.1	7	4.3	8	3.2	9	2.9	9	3.8	4	2.2
	80-99	613	58.7	44	72.1	114	78.6	84	71.2	109	67.3	132	53.4	130	41.7	119	50.9	62	34.6
	100-119	264	25.3	10	16.4	21	14.5	17	14.4	34	21.0	73	29.6	109	34.9	70	29.9	68	38.0
	120-140	74	7.1	0	0.0	1	0.7	5	4.2	8	4.9	24	9.7	36	11.5	25	10.7	24	13.4
	140-160	23	2.2	0	0.0	1	0.7	3	2.5	1	0.6	5	2.0	13	4.2	5	2.1	11	6.1
	160-180	7	0.7	0	0.0	0	0.0	0	0.0	1	0.6	1	0.4	5	1.6	2	0.9	4	2.2
	180-200	8	0.8	0	0.0	0	0.0	2	1.7	0	0.0	1	0.4	5	1.6	1	0.4	4	2.2
	200-220	3	0.3	0	0.0	0	0.0	0	0.0	1	0.6	1	0.4	1	0.3	1	0.4	0	0.0
	220-240	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	1	0.4	0	0.0
	240-260	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	260mg/dL以上	3	0.3	0	0.0	0	0.0	0	0.0	0	0.0	2	0.8	1	0.3	1	0.4	0	0.0
	(再掲)110以上	207	19.8	5	8.2	6	4.1	13	11.0	22	13.6	56	22.7	105	33.7	62	26.5	69	38.5
	総 数	1,045	100.0	61	100.0	145	100.0	118	100.0	162	100.0	247	100.0	312	100.0	234	100.0	179	100.0
	平均値(mg/dL) 標準偏差	100.4 21.6		92.1 10.4		92.5 10.0		96.0 18.4		96.6 15.9		102.6 22.9		107.5 26.6		103.6 24.8		109.8 24.3	
女 性	40mg/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	40-59	1	0.1	1	1.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	60-79	61	4.0	7	6.8	17	8.5	8	4.0	12	4.8	12	3.4	5	1.2	5	1.5	4	1.5
	80-99	912	59.1	82	79.6	156	78.0	163	80.7	164	65.3	183	51.4	164	38.1	139	42.6	99	37.6
	100-119	414	26.8	12	11.7	21	10.5	28	13.9	57	22.7	114	32.0	182	42.2	126	38.7	111	42.2
	120-140	109	7.1	1	1.0	6	3.0	2	1.0	14	5.6	31	8.7	55	12.8	41	12.6	31	11.8
	140-160	26	1.7	0	0.0	0	0.0	0	0.0	2	0.8	9	2.5	15	3.5	7	2.1	10	3.8
	160-180	11	0.7	0	0.0	0	0.0	0	0.0	1	0.4	2	0.6	8	1.9	2	0.6	8	3.0
	180-200	4	0.3	0	0.0	0	0.0	0	0.0	1	0.4	3	0.8	0	0.0	2	0.6	0	0.0
	200-220	2	0.1	0	0.0	0	0.0	1	0.5	0	0.0	0	0.0	1	0.2	1	0.3	0	0.0
	220-240	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	1	0.3	0	0.0
	240-260	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	260mg/dL以上	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	1	0.2	2	0.6	0	0.0
	(再掲)110以上	292	18.9	5	4.9	12	6.0	13	6.4	30	12.0	80	22.5	152	35.3	95	29.1	98	37.3
	総 数	1,543	100.0	103	100.0	200	100.0	202	100.0	251	100.0	356	100.0	431	100.0	326	100.0	263	100.0
	平均値(mg/dL) 標準偏差	99.7 19.5		90.2 10.0		91.3 10.2		92.9 12.0		97.2 14.0		103.0 24.5		107.7 21.6		107.0 28.4		107.8 18.4	

注) インスリン注射または血糖を下げる薬服用者除外

(23年)

第 32 表の 1 血清総コレステロール値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-79歳		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	120mg/dL未満	11	0.8	0	0.0	1	0.6	0	0.0	2	0.9	2	0.6	6	1.2	9	0.8	2	0.6	6	2.1
	120-139	56	3.8	2	2.7	3	1.7	5	3.2	2	0.9	11	3.1	33	6.8	38	3.6	18	5.0	25	8.6
	140-159	134	9.2	19	26.0	14	7.7	5	3.2	7	3.3	28	7.9	61	12.6	75	7.0	33	9.2	41	14.1
	160-179	259	17.7	18	24.7	34	18.8	14	9.1	28	13.0	60	16.9	105	21.7	177	16.5	64	17.8	71	24.5
	180-199	318	21.7	13	17.8	41	22.7	28	18.2	39	18.1	74	20.8	123	25.4	230	21.5	88	24.4	72	24.8
	200-219	338	23.1	13	17.8	41	22.7	43	27.9	55	25.6	106	29.8	80	16.5	264	24.7	83	23.1	44	15.2
	220-239	198	13.5	4	5.5	23	12.7	32	20.8	42	19.5	50	14.0	47	9.7	163	15.2	48	13.3	19	6.6
	240-259	96	6.6	4	5.5	17	9.4	19	12.3	24	11.2	14	3.9	18	3.7	69	6.4	14	3.9	8	2.8
	260-279	37	2.5	0	0.0	5	2.8	5	3.2	11	5.1	7	2.0	9	1.9	31	2.9	7	1.9	4	1.4
	280-299	12	0.8	0	0.0	2	1.1	3	1.9	4	1.9	2	0.6	1	0.2	10	0.9	1	0.3	0	0.0
	300mg/dL以上	4	0.3	0	0.0	0	0.0	0	0.0	1	0.5	2	0.6	1	0.2	4	0.4	2	0.6	0	0.0
	(再掲)220以上	347	23.7	8	11.0	47	26.0	59	38.3	82	38.1	75	21.1	76	15.7	277	25.9	72	20.0	31	10.7
	(再掲)240以上	149	10.2	4	5.5	24	13.3	27	17.5	40	18.6	25	7.0	29	6.0	114	10.7	24	6.7	12	4.1
総数	1,463	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	1,070	100.0	360	100.0	290	100.0	
平均値(mg/dL)	196.7		181.2		199.8		210.4		210.1		198.1		186.4		199.3		194.1		180.4		
標準偏差	34.1		30.8		32.6		31.3		34.2		32.4		33.1		34.0		33.5		31.5		
女 性	120mg/dL未満	4	0.2	1	0.8	2	0.7	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	1	0.3
	120-139	38	1.8	9	7.1	8	2.9	5	1.8	1	0.3	4	0.8	11	1.8	18	1.2	5	1.1	7	1.9
	140-159	138	6.6	29	22.8	37	13.6	16	5.8	6	1.8	12	2.4	38	6.4	54	3.6	17	3.7	26	7.2
	160-179	327	15.6	38	29.9	73	26.7	55	19.9	27	8.2	49	9.9	85	14.3	186	12.3	52	11.2	58	16.1
	180-199	406	19.4	15	11.8	64	23.4	65	23.5	56	17.1	80	16.2	126	21.2	285	18.8	89	19.2	76	21.1
	200-219	496	23.7	18	14.2	56	20.5	72	26.0	79	24.1	117	23.7	154	25.9	388	25.6	115	24.8	93	25.8
	220-239	341	16.3	11	8.7	16	5.9	38	13.7	66	20.1	100	20.2	110	18.5	277	18.3	88	19.0	69	19.2
	240-259	189	9.0	5	3.9	12	4.4	14	5.1	51	15.5	69	14.0	38	6.4	165	10.9	55	11.9	17	4.7
	260-279	83	4.0	1	0.8	2	0.7	7	2.5	24	7.3	31	6.3	18	3.0	75	5.0	21	4.5	8	2.2
	280-299	47	2.2	0	0.0	3	1.1	4	1.4	13	4.0	18	3.6	9	1.5	42	2.8	13	2.8	3	0.8
	300mg/dL以上	25	1.2	0	0.0	0	0.0	1	0.4	5	1.5	14	2.8	5	0.8	25	1.7	9	1.9	2	0.6
	(再掲)220以上	685	32.7	17	13.4	33	12.1	64	23.1	159	48.5	232	47.0	180	30.3	584	38.5	186	40.1	99	27.5
	(再掲)240以上	344	16.4	6	4.7	17	6.2	26	9.4	93	28.4	132	26.7	70	11.8	307	20.3	98	21.1	30	8.3
総数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	1,515	100.0	464	100.0	360	100.0	
平均値(mg/dL)	206.2		179.1		187.2		199.7		221.0		219.9		204.3		212.8		213.7		201.2		
標準偏差	35.9		32.2		30.1		31.1		34.2		36.4		33.0		35.0		34.8		31.5		

(23年)

第 32 表の 2 血清総コレステロール値の分布（性・年齢階級別）〔服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-79歳		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	120mg/dL未満	9	0.7	0	0.0	1	0.6	0	0.0	2	1.1	1	0.4	5	1.3	7	0.8	1	0.4	5	2.2
	120-139	44	3.6	2	2.7	3	1.7	5	3.5	2	1.1	7	2.5	25	6.7	28	3.2	14	5.1	18	7.9
	140-159	103	8.3	19	26.0	14	7.8	5	3.5	4	2.1	20	7.0	41	11.1	50	5.7	19	7.0	31	13.7
	160-179	196	15.8	18	24.7	33	18.4	12	8.4	21	11.2	34	11.9	78	21.0	120	13.7	38	14.0	57	25.1
	180-199	269	21.7	13	17.8	40	22.3	27	18.9	33	17.6	62	21.8	94	25.3	193	22.0	70	25.7	55	24.2
	200-219	304	24.5	13	17.8	41	22.9	39	27.3	51	27.1	93	32.6	67	18.1	233	26.6	68	25.0	37	16.3
	220-239	174	14.0	4	5.5	23	12.8	28	19.6	39	20.7	44	15.4	36	9.7	141	16.1	40	14.7	14	6.2
	240-259	94	7.6	4	5.5	17	9.5	19	13.3	23	12.2	14	4.9	17	4.6	67	7.6	14	5.1	7	3.1
	260-279	33	2.7	0	0.0	5	2.8	5	3.5	10	5.3	7	2.5	6	1.6	27	3.1	5	1.8	3	1.3
	280-299	11	0.9	0	0.0	2	1.1	3	2.1	3	1.6	2	0.7	1	0.3	9	1.0	1	0.4	0	0.0
	300mg/dL以上	2	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.4	1	0.3	2	0.2	2	0.7	0	0.0
	(再掲)220以上	314	25.3	8	11.0	47	26.3	55	38.5	75	39.9	68	23.9	61	16.4	246	28.1	62	22.8	24	10.6
	(再掲)240以上	140	11.3	4	5.5	24	13.4	27	18.9	36	19.1	24	8.4	25	6.7	105	12.0	22	8.1	10	4.4
総数	1,239	100.0	73	100.0	179	100.0	143	100.0	188	100.0	285	100.0	371	100.0	877	100.0	272	100.0	227	100.0	
平均値(mg/dL)	198.7		181.2		200.1		210.9		211.4		202.2		187.8		202.2		197.9		180.9		
標準偏差	33.8		30.8		32.7		31.9		33.1		31.3		33.3		33.3		33.5		31.3		
女 性	120mg/dL未満	4	0.2	1	0.8	2	0.7	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	1	0.4
	120-139	32	1.8	9	7.1	7	2.6	5	1.9	1	0.3	1	0.3	9	2.1	14	1.1	3	0.9	6	2.4
	140-159	118	6.7	29	22.8	37	13.8	16	6.0	4	1.4	5	1.3	27	6.3	36	2.9	7	2.1	22	8.6
	160-179	257	14.6	38	29.9	73	27.1	53	19.8	20	6.8	25	6.7	48	11.3	129	10.5	27	8.0	33	12.9
	180-199	324	18.4	15	11.8	63	23.4	63	23.5	45	15.3	52	13.9	86	20.2	215	17.5	59	17.5	52	20.4
	200-219	403	22.9	18	14.2	55	20.4	69	25.7	70	23.8	85	22.7	106	24.9	308	25.1	83	24.6	59	23.1
	220-239	295	16.8	11	8.7	16	5.9	37	13.8	62	21.1	81	21.7	88	20.7	234	19.1	66	19.6	57	22.4
	240-259	179	10.2	5	3.9	11	4.1	14	5.2	50	17.0	66	17.6	33	7.7	156	12.7	50	14.8	15	5.9
	260-279	76	4.3	1	0.8	2	0.7	6	2.2	24	8.2	29	7.8	14	3.3	71	5.8	20	5.9	5	2.0
	280-299	45	2.6	0	0.0	3	1.1	4	1.5	13	4.4	16	4.3	9	2.1	40	3.3	13	3.9	3	1.2
	300mg/dL以上	25	1.4	0	0.0	0	0.0	1	0.4	5	1.7	14	3.7	5	1.2	25	2.0	9	2.7	2	0.8
	(再掲)220以上	620	35.3	17	13.4	32	11.9	62	23.1	154	52.4	206	55.1	149	35.0	526	42.8	158	46.9	82	32.2
	(再掲)240以上	325	18.5	6	4.7	16	5.9	25	9.3	92	31.3	125	33.4	61	14.3	292	23.8	92	27.3	25	9.8
総数	1,758	100.0	127	100.0	269	100.0	268	100.0	294	100.0	374	100.0	426	100.0	1,228	100.0	337	100.0	255	100.0	
平均値(mg/dL)	207.9		179.1		187.1		199.5		224.0		226.8		207.4		216.3		220.0		202.8		
標準偏差	36.9		32.2		29.8		31.1		33.9		35.7		34.4		35.7		35.2		33.3		

第33表の1 血清HDL- コレステロール値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	30mg/dL未満	30	2.1	0	0.0	1	0.6	1	0.6	4	1.9	6	1.7	18	3.7	8	2.2	14	4.8
	30-39	205	14.0	8	11.0	24	13.3	21	13.6	20	9.3	40	11.2	92	19.0	52	14.4	58	20.0
	40-49	379	25.9	19	26.0	58	32.0	33	21.4	54	25.1	82	23.0	133	27.5	90	25.0	84	29.0
	50-59	383	26.2	23	31.5	47	26.0	36	23.4	57	26.5	108	30.3	112	23.1	97	26.9	64	22.1
	60-69	261	17.8	11	15.1	26	14.4	36	23.4	37	17.2	74	20.8	77	15.9	72	20.0	42	14.5
	70-79	121	8.3	9	12.3	14	7.7	20	13.0	22	10.2	27	7.6	29	6.0	23	6.4	17	5.9
	80-89	48	3.3	2	2.7	5	2.8	4	2.6	11	5.1	10	2.8	16	3.3	11	3.1	8	2.8
	90-99	27	1.8	0	0.0	4	2.2	2	1.3	8	3.7	9	2.5	4	0.8	4	1.1	3	1.0
	100mg/dL以上	9	0.6	1	1.4	2	1.1	1	0.6	2	0.9	0	0.0	3	0.6	3	0.8	0	0.0
	総数	1,463	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	360	100.0	290	100.0
平均値(mg/dL)		54.1		55.6		54.0		56.1		57.0		54.8		51.5		53.6		50.4	
標準偏差		14.9		13.7		15.0		14.7		16.1		14.0		14.7		14.5		14.2	
女 性	30mg/dL未満	6	0.3	0	0.0	0	0.0	1	0.4	1	0.3	1	0.2	3	0.5	4	0.9	0	0.0
	30-39	66	3.2	2	1.6	6	2.2	5	1.8	3	0.9	17	3.4	33	5.5	18	3.9	22	6.1
	40-49	293	14.0	7	5.5	24	8.8	28	10.1	30	9.1	79	16.0	125	21.0	71	15.3	90	25.0
	50-59	529	25.3	34	26.8	63	23.1	58	20.9	82	25.0	127	25.7	165	27.7	133	28.7	86	23.9
	60-69	526	25.1	40	31.5	77	28.2	84	30.3	76	23.2	107	21.7	142	23.9	111	23.9	87	24.2
	70-79	369	17.6	24	18.9	59	21.6	52	18.8	66	20.1	86	17.4	82	13.8	75	16.2	51	14.2
	80-89	189	9.0	17	13.4	28	10.3	34	12.3	38	11.6	47	9.5	25	4.2	30	6.5	16	4.4
	90-99	78	3.7	1	0.8	12	4.4	10	3.6	20	6.1	22	4.5	13	2.2	14	3.0	5	1.4
	100mg/dL以上	38	1.8	2	1.6	4	1.5	5	1.8	12	3.7	8	1.6	7	1.2	8	1.7	3	0.8
	総数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	360	100.0
平均値(mg/dL)		63.7		65.7		65.8		66.0		67.5		63.4		59.2		61.8		58.5	
標準偏差		15.2		12.3		13.9		14.4		16.3		16.1		14.4		15.3		14.0	

(23年)

第33表の2 血清HDL- コレステロール値の分布（性・年齢階級別）〔服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	30mg/dL未満	27	2.2	0	0.0	1	0.6	1	0.7	3	1.6	5	1.8	17	4.6	7	2.6	13	5.7
	30-39	165	13.3	8	11.0	24	13.4	19	13.3	19	10.1	33	11.6	62	16.7	37	13.6	40	17.6
	40-49	314	25.3	19	26.0	57	31.8	30	21.0	44	23.4	60	21.1	104	28.0	68	25.0	66	29.1
	50-59	335	27.0	23	31.5	47	26.3	34	23.8	52	27.7	89	31.2	90	24.3	76	27.9	53	23.3
	60-69	214	17.3	11	15.1	25	14.0	34	23.8	29	15.4	59	20.7	56	15.1	50	18.4	32	14.1
	70-79	110	8.9	9	12.3	14	7.8	18	12.6	21	11.2	23	8.1	25	6.7	19	7.0	15	6.6
	80-89	43	3.5	2	2.7	5	2.8	4	2.8	10	5.3	10	3.5	12	3.2	8	2.9	7	3.1
	90-99	22	1.8	0	0.0	4	2.2	2	1.4	8	4.3	6	2.1	2	0.5	4	1.5	1	0.4
	100mg/dL以上	9	0.7	1	1.4	2	1.1	1	0.7	2	1.1	0	0.0	3	0.8	3	1.1	0	0.0
	総数	1,239	1.0	73	100.0	179	100.0	143	100.0	188	100.0	285	100.0	371	100.0	272	100.0	227	100.0
平均値(mg/dL)		54.4		55.6		54.0		56.2		57.4		55.0		51.7		53.8		50.5	
標準偏差		15.0		13.7		15.1		14.7		16.5		14.1		14.8		14.9		14.0	
女 性	30mg/dL未満	5	0.3	0	0.0	0	0.0	1	0.4	1	0.3	1	0.3	2	0.5	3	0.9	0	0.0
	30-39	56	3.2	2	1.6	6	2.2	5	1.9	2	0.7	12	3.2	29	6.8	14	4.2	19	7.5
	40-49	207	11.8	7	5.5	23	8.6	26	9.7	23	7.8	52	13.9	76	17.8	46	13.6	55	21.6
	50-59	439	25.0	34	26.8	63	23.4	53	19.8	73	24.8	91	24.3	125	29.3	92	27.3	67	26.3
	60-69	450	25.6	40	31.5	76	28.3	83	31.0	70	23.8	86	23.0	95	22.3	81	24.0	58	22.7
	70-79	325	18.5	24	18.9	59	21.9	51	19.0	59	20.1	70	18.7	62	14.6	60	17.8	38	14.9
	80-89	165	9.4	17	13.4	26	9.7	34	12.7	36	12.2	34	9.1	18	4.2	20	5.9	10	3.9
	90-99	75	4.3	1	0.8	12	4.5	10	3.7	19	6.5	21	5.6	12	2.8	13	3.9	5	2.0
	100mg/dL以上	36	2.0	2	1.6	4	1.5	5	1.9	11	3.7	7	1.9	7	1.6	8	2.4	3	1.2
	総数	1,758	100.0	127	100.0	269	100.0	268	100.0	294	100.0	374	100.0	426	100.0	337	100.0	255	100.0
平均値(mg/dL)		64.5		65.7		65.7		66.3		68.2		64.5		59.7		62.6		58.7	
標準偏差		15.4		12.3		13.9		14.4		16.3		16.4		15.2		16.0		14.7	

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

(23年)

第 34 表の 1 血清LDL-コレステロール値 (Friedewaldの式) の分布 (性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	40mg/dL未満	2	0.1	0	0.0	0	0.0	0	0.0	1	0.5	0	0.0	1	0.2	0	0.0	1	0.4
	40-50未満	16	1.1	0	0.0	2	1.2	2	1.4	2	1.0	6	1.7	4	0.8	7	2.0	1	0.4
	50-60未満	35	2.5	2	2.8	2	1.2	1	0.7	4	2.0	11	3.2	15	3.2	7	2.0	15	5.3
	60-70未満	56	4.0	6	8.3	3	1.8	3	2.1	4	2.0	15	4.4	25	5.3	17	4.9	14	4.9
	70-80未満	89	6.3	7	9.7	9	5.3	1	0.7	4	2.0	23	6.7	45	9.5	31	8.9	27	9.5
	80-90未満	132	9.4	10	13.9	17	9.9	11	7.7	15	7.5	26	7.6	53	11.2	28	8.0	38	13.3
	90-100未満	147	10.5	10	13.9	18	10.5	12	8.4	19	9.5	30	8.7	58	12.2	36	10.3	42	14.7
	100-110未満	206	14.7	12	16.7	29	17.0	22	15.4	27	13.6	45	13.1	71	15.0	46	13.2	44	15.4
	110-120未満	184	13.1	6	8.3	22	12.9	15	10.5	32	16.1	41	12.0	68	14.3	53	15.2	33	11.6
	120-130未満	164	11.7	10	13.9	18	10.5	24	16.8	21	10.6	51	14.9	40	8.4	36	10.3	25	8.8
	130-140未満	126	9.0	4	5.6	18	10.5	13	9.1	20	10.1	37	10.8	34	7.2	33	9.5	15	5.3
	140-150未満	115	8.2	1	1.4	12	7.0	25	17.5	16	8.0	36	10.5	25	5.3	24	6.9	15	5.3
	150-160未満	49	3.5	1	1.4	10	5.8	6	4.2	9	4.5	6	1.7	17	3.6	12	3.4	7	2.5
	160-170未満	33	2.4	2	2.8	8	4.7	3	2.1	8	4.0	6	1.7	6	1.3	7	2.0	3	1.1
	170-180未満	20	1.4	0	0.0	2	1.2	3	2.1	7	3.5	3	0.9	5	1.1	5	1.4	3	1.1
	180-190未満	15	1.1	1	1.4	1	0.6	2	1.4	5	2.5	4	1.2	2	0.4	2	0.6	0	0.0
	190-200未満	7	0.5	0	0.0	0	0.0	0	0.0	4	2.0	1	0.3	2	0.4	1	0.3	2	0.7
	200-210未満	4	0.3	0	0.0	0	0.0	0	0.0	1	0.5	1	0.3	2	0.4	2	0.6	0	0.0
	210-220未満	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	1	0.2	1	0.3	0	0.0
	220mg/dL以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	総数	1,402	100.0	72	100.0	171	100.0	143	100.0	199	100.0	343	100.0	474	100.0	348	100.0	285	100.0
	平均値 (mg/dL)	112.0		102.8		114.7		120.4		120.7		112.0		106.4		110.1		102.7	
	標準偏差	29.8		26.2		27.4		26.7		32.6		29.8		29.2		30.2		27.8	
女 性	40mg/dL未満	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	0	0.0	0	0.0	0	0.0
	40-50未満	6	0.3	0	0.0	2	0.7	0	0.0	0	0.0	1	0.2	3	0.5	1	0.2	2	0.6
	50-60未満	24	1.2	5	3.9	6	2.2	2	0.7	2	0.6	4	0.8	5	0.8	5	1.1	2	0.6
	60-70未満	57	2.8	9	7.1	18	6.6	5	1.8	6	1.9	2	0.4	17	2.9	11	2.4	7	2.0
	70-80未満	104	5.0	15	11.8	24	8.9	19	7.0	8	2.5	16	3.3	22	3.7	12	2.6	15	4.2
	80-90未満	182	8.8	25	19.7	39	14.4	29	10.7	14	4.3	29	6.0	46	7.8	30	6.6	31	8.7
	90-100未満	231	11.2	18	14.2	49	18.1	38	14.0	24	7.5	40	8.2	62	10.5	47	10.3	36	10.1
	100-110未満	266	12.9	20	15.7	39	14.4	40	14.7	36	11.2	51	10.5	80	13.6	55	12.0	50	14.0
	110-120未満	266	12.9	15	11.8	31	11.4	36	13.2	46	14.3	52	10.7	86	14.6	54	11.8	59	16.5
	120-130未満	263	12.7	6	4.7	33	12.2	41	15.1	46	14.3	63	12.9	74	12.6	51	11.1	50	14.0
	130-140未満	214	10.3	3	2.4	10	3.7	23	8.5	39	12.1	66	13.6	73	12.4	66	14.4	41	11.5
	140-150未満	145	7.0	4	3.1	9	3.3	17	6.3	24	7.5	47	9.7	44	7.5	37	8.1	26	7.3
	150-160未満	109	5.3	3	2.4	6	2.2	6	2.2	26	8.1	39	8.0	29	4.9	33	7.2	17	4.7
	160-170未満	64	3.1	4	3.1	3	1.1	6	2.2	14	4.3	21	4.3	16	2.7	14	3.1	8	2.2
	170-180未満	51	2.5	0	0.0	1	0.4	3	1.1	16	5.0	17	3.5	14	2.4	18	3.9	5	1.4
	180-190未満	40	1.9	0	0.0	0	0.0	2	0.7	9	2.8	18	3.7	11	1.9	12	2.6	7	2.0
	190-200未満	22	1.1	0	0.0	0	0.0	3	1.1	8	2.5	9	1.8	2	0.3	6	1.3	0	0.0
	200-210未満	11	0.5	0	0.0	1	0.4	1	0.4	3	0.9	4	0.8	2	0.3	3	0.7	1	0.3
	210-220未満	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	0	0.0	0	0.0	0	0.0
	220mg/dL以上	9	0.4	0	0.0	0	0.0	1	0.4	1	0.3	4	0.8	3	0.5	3	0.7	1	0.3
	総数	2,068	100.0	127	100.0	271	100.0	272	100.0	322	100.0	487	100.0	589	100.0	458	100.0	358	100.0
	平均値 (mg/dL)	118.3		98.3		101.9		113.1		128.3		128.5		118.6		124.0		117.3	
	標準偏差	31.2		25.2		24.8		27.5		31.1		32.9		29.4		31.7		27.5	

(23年)

Friedewald の式 LDL コレステロール (mg/dL) = 総コレステロール (mg/dL) - HDL コレステロール (mg/dL) - 中性脂肪 (mg/dL) / 5
 (中性脂肪 (トリグリセライド) が 400mg/dL 未満の場合)

第34表の2 血清LDL-コレステロール値(Friedewaldの式)の分布(性・年齢階級別)[服薬者除外]

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	40mg/dL未満	2	0.2	0	0.0	0	0.0	0	0.0	1	0.6	0	0.0	1	0.3	0	0.0	1	0.4
	40-50未満	10	0.8	0	0.0	2	1.2	2	1.5	2	1.2	3	1.1	1	0.3	3	1.1	0	0.0
	50-60未満	27	2.3	2	2.8	2	1.2	1	0.8	2	1.2	7	2.6	13	3.6	4	1.5	13	5.8
	60-70未満	38	3.2	6	8.3	2	1.2	3	2.3	3	1.7	9	3.3	15	4.1	9	3.4	10	4.5
	70-80未満	67	5.6	7	9.7	9	5.3	1	0.8	3	1.7	16	5.9	31	8.5	18	6.8	22	9.8
	80-90未満	104	8.8	10	13.9	17	10.0	11	8.3	12	6.9	16	5.9	38	10.4	19	7.2	27	12.1
	90-100未満	109	9.2	10	13.9	18	10.6	11	8.3	16	9.2	15	5.5	39	10.7	23	8.7	28	12.5
	100-110未満	180	15.2	12	16.7	29	17.1	19	14.3	23	13.3	35	12.8	62	17.0	37	14.1	39	17.4
	110-120未満	165	13.9	6	8.3	22	12.9	14	10.5	28	16.2	39	14.3	56	15.3	47	17.9	26	11.6
	120-130未満	146	12.3	10	13.9	18	10.6	22	16.5	21	12.1	45	16.5	30	8.2	28	10.6	19	8.5
	130-140未満	113	9.5	4	5.6	18	10.6	11	8.3	18	10.4	32	11.7	30	8.2	27	10.3	15	6.7
	140-150未満	107	9.0	1	1.4	12	7.1	24	18.0	15	8.7	35	12.8	20	5.5	22	8.4	11	4.9
	150-160未満	46	3.9	1	1.4	10	5.9	6	4.5	8	4.6	6	2.2	15	4.1	10	3.8	7	3.1
	160-170未満	32	2.7	2	2.8	8	4.7	3	2.3	8	4.6	6	2.2	5	1.4	7	2.7	2	0.9
	170-180未満	17	1.4	0	0.0	2	1.2	3	2.3	6	3.5	3	1.1	3	0.8	4	1.5	2	0.9
	180-190未満	13	1.1	1	1.4	1	0.6	2	1.5	4	2.3	3	1.1	2	0.5	2	0.8	0	0.0
	190-200未満	5	0.4	0	0.0	0	0.0	0	0.0	2	1.2	1	0.4	2	0.5	1	0.4	2	0.9
	200-210未満	4	0.3	0	0.0	0	0.0	0	0.0	1	0.6	1	0.4	2	0.5	2	0.8	0	0.0
	210-220未満	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0
	220mg/dL以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	総数	1,186	100.0	72	100.0	170	100.0	133	100.0	173	100.0	273	100.0	365	100.0	263	100.0	224	100.0
	平均値(mg/dL)	114.2		102.8		115.0		120.5		121.6		116.7		108.3		114.3		103.8	
	標準偏差	29.2		26.2		27.2		27.3		31.2		28.9		28.9		29.0		28.2	
女 性	40mg/dL未満	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0
	40-50未満	6	0.3	0	0.0	2	0.7	0	0.0	0	0.0	1	0.3	3	0.7	1	0.3	2	0.8
	50-60未満	16	0.9	5	3.9	6	2.2	2	0.8	1	0.3	1	0.3	1	0.2	2	0.6	0	0.0
	60-70未満	47	2.7	9	7.1	17	6.4	5	1.9	5	1.7	1	0.3	10	2.4	6	1.8	5	2.0
	70-80未満	85	4.9	15	11.8	24	9.0	18	6.8	6	2.1	10	2.7	12	2.8	5	1.5	9	3.5
	80-90未満	141	8.1	25	19.7	39	14.6	28	10.6	9	3.1	14	3.8	26	6.1	14	4.2	19	7.5
	90-100未満	189	10.9	18	14.2	49	18.4	38	14.4	20	6.9	21	5.7	43	10.2	27	8.1	28	11.0
	100-110未満	207	11.9	20	15.7	37	13.9	38	14.4	30	10.4	32	8.7	50	11.8	34	10.2	29	11.4
	110-120未満	220	12.7	15	11.8	31	11.6	35	13.3	39	13.5	36	9.8	64	15.1	39	11.7	43	16.9
	120-130未満	217	12.5	6	4.7	33	12.4	39	14.8	42	14.5	43	11.7	54	12.8	38	11.4	34	13.4
	130-140未満	185	10.6	3	2.4	10	3.7	23	8.7	37	12.8	56	15.2	56	13.2	50	15.0	32	12.6
	140-150未満	130	7.5	4	3.1	9	3.4	16	6.1	24	8.3	44	11.9	33	7.8	33	9.9	18	7.1
	150-160未満	105	6.0	3	2.4	6	2.2	5	1.9	25	8.7	38	10.3	28	6.6	32	9.6	16	6.3
	160-170未満	60	3.5	4	3.1	2	0.7	6	2.3	14	4.8	19	5.1	15	3.5	12	3.6	8	3.1
	170-180未満	49	2.8	0	0.0	1	0.4	3	1.1	16	5.5	16	4.3	13	3.1	18	5.4	4	1.6
	180-190未満	37	2.1	0	0.0	0	0.0	2	0.8	9	3.1	17	4.6	9	2.1	12	3.6	5	2.0
	190-200未満	22	1.3	0	0.0	0	0.0	3	1.1	8	2.8	9	2.4	2	0.5	6	1.8	0	0.0
	200-210未満	10	0.6	0	0.0	1	0.4	1	0.4	3	1.0	4	1.1	1	0.2	2	0.6	1	0.4
	210-220未満	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	2	0.5	0	0.0	0	0.0	0	0.0
	220mg/dL以上	9	0.5	0	0.0	0	0.0	1	0.4	1	0.3	4	1.1	3	0.7	3	0.9	1	0.4
	総数	1,738	100.0	127	100.0	267	100.0	263	100.0	289	100.0	369	100.0	423	100.0	334	100.0	254	100.0
	平均値(mg/dL)	120.2		98.3		101.8		113.0		131.1		135.1		122.4		130.4		119.7	
	標準偏差	31.9		25.2		24.6		27.6		30.8		32.6		29.9		31.6		28.3	

(23年)

Friedewaldの式 LDL コレステロール (mg/dL) = 総コレステロール (mg/dL) - HDL コレステロール (mg/dL) - 中性脂肪 (mg/dL) / 5
 (中性脂肪 (トリグリセライド) が 400mg/dL 未満の場合)

注) コレステロールを下げる薬または中性脂肪 (トリグリセライド) を下げる薬服用者除外

(参考) 第 34 表の 3 血清 LDL- コレステロール値 (直接測定法) の分布 (性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	40mg/dL未満	2	0.1	0	0.0	0	0.0	0	0.0	1	0.5	0	0.0	1	0.2	0	0.0	1	0.3
	40-49	8	0.5	0	0.0	0	0.0	1	0.6	1	0.5	4	1.1	2	0.4	2	0.6	2	0.7
	50-59	28	1.9	1	1.4	1	0.6	2	1.3	5	2.3	6	1.7	13	2.7	7	1.9	10	3.4
	60-69	54	3.7	4	5.5	5	2.8	3	1.9	4	1.9	14	3.9	24	5.0	18	5.0	14	4.8
	70-79	88	6.0	8	11.0	10	5.5	3	1.9	7	3.3	18	5.1	42	8.7	24	6.7	28	9.7
	80-89	107	7.3	6	8.2	10	5.5	6	3.9	11	5.1	29	8.1	45	9.3	26	7.2	33	11.4
	90-99	145	9.9	10	13.7	13	7.2	7	4.5	17	7.9	39	11.0	59	12.2	45	12.5	38	13.1
	100-109	168	11.5	9	12.3	26	14.4	14	9.1	19	8.8	33	9.3	67	13.8	37	10.3	44	15.2
	110-119	214	14.6	10	13.7	24	13.3	23	14.9	31	14.4	51	14.3	75	15.5	59	16.4	43	14.8
	120-129	168	11.5	7	9.6	22	12.2	17	11.0	28	13.0	43	12.1	51	10.5	46	12.8	27	9.3
	130-139	160	10.9	8	11.0	21	11.6	20	13.0	28	13.0	48	13.5	35	7.2	29	8.1	18	6.2
	140-149	124	8.5	4	5.5	15	8.3	25	16.2	22	10.2	32	9.0	26	5.4	28	7.8	12	4.1
	150-159	79	5.4	1	1.4	11	6.1	15	9.7	12	5.6	19	5.3	21	4.3	16	4.4	10	3.4
	160-169	53	3.6	3	4.1	9	5.0	10	6.5	11	5.1	8	2.2	12	2.5	14	3.9	3	1.0
	170-179	28	1.9	2	2.7	8	4.4	4	2.6	6	2.8	4	1.1	4	0.8	3	0.8	4	1.4
	180-189	13	0.9	0	0.0	5	2.8	1	0.6	3	1.4	3	0.8	1	0.2	2	0.6	0	0.0
	190-199	16	1.1	0	0.0	1	0.6	3	1.9	6	2.8	3	0.8	3	0.6	2	0.6	2	0.7
	200-209	7	0.5	0	0.0	0	0.0	0	0.0	3	1.4	1	0.3	3	0.6	2	0.6	1	0.3
	210-219	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	220mg/dL以上	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0
	総数	1,463	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	360	100.0	290	100.0
	平均値 (mg/dL) 標準偏差	116.3 30.5		108.8 28.9		122.2 29.7		127.7 28.4		124.0 32.4		115.5 29.8		108.8 29.1		113.1 29.7		105.1 28.1	
女 性	40mg/dL未満	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.3	0	0.0	2	0.6
	40-49	3	0.1	1	0.8	1	0.4	0	0.0	0	0.0	1	0.2	0	0.0	1	0.2	0	0.0
	50-59	22	1.1	4	3.1	5	1.8	3	1.1	1	0.3	2	0.4	7	1.2	3	0.6	4	1.1
	60-69	56	2.7	9	7.1	17	6.2	6	2.2	4	1.2	5	1.0	15	2.5	8	1.7	8	2.2
	70-79	101	4.8	15	11.8	20	7.3	14	5.1	14	4.3	16	3.2	22	3.7	15	3.2	14	3.9
	80-89	163	7.8	19	15.0	30	11.0	24	8.7	13	4.0	28	5.7	49	8.2	34	7.3	29	8.1
	90-99	210	10.0	19	15.0	47	17.2	38	13.7	15	4.6	34	6.9	57	9.6	37	8.0	36	10.0
	100-109	250	11.9	17	13.4	41	15.0	38	13.7	35	10.7	45	9.1	74	12.4	48	10.3	50	13.9
	110-119	276	13.2	17	13.4	45	16.5	34	12.3	46	14.0	59	11.9	75	12.6	55	11.9	48	13.3
	120-129	281	13.4	8	6.3	15	5.5	45	16.2	45	13.7	75	15.2	93	15.6	69	14.9	57	15.8
	130-139	235	11.2	4	3.1	22	8.1	30	10.8	44	13.4	63	12.8	72	12.1	66	14.2	40	11.1
	140-149	170	8.1	6	4.7	15	5.5	16	5.8	30	9.1	49	9.9	54	9.1	43	9.3	33	9.2
	150-159	97	4.6	2	1.6	7	2.6	8	2.9	19	5.8	30	6.1	31	5.2	25	5.4	20	5.6
	160-169	82	3.9	6	4.7	5	1.8	8	2.9	20	6.1	31	6.3	12	2.0	20	4.3	6	1.7
	170-179	57	2.7	0	0.0	2	0.7	4	1.4	17	5.2	22	4.5	12	2.0	16	3.4	3	0.8
	180-189	39	1.9	0	0.0	0	0.0	5	1.8	8	2.4	16	3.2	10	1.7	11	2.4	5	1.4
	190-199	22	1.1	0	0.0	0	0.0	1	0.4	8	2.4	7	1.4	6	1.0	6	1.3	4	1.1
	200-209	12	0.6	0	0.0	0	0.0	1	0.4	7	2.1	3	0.6	1	0.2	3	0.6	0	0.0
	210-219	6	0.3	0	0.0	0	0.0	1	0.4	1	0.3	3	0.6	1	0.2	2	0.4	0	0.0
	220mg/dL以上	10	0.5	0	0.0	1	0.4	1	0.4	1	0.3	5	1.0	2	0.3	2	0.4	1	0.3
	総数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	360	100.0
	平均値 (mg/dL) 標準偏差	119.8 31.5		100.3 27.0		105.3 26.5		115.6 28.8		130.3 31.9		129.5 32.7		118.6 29.2		124.7 31.0		117.0 28.0	

(23年)

(参考) 第34表の4 血清LDL- コレステロール値(直接測定法)の分布(性・年齢階級別) [服薬者除外]

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	40mg/dL未満	2	0.2	0	0.0	0	0.0	0	0.0	1	0.5	0	0.0	1	0.3	0	0.0	1	0.4
	40-49	4	0.3	0	0.0	0	0.0	1	0.7	1	0.5	1	0.4	1	0.3	1	0.4	1	0.4
	50-59	22	1.8	1	1.4	1	0.6	2	1.4	3	1.6	5	1.8	10	2.7	5	1.8	8	3.5
	60-69	42	3.4	4	5.5	4	2.2	3	2.1	4	2.1	9	3.2	18	4.9	10	3.7	12	5.3
	70-79	66	5.3	8	11.0	9	5.0	3	2.1	5	2.7	13	4.6	28	7.5	16	5.9	20	8.8
	80-89	83	6.7	6	8.2	10	5.6	4	2.8	10	5.3	20	7.0	33	8.9	17	6.3	25	11.0
	90-99	115	9.3	10	13.7	13	7.3	7	4.9	15	8.0	25	8.8	45	12.1	33	12.1	28	12.3
	100-109	127	10.3	9	12.3	26	14.5	13	9.1	13	6.9	20	7.0	46	12.4	21	7.7	34	15.0
	110-119	189	15.3	10	13.7	24	13.4	21	14.7	28	14.9	45	15.8	61	16.4	49	18.0	35	15.4
	120-129	148	11.9	7	9.6	22	12.3	16	11.2	25	13.3	40	14.0	38	10.2	36	13.2	21	9.3
	130-139	146	11.8	8	11.0	21	11.7	18	12.6	27	14.4	40	14.0	32	8.6	25	9.2	16	7.0
	140-149	114	9.2	4	5.5	15	8.4	22	15.4	21	11.2	31	10.9	21	5.7	25	9.2	9	4.0
	150-159	72	5.8	1	1.4	11	6.1	15	10.5	10	5.3	17	6.0	18	4.9	13	4.8	9	4.0
	160-169	52	4.2	3	4.1	9	5.0	10	7.0	11	5.9	8	2.8	11	3.0	13	4.8	3	1.3
	170-179	26	2.1	2	2.7	8	4.5	4	2.8	5	2.7	4	1.4	3	0.8	3	1.1	3	1.3
	180-189	12	1.0	0	0.0	5	2.8	1	0.7	3	1.6	2	0.7	1	0.3	2	0.7	0	0.0
	190-199	13	1.0	0	0.0	1	0.6	3	2.1	4	2.1	3	1.1	2	0.5	1	0.4	2	0.9
	200-209	5	0.4	0	0.0	0	0.0	0	0.0	2	1.1	1	0.4	2	0.5	2	0.7	0	0.0
	210-219	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	220mg/dL以上	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0
総 数		1,239	100.0	73	100.0	179	100.0	143	100.0	188	100.0	285	100.0	371	100.0	272	100.0	227	100.0
平均値(mg/dL) 標準偏差		118.3 30.2		108.8 28.9		122.8 29.3		128.3 28.9		124.5 31.5		119.3 29.3		110.3 29.2		116.4 29.5		105.8 28.1	
女 性	40mg/dL未満	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.5	0	0.0	2	0.8
	40-49	2	0.1	1	0.8	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	50-59	17	1.0	4	3.1	4	1.5	3	1.1	1	0.3	1	0.3	4	0.9	1	0.3	3	1.2
	60-69	49	2.8	9	7.1	17	6.3	6	2.2	3	1.0	4	1.1	10	2.3	5	1.5	6	2.4
	70-79	85	4.8	15	11.8	20	7.4	14	5.2	12	4.1	10	2.7	14	3.3	9	2.7	9	3.5
	80-89	124	7.1	19	15.0	30	11.2	24	9.0	8	2.7	13	3.5	30	7.0	17	5.0	19	7.5
	90-99	165	9.4	19	15.0	47	17.5	36	13.4	12	4.1	16	4.3	35	8.2	21	6.2	22	8.6
	100-109	203	11.5	17	13.4	40	14.9	38	14.2	29	9.9	27	7.2	52	12.2	30	8.9	35	13.7
	110-119	225	12.8	17	13.4	44	16.4	33	12.3	39	13.3	41	11.0	51	12.0	38	11.3	33	12.9
	120-129	228	13.0	8	6.3	15	5.6	44	16.4	40	13.6	56	15.0	65	15.3	50	14.8	38	14.9
	130-139	190	10.8	4	3.1	22	8.2	27	10.1	40	13.6	49	13.1	48	11.3	47	13.9	25	9.8
	140-149	159	9.0	6	4.7	15	5.6	15	5.6	30	10.2	46	12.3	47	11.0	38	11.3	29	11.4
	150-159	92	5.2	2	1.6	6	2.2	8	3.0	19	6.5	28	7.5	29	6.8	23	6.8	19	7.5
	160-169	80	4.6	6	4.7	5	1.9	8	3.0	20	6.8	30	8.0	11	2.6	19	5.6	5	2.0
	170-179	52	3.0	0	0.0	2	0.7	4	1.5	16	5.4	20	5.3	10	2.3	16	4.7	1	0.4
	180-189	37	2.1	0	0.0	0	0.0	5	1.9	8	2.7	15	4.0	9	2.1	11	3.3	4	1.6
	190-199	20	1.1	0	0.0	0	0.0	0	0.0	8	2.7	7	1.9	5	1.2	5	1.5	4	1.6
	200-209	12	0.7	0	0.0	0	0.0	1	0.4	7	2.4	3	0.8	1	0.2	3	0.9	0	0.0
	210-219	6	0.3	0	0.0	0	0.0	1	0.4	1	0.3	3	0.8	1	0.2	2	0.6	0	0.0
	220mg/dL以上	10	0.6	0	0.0	1	0.4	1	0.4	1	0.3	5	1.3	2	0.5	2	0.6	1	0.4
総 数		1,758	100.0	127	100.0	269	100.0	268	100.0	294	100.0	374	100.0	426	100.0	337	100.0	255	100.0
平均値(mg/dL) 標準偏差		121.5 32.3		100.3 27.0		105.2 26.3		115.1 28.6		132.9 31.9		135.7 32.7		121.7 30.3		130.4 31.3		118.8 29.3	

注) コレステロールを下げる薬または中性脂肪(トリグリセライド)を下げる薬服用者除外

(23年)

第 35 表の 1 血清 non HDL- コレステロール値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	70mg/dL未満	14	1.0	0	0.0	1	0.6	1	0.6	2	0.9	4	1.1	6	1.2	2	0.6	6	2.1
	70-80未満	20	1.4	4	5.5	0	0.0	3	1.9	4	1.9	3	0.8	6	1.2	1	0.3	6	2.1
	80-90未満	50	3.4	7	9.6	6	3.3	2	1.3	3	1.4	10	2.8	22	4.5	19	5.3	11	3.8
	90-100未満	71	4.9	7	9.6	7	3.9	3	1.9	7	3.3	15	4.2	32	6.6	17	4.7	23	7.9
	100-110未満	83	5.7	7	9.6	12	6.6	4	2.6	10	4.7	18	5.1	32	6.6	20	5.6	24	8.3
	110-120未満	151	10.3	11	15.1	22	12.2	9	5.8	14	6.5	31	8.7	64	13.2	35	9.7	42	14.5
	120-130未満	140	9.6	7	9.6	16	8.8	9	5.8	14	6.5	31	8.7	63	13.0	39	10.8	40	13.8
	130-140未満	164	11.2	6	8.2	17	9.4	20	13.0	24	11.2	43	12.1	54	11.2	42	11.7	35	12.1
	140-150未満	175	12.0	9	12.3	15	8.3	13	8.4	26	12.1	52	14.6	60	12.4	53	14.7	31	10.7
	150-160未満	157	10.7	6	8.2	23	12.7	17	11.0	20	9.3	48	13.5	43	8.9	42	11.7	24	8.3
	160-170未満	148	10.1	3	4.1	16	8.8	29	18.8	27	12.6	38	10.7	35	7.2	29	8.1	20	6.9
	170-180未満	91	6.2	0	0.0	19	10.5	11	7.1	14	6.5	25	7.0	22	4.5	22	6.1	10	3.4
	180-190未満	72	4.9	1	1.4	8	4.4	10	6.5	18	8.4	15	4.2	20	4.1	12	3.3	10	3.4
	190-200未満	47	3.2	2	2.7	7	3.9	8	5.2	9	4.2	11	3.1	10	2.1	14	3.9	2	0.7
	200-210未満	27	1.8	1	1.4	3	1.7	8	5.2	4	1.9	3	0.8	8	1.7	5	1.4	3	1.0
	210-220未満	22	1.5	2	2.7	4	2.2	4	2.6	8	3.7	3	0.8	1	0.2	3	0.8	1	0.3
	220-230未満	12	0.8	0	0.0	3	1.7	1	0.6	5	2.3	2	0.6	1	0.2	1	0.3	0	0.0
	230-240未満	13	0.9	0	0.0	1	0.6	2	1.3	3	1.4	2	0.6	5	1.0	3	0.8	2	0.7
	240-250未満	5	0.3	0	0.0	1	0.6	0	0.0	3	1.4	1	0.3	0	0.0	0	0.0	0	0.0
	250mg/dL以上	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	1	0.3	0	0.0
	総数	1,463	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	360	100.0	290	100.0
	平均値(mg/dL)	142.5		125.7		145.8		154.3		153.1		143.2		134.9		140.5		130.0	
	標準偏差	34.3		33.5		34.7		34.0		37.1		32.0		31.9		32.3		30.6	
女 性	70mg/dL未満	8	0.4	2	1.6	3	1.1	0	0.0	0	0.0	1	0.2	2	0.3	1	0.2	2	0.6
	70-80未満	35	1.7	12	9.4	12	4.4	5	1.8	1	0.3	1	0.2	4	0.7	3	0.6	1	0.3
	80-90未満	54	2.6	11	8.7	16	5.9	14	5.1	3	0.9	3	0.6	7	1.2	1	0.2	7	1.9
	90-100未満	108	5.2	24	18.9	32	11.7	13	4.7	10	3.0	8	1.6	21	3.5	8	1.7	15	4.2
	100-110未満	160	7.6	17	13.4	41	15.0	29	10.5	12	3.7	24	4.9	37	6.2	28	6.0	19	5.3
	110-120未満	192	9.2	17	13.4	33	12.1	33	11.9	17	5.2	36	7.3	56	9.4	42	9.1	36	10.0
	120-130未満	229	10.9	13	10.2	44	16.1	31	11.2	41	12.5	31	6.3	69	11.6	36	7.8	47	13.1
	130-140未満	240	11.5	7	5.5	26	9.5	49	17.7	37	11.3	52	10.5	69	11.6	54	11.6	41	11.4
	140-150未満	241	11.5	5	3.9	23	8.4	32	11.6	51	15.5	59	11.9	71	11.9	50	10.8	44	12.2
	150-160未満	227	10.8	7	5.5	15	5.5	25	9.0	32	9.8	63	12.8	85	14.3	65	14.0	51	14.2
	160-170未満	166	7.9	6	4.7	10	3.7	13	4.7	28	8.5	56	11.3	53	8.9	46	9.9	32	8.9
	170-180未満	147	7.0	4	3.1	9	3.3	10	3.6	24	7.3	56	11.3	44	7.4	42	9.1	28	7.8
	180-190未満	80	3.8	0	0.0	4	1.5	9	3.2	18	5.5	24	4.9	25	4.2	28	6.0	14	3.9
	190-200未満	75	3.6	2	1.6	3	1.1	4	1.4	16	4.9	25	5.1	25	4.2	25	5.4	10	2.8
	200-210未満	49	2.3	0	0.0	1	0.4	3	1.1	15	4.6	20	4.0	10	1.7	13	2.8	4	1.1
	210-220未満	29	1.4	0	0.0	0	0.0	2	0.7	8	2.4	12	2.4	7	1.2	5	1.1	5	1.4
	220-230未満	19	0.9	0	0.0	0	0.0	1	0.4	8	2.4	6	1.2	4	0.7	6	1.3	2	0.6
	230-240未満	19	0.9	0	0.0	1	0.4	3	1.1	5	1.5	8	1.6	2	0.3	3	0.6	1	0.3
	240-250未満	4	0.2	0	0.0	0	0.0	0	0.0	1	0.3	2	0.4	1	0.2	3	0.6	0	0.0
	250mg/dL以上	12	0.6	0	0.0	0	0.0	1	0.4	1	0.3	7	1.4	3	0.5	5	1.1	1	0.3
	総数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	360	100.0
	平均値(mg/dL)	142.6		113.4		121.5		133.8		153.5		156.4		145.1		151.9		142.7	
	標準偏差	35.0		28.9		28.3		31.4		34.3		35.2		31.4		33.5		30.3	

(23年)

non LDL コレステロール (mg/dL) = 総コレステロール (mg/dL) - HDL コレステロール (mg/dL)

第35表の2 血清 non HDL- コレステロール値の分布（性・年齢階級別）〔服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	70mg/dL未満	11	0.9	0	0.0	1	0.6	1	0.7	2	1.1	3	1.1	4	1.1	2	0.7	4	1.8
	70-80未満	17	1.4	4	5.5	0	0.0	3	2.1	3	1.6	2	0.7	5	1.3	1	0.4	5	2.2
	80-90未満	40	3.2	7	9.6	6	3.4	2	1.4	2	1.1	5	1.8	18	4.9	11	4.0	10	4.4
	90-100未満	56	4.5	7	9.6	7	3.9	3	2.1	7	3.7	10	3.5	22	5.9	11	4.0	16	7.0
	100-110未満	66	5.3	7	9.6	12	6.7	4	2.8	8	4.3	13	4.6	22	5.9	11	4.0	19	8.4
	110-120未満	117	9.4	11	15.1	21	11.7	8	5.6	12	6.4	19	6.7	46	12.4	24	8.8	32	14.1
	120-130未満	109	8.8	7	9.6	15	8.4	7	4.9	9	4.8	21	7.4	50	13.5	28	10.3	31	13.7
	130-140未満	134	10.8	6	8.2	17	9.5	19	13.3	19	10.1	34	11.9	39	10.5	32	11.8	26	11.5
	140-150未満	148	11.9	9	12.3	15	8.4	11	7.7	26	13.8	42	14.7	45	12.1	37	13.6	26	11.5
	150-160未満	140	11.3	6	8.2	23	12.8	16	11.2	18	9.6	43	15.1	34	9.2	36	13.2	18	7.9
	160-170未満	136	11.0	3	4.1	16	8.9	27	18.9	25	13.3	33	11.6	32	8.6	26	9.6	18	7.9
	170-180未満	85	6.9	0	0.0	19	10.6	10	7.0	13	6.9	24	8.4	19	5.1	19	7.0	9	4.0
	180-190未満	65	5.2	1	1.4	8	4.5	9	6.3	17	9.0	15	5.3	15	4.0	10	3.7	7	3.1
	190-200未満	41	3.3	2	2.7	7	3.9	8	5.6	7	3.7	10	3.5	7	1.9	12	4.4	1	0.4
	200-210未満	26	2.1	1	1.4	3	1.7	8	5.6	4	2.1	2	0.7	8	2.2	5	1.8	3	1.3
	210-220未満	21	1.7	2	2.7	4	2.2	4	2.8	7	3.7	3	1.1	1	0.3	3	1.1	1	0.4
	220-230未満	11	0.9	0	0.0	3	1.7	1	0.7	4	2.1	2	0.7	1	0.3	1	0.4	0	0.0
	230-240未満	11	0.9	0	0.0	1	0.6	2	1.4	3	1.6	2	0.7	3	0.8	2	0.7	1	0.4
	240-250未満	4	0.3	0	0.0	1	0.6	0	0.0	2	1.1	1	0.4	0	0.0	0	0.0	0	0.0
	250mg/dL以上	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.4	0	0.0
	総数	1,239	100.0	73	100.0	179	100.0	143	100.0	188	100.0	285	100.0	371	100.0	272	100.0	227	100.0
	平均値(mg/dL)	144.3		125.7		146.0		154.7		153.9		147.2		136.1		144.1		130.4	
	標準偏差	34.5		33.5		34.7		34.8		36.5		31.9		32.0		32.5		30.4	
女 性	70mg/dL未満	7	0.4	2	1.6	3	1.1	0	0.0	0	0.0	0	0.0	2	0.5	0	0.0	2	0.8
	70-80未満	32	1.8	12	9.4	12	4.5	5	1.9	1	0.3	0	0.0	2	0.5	1	0.3	1	0.4
	80-90未満	50	2.8	11	8.7	15	5.6	14	5.2	3	1.0	2	0.5	5	1.2	1	0.3	5	2.0
	90-100未満	98	5.6	24	18.9	32	11.9	13	4.9	8	2.7	6	1.6	15	3.5	5	1.5	11	4.3
	100-110未満	130	7.4	17	13.4	41	15.2	29	10.8	8	2.7	13	3.5	22	5.2	13	3.9	13	5.1
	110-120未満	154	8.8	17	13.4	33	12.3	32	11.9	14	4.8	16	4.3	42	9.9	24	7.1	27	10.6
	120-130未満	185	10.5	13	10.2	42	15.6	30	11.2	36	12.2	18	4.8	46	10.8	23	6.8	32	12.5
	130-140未満	191	10.9	7	5.5	26	9.7	48	17.9	30	10.2	38	10.2	42	9.9	40	11.9	23	9.0
	140-150未満	197	11.2	5	3.9	23	8.6	30	11.2	44	15.0	42	11.2	53	12.4	38	11.3	32	12.5
	150-160未満	175	10.0	7	5.5	15	5.6	23	8.6	29	9.9	47	12.6	54	12.7	42	12.5	31	12.2
	160-170未満	138	7.8	6	4.7	10	3.7	12	4.5	27	9.2	47	12.6	36	8.5	33	9.8	22	8.6
	170-180未満	134	7.6	4	3.1	8	3.0	10	3.7	24	8.2	48	12.8	40	9.4	36	10.7	25	9.8
	180-190未満	75	4.3	0	0.0	4	1.5	9	3.4	18	6.1	21	5.6	23	5.4	24	7.1	13	5.1
	190-200未満	71	4.0	2	1.6	3	1.1	4	1.5	15	5.1	25	6.7	22	5.2	24	7.1	8	3.1
	200-210未満	46	2.6	0	0.0	1	0.4	3	1.1	15	5.1	19	5.1	8	1.9	12	3.6	3	1.2
	210-220未満	23	1.3	0	0.0	0	0.0	2	0.7	7	2.4	10	2.7	4	0.9	4	1.2	3	1.2
	220-230未満	19	1.1	0	0.0	0	0.0	1	0.4	8	2.7	6	1.6	4	0.9	6	1.8	2	0.8
	230-240未満	17	1.0	0	0.0	1	0.4	2	0.7	5	1.7	7	1.9	2	0.5	3	0.9	1	0.4
	240-250未満	4	0.2	0	0.0	0	0.0	0	0.0	1	0.3	2	0.5	1	0.2	3	0.9	0	0.0
	250mg/dL以上	12	0.7	0	0.0	0	0.0	1	0.4	1	0.3	7	1.9	3	0.7	5	1.5	1	0.4
	総数	1,758	100.0	127	100.0	269	100.0	268	100.0	294	100.0	374	100.0	426	100.0	337	100.0	255	100.0
	平均値(mg/dL)	143.4		113.4		121.4		133.1		155.8		162.3		147.7		157.4		144.1	
	標準偏差	36.3		28.9		28.3		31.2		34.4		35.3		32.8		34.1		31.8	

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

(23年)

non LDL コレステロール (mg/dL) = 総コレステロール (mg/dL) - HDL コレステロール (mg/dL)

第 36 表の 1 血清中性脂肪（トリグリセライド）値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	20mg/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	20-49	53	3.6	8	11.0	10	5.5	5	3.2	7	3.3	4	1.1	19	3.9	7	1.9	14	4.8
	50-79	216	14.8	20	27.4	30	16.6	17	11.0	31	14.4	45	12.6	73	15.1	46	12.8	52	17.9
	80-109	284	19.4	16	21.9	27	14.9	32	20.8	37	17.2	67	18.8	105	21.7	77	21.4	60	20.7
	110-139	247	16.9	8	11.0	35	19.3	22	14.3	27	12.6	66	18.5	89	18.4	62	17.2	57	19.7
	140-169	182	12.4	6	8.2	18	9.9	18	11.7	26	12.1	50	14.0	64	13.2	53	14.7	33	11.4
	170-199	127	8.7	4	5.5	15	8.3	10	6.5	27	12.6	29	8.1	42	8.7	33	9.2	24	8.3
	200mg/dL以上	354	24.2	11	15.1	46	25.4	50	32.5	60	27.9	95	26.7	92	19.0	82	22.8	50	17.2
	150以上(再掲)	603	41.2	17	23.3	76	42.0	69	44.8	106	49.3	158	44.4	177	36.6	149	41.4	98	33.8
	総数	1,463	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	360	100.0	290	100.0
	平均値(mg/dL)	160.1		120.1		165.0		182.5		178.9		165.0		145.3		158.2		137.9	
	標準偏差	109.9		83.1		125.2		130.3		131.8		106.3		86.5		100.3		81.3	
女 性	20mg/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	20-49	175	8.4	39	30.7	47	17.2	45	16.2	16	4.9	10	2.0	18	3.0	16	3.4	10	2.8
	50-79	502	24.0	48	37.8	84	30.8	88	31.8	74	22.6	101	20.4	107	18.0	76	16.4	69	19.2
	80-109	476	22.7	17	13.4	68	24.9	52	18.8	75	22.9	117	23.7	147	24.7	113	24.4	93	25.8
	110-139	321	15.3	13	10.2	28	10.3	34	12.3	66	20.1	88	17.8	92	15.5	73	15.7	61	16.9
	140-169	212	10.1	4	3.1	15	5.5	13	4.7	32	9.8	53	10.7	95	16.0	54	11.6	59	16.4
	170-199	142	6.8	4	3.1	10	3.7	22	7.9	22	6.7	38	7.7	46	7.7	47	10.1	24	6.7
	200mg/dL以上	266	12.7	2	1.6	21	7.7	23	8.3	43	13.1	87	17.6	90	15.1	85	18.3	44	12.2
	150以上(再掲)	542	25.9	8	6.3	38	13.9	55	19.9	82	25.0	157	31.8	202	33.9	171	36.9	107	29.7
	総数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	360	100.0
	平均値(mg/dL)	122.9		75.6		98.5		107.4		128.0		140.1		134.4		141.6		128.1	
	標準偏差	77.5		41.6		64.4		81.5		77.2		83.8		73.7		81.3		64.2	

(23年)

第 36 表の 2 血清中性脂肪（トリグリセライド）値の分布（性・年齢階級別）[服薬者除外]

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	20mg/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	20-49	51	4.1	8	11.0	10	5.6	5	3.5	7	3.7	4	1.4	17	4.6	6	2.2	13	5.7
	50-79	197	15.9	20	27.4	30	16.8	15	10.5	30	16.0	40	14.0	62	16.7	40	14.7	44	19.4
	80-109	247	19.9	16	21.9	27	15.1	30	21.0	32	17.0	60	21.1	82	22.1	62	22.8	50	22.0
	110-139	209	16.9	8	11.0	35	19.6	22	15.4	23	12.2	52	18.2	69	18.6	48	17.6	42	18.5
	140-169	137	11.1	6	8.2	18	10.1	15	10.5	21	11.2	36	12.6	41	11.1	34	12.5	20	8.8
	170-199	108	8.7	4	5.5	15	8.4	10	7.0	24	12.8	20	7.0	35	9.4	23	8.5	21	9.3
	200mg/dL以上	290	23.4	11	15.1	44	24.6	46	32.2	51	27.1	73	25.6	65	17.5	59	21.7	37	16.3
	150以上(再掲)	490	39.5	17	23.3	74	41.3	62	43.4	91	48.4	116	40.7	130	35.0	103	37.9	74	32.6
	総数	1,239	100.0	73	100.0	179	100.0	143	100.0	188	100.0	285	100.0	371	100.0	272	100.0	227	100.0
	平均値(mg/dL)	158.4		120.1		163.1		181.5		179.5		164.0		139.9		154.8		133.1	
	標準偏差	112.9		83.1		124.1		130.1		137.4		114.0		82.5		104.1		93.3	
女 性	20mg/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	20-49	171	9.7	39	30.7	47	17.5	44	16.4	16	5.4	9	2.4	16	3.8	15	4.5	8	3.1
	50-79	463	26.3	48	37.8	82	30.5	88	32.8	66	22.4	83	22.2	96	22.5	61	18.1	62	24.3
	80-109	395	22.5	17	13.4	66	24.5	52	19.4	66	22.4	89	23.8	105	24.6	88	26.1	60	23.5
	110-139	260	14.8	13	10.2	28	10.4	32	11.9	61	20.7	65	17.4	61	14.3	50	14.8	42	16.5
	140-169	165	9.4	4	3.1	15	5.6	12	4.5	31	10.5	42	11.2	61	14.3	34	10.1	41	16.1
	170-199	112	6.4	4	3.1	10	3.7	20	7.5	20	6.8	28	7.5	30	7.0	35	10.4	14	5.5
	200mg/dL以上	192	10.9	2	1.6	21	7.8	20	7.5	34	11.6	58	15.5	57	13.4	54	16.0	28	11.0
	150以上(再掲)	404	23.0	8	6.3	38	14.1	49	18.3	70	23.8	112	29.9	127	29.8	113	33.5	68	26.7
	総数	1,758	100.0	127	100.0	269	100.0	268	100.0	294	100.0	374	100.0	426	100.0	337	100.0	255	100.0
	平均値(mg/dL)	117.1		75.6		98.6		105.0		125.0		135.9		126.8		134.8		122.0	
	標準偏差	74.7		41.6		64.8		80.3		73.7		83.2		68.9		76.1		61.9	

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

(23年)

第37表 血清クレアチニン値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	0.40mg/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	0.40-0.49	3	0.2	0	0.0	1	0.6	0	0.0	1	0.5	0	0.0	1	0.2	0	0.0	1	0.3
	0.50-0.59	38	2.6	2	2.7	4	2.2	3	1.9	4	1.9	13	3.7	12	2.5	13	3.6	5	1.7
	0.60-0.69	216	14.8	8	11.0	17	9.4	24	15.6	43	20.0	62	17.4	62	12.8	63	17.5	32	11.0
	0.70-0.79	446	30.5	33	45.2	69	38.1	52	33.8	68	31.6	108	30.3	116	24.0	89	24.7	71	24.5
	0.80-0.89	383	26.2	22	30.1	65	35.9	50	32.5	59	27.4	85	23.9	102	21.1	72	20.0	64	22.1
	0.90-0.99	199	13.6	7	9.6	20	11.0	19	12.3	24	11.2	48	13.5	81	16.7	62	17.2	45	15.5
	1.00-1.19	113	7.7	1	1.4	4	2.2	6	3.9	13	6.0	23	6.5	66	13.6	42	11.7	39	13.4
	1.20-1.39	38	2.6	0	0.0	1	0.6	0	0.0	3	1.4	11	3.1	23	4.8	10	2.8	19	6.6
	1.40-1.59	11	0.8	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	10	2.1	3	0.8	8	2.8
	1.60-1.79	4	0.3	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	3	0.6	3	0.8	1	0.3
	1.80-1.99	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	0	0.0	2	0.7
	2.00mg/dL以上	10	0.7	0	0.0	0	0.0	0	0.0	0	0.0	4	1.1	6	1.2	3	0.8	3	1.0
総数		1,463	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	360	100.0	290	100.0
平均値(mg/dL)		0.8		0.8		0.8		0.8		0.8		0.9		0.9		0.9		0.9	
標準偏差		0.4		0.1		0.1		0.1		0.1		0.5		0.4		0.4		0.3	
女 性	0.40mg/dL未満	20	1.0	2	1.6	7	2.6	2	0.7	1	0.3	3	0.6	5	0.8	5	1.1	2	0.6
	0.40-0.49	264	12.6	17	13.4	46	16.8	33	11.9	42	12.8	59	11.9	67	11.3	57	12.3	29	8.1
	0.50-0.59	866	41.4	76	59.8	133	48.7	121	43.7	148	45.1	209	42.3	179	30.1	178	38.4	94	26.1
	0.60-0.69	618	29.5	28	22.0	72	26.4	92	33.2	99	30.2	147	29.8	180	30.3	137	29.5	109	30.3
	0.70-0.79	216	10.3	4	3.1	13	4.8	25	9.0	35	10.7	54	10.9	85	14.3	60	12.9	58	16.1
	0.80-0.89	56	2.7	0	0.0	1	0.4	3	1.1	0	0.0	13	2.6	39	6.6	16	3.4	33	9.2
	0.90-0.99	14	0.7	0	0.0	0	0.0	1	0.4	0	0.0	4	0.8	9	1.5	4	0.9	8	2.2
	1.00-1.19	22	1.1	0	0.0	0	0.0	0	0.0	0	0.0	5	1.0	17	2.9	5	1.1	15	4.2
	1.20-1.39	6	0.3	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	5	0.8	0	0.0	5	1.4
	1.40-1.59	5	0.2	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0	4	0.7	2	0.4	2	0.6
	1.60-1.79	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.3
	1.80-1.99	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.3	0	0.0	2	0.6
	2.00mg/dL以上	4	0.2	0	0.0	0	0.0	0	0.0	2	0.6	0	0.0	2	0.3	0	0.0	2	0.6
総数		2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	360	100.0
平均値(mg/dL)		0.6		0.6		0.6		0.6		0.6		0.6		0.7		0.6		0.7	
標準偏差		0.2		0.1		0.1		0.1		0.2		0.1		0.4		0.1		0.4	

(23年)

第38表 推算 GFR 値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	15ml/min/1.73m ² 未満	3	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	2	0.4	1	0.3	1	0.3
	15-29未満	9	0.6	0	0.0	0	0.0	0	0.0	0	0.0	3	0.8	6	1.2	2	0.6	4	1.4
	30-59未満	192	13.1	0	0.0	1	0.6	2	1.3	11	5.1	40	11.2	138	28.5	76	21.1	89	30.7
	60-89未満	894	61.1	14	19.2	85	47.0	97	63.0	150	69.8	248	69.7	300	62.0	227	63.1	180	62.1
	90ml/min/1.73m ² 以上	365	24.9	59	80.8	95	52.5	55	35.7	54	25.1	64	18.0	38	7.9	54	15.0	16	5.5
	総数	1,463	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	360	100.0	290	100.0
	平均値	78.3		103.3		91.7		85.8		81.5		75.8		67.5		72.3		65.7	
標準偏差		18.5		14.9		14.1		13.2		14.8		15.8		16.9		16.6		16.5	
女 性	15ml/min/1.73m ² 未満	3	0.1	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	2	0.3	0	0.0	2	0.6
	15-29未満	9	0.4	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	8	1.3	2	0.4	6	1.7
	30-59未満	199	9.5	0	0.0	1	0.4	2	0.7	5	1.5	46	9.3	145	24.4	62	13.4	118	32.8
	60-89未満	1,207	57.6	12	9.4	85	31.1	152	54.9	234	71.3	358	72.5	366	61.5	323	69.6	206	57.2
	90ml/min/1.73m ² 以上	676	32.3	115	90.6	187	68.5	123	44.4	87	26.5	90	18.2	74	12.4	77	16.6	28	7.8
	総数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	360	100.0
	平均値	82.8		109.7		99.6		88.7		82.6		78.2		70.4		75.8		66.4	
標準偏差		19.6		15.6		18.6		14.1		14.5		14.1		17.5		15.2		17.3	

(23年)

推算 GFR (ml/min/1.73m²) = 194 × 血清クレアチニン値 (mg/dL)^{-1.094} × 年齢 (歳)^{-0.287} (男性)
 = 194 × 血清クレアチニン値 (mg/dL)^{-1.094} × 年齢 (歳)^{-0.287} × 0.739 (女性)

* CKD ガイドライン 2009 年版の推算式

第 39 表の 1 血清鉄の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	40 μ g/dL未満	49	3.4	0	0.0	4	2.2	2	1.3	4	1.9	12	3.4	27	5.6	13	3.6	21	7.2
	40-59	169	11.6	4	5.5	24	13.3	15	9.7	17	7.9	36	10.1	73	15.1	44	12.3	48	16.6
	60-79	342	23.4	15	20.5	52	28.7	33	21.4	40	18.6	83	23.4	119	24.6	84	23.4	75	25.9
	80-99	382	26.1	17	23.3	35	19.3	43	27.9	54	25.1	96	27.0	137	28.3	104	29.0	78	26.9
	100-119	231	15.8	19	26.0	32	17.7	27	17.5	37	17.2	63	17.7	53	11.0	50	13.9	33	11.4
	120-139	144	9.8	7	9.6	15	8.3	21	13.6	36	16.7	31	8.7	34	7.0	26	7.2	17	5.9
	140-159	80	5.5	4	5.5	10	5.5	8	5.2	13	6.0	20	5.6	25	5.2	22	6.1	11	3.8
	160-179	33	2.3	2	2.7	3	1.7	3	1.9	11	5.1	6	1.7	8	1.7	9	2.5	2	0.7
	180-199	19	1.3	2	2.7	3	1.7	1	0.6	1	0.5	5	1.4	7	1.4	4	1.1	5	1.7
	200 μ g/dL以上	13	0.9	3	4.1	3	1.7	1	0.6	2	0.9	3	0.8	1	0.2	3	0.8	0	0.0
	総数	1,462	100.0	73	100.0	181	100.0	154	100.0	215	100.0	355	100.0	484	100.0	359	100.0	290	100.0
	平均値(μ g/dL)	92.9		106.3		92.7		95.8		101.0		93.5		85.9		91.6		82.1	
	標準偏差	35.2		40.8		36.1		30.6		35.2		35.3		34.0		36.4		32.4	
女 性	40 μ g/dL未満	237	11.3	20	15.7	54	19.8	62	22.4	22	6.7	26	5.3	53	8.9	25	5.4	38	10.6
	40-59	492	23.5	28	22.0	61	22.3	58	20.9	68	20.7	104	21.1	173	29.1	114	24.6	111	30.8
	60-79	612	29.2	31	24.4	60	22.0	57	20.6	108	32.9	174	35.2	182	30.6	160	34.5	108	30.0
	80-99	412	19.7	23	18.1	45	16.5	48	17.3	78	23.8	106	21.5	112	18.8	96	20.7	66	18.3
	100-119	189	9.0	12	9.4	22	8.1	27	9.7	31	9.5	51	10.3	46	7.7	46	9.9	18	5.0
	120-139	74	3.5	3	2.4	13	4.8	13	4.7	11	3.4	22	4.5	12	2.0	13	2.8	7	1.9
	140-159	45	2.1	6	4.7	9	3.3	10	3.6	5	1.5	7	1.4	8	1.3	8	1.7	5	1.4
	160-179	20	1.0	2	1.6	5	1.8	1	0.4	3	0.9	4	0.8	5	0.8	1	0.2	4	1.1
	180-199	6	0.3	1	0.8	2	0.7	1	0.4	1	0.3	0	0.0	1	0.2	0	0.0	1	0.3
	200 μ g/dL以上	7	0.3	1	0.8	2	0.7	0	0.0	1	0.3	0	0.0	3	0.5	1	0.2	2	0.6
	総数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	360	100.0
	平均値(μ g/dL)	73.2		74.7		72.5		68.6		76.1		76.2		71.4		74.5		69.7	
	標準偏差	32.5		37.5		40.6		36.8		29.4		25.9		31.4		27.2		32.5	

(23年)

第 39 表の 2 血清鉄の分布（性・年齢階級別）〔服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	40 μ g/dL未満	47	3.3	0	0.0	4	2.2	2	1.3	4	1.9	11	3.1	26	5.5	12	3.4	20	7.2
	40-59	164	11.4	4	5.5	24	13.3	15	9.7	16	7.5	36	10.3	69	14.7	43	12.1	45	16.2
	60-79	339	23.5	15	20.5	52	28.7	33	21.4	40	18.7	82	23.4	117	24.9	84	23.7	73	26.3
	80-99	377	26.1	17	23.3	35	19.3	43	27.9	54	25.2	96	27.4	132	28.1	103	29.0	74	26.6
	100-119	230	15.9	19	26.0	32	17.7	27	17.5	37	17.3	62	17.7	53	11.3	49	13.8	33	11.9
	120-139	142	9.8	7	9.6	15	8.3	21	13.6	36	16.8	30	8.5	33	7.0	26	7.3	16	5.8
	140-159	80	5.5	4	5.5	10	5.5	8	5.2	13	6.1	20	5.7	25	5.3	22	6.2	11	4.0
	160-179	33	2.3	2	2.7	3	1.7	3	1.9	11	5.1	6	1.7	8	1.7	9	2.5	2	0.7
	180-199	18	1.2	2	2.7	3	1.7	1	0.6	1	0.5	5	1.4	6	1.3	4	1.1	4	1.4
	200 μ g/dL以上	13	0.9	3	4.1	3	1.7	1	0.6	2	0.9	3	0.9	1	0.2	3	0.8	0	0.0
	総数	1,443	100.0	73	100.0	181	100.0	154	100.0	214	100.0	351	100.0	470	100.0	355	100.0	278	100.0
	平均値(μ g/dL)	93.0		106.3		92.7		95.8		101.2		93.6		86.0		91.9		82.0	
	標準偏差	35.2		40.8		36.1		30.6		35.0		35.2		33.9		36.4		32.0	
女 性	40 μ g/dL未満	229	11.2	20	15.7	52	19.6	57	21.4	22	6.8	26	5.3	52	9.0	24	5.3	38	10.9
	40-59	482	23.5	28	22.0	59	22.3	56	21.1	68	20.9	104	21.4	167	28.9	113	24.8	106	30.5
	60-79	600	29.3	31	24.4	59	22.3	57	21.4	106	32.6	170	34.9	177	30.6	155	34.1	105	30.2
	80-99	405	19.8	23	18.1	44	16.6	46	17.3	78	24.0	103	21.1	111	19.2	95	20.9	65	18.7
	100-119	185	9.0	12	9.4	20	7.5	26	9.8	30	9.2	51	10.5	46	8.0	46	10.1	18	5.2
	120-139	73	3.6	3	2.4	13	4.9	12	4.5	11	3.4	22	4.5	12	2.1	13	2.9	7	2.0
	140-159	44	2.1	6	4.7	9	3.4	10	3.8	5	1.5	7	1.4	7	1.2	8	1.8	4	1.1
	160-179	19	0.9	2	1.6	5	1.9	1	0.4	3	0.9	4	0.8	4	0.7	1	0.2	3	0.9
	180-199	6	0.3	1	0.8	2	0.8	1	0.4	1	0.3	0	0.0	1	0.2	0	0.0	1	0.3
	200 μ g/dL以上	5	0.2	1	0.8	2	0.8	0	0.0	1	0.3	0	0.0	1	0.2	0	0.0	1	0.3
	総数	2,048	100.0	127	100.0	265	100.0	266	100.0	325	100.0	487	100.0	578	100.0	455	100.0	348	100.0
	平均値(μ g/dL)	73.1		74.7		72.7		69.2		76.0		76.1		70.5		74.1		68.5	
	標準偏差	31.5		37.5		40.9		36.5		29.5		26.1		27.3		25.5		27.9	

注) 貧血治療薬服薬者除外

(23年)

第40表の1 総鉄結合能（TIBC）の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	200 μ g/dL未満	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.3	0	0.0
	200-249	38	2.6	1	1.4	1	0.6	1	0.6	2	0.9	7	2.0	26	5.4	9	2.5	20	6.9
	250-299	340	23.3	12	16.4	26	14.4	19	12.3	35	16.3	89	25.1	159	32.9	106	29.5	105	36.2
	300-349	634	43.4	33	45.2	93	51.4	76	49.4	104	48.4	151	42.5	177	36.6	135	37.6	102	35.2
	350-399	349	23.9	22	30.1	52	28.7	46	29.9	57	26.5	83	23.4	89	18.4	77	21.4	48	16.6
	400-449	85	5.8	4	5.5	8	4.4	11	7.1	15	7.0	21	5.9	26	5.4	25	7.0	12	4.1
	450-499	14	1.0	1	1.4	1	0.6	1	0.6	2	0.9	4	1.1	5	1.0	5	1.4	3	1.0
	500 μ g/dL以上	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.3	0	0.0
	総数	1,462	100.0	73	100.0	181	100.0	154	100.0	215	100.0	355	100.0	484	100.0	359	100.0	290	100.0
女 性	200 μ g/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	200-249	28	1.3	1	0.8	2	0.7	2	0.7	2	0.6	6	1.2	15	2.5	7	1.5	11	3.1
	250-299	370	17.7	7	5.5	33	12.1	25	9.0	52	15.9	110	22.3	143	24.0	110	23.7	87	24.2
	300-349	908	43.4	51	40.2	100	36.6	90	32.5	166	50.6	234	47.4	267	44.9	237	51.1	145	40.3
	350-399	499	23.8	51	40.2	71	26.0	74	26.7	75	22.9	103	20.9	125	21.0	82	17.7	85	23.6
	400-449	213	10.2	14	11.0	44	16.1	59	21.3	24	7.3	35	7.1	37	6.2	26	5.6	25	6.9
	450-499	66	3.2	3	2.4	19	7.0	23	8.3	8	2.4	6	1.2	7	1.2	2	0.4	6	1.7
	500 μ g/dL以上	10	0.5	0	0.0	4	1.5	4	1.4	1	0.3	0	0.0	1	0.2	0	0.0	1	0.3
	総数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	360	100.0
	平均値(μ g/dL) 標準偏差	329.4 44.9		335.0 41.4		335.6 37.4		342.2 37.6		335.9 40.3		328.5 43.9		320.0 50.6		327.8 49.9		314.2 49.2	

(23年)

第40表の2 総鉄結合能（TIBC）の分布（性・年齢階級別）[服薬者除外]

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	200 μ g/dL未満	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.3	0	0.0
	200-249	33	2.3	1	1.4	1	0.6	1	0.6	2	0.9	7	2.0	21	4.5	9	2.5	15	5.4
	250-299	334	23.1	12	16.4	26	14.4	19	12.3	35	16.4	87	24.8	155	33.0	104	29.3	102	36.7
	300-349	630	43.7	33	45.2	93	51.4	76	49.4	103	48.1	150	42.7	175	37.2	135	38.0	100	36.0
	350-399	346	24.0	22	30.1	52	28.7	46	29.9	57	26.6	83	23.6	86	18.3	76	21.4	46	16.5
	400-449	84	5.8	4	5.5	8	4.4	11	7.1	15	7.0	20	5.7	26	5.5	24	6.8	12	4.3
	450-499	14	1.0	1	1.4	1	0.6	1	0.6	2	0.9	4	1.1	5	1.1	5	1.4	3	1.1
	500 μ g/dL以上	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.3	0	0.0
	総数	1,443	100.0	73	100.0	181	100.0	154	100.0	214	100.0	351	100.0	470	100.0	355	100.0	278	100.0
女 性	200 μ g/dL未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	200-249	27	1.3	1	0.8	2	0.8	2	0.8	2	0.6	6	1.2	14	2.4	6	1.3	11	3.2
	250-299	364	17.8	7	5.5	32	12.1	25	9.4	51	15.7	108	22.2	141	24.4	108	23.7	85	24.4
	300-349	890	43.5	51	40.2	98	37.0	84	31.6	165	50.8	232	47.6	260	45.0	234	51.4	140	40.2
	350-399	487	23.8	51	40.2	68	25.7	73	27.4	75	23.1	101	20.7	119	20.6	79	17.4	81	23.3
	400-449	207	10.1	14	11.0	43	16.2	57	21.4	23	7.1	34	7.0	36	6.2	26	5.7	24	6.9
	450-499	63	3.1	3	2.4	18	6.8	21	7.9	8	2.5	6	1.2	7	1.2	2	0.4	6	1.7
	500 μ g/dL以上	10	0.5	0	0.0	4	1.5	4	1.5	1	0.3	0	0.0	1	0.2	0	0.0	1	0.3
	総数	2,048	100.0	127	100.0	265	100.0	266	100.0	325	100.0	487	100.0	578	100.0	455	100.0	348	100.0
	平均値(μ g/dL) 標準偏差	341.9 51.6		355.1 45.1		361.2 60.5		370.3 58.7		339.7 45.1		330.8 43.7		327.8 45.9		326.4 41.1		329.5 48.8	

注) 貧血治療薬服用者除外

(23年)

第 41 表 AST (GOT) の分布 (性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	5U/L未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	5-9	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	1	0.2	0	0.0	1	0.3
	10-19	399	27.3	37	50.7	64	35.4	49	31.8	61	28.4	67	18.8	121	25.0	83	23.1	69	23.8
	20-29	749	51.2	30	41.1	91	50.3	75	48.7	109	50.7	200	56.2	244	50.4	193	53.6	147	50.7
	30-39	206	14.1	4	5.5	15	8.3	21	13.6	30	14.0	56	15.7	80	16.5	53	14.7	51	17.6
	40-49	60	4.1	0	0.0	6	3.3	5	3.2	9	4.2	22	6.2	18	3.7	21	5.8	10	3.4
	50U/L以上	47	3.2	2	2.7	5	2.8	4	2.6	6	2.8	10	2.8	20	4.1	10	2.8	12	4.1
	総数	1,463	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	360	100.0	290	100.0
	平均値(U/L) 標準偏差	25.6 11.5		21.4 11.2		23.4 8.7		24.8 9.7		25.5 10.5		26.7 12.0		26.6 12.6		26.5 11.3		26.6 12.5	
女 性	5U/L未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	5-9	3	0.1	1	0.8	1	0.4	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0
	10-19	868	41.5	104	81.9	195	71.4	174	62.8	114	34.8	136	27.5	145	24.4	112	24.1	83	23.1
	20-29	989	47.2	20	15.7	72	26.4	86	31.0	179	54.6	285	57.7	347	58.3	284	61.2	211	58.6
	30-39	168	8.0	2	1.6	4	1.5	11	4.0	24	7.3	50	10.1	77	12.9	52	11.2	51	14.2
	40-49	33	1.6	0	0.0	1	0.4	4	1.4	2	0.6	13	2.6	13	2.2	7	1.5	8	2.2
	50U/L以上	33	1.6	0	0.0	0	0.0	2	0.7	8	2.4	10	2.0	13	2.2	9	1.9	7	1.9
	総数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	360	100.0
	平均値(U/L) 標準偏差	22.2 8.1		16.9 4.2		17.8 4.5		19.6 6.3		22.9 9.3		24.1 8.1		24.8 8.5		24.3 7.8		7.8 8.6	

(23年)

第 42 表 ALT (GPT) の分布 (性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	5U/L未満	1	0.1	1	1.4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	5-9	40	2.7	5	6.8	3	1.7	1	0.6	3	1.4	8	2.2	20	4.1	12	3.3	13	4.5
	10-19	618	42.2	34	46.6	70	38.7	50	32.5	74	34.4	140	39.3	250	51.7	151	41.9	164	56.6
	20-29	427	29.2	18	24.7	42	23.2	44	28.6	69	32.1	116	32.6	138	28.5	123	34.2	72	24.8
	30-39	182	12.4	6	8.2	25	13.8	21	13.6	39	18.1	49	13.8	42	8.7	43	11.9	23	7.9
	40-49	97	6.6	5	6.8	16	8.8	20	13.0	17	7.9	19	5.3	20	4.1	17	4.7	10	3.4
	50U/L以上	98	6.7	4	5.5	25	13.8	18	11.7	13	6.0	24	6.7	14	2.9	14	3.9	8	2.8
	総数	1,463	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	360	100.0	290	100.0
	平均値(U/L) 標準偏差	25.4 16.3		23.2 14.4		30.3 21.6		31.0 20.7		26.6 13.6		25.9 16.3		21.3 12.1		23.4 12.0		20.4 12.1	
女 性	5U/L未満	1	0.0	1	0.8	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	5-9	193	9.2	40	31.5	52	19.0	32	11.6	11	3.4	20	4.0	38	6.4	18	3.9	29	8.1
	10-19	1,264	60.4	73	57.5	177	64.8	178	64.3	204	62.2	266	53.8	366	61.5	261	56.3	231	64.2
	20-29	431	20.6	10	7.9	34	12.5	40	14.4	68	20.7	145	29.4	134	22.5	133	28.7	72	20.0
	30-39	108	5.2	2	1.6	7	2.6	11	4.0	24	7.3	29	5.9	35	5.9	28	6.0	17	4.7
	40-49	38	1.8	1	0.8	0	0.0	8	2.9	8	2.4	13	2.6	8	1.3	9	1.9	5	1.4
	50U/L以上	59	2.8	0	0.0	3	1.1	8	2.9	13	4.0	21	4.3	14	2.4	15	3.2	6	1.7
	総数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	360	100.0
	平均値(U/L) 標準偏差	18.6 12.0		12.2 5.8		14.6 7.2		17.8 13.3		20.9 15.3		21.4 12.8		18.7 10.1		20.5 11.6		17.6 8.9	

(23年)

第43表 γ -GT (γ -GTP) の分布 (性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	10IU/L未満	9	0.6	1	1.4	1	0.6	0	0.0	0	0.0	2	0.6	5	1.0	1	0.3	5	1.7
	10-19	332	22.7	33	45.2	39	21.5	29	18.8	27	12.6	58	16.3	146	30.2	70	19.4	101	34.8
	20-29	382	26.1	20	27.4	46	25.4	29	18.8	56	26.0	88	24.7	143	29.5	99	27.5	89	30.7
	30-39	223	15.2	9	12.3	23	12.7	29	18.8	36	16.7	53	14.9	73	15.1	60	16.7	41	14.1
	40-49	132	9.0	3	4.1	21	11.6	13	8.4	25	11.6	37	10.4	33	6.8	32	8.9	16	5.5
	50-59	81	5.5	1	1.4	13	7.2	13	8.4	10	4.7	20	5.6	24	5.0	22	6.1	12	4.1
	60-69	66	4.5	1	1.4	9	5.0	6	3.9	18	8.4	18	5.1	14	2.9	17	4.7	6	2.1
	70-79	49	3.3	2	2.7	6	3.3	5	3.2	5	2.3	14	3.9	17	3.5	14	3.9	8	2.8
	80IU/L以上	189	12.9	3	4.1	23	12.7	30	19.5	38	17.7	66	18.5	29	6.0	45	12.5	12	4.1
	総数	1,463	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	360	100.0	290	100.0
女 性	10IU/L未満	101	4.8	23	18.1	28	10.3	12	4.3	10	3.0	3	0.6	25	4.2	5	1.1	20	5.6
	10-19	1,177	56.2	88	69.3	185	67.8	179	64.6	148	45.1	243	49.2	334	56.1	228	49.1	220	61.1
	20-29	403	19.2	11	8.7	35	12.8	45	16.2	78	23.8	116	23.5	118	19.8	108	23.3	62	17.2
	30-39	174	8.3	3	2.4	13	4.8	19	6.9	34	10.4	55	11.1	50	8.4	52	11.2	27	7.5
	40-49	84	4.0	1	0.8	7	2.6	5	1.8	15	4.6	25	5.1	31	5.2	30	6.5	14	3.9
	50-59	44	2.1	1	0.8	3	1.1	7	2.5	10	3.0	13	2.6	10	1.7	11	2.4	5	1.4
	60-69	29	1.4	0	0.0	1	0.4	0	0.0	15	4.6	4	0.8	9	1.5	7	1.5	3	0.8
	70-79	19	0.9	0	0.0	1	0.4	1	0.4	5	1.5	8	1.6	4	0.7	4	0.9	3	0.8
	80IU/L以上	63	3.0	0	0.0	0	0.0	9	3.2	13	4.0	27	5.5	14	2.4	19	4.1	6	1.7
	総数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	360	100.0
	平均値(IU/L) 標準偏差	23.9 66.5		14.3 20.7		17.0 61.8		22.6 50.7		28.5 69.8		28.4 60.2		23.3 77.8		26.8 83.0		21.4 51.1	

(23年)

第44表 血清尿酸値の分布 (性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	2.0mg/dL未満	5	0.3	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	4	0.8	1	0.3	3	1.0
	2.0-2.9	16	1.1	0	0.0	0	0.0	1	0.6	3	1.4	3	0.8	9	1.9	2	0.6	8	2.8
	3.0-3.9	64	4.4	4	5.5	4	2.2	3	1.9	7	3.3	19	5.3	27	5.6	23	6.4	18	6.2
	4.0-4.9	251	17.2	8	11.0	22	12.2	25	16.2	40	18.6	57	16.0	99	20.5	61	16.9	61	21.0
	5.0-5.9	475	32.5	28	38.4	52	28.7	50	32.5	60	27.9	122	34.3	163	33.7	110	30.6	103	35.5
	6.0-6.9	384	26.2	19	26.0	60	33.1	41	26.6	66	30.7	95	26.7	103	21.3	99	27.5	54	18.6
	7.0-7.9	193	13.2	12	16.4	28	15.5	26	16.9	23	10.7	43	12.1	61	12.6	49	13.6	32	11.0
	8.0-8.9	63	4.3	1	1.4	12	6.6	8	5.2	12	5.6	15	4.2	15	3.1	13	3.6	10	3.4
	9.0-9.9	9	0.6	1	1.4	1	0.6	0	0.0	3	1.4	1	0.3	3	0.6	2	0.6	1	0.3
	10mg/dL以上	3	0.2	0	0.0	2	1.1	0	0.0	1	0.5	0	0.0	0	0.0	0	0.0	0	0.0
	総数	1,463	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	360	100.0	290	100.0
	平均値(mg/dL) 標準偏差	5.8 1.3		5.9 1.1		6.2 1.2		6.0 1.2		6.0 1.3		5.8 1.2		5.6 1.3		5.8 1.2		5.5 1.3	
女 性	2.0mg/dL未満	3	0.1	0	0.0	0	0.0	0	0.0	1	0.3	1	0.2	1	0.2	1	0.2	1	0.3
	2.0-2.9	115	5.5	8	6.3	23	8.4	32	11.6	7	2.1	13	2.6	32	5.4	24	5.2	13	3.6
	3.0-3.9	568	27.1	48	37.8	116	42.5	109	39.4	85	25.9	100	20.2	110	18.5	94	20.3	65	18.1
	4.0-4.9	839	40.1	57	44.9	104	38.1	88	31.8	137	41.8	214	43.3	239	40.2	197	42.5	135	37.5
	5.0-5.9	397	19.0	12	9.4	21	7.7	36	13.0	71	21.6	121	24.5	136	22.9	99	21.3	91	25.3
	6.0-6.9	122	5.8	2	1.6	7	2.6	9	3.2	22	6.7	35	7.1	47	7.9	32	6.9	34	9.4
	7.0-7.9	38	1.8	0	0.0	1	0.4	2	0.7	5	1.5	8	1.6	22	3.7	14	3.0	14	3.9
	8.0-8.9	8	0.4	0	0.0	0	0.0	1	0.4	0	0.0	1	0.2	6	1.0	2	0.4	5	1.4
	9.0-9.9	4	0.2	0	0.0	1	0.4	0	0.0	0	0.0	1	0.2	2	0.3	1	0.2	2	0.6
	10mg/dL以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	総数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	464	100.0	360	100.0
	平均値(mg/dL) 標準偏差	4.5 1.1		4.1 0.8		4.0 0.9		4.1 1.0		4.5 0.9		4.6 1.0		4.7 1.2		4.6 1.1		4.8 1.2	

(23年)

第 45 表 メタボリックシンドローム（内臓脂肪症候群）の状況（性・年齢階級別）

	総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上																					
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%																				
総 数	メタボリックシンドローム(内臓脂肪症候群)が強く疑われる者																																							
	635		18.0		4		2.1		21		4.8		28		6.5		83		15.3		198		23.4		301		28.1		425		18.9		220		26.8		185		28.7	
	461		13.1		10		5.1		39		8.9		55		12.8		79		14.6		131		15.5		147		13.7		332		14.8		130		15.8		80		12.4	
	2,430		68.9		181		92.8		380		86.4		347		80.7		380		70.1		518		61.2		624		58.2		1,490		66.3		471		57.4		379		58.9	
	3,526		100.0		195		100.0		440		100.0		430		100.0		542		100.0		847		100.0		1,072		100.0		2,247		100.0		821		100.0		644		100.0	
男 性	メタボリックシンドローム(内臓脂肪症候群)が強く疑われる者																																							
	421		28.8		2		2.7		18		9.9		19		12.4		53		24.7		130		36.5		199		41.3		274		29.9		140		39.2		127		43.8	
	312		21.4		10		13.7		37		20.4		38		24.8		54		25.1		89		25.0		84		17.4		222		24.2		80		22.4		43		14.8	
	727		49.8		61		83.6		126		69.6		96		62.7		108		50.2		137		38.5		199		41.3		420		45.9		137		38.4		120		41.4	
	1,460		100.0		73		100.0		181		100.0		153		100.0		215		100.0		356		100.0		482		100.0		916		100.0		357		100.0		290		100.0	
女 性	メタボリックシンドローム(内臓脂肪症候群)が強く疑われる者																																							
	214		10.4		2		1.6		3		1.2		9		3.2		30		9.2		68		13.8		102		17.3		151		11.3		80		17.2		58		16.4	
	149		7.2		0		0.0		2		0.8		17		6.1		25		7.6		42		8.6		63		10.7		110		8.3		50		10.8		37		10.5	
	1,703		82.4		120		98.4		254		98.1		251		90.6		272		83.2		381		77.6		425		72.0		1,070		80.4		334		72.0		259		73.2	
	2,066		100.0		122		100.0		259		100.0		277		100.0		327		100.0		491		100.0		590		100.0		1,331		100.0		464		100.0		354		100.0	

注) 妊婦除外, ヘモグロビン Alc ≧ 5.5% の場合。項目の基準は p.18 参照

(23年)

第46表の1 腹囲区分別、血中脂質、血圧、血糖のいずれかのリスクを有する割合（性・年齢階級別）

			総 数		リ ス ク 保 有 数							
					0		1つ		2つ		3つ	
			人数	%	人数	%	人数	%	人数	%	人数	%
総 数	20-29歳	<85cm(男)/90cm(女)	170	100.0	152	89.4	16	9.4	2	1.2	0	0.0
		≥85cm(男)/90cm(女)	25	100.0	11	44.0	10	40.0	4	16.0	0	0.0
	30-39歳	<85cm(男)/90cm(女)	342	100.0	270	78.9	63	18.4	9	2.6	0	0.0
		≥85cm(男)/90cm(女)	98	100.0	38	38.8	39	39.8	19	19.4	2	2.0
	40-49歳	<85cm(男)/90cm(女)	313	100.0	213	68.1	82	26.2	16	5.1	2	0.6
		≥85cm(男)/90cm(女)	117	100.0	34	29.1	55	47.0	23	19.7	5	4.3
	50-59歳	<85cm(男)/90cm(女)	367	100.0	150	40.9	154	42.0	57	15.5	6	1.6
		≥85cm(男)/90cm(女)	175	100.0	13	7.4	79	45.1	60	34.3	23	13.1
	60-69歳	<85cm(男)/90cm(女)	505	100.0	109	21.6	226	44.8	123	24.4	47	9.3
		≥85cm(男)/90cm(女)	342	100.0	13	3.8	131	38.3	128	37.4	70	20.5
	70歳以上	<85cm(男)/90cm(女)	606	100.0	48	7.9	284	46.9	204	33.7	70	11.6
		≥85cm(男)/90cm(女)	466	100.0	18	3.9	147	31.5	183	39.3	118	25.3
	総 数	<85cm(男)/90cm(女)	2,303	100.0	948	41.2	825	35.8	411	17.8	125	5.4
		≥85cm(男)/90cm(女)	1,223	100.0	131	10.7	461	37.7	417	34.1	218	17.8
	(再掲) 40-74歳	<85cm(男)/90cm(女)	1,417	100.0	494	34.9	574	40.5	271	19.1	78	5.5
		≥85cm(男)/90cm(女)	830	100.0	73	8.8	332	40.0	277	33.4	148	17.8
男 性	20-29歳	<85cm	55	100.0	45	81.8	8	14.5	2	3.6	0	0.0
		≥85cm	18	100.0	6	33.3	10	55.6	2	11.1	0	0.0
	30-39歳	<85cm	100	100.0	66	66.0	28	28.0	6	6.0	0	0.0
		≥85cm	81	100.0	26	32.1	37	45.7	17	21.0	1	1.2
	40-49歳	<85cm	71	100.0	38	53.5	25	35.2	7	9.9	1	1.4
		≥85cm	82	100.0	25	30.5	38	46.3	16	19.5	3	3.7
	50-59歳	<85cm	99	100.0	27	27.3	50	50.5	21	21.2	1	1.0
		≥85cm	116	100.0	9	7.8	54	46.6	37	31.9	16	13.8
	60-69歳	<85cm	133	100.0	28	21.1	65	48.9	28	21.1	12	9.0
		≥85cm	223	100.0	4	1.8	89	39.9	84	37.7	46	20.6
	70歳以上	<85cm	188	100.0	13	6.9	84	44.7	65	34.6	26	13.8
		≥85cm	294	100.0	11	3.7	84	28.6	115	39.1	84	28.6
	総 数	<85cm	646	100.0	217	33.6	260	40.2	129	20.0	40	6.2
		≥85cm	814	100.0	81	10.0	312	38.3	271	33.3	150	18.4
	(再掲) 40-74歳	<85cm	374	100.0	97	25.9	171	45.7	83	22.2	23	6.1
		≥85cm	542	100.0	46	8.5	222	41.0	174	32.1	100	18.5
女 性	20-29歳	<90cm	115	100.0	107	93.1	8	7.0	0	0.0	0	0.0
		≥90cm	7	100.0	5	75.0	0	0.0	2	28.6	0	0.0
	30-39歳	<90cm	242	100.0	204	84.3	35	14.5	3	1.2	0	0.0
		≥90cm	17	100.0	12	70.6	2	11.8	2	11.8	1	5.9
	40-49歳	<90cm	242	100.0	175	72.3	57	23.6	9	3.7	1	0.4
		≥90cm	35	100.0	9	25.7	17	48.6	7	20.0	2	5.7
	50-59歳	<90cm	268	100.0	123	45.9	104	38.8	36	13.4	5	1.9
		≥90cm	59	100.0	4	6.8	25	42.4	23	39.0	7	11.9
	60-69歳	<90cm	372	100.0	81	21.8	161	43.3	95	25.5	35	9.4
		≥90cm	119	100.0	9	7.6	42	35.3	44	37.0	24	20.2
	70歳以上	<90cm	418	100.0	35	8.4	200	47.8	139	33.3	44	10.5
		≥90cm	172	100.0	7	4.1	63	36.6	68	39.5	34	19.8
	総 数	<85cm(男)/90cm(女)	1,657	100.0	725	43.8	565	34.1	282	17.0	85	5.1
		≥85cm(男)/90cm(女)	409	100.0	46	11.2	149	36.4	146	35.7	68	16.6
	(再掲) 40-74歳	<90cm	1,043	100.0	397	38.1	403	38.6	188	18.0	55	5.3
		≥90cm	288	100.0	27	9.4	110	38.2	103	35.8	48	16.7
(再掲) 65-74歳	<90cm	<90cm	324	100.0	40	12.3	156	48.1	92	28.4	36	11.1
		≥90cm	140	100.0	10	7.1	50	35.7	52	37.1	28	20.0
	<90cm	<90cm	257	100.0	17	6.6	119	46.3	91	35.4	30	11.7
		≥90cm	97	100.0	2	2.1	37	38.1	39	40.2	19	19.6

注) 妊婦除外, ヘモグロビン A1c ≥ 5.5% の場合。リスクの項目は基準は p.18 参照

(23年)

第 46 表の 2 腹囲区分別, 血中脂質, 血圧, 血糖のリスク保有状況〔すべての組み合わせ (①血中脂質②血圧③血糖)〕(性・年齢階級別)

年齢(歳)			総 数		リ ス ク 保 有 数															
					①有②有③有		①有②有③無		①無②有③有		①有②無③有		①有②無③無		①無②有③無		①無②無③有		①無②無③無	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	20-29歳	<85cm(男)/90cm(女)	170	100.0	0	0.0	1	0.6	1	0.6	0	0.0	4	2.4	11	6.5	1	0.6	152	89.4
		≥85cm(男)/90cm(女)	25	100.0	0	0.0	2	8.0	2	8.0	0	0.0	3	12.0	7	28.0	0	0.0	11	44.0
	30-39歳	<85cm(男)/90cm(女)	342	100.0	0	0.0	2	0.6	6	1.8	1	0.3	9	2.6	47	13.7	7	2.0	270	78.9
		≥85cm(男)/90cm(女)	98	100.0	2	2.0	8	8.2	9	9.2	2	2.0	10	10.2	27	27.6	2	2.0	38	38.8
	40-49歳	<85cm(男)/90cm(女)	313	100.0	2	0.6	9	2.9	6	1.9	1	0.3	5	1.6	60	19.2	17	5.4	213	68.1
		≥85cm(男)/90cm(女)	117	100.0	5	4.3	14	12.0	6	5.1	3	2.6	7	6.0	44	37.6	4	3.4	34	29.1
	50-59歳	<85cm(男)/90cm(女)	367	100.0	6	1.6	19	5.2	34	9.3	4	1.1	6	1.6	129	35.1	19	5.2	150	40.9
		≥85cm(男)/90cm(女)	175	100.0	23	13.1	17	9.7	39	22.3	4	2.3	6	3.4	64	36.6	9	5.1	13	7.4
	60-69歳	<85cm(男)/90cm(女)	505	100.0	47	9.3	46	9.1	69	13.7	8	1.6	14	2.8	186	36.8	26	5.1	109	21.6
		≥85cm(男)/90cm(女)	342	100.0	70	20.5	47	13.7	77	22.5	4	1.2	5	1.5	119	34.8	7	2.0	13	3.8
	70歳以上	<85cm(男)/90cm(女)	606	100.0	70	11.6	88	14.5	110	18.2	6	1.0	20	3.3	241	39.8	23	3.8	48	7.9
		≥85cm(男)/90cm(女)	466	100.0	118	25.3	73	15.7	101	21.7	9	1.9	5	1.1	132	28.3	10	2.1	18	3.9
	総 数	<85cm(男)/90cm(女)	2,303	100.0	125	5.4	165	7.2	226	9.8	20	0.9	58	2.5	674	29.3	93	4.0	942	40.9
		≥85cm(男)/90cm(女)	1,223	100.0	218	17.8	161	13.2	234	19.1	22	1.8	36	2.9	393	32.1	32	2.6	127	10.4
	(再掲) 40-74歳	<85cm(男)/90cm(女)	1,417	100.0	78	5.5	106	7.5	148	10.4	17	1.2	36	2.5	469	33.1	69	4.9	494	34.9
		≥85cm(男)/90cm(女)	830	100.0	148	17.8	102	12.3	159	19.2	16	1.9	20	2.4	287	34.6	25	3.0	73	8.8
	(再掲) 65-74歳	<85cm(男)/90cm(女)	452	100.0	54	11.9	53	11.7	67	14.8	8	1.8	17	3.8	177	39.2	21	4.6	55	12.2
		≥85cm(男)/90cm(女)	369	100.0	88	23.8	44	11.9	81	22.0	7	1.9	4	1.1	118	32.0	8	2.2	19	5.1
	(再掲) 75歳以上	<85cm(男)/90cm(女)	374	100.0	47	12.6	56	15.0	71	19.0	2	0.5	9	2.4	147	39.3	16	4.3	26	7.0
		≥85cm(男)/90cm(女)	270	100.0	68	25.2	49	18.1	64	23.7	4	1.5	3	1.1	72	26.7	5	1.9	5	1.9
男 性	20-29歳	<85cm	55	100.0	0	0.0	1	1.8	1	1.8	0	0.0	2	3.6	6	10.9	0	0.0	45	81.8
		≥85cm	18	100.0	0	0.0	2	11.1	0	0.0	0	0.0	3	16.7	7	38.9	0	0.0	6	33.3
	30-39歳	<85cm	100	100.0	0	0.0	2	2.0	3	3.0	1	1.0	5	5.0	22	22.0	1	1.0	66	66.0
		≥85cm	81	100.0	1	1.2	8	9.9	7	8.6	2	2.5	8	9.9	27	33.3	2	2.5	26	32.1
	40-49歳	<85cm	71	100.0	1	1.4	4	5.6	2	2.8	1	1.4	4	5.6	19	26.8	2	2.8	38	53.5
		≥85cm	82	100.0	3	3.7	9	11.0	4	4.9	3	3.7	6	7.3	30	36.6	2	2.4	25	30.5
	50-59歳	<85cm	99	100.0	1	1.0	7	7.1	13	13.1	1	1.0	2	2.0	43	43.4	5	5.1	27	27.3
		≥85cm	116	100.0	16	13.8	14	12.1	19	16.4	4	3.4	4	3.4	44	37.9	6	5.2	9	7.8
	60-69歳	<85cm	133	100.0	12	9.0	8	6.0	16	12.0	4	3.0	4	3.0	54	40.6	7	5.3	28	21.1
		≥85cm	223	100.0	46	20.6	30	13.5	51	22.9	3	1.3	3	1.3	81	36.3	5	2.2	4	1.8
	70歳以上	<85cm	188	100.0	26	13.8	21	11.2	41	21.8	3	1.6	4	2.1	68	36.2	12	6.4	13	6.9
		≥85cm	294	100.0	84	28.6	46	15.6	65	22.1	4	1.4	2	0.7	76	25.9	6	2.0	11	3.7
	総 数	<85cm(男)/90cm(女)	646	100.0	40	6.2	43	6.7	76	11.8	10	1.5	21	3.3	212	32.8	27	4.2	217	33.6
		≥85cm(男)/90cm(女)	814	100.0	150	18.4	109	13.4	146	17.9	16	2.0	26	3.2	265	32.6	21	2.6	81	10.0
	(再掲) 40-74歳	<85cm	374	100.0	23	6.1	30	8.0	45	12.0	8	2.1	12	3.2	142	38.0	17	4.5	97	25.9
		≥85cm	542	100.0	100	18.5	66	12.2	97	17.9	11	2.0	14	2.6	192	35.4	16	3.0	46	8.5
	(再掲) 65-74歳	<85cm	128	100.0	18	14.1	13	10.2	19	14.8	4	3.1	5	3.9	49	38.3	5	3.9	15	11.7
		≥85cm	229	100.0	60	26.2	25	10.9	52	22.7	3	1.3	2	0.9	74	32.3	4	1.7	9	3.9
	(再掲) 75歳以上	<85cm	117	100.0	17	14.5	10	8.5	27	23.1	1	0.9	2	1.7	42	35.9	9	7.7	9	7.7
		≥85cm	173	100.0	49	28.3	33	19.1	42	24.3	3	1.7	1	0.6	39	22.5	3	1.7	3	1.7
女 性	20-29歳	<90cm	115	100.0	0	0.0	0	0.0	0	0.0	0	0.0	2	1.7	5	4.3	1	0.9	107	93.0
		≥90cm	7	100.0	0	0.0	0	0.0	2	28.6	0	0.0	0	0.0	0	0.0	0	0.0	5	71.4
	30-39歳	<90cm	242	100.0	0	0.0	0	0.0	3	1.2	0	0.0	4	1.7	25	10.1	6	2.5	204	84.3
		≥90cm	17	100.0	1	5.9	0	0.0	2	11.8	0	0.0	2	11.8	0	0.0	0	0.0	12	70.6
	40-49歳	<90cm	242	100.0	1	0.4	5	2.1	4	1.7	0	0.0	1	0.4	41	16.9	15	6.2	175	72.3
		≥90cm	35	100.0	2	5.7	5	14.3	2	5.7	0	0.0	1	2.9	14	40.0	2	5.7	9	25.7
	50-59歳	<90cm	268	100.0	5	1.9	12	4.5	21	7.8	3	1.1	4	1.5	86	32.1	14	5.2	123	45.9
		≥90cm	59	100.0	7	11.9	3	5.1	20	33.9	0	0.0	2	3.4	20	33.9	3	5.1	4	6.8
	60-69歳	<90cm	372	100.0	35	9.4	38	10.2	53	14.2	4	1.1	10	2.7	132	35.5	19	5.1	81	21.8
		≥90cm	119	100.0	24	20.2	17	14.3	26	21.8	1	0.8	2	1.7	38	31.9	2	1.7	9	7.6
	70歳以上	<90cm	418	100.0	44	10.5	67	16.0	69	16.5	3	0.7	16	3.8	173	41.4	11	2.6	35	8.4
		≥90cm	172	100.0	34	19.8	27	15.7	36	20.9	5	2.9	3	1.7	56	32.6	4	2.3	7	4.1
	総 数	<85cm(男)/90cm(女)	1,657	100.0	85	5.1	122	7.4	150	9.1	10	0.6	37	2.2	462	27.9	66	4.0	725	43.8
		≥85cm(男)/90cm(女)	409	100.0	68	16.6	52	12.7	88	21.5	6	1.5	10	2.4	128	31.3	11	2.7	46	11.2
	(再掲) 40-74歳	<90cm	1,043	100.0	55	5.3	76	7.3	103	9.9	9	0.9	24	2.3	327	31.4	52	5.0	397	38.1
		≥90cm	288	100.0	48	16.7	36	12.5	62	21.5	5	1.7	6	2.1	95	33.0	9	3.1	27	9.4
	(再掲) 65-74歳	<90cm	452	100.0	54	11.9	53	11.7	67	14.8	8	1.8	17	3.8	177	39.2	21	4.6	55	12.2
		≥90cm	369	100.0	88	23.8	44	11.9	81	22.0	7	1.9	4	1.1	118	32.0	8	2.2	19	5.1
	(再掲) 75歳以上	<90cm	374	100.0	47	12.6	56	15.0	71	19.0	2	0.5	9	2.4	147	39.3	16	4.3	26	7.0
		≥90cm	270	100.0	68	25.2	49	18.1	64	23.7	4	1.5	3	1.1	72	26.7	5	1.9	5	1.9

注) 妊婦除外, ヘモグロビン A1c ≥ 5.5% の場合。リスクの項目は基準は p.18 参照

(23年)

第 47 表 「糖尿病が強く疑われる人」及び「糖尿病の可能性を否定できない人」の状況（性・年齢階級別）

	総 数	20～29歳		30～39歳		40～49歳		50～59歳		60～69歳		70歳以上		(再掲) 40～74歳		(再掲) 65～74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	糖尿病が強く疑われる人	384	10.9	1	0.5	8	1.8	15	3.5	41	7.6	141	16.8	264	11.9	141	17.4	111	17.5
	(再掲)糖尿病が強く疑われる人のうち服薬者*	201	52.3	1	100.0	1	12.5	8	53.3	20	48.8	74	52.5	136	51.5	86	61.0	63	56.8
	糖尿病の可能性が否定できない人	366	10.4	2	1.0	12	2.7	19	4.4	53	9.9	107	12.7	248	11.1	188	14.5	104	16.4
	上記以外	2,758	78.6	196	98.5	429	95.5	393	92.0	443	82.5	593	70.5	1,714	77.0	552	68.1	419	66.1
	総 数	3,508	100.0	199	100.0	449	100.0	427	100.0	537	100.0	841	100.0	2,226	100.0	811	100.0	634	100.0
男 性	糖尿病が強く疑われる人	226	15.7	0	0.0	4	2.2	10	6.7	23	10.8	80	22.7	153	16.9	86	24.3	69	24.4
	(再掲)糖尿病が強く疑われる人のうち服薬者*	124	54.9	0	0.0	1	25.0	4	40.0	12	52.2	48	60.0	83	54.2	54	62.8	40	58.0
	糖尿病の可能性が否定できない人	166	11.5	1	1.4	7	3.9	6	4.0	22	10.3	39	11.0	106	11.7	56	15.8	52	18.4
	上記以外	1,049	72.8	72	98.6	169	93.9	134	89.3	168	78.9	234	66.3	646	71.4	212	59.9	162	57.2
	総 数	1,441	100.0	73	100.0	180	100.0	150	100.0	213	100.0	353	100.0	905	100.0	354	100.0	283	100.0
女 性	糖尿病が強く疑われる人	158	7.6	1	0.8	4	1.5	5	1.8	18	5.6	61	12.5	111	8.4	55	12.0	42	12.0
	(再掲)糖尿病が強く疑われる人のうち服薬者*	77	48.7	1	100.0	0	0.0	4	80.0	8	44.4	26	42.6	53	47.7	32	58.2	23	54.8
	糖尿病の可能性が否定できない人	200	9.7	1	0.8	5	1.9	13	4.7	31	9.6	68	13.9	142	10.7	62	13.6	52	14.8
	上記以外	1,709	82.7	124	98.4	260	96.7	259	93.5	275	84.9	432	74.1	1,068	80.8	340	74.4	257	73.2
	総 数	2,067	100.0	126	100.0	269	100.0	277	100.0	324	100.0	488	100.0	1,321	100.0	457	100.0	351	100.0

(23年)

注) 判定基準は p18 参照
* 「服薬者」とは、質問票で「インスリン注射または血糖値を下げる薬」の使用有と回答した者

「糖尿病が強く疑われる人」の判定
ヘモグロビン A1 c 値が 6.1% 以上、または、質問票で「現在糖尿病の治療を受けている」と答えた人
「糖尿病の可能性を否定できない人」の判定
ヘモグロビン A1 c 値が 5.6% 以上、6.1% 未満で、“糖尿病が強く疑われる人”以外の人

第48表 高血圧症有病者の状況（性・年齢階級別）

総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)40-74歳		(再掲)65-74歳		(再掲)75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	高血圧症有病者	1,858	48.3	11	5.0	53	11.0	90	19.7	268	45.2	543	59.8	893	75.4	1,221	50.4	583	65.5
	(再掲)高血圧症有病者のうち服薬者*	1,073	57.8	0	0.0	5	9.4	15	16.7	120	44.8	318	58.6	615	68.9	654	53.6	365	62.6
	正常高値血圧者	517	13.4	14	6.4	52	10.8	67	14.7	90	15.2	156	17.2	138	11.7	381	15.7	150	16.9
	上記以外	1,469	38.2	194	88.6	378	78.3	300	65.6	235	39.6	209	23.0	153	12.9	820	33.9	157	17.6
総数		3,844	100.0	219	100.0	483	100.0	457	100.0	593	100.0	908	100.0	1,184	100.0	2,422	100.0	890	100.0
男性	高血圧症有病者	895	56.5	8	10.1	40	21.3	46	28.0	132	55.9	260	67.0	409	77.5	588	59.1	275	70.9
	(再掲)高血圧症有病者のうち服薬者*	499	55.8	0	0.0	4	10.0	8	17.4	54	40.9	139	53.5	294	71.9	304	51.7	181	65.8
	正常高値血圧者	229	14.5	10	12.7	32	17.0	33	20.1	38	16.1	60	15.5	56	10.6	160	16.1	56	14.4
	上記以外	459	29.0	61	77.2	116	61.7	85	51.8	66	28.0	68	17.5	63	11.9	247	24.8	57	14.7
総数		1,583	100.0	79	100.0	188	100.0	164	100.0	236	100.0	388	100.0	528	100.0	995	100.0	388	100.0
女性	高血圧症有病者	963	42.6	3	2.1	13	4.4	44	15.0	136	38.1	283	54.4	484	73.8	633	44.4	308	61.4
	(再掲)高血圧症有病者のうち服薬者*	574	59.6	0	0.0	1	7.7	7	15.9	66	48.5	179	63.3	321	66.3	350	55.3	184	59.7
	正常高値血圧者	288	12.7	4	2.9	20	6.8	34	11.6	52	14.6	96	18.5	82	12.5	221	15.5	94	18.7
	上記以外	1,010	44.7	133	95.0	262	88.8	215	73.4	169	47.3	141	27.1	90	13.7	573	40.2	100	19.9
総数		2,261	100.0	140	100.0	295	100.0	293	100.0	357	100.0	520	100.0	656	100.0	1,427	100.0	502	100.0

* 「服薬者」とは、質問票で「血圧を下げる薬」の使用有と回答した者

「高血圧症有病者」の判定
収縮期血圧140mmHg以上または拡張期血圧90mmHg以上、もしくは血圧を下げる薬を服用している者。

(23年)

第49表 「脂質異常症が疑われる人」の状況（採血時間によらず、性・年齢階級別）

	総数	20～29歳		30～39歳		40～49歳		50～59歳		60～69歳		70歳以上		(再掲) 40～74歳		(再掲) 65～74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	脂質異常症が疑われる人	807	22.7	10	5.0	34	7.5	46	10.7	86	15.8	242	28.5	389	36.1	524	23.3	274	33.3
	(再掲)脂質異常症が疑われる人のうち服薬者*	552	68.4	0	0.0	3	8.8	20	43.5	61	70.9	191	78.9	277	71.2	384	73.3	213	77.7
	上記以外	2,750	77.3	190	95.0	420	92.5	385	89.3	457	84.2	608	71.5	690	63.9	1,729	76.7	550	66.7
男性	総数	3,557	100.0	200	100.0	454	100.0	431	100.0	543	100.0	850	100.0	1,079	100.0	2,253	100.0	824	100.0
	脂質異常症が疑われる人	415	28.4	8	11.0	27	14.9	31	20.1	49	22.8	109	30.6	191	39.5	264	28.7	131	36.4
	(再掲)脂質異常症が疑われる人のうち服薬者*	221	53.3	0	0.0	2	7.4	11	35.5	27	55.1	71	65.1	110	57.6	158	59.8	87	66.4
女性	上記以外	1,048	71.6	65	89.0	154	85.1	123	79.9	166	77.2	247	69.4	293	60.5	655	71.3	229	63.6
	総数	1,463	100.0	73	100.0	181	100.0	154	100.0	215	100.0	356	100.0	484	100.0	919	100.0	360	100.0
	脂質異常症が疑われる人	392	18.7	2	1.6	7	2.6	15	5.4	37	11.3	133	26.9	198	33.3	260	19.5	143	30.8
	(再掲)脂質異常症が疑われる人のうち服薬者*	331	84.4	0	0.0	1	14.3	9	60.0	34	91.9	120	90.2	167	84.3	226	86.9	126	88.1
	上記以外	1,702	81.3	125	98.4	266	97.4	262	94.6	291	88.7	361	73.1	397	66.7	1,074	80.5	321	69.2
	総数	2,094	100.0	127	100.0	273	100.0	277	100.0	328	100.0	494	100.0	595	100.0	1,334	100.0	464	100.0

* 「服薬者」とは、質問票で「コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬」の使用有と回答した者

(23年)

「脂質異常症が疑われる人」の判定
国民健康・栄養調査の血液検査では、空腹時採血が困難であるため、脂質異常症の診断基準項目である中性脂肪による判定は行わず、下記の通りとした。
・「脂質異常症が疑われる人」
HDL コレステロールが 40mg/dL 未満、もしくはコレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬を服用している者。

第 50 表 運動習慣の状況（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 20-64歳		(再掲) 65歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	習慣無(健康上の理由)	330	7.9	6	2.1	12	2.2	10	1.9	31	4.7	64	6.7	207	16.6	79	3.2	251	14.8	96	10.4	155	20.1
	習慣無(その他)	2,532	60.5	232	82.9	437	81.5	387	74.9	444	67.8	497	52.0	535	43.0	1,787	71.6	745	44.0	406	44.1	339	43.9
	運動の習慣有	1,326	31.7	42	15.0	87	16.2	120	23.2	180	27.5	394	41.3	503	40.4	630	25.2	696	41.1	418	45.4	278	36.0
	総 数	4,188	100.0	280	100.0	536	100.0	517	100.0	655	100.0	955	100.0	1,245	100.0	2,496	100.0	1,692	100.0	920	100.0	772	100.0
男性	習慣無(健康上の理由)	111	6.3	3	2.7	6	2.8	5	2.5	12	4.4	19	4.6	66	11.9	30	2.9	81	10.8	28	6.9	53	15.5
	習慣無(その他)	1,039	58.7	83	74.1	175	80.3	141	70.9	196	71.5	221	53.5	223	40.3	727	71.1	312	41.8	166	41.1	146	42.6
	運動の習慣有	619	35.0	26	23.2	37	17.0	53	26.6	66	24.1	173	41.9	264	47.7	265	25.9	354	47.4	210	52.0	144	42.0
	総 数	1,769	100.0	112	100.0	218	100.0	199	100.0	274	100.0	413	100.0	553	100.0	1,022	100.0	747	100.0	404	100.0	343	100.0
女性	習慣無(健康上の理由)	219	9.1	3	1.8	6	1.9	5	1.6	19	5.0	45	8.3	141	20.4	49	3.3	170	18.0	68	13.2	102	23.8
	習慣無(その他)	1,493	61.7	149	88.7	262	82.4	246	77.4	248	65.1	276	50.9	312	45.1	1,060	71.9	433	45.8	240	46.5	193	45.0
	運動の習慣有	707	29.2	16	9.5	50	15.7	67	21.1	114	29.9	221	40.8	239	34.5	365	24.8	342	36.2	208	40.3	134	31.2
	総 数	2,419	100.0	168	100.0	318	100.0	318	100.0	381	100.0	542	100.0	692	100.0	1,474	100.0	945	100.0	516	100.0	429	100.0

(23年)

第 51 表 1 週間の運動日数（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	運動無し	2,862	68.3	238	85.0	449	83.8	397	76.8	475	72.5	561	58.7	742	59.6	502	54.6	494	64.0
	2日/週	284	6.8	15	5.4	31	5.8	38	7.4	50	7.6	68	7.1	82	6.6	73	7.9	41	5.3
	3日/週	232	5.5	12	4.3	15	2.8	20	3.9	34	5.2	69	7.2	82	6.6	68	7.4	47	6.1
	4日/週	141	3.4	3	1.1	5	0.9	10	1.9	15	2.3	53	5.5	55	4.4	53	5.8	25	3.2
	5日/週	175	4.2	5	1.8	18	3.4	21	4.1	24	3.7	58	6.1	49	3.9	53	5.8	26	3.4
	6日/週	80	1.9	1	0.4	4	0.7	6	1.2	6	0.9	26	2.7	37	3.0	31	3.4	21	2.7
	7日/週	414	9.9	6	2.1	14	2.6	25	4.8	51	7.8	120	12.6	198	15.9	140	15.2	118	15.3
	総 数	4,188	100.0	280	100.0	536	100.0	517	100.0	655	100.0	955	100.0	1,245	100.0	920	100.0	772	100.0
男性	運動無し	1,150	65.0	86	76.8	181	83.0	146	73.4	208	75.9	240	58.1	289	52.3	194	48.0	199	58.0
	2日/週	131	7.4	7	6.3	12	5.5	18	9.0	22	8.0	31	7.5	41	7.4	32	7.9	21	6.1
	3日/週	94	5.3	8	7.1	6	2.8	8	4.0	11	4.0	25	6.1	36	6.5	31	7.7	19	5.5
	4日/週	66	3.7	2	1.8	2	0.9	7	3.5	1	0.4	25	6.1	29	5.2	28	6.9	12	3.5
	5日/週	86	4.9	2	1.8	10	4.6	11	5.5	10	3.6	26	6.3	27	4.9	27	6.7	15	4.4
	6日/週	41	2.3	1	0.9	2	0.9	4	2.0	4	1.5	13	3.1	17	3.1	14	3.5	10	2.9
	7日/週	201	11.4	6	5.4	5	2.3	5	2.5	18	6.6	53	12.8	114	20.6	78	19.3	67	19.5
	総 数	1,769	100.0	112	100.0	218	100.0	199	100.0	274	100.0	413	100.0	553	100.0	404	100.0	343	100.0
女性	運動無し	1,712	70.8	152	90.5	268	84.3	251	78.9	267	70.1	321	59.2	453	65.5	308	59.7	295	68.8
	2日/週	153	6.3	8	4.8	19	6.0	20	6.3	28	7.3	37	6.8	41	5.9	41	7.9	20	4.7
	3日/週	138	5.7	4	2.4	9	2.8	12	3.8	23	6.0	44	8.1	46	6.6	37	7.2	28	6.5
	4日/週	75	3.1	1	0.6	3	0.9	3	0.9	14	3.7	28	5.2	26	3.8	25	4.8	13	3.0
	5日/週	89	3.7	3	1.8	8	2.5	10	3.1	14	3.7	32	5.9	22	3.2	26	5.0	11	2.6
	6日/週	39	1.6	0	0.0	2	0.6	2	0.6	2	0.5	13	2.4	20	2.9	17	3.3	11	2.6
	7日/週	213	8.8	0	0.0	9	2.8	20	6.3	33	8.7	67	12.4	84	12.1	62	12.0	51	11.9
	総 数	2,419	100.0	168	100.0	318	100.0	318	100.0	381	100.0	542	100.0	692	100.0	516	100.0	429	100.0

(23年)

第 52 表 運動を行う日の平均運動時間（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	運動無し	2,862	68.3	238	85.0	449	83.8	397	76.8	475	72.5	561	58.7	742	59.6	502	54.6	494	64.0
	30分以上60分未満	534	12.8	16	5.7	30	5.6	44	8.5	79	12.1	159	16.6	206	16.5	160	17.4	130	16.8
	60分以上120分未満	509	12.2	10	3.6	37	6.9	47	9.1	69	10.5	155	16.2	191	15.3	165	17.9	100	13.0
	120分以上180分未満	189	4.5	13	4.6	16	3.0	17	3.3	18	2.7	56	5.9	69	5.5	61	6.6	32	4.1
	180分以上	94	2.2	3	1.1	4	0.7	12	2.3	14	2.1	24	2.5	37	3.0	32	3.5	16	2.1
	総 数	4,188	100.0	280	100.0	536	100.0	517	100.0	655	100.0	955	100.0	1,245	100.0	920	100.0	772	100.0
男性	運動無し	1,150	65.0	86	76.8	181	83.0	146	73.4	208	75.9	240	58.1	289	52.3	194	48.0	199	58.0
	30分以上60分未満	222	12.5	11	9.8	13	6.0	19	9.5	28	10.2	55	13.3	96	17.4	70	17.3	60	17.5
	60分以上120分未満	244	13.8	3	2.7	14	6.4	16	8.0	24	8.8	83	20.1	104	18.8	93	23.0	51	14.9
	120分以上180分未満	93	5.3	9	8.0	6	2.8	11	5.5	6	2.2	22	5.3	39	7.1	28	6.9	21	6.1
	180分以上	60	3.4	3	2.7	4	1.8	7	3.5	8	2.9	13	3.1	25	4.5	19	4.7	12	3.5
	総 数	1,769	100.0	112	100.0	218	100.0	199	100.0	274	100.0	413	100.0	553	100.0	404	100.0	343	100.0
女性	運動無し	1,712	70.8	152	90.5	268	84.3	251	78.9	267	70.1	321	59.2	453	65.5	308	59.7	295	68.8
	30分以上60分未満	312	12.9	5	3.0	17	5.3	25	7.9	51	13.4	104	19.2	110	15.9	90	17.4	70	16.3
	60分以上120分未満	265	11.0	7	4.2	23	7.2	31	9.7	45	11.8	72	13.3	87	12.6	72	14.0	49	11.4
	120分以上180分未満	96	4.0	4	2.4	10	3.1	6	1.9	12	3.1	34	6.3	30	4.3	33	6.4	11	2.6
	180分以上	34	1.4	0	0.0	0	0.0	5	1.6	6	1.6	11	2.0	12	1.7	13	2.5	4	0.9
	総 数	2,419	100.0	168	100.0	318	100.0	318	100.0	381	100.0	542	100.0	692	100.0	516	100.0	429	100.0

(23年)

第53表 運動強度の状況（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	運動無し	2,862	68.3	238	85.0	449	83.8	397	76.8	475	72.5	561	58.7	742	59.6	502	54.6	494	64.0
	高強度	96	2.3	12	4.3	17	3.2	22	4.3	11	1.7	21	2.2	13	1.0	15	1.6	5	0.6
	中強度	414	9.9	14	5.0	34	6.3	52	10.1	77	11.8	133	13.9	104	8.4	121	13.2	49	6.3
	低強度	816	19.5	16	5.7	36	6.7	46	8.9	92	14.0	240	25.1	386	31.0	282	30.7	224	29.0
	総 数	4,188	100.0	280	100.0	536	100.0	517	100.0	655	100.0	955	100.0	1,245	100.0	920	100.0	772	100.0
男性	運動無し	1,150	65.0	86	76.8	181	83.0	146	73.4	208	75.9	240	58.1	289	52.3	194	48.0	199	58.0
	高強度	61	3.4	10	8.9	9	4.1	15	7.5	8	2.9	10	2.4	9	1.6	9	2.2	4	1.2
	中強度	195	11.0	8	7.1	15	6.9	23	11.6	25	9.1	59	14.3	65	11.8	64	15.8	30	8.7
	低強度	363	20.5	8	7.1	13	6.0	15	7.5	33	12.0	104	25.2	190	34.4	137	33.9	110	32.1
	総 数	1,769	100.0	112	100.0	218	100.0	199	100.0	274	100.0	413	100.0	553	100.0	404	100.0	343	100.0
女性	運動無し	1,712	70.8	152	90.5	268	84.3	251	78.9	267	70.1	321	59.2	453	65.5	308	59.7	295	68.8
	高強度	35	1.4	2	1.2	8	2.5	7	2.2	3	0.8	11	2.0	4	0.6	6	1.2	1	0.2
	中強度	219	9.1	6	3.6	19	6.0	29	9.1	52	13.6	74	13.7	39	5.6	57	11.0	19	4.4
	低強度	453	18.7	8	4.8	23	7.2	31	9.7	59	15.5	136	25.1	196	28.3	145	28.1	114	26.6
	総 数	2,419	100.0	168	100.0	318	100.0	318	100.0	381	100.0	542	100.0	692	100.0	516	100.0	429	100.0

(23年)

第54表 1日の歩行数の分布（性・年齢階級別）

		総 数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 20歳以上		(再掲) 20-64歳		(再掲) 65歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	2,000歩未満	610	9.1	8	2.3	27	4.8	46	4.9	22	2.2	54	5.2	92	7.1	361	23.7	602	9.5	192	4.5	410	19.4	122	10.4	288	30.7
	2,000-3,999	1,193	17.8	36	10.5	86	15.4	149	15.8	138	13.9	166	15.9	229	17.6	389	25.5	1,157	18.2	654	15.3	503	23.9	235	20.1	268	28.6
	4,000-5,999	1,367	20.4	59	17.2	106	19.0	201	21.3	226	22.7	207	19.8	256	19.7	312	20.4	1,308	20.5	880	20.7	428	20.3	257	21.9	171	18.2
	6,000-7,999	1,236	18.4	72	21.0	115	20.6	172	18.2	205	20.6	205	19.6	263	20.2	204	13.4	1,164	18.3	845	19.8	319	15.1	212	18.1	107	11.4
	8,000-9,999	966	14.4	54	15.7	81	14.5	154	16.3	159	16.0	170	16.2	217	16.7	131	8.6	912	14.3	689	16.2	223	10.6	163	13.9	60	6.4
	10,000歩以上	1,340	20.0	114	33.2	142	25.5	223	23.6	245	24.6	245	23.4	242	18.6	129	8.5	1,226	19.2	1,001	23.5	225	10.7	182	15.5	43	4.6
	9,200歩(男)/8,300歩(女)以上	1,905	28.4	142	41.4	189	33.9	312	33.0	339	34.1	352	33.6	371	28.6	200	13.1	1,763	27.7	1,415	33.2	348	16.5	273	23.3	75	8.0
	総 数	6,712	100.0	343	100.0	557	100.0	945	100.0	995	100.0	1,047	100.0	1,299	100.0	1,526	100.0	6,369	100.0	4,261	100.0	2,108	100.0	1,171	100.0	937	100.0
	平均値(歩/日)	6,895		8,721		7,802		7,537		7,692		7,524		6,970		4,740		6,797		7,554		5,266		6,309		3,963	
	標準偏差	4,293		4,824		4,512		4,271		4,049		4,295		4,081		3,609		4,241		4,225		3,841		3,918		3,312	
男性	2,000歩未満	271	8.9	5	2.9	13	5.3	22	5.0	11	2.5	31	6.3	45	7.9	144	21.3	266	9.2	101	5.2	165	17.7	58	11.1	107	26.2
	2,000-3,999	516	16.9	25	14.5	42	17.1	54	12.3	58	12.9	84	16.9	90	15.7	163	24.1	491	17.1	287	14.7	204	21.9	94	18.0	110	26.9
	4,000-5,999	567	18.6	31	17.9	44	17.9	85	19.4	91	20.3	83	16.7	110	19.2	123	18.2	536	18.6	366	18.8	170	18.3	103	19.7	67	16.4
	6,000-7,999	552	18.1	32	18.5	41	16.7	80	18.2	98	21.9	95	19.2	106	18.5	100	14.8	520	18.1	372	19.1	148	15.9	89	17.0	59	14.4
	8,000-9,999	429	14.1	26	15.0	33	13.4	69	15.7	63	14.1	75	15.1	96	16.8	67	9.9	403	14.0	292	15.0	111	11.9	82	15.7	29	7.1
	10,000歩以上	717	23.5	54	31.2	73	29.7	129	29.4	127	28.3	128	25.8	126	22.0	80	11.8	663	23.0	530	27.2	133	14.3	96	18.4	37	9.0
	(再掲)9,200歩以上	855	28.0	60	34.7	84	34.1	150	34.2	147	32.8	158	31.9	156	27.2	100	14.8	795	27.6	628	32.2	167	17.9	123	23.6	44	10.8
	総 数	3,052	100.0	173	100.0	246	100.0	439	100.0	448	100.0	496	100.0	573	100.0	677	100.0	2,879	100.0	1,948	100.0	931	100.0	522	100.0	409	100.0
	平均値(歩/日)	7,301		8,433		8,199		8,238		8,090		7,693		7,307		5,263		7,233		7,935		5,763		6,599		4,696	
	標準偏差	4,679		4,895		5,151		4,725		4,571		4,678		4,486		3,983		4,657		4,730		4,135		4,090		3,948	
女性	2,000歩未満	339	9.3	3	1.8	14	4.5	24	4.7	11	2.0	23	4.2	47	6.5	217	25.6	336	9.6	91	3.9	245	20.8	64	9.9	181	34.3
	2,000-3,999	677	18.5	11	6.5	44	14.1	95	18.8	80	14.6	82	14.9	139	19.1	226	26.6	666	19.1	367	15.9	299	25.4	141	21.7	158	29.9
	4,000-5,999	800	21.9	28	16.5	62	19.9	116	22.9	135	24.7	124	22.5	146	20.1	189	22.3	772	22.1	514	22.2	258	21.9	154	23.7	104	19.7
	6,000-7,999	684	18.7	40	23.5	74	23.8	92	18.2	107	19.6	110	20.0	157	21.6	104	12.2	644	18.5	473	20.4	171	14.5	123	19.0	48	9.1
	8,000-9,999	537	14.7	28	16.5	48	15.4	85	16.8	96	17.6	95	17.2	121	16.7	64	7.5	509	14.6	397	17.2	112	9.5	81	12.5	31	5.9
	10,000歩以上	623	17.0	60	35.3	69	22.2	94	18.6	118	21.6	117	21.2	116	16.0	49	5.8	563	16.1	471	20.4	92	7.8	86	13.3	6	1.1
	(再掲)8,300歩以上	1,050	28.7	82	48.2	105	33.8	162	32.0	192	35.1	194	35.2	215	29.6	100	11.8	968	27.7	787	34.0	181	15.4	150	23.1	31	5.9
	総 数	3,660	100.0	170	100.0	311	100.0	506	100.0	547	100.0	551	100.0	726	100.0	849	100.0	3,490	100.0	2,313	100.0	1,177	100.0	649	100.0	528	100.0
	平均値(歩/日)	6,556		9,015		7,487		6,930		7,367		7,371		6,705		4,323		6,437		7,233		4,873		6,075		3,394	
	標準偏差	3,912		4,747		3,914		3,732		3,535		3,917		3,712		3,222		3,827		3,719		3,545		3,761		2,584	

(23年)

