

第 2 部 身体状況調査の結果

第 11 表 身長・体重の平均値及び標準偏差（性・年齢階級別）

年齢(歳)	男 性						女 性					
	身 長(cm)			体 重(kg)			身 長(cm)			体 重(kg)		
	人 数	平均値	標準偏差	人 数	平均値	標準偏差	人 数	平均値	標準偏差	人 数	平均値	標準偏差
総 数	3,232	163.9	12.4	3,219	62.0	14.8	3,861	152.5	9.0	3,826	51.3	10.6
1	—	—	—	—	—	—	—	—	—	—	—	—
2	—	—	—	—	—	—	—	—	—	—	—	—
3	—	—	—	—	—	—	—	—	—	—	—	—
4	—	—	—	—	—	—	—	—	—	—	—	—
5	—	—	—	—	—	—	—	—	—	—	—	—
6	42	117.1	5.2	42	20.9	2.7	38	115.1	4.0	38	20.1	2.4
7	44	120.8	6.2	43	23.6	4.5	34	121.8	5.2	34	23.5	3.6
8	38	129.4	6.1	38	27.2	4.3	38	128.4	5.9	38	26.5	4.4
9	37	132.7	5.5	37	30.8	7.4	39	135.2	5.9	38	30.8	6.7
10	46	138.5	5.8	46	32.9	6.9	36	139.2	7.4	36	33.1	6.4
11	42	143.8	6.9	42	36.2	6.8	41	145.3	6.3	41	36.8	6.7
12	32	151.9	7.7	32	44.5	12.9	28	150.8	5.8	28	40.8	7.8
13	30	159.6	10.1	30	46.9	10.3	29	155.7	4.6	29	46.2	7.1
14	32	164.4	6.6	32	52.6	9.1	31	156.1	4.9	31	48.4	5.8
15	35	169.0	5.8	35	56.7	10.7	39	156.6	5.5	39	51.0	7.6
16	39	168.5	5.8	39	57.7	9.2	28	158.0	4.1	27	51.0	7.2
17	33	170.9	5.1	32	60.9	8.1	34	158.0	4.9	33	51.9	6.0
18	19	172.5	4.8	19	62.5	13.4	26	157.7	5.5	26	56.3	12.8
19	20	169.0	5.7	20	59.4	8.3	21	159.2	5.1	21	55.2	9.9
20	21	171.0	6.3	21	61.8	13.2	20	159.5	7.5	20	55.0	14.1
21	16	170.2	5.7	16	60.7	8.6	23	156.7	3.7	22	49.4	6.5
22	19	170.1	6.1	18	69.5	13.4	22	157.2	6.0	20	50.3	7.4
23	22	171.9	5.3	22	65.7	12.2	21	154.8	4.7	21	48.4	7.6
24	23	171.5	5.6	23	62.9	10.5	34	159.1	4.7	33	49.8	8.8
25	19	169.6	6.9	19	63.5	10.3	25	157.7	5.1	25	50.2	6.8
26-29	98	170.4	6.2	96	66.5	12.3	114	158.7	5.4	111	51.8	9.0
30-39	387	171.5	6.1	385	69.6	11.7	486	158.3	5.3	465	54.0	9.2
40-49	397	170.6	5.8	395	70.4	11.6	489	157.8	5.2	487	54.7	9.3
50-59	453	168.0	6.1	451	68.2	10.4	567	155.2	5.4	564	54.1	8.6
60-69	650	165.3	6.1	649	64.6	9.1	765	151.8	5.5	763	53.4	8.3
70歳以上	638	160.9	6.3	637	59.9	9.4	833	147.7	6.2	836	50.4	8.8
(再掲)												
20歳以上	2,743	166.8	7.2	2,732	65.7	11.1	3,399	153.6	6.9	3,367	52.9	9.0
20-29	218	170.6	6.0	215	65.1	12.0	259	158.1	5.4	252	51.0	9.0
60-64	335	166.4	6.2	334	65.7	9.2	377	152.7	5.5	376	54.2	8.7
65-69	315	164.1	5.8	315	63.4	8.7	388	150.9	5.3	387	52.6	7.9
70-74	261	162.8	6.2	260	62.3	9.5	316	149.9	5.4	316	52.2	8.7
75-79	190	160.2	5.8	190	59.5	9.2	251	147.1	6.0	251	50.5	8.5
80歳以上	187	159.1	6.3	187	56.9	8.4	266	145.6	6.5	269	48.1	8.8

(22年)

注) 体重は妊婦除外。平成 22 年は乳幼児身体発育調査実施のため 1 - 5 歳は除く。

第12表 BMIの平均値及び標準偏差（15歳以上，性・年齢階級別）

年齢(歳)	男 性			女 性		
	人 数	平均値	標準偏差	人 数	平均値	標準偏差
総 数	2,876	23.41	3.40	3,509	22.37	3.57
15-19歳	145	20.45	3.06	146	21.19	3.58
20-29歳	215	22.33	3.68	252	20.37	3.16
30-39歳	385	23.64	3.57	465	21.56	3.34
40-49歳	395	24.18	3.57	487	21.96	3.55
50-59歳	451	24.13	3.30	564	22.50	3.49
60-69歳	648	23.63	2.92	762	23.17	3.41
70歳以上	637	23.09	3.22	833	23.06	3.61
(再掲)						
20歳以上	2,731	23.57	3.35	3,363	22.42	3.56
20-69歳	2,094	23.71	3.37	2,530	22.21	3.52
40-69歳	1,494	23.93	3.23	1,813	22.64	3.51
65-69歳	314	23.53	2.77	387	23.09	3.26
70-74歳	260	23.47	3.28	316	23.23	3.55
75-79歳	190	23.17	3.27	251	23.30	3.44
80-84歳	127	22.67	3.01	173	22.95	3.79
80歳以上	187	22.49	3.01	266	22.63	3.82
85歳以上	60	22.12	2.99	93	22.05	3.83

注) 妊婦除外

(22年)

第13表 BMIの分布（15歳以上，性別）

	総 数		男 性		女 性	
	人 数	%	人 数	%	人 数	%
総 数	6,385	100.0	2,876	100.0	3,509	100.0
15未満	11	0.2	4	0.1	7	0.2
15-16未満	39	0.6	13	0.5	26	0.7
16-17未満	104	1.6	29	1.0	75	2.1
17-18未満	232	3.6	65	2.3	167	4.8
18-19未満	427	6.7	128	4.5	299	8.5
19-20未満	599	9.4	228	7.9	371	10.6
20-21未満	677	10.6	240	8.3	437	12.5
21-22未満	705	11.0	302	10.5	403	11.5
22-23未満	750	11.7	340	11.8	410	11.7
23-24未満	675	10.6	363	12.6	312	8.9
24-25未満	597	9.4	320	11.1	277	7.9
25-26未満	497	7.8	266	9.2	231	6.6
26-27未満	328	5.1	186	6.5	142	4.0
27-28未満	246	3.9	148	5.1	98	2.8
28-29未満	174	2.7	83	2.9	91	2.6
29-30未満	106	1.7	55	1.9	51	1.5
30-31未満	70	1.1	34	1.2	36	1.0
31-32未満	48	0.8	22	0.8	26	0.7
32-33未満	37	0.6	21	0.7	16	0.5
33-34未満	21	0.3	12	0.4	9	0.3
34-35未満	14	0.2	7	0.2	7	0.2
35以上	28	0.4	10	0.3	18	0.5

注) 妊婦除外

(22年)

第14表 BMIの状況（15歳以上、性・年齢階級別）

		総 数		や せ		普 通		肥 満		(再掲) 20以下		(再掲) 25以上30未満		(再掲) 30以上	
				18.5未満		18.5以上25未満		25以上							
		人 数	%	人 数	%	人 数	%	人 数	%	人 数	%	人 数	%	人 数	%
総数	総 数	6,385	100.0	558	8.7	4,258	66.7	1,569	24.6	－	－	1,351	21.2	218	3.4
	15-19歳	291	100.0	61	21.0	203	69.8	27	9.3	－	－	21	7.2	6	2.1
	20-29歳	467	100.0	98	21.0	308	66.0	61	13.1	－	－	44	9.4	17	3.6
	30-39歳	850	100.0	77	9.1	598	70.4	175	20.6	－	－	142	16.7	33	3.9
	40-49歳	882	100.0	67	7.6	587	66.6	228	25.9	－	－	189	21.4	39	4.4
	50-59歳	1,015	100.0	64	6.3	676	66.6	275	27.1	－	－	235	23.2	40	3.9
	60-69歳	1,410	100.0	65	4.6	945	67.0	400	28.4	－	－	361	25.6	39	2.8
	70歳以上	1,470	100.0	126	8.6	941	64.0	403	27.4	－	－	359	24.4	44	3.0
	(再掲)														
	20歳以上	6,094	100.0	497	8.2	4,055	66.5	1,542	25.3	－	－	1,330	21.8	212	3.5
	20-69歳	4,624	100.0	371	8.0	3,114	67.3	1,139	24.6	－	－	971	21.0	168	3.6
	40-69歳	3,307	100.0	196	5.9	2,208	66.8	903	27.3	－	－	785	23.7	118	3.6
	65-69歳	701	100.0	31	4.4	483	68.9	187	26.7	98	14.0	170	24.3	17	2.4
	70-74歳	576	100.0	37	6.4	374	64.9	165	28.6	92	16.0	146	25.3	19	3.3
	75-79歳	441	100.0	34	7.7	279	63.3	128	29.0	72	16.3	114	25.9	14	3.2
	80-84歳	300	100.0	29	9.7	191	63.7	80	26.7	73	24.3	72	24.0	8	2.7
	65歳以上	2,171	100.0	157	7.2	1,424	65.6	590	27.2	377	17.4	529	24.4	61	2.8
	80歳以上	453	100.0	55	12.1	288	63.6	110	24.3	115	25.4	99	21.9	11	2.4
	85歳以上	153	100.0	26	17.0	97	63.4	30	19.6	42	27.5	27	17.6	3	2.0
男性	総 数	2,876	100.0	162	5.6	1,870	65.0	844	29.3	－	－	738	25.7	106	3.7
	15-19歳	145	100.0	36	24.8	96	66.2	13	9.0	－	－	11	7.6	2	1.4
	20-29歳	215	100.0	25	11.6	148	68.8	42	19.5	－	－	32	14.9	10	4.7
	30-39歳	385	100.0	10	2.6	264	68.6	111	28.8	－	－	87	22.6	24	6.2
	40-49歳	395	100.0	12	3.0	244	61.8	139	35.2	－	－	111	28.1	28	7.1
	50-59歳	451	100.0	9	2.0	274	60.8	168	37.3	－	－	147	32.6	21	4.7
	60-69歳	648	100.0	19	2.9	435	67.1	194	29.9	－	－	183	28.2	11	1.7
	70歳以上	637	100.0	51	8.0	409	64.2	177	27.8	－	－	167	26.2	10	1.6
	(再掲)														
	20歳以上	2,731	100.0	126	4.6	1,774	65.0	831	30.4	－	－	727	26.6	104	3.8
	20-69歳	2,094	100.0	75	3.6	1,365	65.2	654	31.2	－	－	560	26.7	94	4.5
	40-69歳	1,494	100.0	40	2.7	953	63.8	501	33.5	－	－	441	29.5	60	4.0
	65-69歳	314	100.0	11	3.5	215	68.5	88	28.0	31	9.9	84	26.8	4	1.3
	70-74歳	260	100.0	17	6.5	162	62.3	81	31.2	37	14.2	76	29.2	5	1.9
	75-79歳	190	100.0	16	8.4	118	62.1	56	29.5	36	18.9	51	26.8	5	2.6
	80-84歳	127	100.0	11	8.7	87	68.5	29	22.8	30	23.6	29	22.8	0	0.0
	65歳以上	951	100.0	62	6.5	624	65.6	265	27.9	148	15.6	251	26.4	14	1.5
	80歳以上	187	100.0	18	9.6	129	69.0	40	21.4	44	23.5	40	21.4	0	0.0
	85歳以上	60	100.0	7	11.7	42	70.0	11	18.3	14	23.3	11	18.3	0	0.0
女性	総 数	3,509	100.0	396	11.3	2,388	68.1	725	20.7	－	－	613	17.5	112	3.2
	15-19歳	146	100.0	25	17.1	107	73.3	14	9.6	－	－	10	6.8	4	2.7
	20-29歳	252	100.0	73	29.0	160	63.5	19	7.5	－	－	12	4.8	7	2.8
	30-39歳	465	100.0	67	14.4	334	71.8	64	13.8	－	－	55	11.8	9	1.9
	40-49歳	487	100.0	55	11.3	343	70.4	89	18.3	－	－	78	16.0	11	2.3
	50-59歳	564	100.0	55	9.8	402	71.3	107	19.0	－	－	88	15.6	19	3.4
	60-69歳	762	100.0	46	6.0	510	66.9	206	27.0	－	－	178	23.4	28	3.7
	70歳以上	833	100.0	75	9.0	532	63.9	226	27.1	－	－	192	23.0	34	4.1
	(再掲)														
	20歳以上	3,363	100.0	371	11.0	2,281	67.8	711	21.1	－	－	603	17.9	108	3.2
	20-69歳	2,530	100.0	296	11.7	1,749	69.1	485	19.2	－	－	411	16.2	74	2.9
	40-69歳	1,813	100.0	156	8.6	1,255	69.2	402	22.2	－	－	344	19.0	58	3.2
	65-69歳	387	100.0	20	5.2	268	69.3	99	25.6	67	17.3	86	22.2	13	3.4
	70-74歳	316	100.0	20	6.3	212	67.1	84	26.6	55	17.4	70	22.2	14	4.4
	75-79歳	251	100.0	18	7.2	161	64.1	72	28.7	36	14.3	63	25.1	9	3.6
	80-84歳	173	100.0	18	10.4	104	60.1	51	29.5	229	18.8	43	24.9	8	4.6
	65歳以上	1,220	100.0	95	7.8	800	65.6	325	26.6	43	24.9	278	22.8	47	3.9
	80歳以上	266	100.0	37	13.9	159	59.8	70	26.3	71	26.7	59	22.2	11	4.1
	85歳以上	93	100.0	19	20.4	55	59.1	19	20.4	28	30.1	16	17.2	3	3.2

注) 妊婦除外。BMIの判定はp. 17 参照

(22年)

第15表 学校保健統計調査方式による肥満区分（6－14歳，性・年齢階級別）

		総 数		6－8歳		9－11歳		12－14歳	
		人 数	%	人 数	%	人 数	%	人 数	%
総数	総 数	655	100.0	233	100.0	240	100.0	182	100.0
	－30%未満	3	0.5	1	0.4	1	0.4	1	0.5
	－30%超～－20%未満	25	3.8	3	1.3	15	6.3	7	3.8
	－20%超～－10%未満	139	21.2	33	14.2	52	21.7	54	29.7
	－10%超～ 0%未満	239	36.5	111	47.6	76	31.7	52	28.6
	0%以上～+10%未満	135	20.6	55	23.6	45	18.8	35	19.2
	+10%以上～+20%未満	75	11.5	19	8.2	33	13.8	23	12.6
	+20%以上～+30%未満	25	3.8	9	3.9	11	4.6	5	2.7
	+30%以上～+40%未満	6	0.9	1	0.4	4	1.7	1	0.5
	+40%以上～+50%未満	4	0.6	1	0.4	1	0.4	2	1.1
	+50%以上	4	0.6	0	0.0	2	0.8	2	1.1
男性	総 数	342	100.0	123	100.0	125	100.0	94	100.0
	－30%未満	1	0.3	1	0.8	0	0.0	0	0.0
	－30%超～－20%未満	17	5.0	2	1.6	9	7.2	6	6.4
	－20%超～－10%未満	66	19.3	19	15.4	24	19.2	23	24.5
	－10%超～ 0%未満	123	36.0	52	42.3	44	35.2	27	28.7
	0%以上～+10%未満	73	21.3	33	26.8	20	16.0	20	21.3
	+10%以上～+20%未満	35	10.2	8	6.5	17	13.6	10	10.6
	+20%以上～+30%未満	18	5.3	6	4.9	8	6.4	4	4.3
	+30%以上～+40%未満	3	0.9	1	0.8	1	0.8	1	1.1
	+40%以上～+50%未満	2	0.6	1	0.8	0	0.0	1	1.1
	+50%以上	4	1.2	0	0.0	2	1.6	2	2.1
女性	総 数	313	100.0	110	100.0	115	100.0	88	100.0
	－30%未満	2	0.6	0	0.0	1	0.9	1	1.1
	－30%超～－20%未満	8	2.6	1	0.9	6	5.2	1	1.1
	－20%超～－10%未満	73	23.3	14	12.7	28	24.3	31	35.2
	－10%超～ 0%未満	116	37.1	59	53.6	32	27.8	25	28.4
	0%以上～+10%未満	62	19.8	22	20.0	25	21.7	15	17.0
	+10%以上～+20%未満	40	12.8	11	10.0	16	13.9	13	14.8
	+20%以上～+30%未満	7	2.2	3	2.7	3	2.6	1	1.1
	+30%以上～+40%未満	3	1.0	0	0.0	3	2.6	0	0.0
	+40%以上～+50%未満	2	0.6	0	0.0	1	0.9	1	1.1
	+50%以上	0	0.0	0	0.0	0	0.0	0	0.0

注) 学校保健統計調査方式による区分は，p.17 参照

(22年)

第16表 腹囲の分布（6歳以上、性・年齢階級別）

総数		6-8歳		9-11歳		12-14歳		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 20歳以上		(再掲) 40-74歳	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総数	6641	100.0	222	100.0	157	100.0	262	100.0	415	100.0	783	100.0	840	100.0	978	100.0	1,353	100.0	1,410	100.0	5,779	100.0	3,735	100.0
	50cm未満	23	0.3	3	1.4	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	50－55cm未満	132	2.0	31	14.0	3	1.9	0	0.0	1	0.2	0	0.0	1	0.1	1	0.1	0	0.0	1	0.1	4	0.1	2	0.1
	55－60	200	3.0	69	31.2	20	12.7	9	3.4	8	1.9	7	0.9	4	0.5	1	0.1	4	0.3	2	0.1	26	0.4	10	0.3
	60－65	297	4.5	27	12.2	36	22.9	27	10.3	37	8.9	26	3.3	37	4.4	22	2.2	13	1.0	21	1.5	156	2.7	79	2.1
	65－70	507	7.6	30	13.5	43	27.4	60	22.9	70	16.9	73	9.3	70	8.3	56	5.7	51	3.8	47	3.3	367	6.4	192	5.1
	70－75	884	13.3	4	1.8	36	22.9	85	32.4	98	23.6	154	19.7	133	15.8	106	10.8	110	8.1	143	10.1	744	12.9	397	10.6
	75－80	988	14.9	1	0.5	12	7.6	44	16.8	87	21.0	168	21.5	160	19.0	149	15.2	198	14.6	160	11.3	922	16.0	576	15.4
	80－85	1,153	17.4	0	0.0	6	2.7	2	1.3	18	6.9	156	19.9	163	19.4	229	23.4	260	19.2	270	19.1	1,127	19.5	763	20.4
	85－90	1,051	15.8	0	0.0	1	0.5	4	2.5	6	2.3	84	10.7	118	14.0	180	18.4	317	23.4	311	22.1	1,040	18.0	750	20.1
	90－95	741	11.2	0	0.0	0	0.0	7	2.7	11	2.7	56	7.2	73	8.7	120	12.3	236	17.4	238	16.9	734	12.7	527	14.1
	95－100	380	5.7	0	0.0	1	0.6	0	0.0	9	2.2	33	4.2	47	5.6	68	7.0	89	6.6	133	9.4	379	6.6	251	6.7
	100－105	176	2.7	0	0.0	0	0.0	3	1.1	7	1.7	16	2.0	14	1.7	29	3.0	53	3.9	54	3.8	173	3.0	117	3.1
	105－110	60	0.9	0	0.0	0	0.0	0	0.0	5	1.2	2	0.3	11	1.3	9	0.9	11	0.8	22	1.6	60	1.0	39	1.0
男性	110－115	31	0.5	0	0.0	0	0.0	2	0.8	2	0.5	2	0.3	6	0.7	6	0.6	7	0.5	6	0.4	29	0.5	22	0.6
	115－120	12	0.2	0	0.0	0	0.0	0	0.0	1	0.2	4	0.5	2	0.2	2	0.2	2	0.1	1	0.1	12	0.2	7	0.2
	120cm以上	6	0.1	0	0.0	0	0.0	0	0.0	0	0.0	2	0.3	1	0.1	0	0.0	2	0.1	1	0.1	6	0.1	3	0.1
	総数	3,032	100.0	115	100.0	82	100.0	132	100.0	192	100.0	350	100.0	375	100.0	435	100.0	623	100.0	613	100.0	2,588	100.0	1,686	100.0
	50cm未満	8	0.3	1	0.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	50－55cm未満	68	2.2	50	43.5	16	13.9	1	1.2	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	1	0.0	1	0.1
	55－60	88	2.9	37	32.2	42	36.5	9	11.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	60－65	78	2.6	13	11.3	25	21.7	16	19.5	4	2.1	1	0.3	2	0.5	1	0.2	2	0.3	4	0.7	14	0.5	6	0.4
	65－70	146	4.8	4	3.5	16	13.9	23	28.0	35	26.5	13	3.7	6	1.6	7	1.6	9	1.4	17	2.8	68	2.6	27	1.6
	70－75	312	10.3	3	2.6	6	5.2	22	26.8	40	30.3	46	13.1	32	8.5	28	6.4	32	5.1	60	9.8	241	9.3	113	6.7
	75－80	392	12.9	1	0.9	5	4.3	27	20.5	46	24.0	70	20.0	63	16.8	38	8.7	76	12.2	61	10.0	354	13.7	204	12.1
	80－85	591	19.5	0	0.0	0	0.0	6	4.5	8	4.2	39	11.1	47	12.5	109	25.1	132	21.2	128	20.9	575	22.2	381	22.6
	85－90	586	19.3	0	0.0	1	0.9	3	3.7	21	10.9	53	15.1	78	20.8	109	25.1	174	27.9	146	23.8	581	22.4	425	25.2
女性	90－95	409	13.5	0	0.0	0	0.0	0	0.0	8	4.2	24	6.9	35	9.3	44	10.1	38	6.1	63	10.3	403	15.6	291	17.3
	95－100	213	7.0	0	0.0	1	1.2	0	0.0	8	4.2	24	6.9	35	9.3	44	10.1	38	6.1	63	10.3	212	8.2	140	8.3
	100－105	93	3.1	0	0.0	0	0.0	0	0.0	2	1.5	7	3.6	12	3.2	16	3.7	23	3.7	20	3.3	91	3.5	63	3.7
	105－110	30	1.0	0	0.0	0	0.0	0	0.0	2	1.0	2	1.0	7	1.9	7	1.6	4	0.6	8	1.3	30	1.2	23	1.4
	110－115	11	0.4	0	0.0	0	0.0	0	0.0	1	0.5	2	0.6	4	1.1	4	0.9	0	0.0	0	0.0	11	0.4	8	0.5
	115－120	5	0.2	0	0.0	0	0.0	0	0.0	0	0.0	3	0.9	1	0.3	0	0.0	1	0.2	0	0.0	5	0.2	2	0.1
	120cm以上	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	1	0.2	0	0.0	2	0.1	2	0.1
	総数	3,609	100.0	107	100.0	75	100.0	130	100.0	223	100.0	433	100.0	465	100.0	543	100.0	730	100.0	797	100.0	3,191	100.0	2,049	100.0
	50cm未満	15	0.4	2	1.9	0	0.0	1	0.8	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	50－55cm未満	64	1.8	44	41.5	15	14.0	2	2.7	1	0.4	0	0.0	0	0.0	1	0.2	0	0.0	1	0.1	3	0.1	1	0.0
	55－60	112	3.1	32	30.2	34	31.8	9	6.9	8	3.6	7	1.6	4	0.9	1	0.2	4	0.5	2	0.3	26	0.8	10	0.5
	60－65	219	6.1	14	13.2	26	24.3	17	13.1	33	14.8	25	5.8	35	7.5	21	3.9	11	1.5	17	2.1	142	4.5	73	3.6
	65－70	361	10.0	3	2.8	14	13.1	20	26.7	54	24.2	60	13.9	64	13.8	49	9.0	42	5.8	30	3.8	299	9.4	165	8.1
	70－75	572	15.8	1	0.9	9	8.4	14	18.7	55	24.7	108	24.9	101	21.7	78	14.4	78	10.7	83	10.4	503	15.8	284	13.9
	75－80	596	16.5	0	0.0	4	3.7	7	9.3	41	18.4	98	22.6	97	20.9	111	20.4	122	16.7	99	12.4	568	17.8	372	18.2
	80－85	562	15.6	0	0.0	3	2.8	0	0.0	13	5.8	72	16.6	77	16.6	120	22.1	128	17.5	142	17.8	552	17.3	382	18.6
	85－90	465	12.9	0	0.0	0	0.0	5	3.8	9	4.0	31	7.2	40	8.6	71	13.1	143	19.6	165	20.7	459	14.4	325	15.9
	90－95	332	9.2	0	0.0	0	0.0	1	0.8	3	1.3	17	3.9	26	5.6	48	8.8	105	14.4	132	16.6	331	10.4	236	11.5
	95－100	167	4.6	0	0.0	0	0.0	0	0.0	1	0.4	9	2.1	12	2.6	24	4.4	51	7.0	70	8.8	167	5.2	111	5.4
	100－105	83	2.3	0	0.0	0	0.0	1	0.8	0	0.0	3	0.7	2	0.4	13	2.4	30	4.1	34	4.3	82	2.6	54	2.6
	105－110	30	0.8	0	0.0	0	0.0	0	0.0	3	1.3	0	0.0	4	0.9	2	0.4	7	1.0	14	1.8	30	0.9	16	0.8
	110－115	20	0.6	0	0.0	0	0.0	2	1.5	1	0.4	0	0.0	2	0.4	2	0.4	7	1.0	6	0.8	18	0.6	14	0.7
	115－120	7	0.2	0	0.0	0	0.0	0	0.0	1	0.4	1	0.2	1	0.2	2	0.4	1	0.1	1	0.1	7	0.2	5	0.2
	120cm以上	4	0.1	0	0.0	0	0.0	0	0.0	0	0.0	2	0.5	0	0.0	0	0.0	1	0.1	1	0.1	4	0.1	1	0.0

注）妊婦除外

(22年)

第 17 表 腹囲の区分（20 歳以上，性・年齢階級別）

	総 数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	5,779	100.0	415	100.0	783	100.0	840	100.0	978	100.0	1,353	100.0	1,410	100.0
	<85cm(男)/<90cm(女) ≥85cm(男)/≥90cm(女)	3,805 1,974	65.8 34.2	359 56	86.5 13.5	615 168	78.5 21.5	608 232	72.4 27.6	635 343	64.9 35.1	779 574	57.6 42.4	809 601	57.4 42.6
男性	総 数	2,588	100.0	192	100.0	350	100.0	375	100.0	435	100.0	623	100.0	613	100.0
	<85cm ≥85cm	1,253 1,335	48.4 51.6	145 47	75.5 24.5	214 136	61.1 38.9	190 185	50.7 49.3	183 252	42.1 57.9	251 372	40.3 59.7	270 343	44.0 56.0
女性	総 数	3,191	100.0	223	100.0	433	100.0	465	100.0	543	100.0	730	100.0	797	100.0
	<90cm ≥90cm	2,552 639	80.0 20.0	214 9	96.0 4.0	401 32	92.6 7.4	418 47	89.9 10.1	452 91	83.2 16.8	528 202	72.3 27.7	539 258	67.6 32.4

注) 妊婦除外

(22年)

第 18 表 BMI・腹囲の区分（20 歳以上，性・年齢階級別）

	総 数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	5,721	100.0	408	100.0	779	100.0	832	100.0	967	100.0	1,343	100.0	1,392	100.0
	BMI<25,腹囲<85cm(男)/90cm(女)	3,488	61.0	338	82.8	563	72.3	546	65.6	591	61.1	707	52.6	743	53.4
	BMIのみ超(BMI≥25)	285	5.0	15	3.7	49	6.3	57	6.9	42	4.3	68	5.1	54	3.9
	腹囲のみ超(腹囲≥85cm(男)/90cm(女))	766	13.4	14	3.4	54	6.9	71	8.5	110	11.4	252	18.8	265	19.0
男性	BMI≥25,腹囲≥85cm(男)/90cm(女)	1,182	20.7	41	10.0	113	14.5	158	19.0	224	23.2	316	23.5	330	23.7
	総 数	2,559	100.0	189	100.0	347	100.0	370	100.0	428	100.0	619	100.0	606	100.0
	BMI<25,腹囲<85cm	1,180	46.1	136	72.0	198	57.1	175	47.3	177	41.4	237	38.3	257	42.4
	BMIのみ超(BMI≥25)	61	2.4	7	3.7	14	4.0	13	3.5	6	1.4	12	1.9	9	1.5
女性	腹囲のみ超(腹囲≥85cm)	588	23.0	14	7.4	49	14.1	65	17.6	87	20.3	197	31.8	176	29.0
	BMI≥25,腹囲≥85cm	730	28.5	32	16.9	86	24.8	117	31.6	158	36.9	173	27.9	164	27.1
	総 数	3,162	100.0	219	100.0	432	100.0	462	100.0	539	100.0	724	100.0	786	100.0
	BMI<25,腹囲<90cm	2,308	73.0	202	92.2	365	84.5	371	80.3	414	76.8	470	64.9	486	61.8
妊婦除外	BMIのみ超(BMI≥25)	224	7.1	8	3.7	35	8.1	44	9.5	36	6.7	56	7.7	45	5.7
	腹囲のみ超(腹囲≥90cm)	178	5.6	0	0.0	5	1.2	6	1.3	23	4.3	55	7.6	89	11.3
	BMI≥25,腹囲≥90cm	452	14.3	9	4.1	27	6.3	41	8.9	66	12.2	143	19.8	166	21.1

注) 妊婦除外

(22年)

〔参考〕 内臓脂肪型肥満の診断基準：

- BMI 25 以上で，男性のウエスト周囲径 85cm 以上，女性のウエスト周囲径 90cm 以上を上半身肥満の疑いとする。
- 上半身肥満の疑いと判定され，腹部CT法による内臓脂肪面積 100cm²以上（男女とも）を内臓脂肪型肥満と診断する。（日本肥満学会肥満症診断基準検討委員会，2000 年）

※国民健康・栄養調査の「腹囲」は，「立位のへその高さ」で計測したが，計測位置は「ウエスト周囲径」と同じである。

第19表 薬の服用状況（性・年齢階級別）

		総数		20－29歳		30－39歳		40－49歳		50－59歳		60－69歳		70歳以上		
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
総数	総 数	4,647	100.0	289	100.0	595	100.0	602	100.0	786	100.0	1,160	100.0	1,215	100.0	
	使用なし	3,050	65.6	286	99.0	583	98.0	549	91.2	579	73.7	623	53.7	430	35.4	
	使用あり(一つ以上)	1,597	34.4	3	1.0	12	2.0	53	8.8	207	26.3	537	46.3	785	64.6	
	血圧を下げる薬	1,213	26.1	0	0.0	4	0.7	31	5.1	143	18.2	395	34.1	640	52.7	
	脈の乱れを治す薬	165	3.6	0	0.0	2	0.3	3	0.5	14	1.8	44	3.8	102	8.4	
	インスリン注射又は血糖を下げる薬	289	6.2	1	0.3	2	0.3	9	1.5	35	4.5	106	9.1	136	11.2	
	コレステロールを下げる薬	609	13.1	0	0.0	4	0.7	14	2.3	84	10.7	213	18.4	294	24.2	
	中性脂肪(トリグリセライド)を下げる薬	147	3.2	0	0.0	0	0.0	8	1.3	23	2.9	47	4.1	69	5.7	
	貧血治療のための薬(鉄剤)	68	1.5	2	0.7	3	0.5	6	1.0	10	1.3	10	0.9	37	3.0	
男性	総 数	1,977	100.0	128	100.0	219	100.0	241	100.0	327	100.0	533	100.0	529	100.0	
	使用なし	1,253	63.4	128	100.0	214	97.7	210	87.1	219	67.0	282	52.9	200	37.8	
	使用あり(一つ以上)	724	36.6	0	0.0	5	2.3	31	12.9	108	33.0	251	47.1	329	62.2	
	血圧を下げる薬	577	29.2	0	0.0	3	1.4	21	8.7	82	25.1	192	36.0	279	52.7	
	脈の乱れを治す薬	91	4.6	0	0.0	0	0.0	2	0.8	8	2.5	27	5.1	54	10.2	
	インスリン注射又は血糖を下げる薬	160	8.1	0	0.0	2	0.9	5	2.1	26	8.0	64	12.0	63	11.9	
	コレステロールを下げる薬	217	11.0	0	0.0	1	0.5	12	5.0	38	11.6	76	14.3	90	17.0	
	中性脂肪(トリグリセライド)を下げる薬	91	4.6	0	0.0	0	0.0	8	3.3	18	5.5	32	6.0	33	6.2	
	貧血治療のための薬(鉄剤)	20	1.0	0	0.0	0	0.0	1	0.4	2	0.6	4	0.8	13	2.5	
女性	総 数	2,670	100.0	161	100.0	376	100.0	361	100.0	459	100.0	627	100.0	686	100.0	
	使用なし	1,797	67.3	158	98.1	369	98.1	339	93.9	360	78.4	341	54.4	230	33.5	
	使用あり(一つ以上)	873	32.7	3	1.9	7	1.9	22	6.1	99	21.6	286	45.6	456	66.5	
	血圧を下げる薬	636	23.8	0	0.0	1	0.3	10	2.8	61	13.3	203	32.4	361	52.6	
	脈の乱れを治す薬	74	2.8	0	0.0	2	0.5	1	0.3	6	1.3	17	2.7	48	7.0	
	インスリン注射又は血糖を下げる薬	129	4.8	1	0.6	0	0.0	4	1.1	9	2.0	42	6.7	73	10.7	
	コレステロールを下げる薬	392	14.7	0	0.0	3	0.8	2	0.6	46	10.0	137	21.9	204	29.8	
	中性脂肪(トリグリセライド)を下げる薬	56	2.1	0	0.0	0	0.0	0	0.0	5	1.1	15	2.4	36	5.3	
	貧血治療のための薬(鉄剤)	48	1.8	2	1.3	3	0.8	5	1.4	8	1.7	6	1.0	24	3.5	
	〔再掲〕妊婦除外〕	総 数	2,650	100.0	155	100.0	363	100.0	360	100.0	459	100.0	627	100.0	686	100.0
		使用なし	1,779	67.1	154	99.4	356	98.1	338	93.9	360	78.4	341	54.4	230	33.5
		使用あり(一つ以上)	871	32.9	1	0.6	7	1.9	22	6.1	99	21.6	286	45.6	456	66.5
		血圧を下げる薬	636	24.0	0	0.0	1	0.3	10	2.8	61	13.3	203	32.4	361	52.6
		脈の乱れを治す薬	74	2.8	0	0.0	2	0.6	1	0.3	6	1.3	17	2.7	48	7.0
		インスリン注射又は血糖を下げる薬	129	4.9	1	0.6	0	0.0	4	1.1	9	2.0	42	6.7	73	10.7
		コレステロールを下げる薬	392	14.8	0	0.0	3	0.8	2	0.6	46	10.0	137	21.9	204	29.8
		中性脂肪(トリグリセライド)を下げる薬	56	2.1	0	0.0	0	0.0	0	0.0	5	1.1	15	2.4	36	5.3
		貧血治療のための薬(鉄剤)	46	1.7	0	0.0	3	0.8	5	1.4	8	1.7	6	1.0	24	3.5

* 複数回答

(22年)

第20表の1 血圧の状況（性・年齢階級別）

		総数		至適血圧		正常血圧		正常高値血圧		Ⅰ度高血圧		Ⅱ度高血圧		Ⅲ度高血圧		(再掲) 収縮期高血圧		
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%*	
総数	総 数	4,220	100.0	1,107	26.2	737	17.5	818	19.4	1,135	26.9	342	8.1	81	1.9	841	19.9	
	15-19歳	101	100.0	80	79.2	16	15.8	4	4.0	1	1.0	0	0.0	0	0.0	0	0.0	
	20-29歳	202	100.0	137	67.8	40	19.8	14	6.9	8	4.0	3	1.5	0	0.0	4	2.0	
	30-39歳	511	100.0	304	59.5	102	20.0	61	11.9	37	7.2	6	1.2	1	0.2	8	1.6	
	40-49歳	509	100.0	212	41.7	108	21.2	92	18.1	77	15.1	15	2.9	5	1.0	22	4.3	
	50-59歳	696	100.0	152	21.8	144	20.7	140	20.1	184	26.4	66	9.5	10	1.4	78	11.2	
	60-69歳	1,076	100.0	130	12.1	171	15.9	249	23.1	388	36.1	104	9.7	34	3.2	266	24.7	
	70歳以上	1,125	100.0	92	8.2	156	13.9	258	22.9	440	39.1	148	13.2	31	2.8	463	41.2	
	(再掲) 65-74歳	1,022	100.0	113	11.1	155	15.2	231	22.6	375	36.7	115	11.3	33	3.2	326	31.9	
	(再掲) 75歳以上	663	100.0	45	6.8	93	14.0	153	23.1	263	39.7	90	13.6	19	2.9	289	43.6	
男性	総 数	1,744	100.0	285	16.3	289	16.6	381	21.8	579	33.2	171	9.8	39	2.2	379	21.7	
	15-19歳	47	100.0	34	72.3	10	21.3	2	4.3	1	2.1	0	0.0	0	0.0	0	0.0	
	20-29歳	84	100.0	38	45.2	25	29.8	11	13.1	8	9.5	2	2.4	0	0.0	4	4.8	
	30-39歳	170	100.0	56	32.9	42	24.7	40	23.5	26	15.3	5	2.9	1	0.6	5	2.9	
	40-49歳	185	100.0	42	22.7	45	24.3	40	21.6	46	24.9	9	4.9	3	1.6	10	5.4	
	50-59歳	277	100.0	34	12.3	47	17.0	62	22.4	94	33.9	34	12.3	6	2.2	30	10.8	
	60-69歳	486	100.0	49	10.1	62	12.8	120	24.7	187	38.5	52	10.7	16	3.3	109	22.4	
	70歳以上	495	100.0	32	6.5	58	11.7	106	21.4	217	43.8	69	13.9	13	2.6	221	44.6	
	(再掲) 65-74歳	447	100.0	38	8.5	51	11.4	102	22.8	182	40.7	59	13.2	15	3.4	147	32.9	
	(再掲) 75歳以上	291	100.0	19	6.5	37	12.7	66	22.7	126	43.3	35	12.0	8	2.7	134	46.0	
女性	総 数	2,476	100.0	822	33.2	448	18.1	437	17.6	556	22.5	171	6.9	42	1.7	462	18.7	
	15-19歳	54	100.0	46	85.2	6	11.1	2	3.7	0	0.0	0	0.0	0	0.0	0	0.0	
	20-29歳	118	100.0	99	83.9	15	12.7	3	2.5	0	0.0	1	0.8	0	0.0	0	0.0	
	30-39歳	341	100.0	248	72.7	60	17.6	21	6.2	11	3.2	1	0.3	0	0.0	3	0.9	
	40-49歳	324	100.0	170	52.5	63	19.4	52	16.0	31	9.6	6	1.9	2	0.6	12	3.7	
	50-59歳	419	100.0	118	28.2	97	23.2	78	18.6	90	21.5	32	7.6	4	1.0	48	11.5	
	60-69歳	590	100.0	81	13.7	109	18.5	129	21.9	201	34.1	52	8.8	18	3.1	157	26.6	
	70歳以上	630	100.0	60	9.5	98	15.6	152	24.1	223	35.4	79	12.5	18	2.9	242	38.4	
	(再掲) 65-74歳	575	100.0	75	13.0	104	18.1	129	22.4	193	33.6	56	9.7	18	3.1	179	31.1	
	(再掲) 75歳以上	372	100.0	26	7.0	56	15.1	87	23.4	137	36.8	55	14.8	11	3.0	155	41.7	
	(再掲) 妊婦除外)	総 数**	2,457	100.0	500	20.4	137	5.6	75	3.1	42	1.7	8	0.3	2	0.1	15	2.0
		20-29歳	112	100.0	94	83.9	15	13.4	2	1.8	0	0.0	1	0.9	0	0.0	0	0.0
		30-39歳	329	100.0	237	72.0	59	17.9	21	6.4	11	3.3	1	0.3	0	0.0	3	0.9
		40-49歳	323	100.0	169	52.3	63	19.5	52	16.1	31	9.6	6	1.9	2	0.6	12	3.7

注) 日本高血圧学会による血圧の分類はp.16を参照。

(22年)

数値は2回の測定値の平均値を用いた。なお、1回しか測定できなかった者については、その値を採用した。

* 総数に対する割合

** 15歳以上の妊婦除外の総数

第 20 表の 2 血圧の状況（性・年齢階級別）〔服薬者除外〕

		総数		至適血圧		正常血圧		正常高値血圧		Ⅰ度高血圧		Ⅱ度高血圧		Ⅲ度高血圧		(再掲) 収縮期高血圧		
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%*	
総数	総 数	3,114	100.0	1,048	33.7	612	19.7	560	18.0	653	21.0	195	6.3	46	1.5	421	13.5	
	15-19歳	101	100.0	80	79.2	16	15.8	4	4.0	1	1.0	0	0.0	0	0.0	0	0.0	
	20-29歳	202	100.0	137	67.8	40	19.8	14	6.9	8	4.0	3	1.5	0	0.0	4	2.0	
	30-39歳	509	100.0	304	59.7	102	20.0	59	11.6	37	7.3	6	1.2	1	0.2	8	1.6	
	40-49歳	488	100.0	211	43.2	107	21.9	88	18.0	66	13.5	12	2.5	4	0.8	21	4.3	
	50-59歳	571	100.0	143	25.0	127	22.2	114	20.0	133	23.3	50	8.8	4	0.7	58	10.2	
	60-69歳	713	100.0	109	15.3	133	18.7	167	23.4	218	30.6	63	8.8	23	3.2	145	20.3	
	70歳以上	530	100.0	64	12.1	87	16.4	114	21.5	190	35.8	61	11.5	14	2.6	185	34.9	
	(再掲) 65-74歳	604	100.0	92	15.2	105	17.4	133	22.0	188	31.1	65	10.8	21	3.5	152	25.2	
	(再掲) 75歳以上	290	100.0	28	9.7	50	17.2	64	22.1	108	37.2	32	11.0	8	2.8	108	37.2	
男性	総 数	1,226	100.0	259	21.1	244	19.9	263	21.5	335	27.3	102	8.3	23	1.9	187	15.3	
	15-19歳	47	100.0	34	72.3	10	21.3	2	4.3	1	2.1	0	0.0	0	0.0	0	0.0	
	20-29歳	84	100.0	38	45.2	25	29.8	11	13.1	8	9.5	2	2.4	0	0.0	4	4.8	
	30-39歳	169	100.0	56	33.1	42	24.9	39	23.1	26	15.4	5	3.0	1	0.6	5	3.0	
	40-49歳	171	100.0	41	24.0	45	26.3	37	21.6	38	22.2	8	4.7	2	1.2	9	5.3	
	50-59歳	207	100.0	31	15.0	39	18.8	47	22.7	65	31.4	22	10.6	3	1.4	21	10.1	
	60-69歳	316	100.0	39	12.3	53	16.8	81	25.6	103	32.6	30	9.5	10	3.2	56	17.7	
	70歳以上	232	100.0	20	8.6	30	12.9	46	19.8	94	40.5	35	15.1	7	3.0	92	39.7	
	(再掲) 65-74歳	247	100.0	29	11.7	35	14.2	53	21.5	88	35.6	32	13.0	10	4.0	66	26.7	
	(再掲) 75歳以上	137	100.0	11	8.0	19	13.9	32	23.4	54	39.4	18	13.1	3	2.2	53	38.7	
女性	総 数	1,888	100.0	789	41.8	368	19.5	297	15.7	318	16.8	93	4.9	23	1.2	234	12.4	
	15-19歳	54	100.0	46	85.2	6	11.1	2	3.7	0	0.0	0	0.0	0	0.0	0	0.0	
	20-29歳	118	100.0	99	83.9	15	12.7	3	2.5	0	0.0	1	0.8	0	0.0	0	0.0	
	30-39歳	340	100.0	248	72.9	60	17.6	20	5.9	11	3.2	1	0.3	0	0.0	3	0.9	
	40-49歳	317	100.0	170	53.6	62	19.6	51	16.1	28	8.8	4	1.3	2	0.6	12	3.8	
	50-59歳	364	100.0	112	30.8	88	24.2	67	18.4	68	18.7	28	7.7	1	0.3	37	10.2	
	60-69歳	397	100.0	70	17.6	80	20.2	86	21.7	115	29.0	33	8.3	13	3.3	89	22.4	
	70歳以上	298	100.0	44	14.8	57	19.1	68	22.8	96	32.2	26	8.7	7	2.3	93	31.2	
	(再掲) 65-74歳	357	100.0	63	17.6	70	19.6	80	22.4	100	28.0	33	9.2	11	3.1	86	24.1	
	(再掲) 75歳以上	153	100.0	17	11.1	31	20.3	32	20.9	54	35.3	14	9.2	5	3.3	55	35.9	
	(再掲) 妊婦除外	総 数**	1,869	100.0	500	26.8	136	7.3	73	3.9	39	2.1	6	0.3	2	0.1	15	0.8
		20-29歳	112	100.0	94	83.9	15	13.4	2	1.8	0	0.0	1	0.9	0	0.0	0	0.0
		30-39歳	328	100.0	237	72.3	59	18.0	20	6.1	11	3.4	1	0.3	0	0.0	3	0.9
		40-49歳	316	100.0	169	53.5	62	19.6	51	16.1	28	8.9	4	1.3	2	0.6	12	3.8

注) 血圧を下げる薬服用者除外。日本高血圧学会による血圧の分類は p.16 を参照。

(22年)

数値は 2 回の測定値の平均値を用いた。なお、1 回しか測定できなかった者については、その値を採用した。

* 総数に対する割合

** 15 歳以上の妊婦除外の総数

第21表の1 収縮期（最高）・拡張期（最低）血圧の分布（性・年齢階級別）

			総 数		15～19歳		20～29歳		30～39歳		40～49歳		50～59歳		60～69歳		70歳以上		(再掲) 40～89歳	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	最高血圧	総 数	1,744	100.0	47	100.0	84	100.0	170	100.0	185	100.0	277	100.0	486	100.0	495	100.0	1,435	100.0
		90mmHg未満	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.1
		90～99	16	0.9	6	12.8	1	1.2	3	1.8	1	0.5	2	0.7	2	0.4	1	0.2	6	0.4
		100～109	100	5.7	15	31.9	18	21.4	19	11.2	20	10.8	10	3.6	11	2.3	7	1.4	47	3.3
		110～119	219	12.6	16	34.0	22	26.2	47	27.6	34	18.4	31	11.2	41	8.4	28	5.7	134	9.3
		120～129	351	20.1	8	17.0	27	32.1	54	31.8	59	31.9	60	21.7	78	16.0	65	13.1	261	18.2
		130～139	380	21.8	2	4.3	9	10.7	31	18.2	37	20.0	74	26.7	124	25.5	103	20.8	337	23.5
		140～149	327	18.8	0	0.0	7	8.3	10	5.9	19	10.3	45	16.2	116	23.9	130	26.3	309	21.5
		150～159	197	11.3	0	0.0	0	0.0	4	2.4	12	6.5	38	13.7	60	12.3	83	16.8	192	13.4
		160～169	77	4.4	0	0.0	0	0.0	1	0.6	1	0.5	9	3.2	20	4.1	46	9.3	74	5.2
		170～179	45	2.6	0	0.0	0	0.0	1	0.6	1	0.5	5	1.8	20	4.1	18	3.6	44	3.1
		180mmHg以上	31	1.8	0	0.0	0	0.0	0	0.0	0	0.0	1	0.5	3	1.1	14	2.9	13	2.6
		平均値 mmHg	135.3		110.7		120.1		123.1		127.2		135.4		139.7		142.9		138.3	
		標準偏差	18.2		9.6		11.3		12.5		14.4		16.2		17.9		17.3		17.6	
	最低血圧	総 数	1,744	100.0	47	100.0	84	100.0	170	100.0	185	100.0	277	100.0	486	100.0	495	100.0	1,435	100.0
		40mmHg未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		40～49	3	0.2	2	4.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.1
		50～59	35	2.0	9	19.1	2	2.4	2	1.2	1	0.5	0	0.0	5	1.0	16	3.2	22	1.5
		60～69	188	10.8	16	34.0	21	25.0	25	14.7	12	6.5	11	4.0	30	6.2	73	14.7	125	8.7
		70～79	533	30.6	16	34.0	31	36.9	65	38.2	57	30.8	70	25.3	141	29.0	153	30.9	420	29.3
		80～89	575	33.0	3	6.4	24	28.6	51	30.0	67	36.2	92	33.2	164	33.7	174	35.2	492	34.3
		90～99	309	17.7	1	2.1	4	4.8	22	12.9	37	20.0	70	25.3	112	23.0	63	12.7	281	19.6
		100～109	87	5.0	0	0.0	2	2.4	4	2.4	8	4.3	30	10.8	30	6.2	13	2.6	81	5.6
		110～119	10	0.6	0	0.0	0	0.0	0	0.0	2	1.1	3	1.1	3	0.6	2	0.4	10	0.7
120～129	4	0.2	0	0.0	0	0.0	1	0.6	1	0.5	1	0.4	1	0.2	0	0.0	3	0.2		
130～139	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0		
140mmHg以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0		
平均値 mmHg	81.6		66.7		76.0		79.7		83.6		86.3		83.8		79.0		82.6			
標準偏差	11.2		9.4		9.7		10.5		10.2		10.9		10.4		10.7		10.9			
女性	最高血圧	総 数	2,476	100.0	54	100.0	118	100.0	341	100.0	324	100.0	419	100.0	590	100.0	630	100.0	1,952	100.0
		90mmHg未満	12	0.5	1	1.9	3	2.5	6	1.8	1	0.3	0	0.0	0	0.0	1	0.2	2	0.1
		90～99	87	3.5	8	14.8	12	10.2	39	11.4	14	4.3	12	2.9	2	0.3	0	0.0	28	1.4
		100～109	355	14.3	20	37.0	55	46.6	118	34.6	86	26.5	37	8.8	20	3.4	19	3.0	162	8.3
		110～119	438	17.7	21	38.9	35	29.7	97	28.4	88	27.2	80	19.1	70	11.9	47	7.5	284	14.5
		120～129	466	18.8	3	5.6	10	8.5	57	16.7	68	21.0	103	24.6	121	20.5	104	16.5	394	20.2
		130～139	409	16.5	1	1.9	3	2.5	15	4.4	36	11.1	76	18.1	125	21.2	153	24.3	388	19.9
		140～149	314	12.7	0	0.0	0	0.0	7	2.1	21	6.5	55	13.1	116	19.7	115	18.3	306	15.7
		150～159	208	8.4	0	0.0	0	0.0	2	0.6	4	1.2	32	7.6	72	12.2	98	15.6	204	10.5
		160～169	112	4.5	0	0.0	0	0.0	0	0.0	2	0.6	16	3.8	33	5.6	61	9.7	110	5.6
		170～179	39	1.6	0	0.0	0	0.0	0	0.0	2	0.6	5	1.2	17	2.9	15	2.4	38	1.9
		180mmHg以上	36	1.5	0	0.0	0	0.0	0	0.0	2	0.6	3	0.7	14	2.4	17	2.7	36	1.8
		平均値 mmHg	128.7		107.2		107.8		111.5		118.6		129.0		137.6		140.4		133.4	
		標準偏差	20.3		8.7		9.3		11.9		15.9		17.6		18.3		18.2		19.4	
	最低血圧	総 数	2,476	100.0	54	100.0	118	100.0	341	100.0	324	100.0	419	100.0	590	100.0	630	100.0	1,952	100.0
		40mmHg未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		40～49	12	0.5	3	5.6	2	1.7	2	0.6	0	0.0	0	0.0	0	0.0	5	0.8	5	0.3
		50～59	99	4.0	12	22.2	16	13.6	33	9.7	11	3.4	6	1.4	3	0.5	18	2.9	37	1.9
		60～69	510	20.6	24	44.4	54	45.8	117	34.3	82	25.3	54	12.9	67	11.4	112	17.8	311	15.9
		70～79	891	36.0	11	20.4	37	31.4	129	37.8	127	39.2	150	35.8	209	35.4	228	36.2	710	36.4
		80～89	657	26.5	4	7.4	8	6.8	51	15.0	77	23.8	131	31.3	197	33.4	189	30.0	592	30.3
		90～99	237	9.6	0	0.0	0	0.0	8	2.3	20	6.2	62	14.8	86	14.6	61	9.7	229	11.7
		100～109	56	2.3	0	0.0	1	0.8	1	0.3	5	1.5	15	3.6	19	3.2	15	2.4	54	2.8
		110～119	9	0.4	0	0.0	0	0.0	0	0.0	2	0.6	1	0.2	5	0.8	1	0.2	9	0.5
120～129	4	0.2	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	3	0.5	1	0.2	4	0.2		
130～139	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0		
140mmHg以上	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.1		
平均値 mmHg	76.8		64.5		67.1		70.6		75.7		80.0		81.3		77.3		78.9			
標準偏差	11.2		8.8		8.4		9.3		10.2		10.4		10.8		10.6		10.7			
〔再掲〕 妊婦除外	最高血圧	総 数	2,457	100.0	54	100.0	112	100.0	329	100.0	323	100.0	419	100.0	590	100.0	630	100.0	1,951	100.0
		90mmHg未満	12	0.5	1	1.9	3	2.7	6	1.8	1	0.3	0	0.0	0	0.0	1	0.2	2	0.1
		90～99	84	3.4	8	14.8	11	9.8	37	11.2	14	4.3	12	2.9	2	0.3	0	0.0	28	1.4
		100～109	348	14.2	20	37.0	53	47.3	114	34.7	85	26.3	37	8.8	20	3.4	19	3.0	161	8.3
		110～119	431	17.5	21	38.9	33	29.5	92	28.0	88	27.2	80	19.1	70	11.9	47	7.5	284	14.6
		120～129	465	18.9	3	5.6	10	8.9	56	17.0	68	21.1	103	24.6	121	20.5	104	16.5	394	20.2
		130～139	408	16.6	1	1.9	2	1.8	15	4.6	36	11.1	76	18.1	125	21.2	153	24.3	388	19.9
		140～149	314	12.8	0	0.0	0	0.0	7	2.1	21	6.5	55	13.1	116	19.7	115	18.3	306	15.7
		150～159	208	8.5	0	0.0	0	0.0	2	0.6	4	1.2	32	7.6	72	12.2	98	15.6	204	10.5
		160～169	112	4.6	0	0.0	0	0.0	0	0.0	2	0.6	16	3.8	33	5.6	61	9.7	110	5.6
		170～179	39	1.6	0	0.0	0	0.0	0	0.0	2	0.6	5	1.2	17	2.9	15	2.4	38	1.9
		180mmHg以上	36	1.5	0	0.0	0	0.0	0	0.0	2	0.6	3	0.7	14	2.4	17	2.7	36	1.8
		平均値 mmHg	128.8		107.2		10													

第 21 表の 2 収縮期（最高）・拡張期（最低）血圧の分布（性・年齢階級別）〔服薬者除外〕

			総 数		15－19歳		20－29歳		30－39歳		40－49歳		50－59歳		60－69歳		70歳以上		(再掲) 40～89歳	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	最高血圧	総 数	1,226	100.0	47	100.0	84	100.0	169	100.0	171	100.0	207	100.0	316	100.0	232	100.0	923	100.0
		90mmHg未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		90－99	16	1.3	6	12.8	1	1.2	3	1.8	1	0.6	2	1.0	2	0.6	1	0.4	6	0.7
		100－109	98	8.0	15	31.9	18	21.4	19	11.2	19	11.1	10	4.8	11	3.5	6	2.6	46	5.0
		110－119	190	15.5	16	34.0	22	26.2	46	27.2	34	19.9	27	13.0	30	9.5	15	6.5	106	11.5
		120－129	290	23.7	8	17.0	27	32.1	54	32.0	56	32.7	49	23.7	64	20.3	32	13.8	201	21.8
		130－139	256	20.9	2	4.3	9	10.7	31	18.3	34	19.9	52	25.1	82	25.9	46	19.8	213	23.1
		140－149	184	15.0	0	0.0	7	8.3	10	5.9	15	8.8	34	16.4	64	20.3	54	23.3	167	18.1
		150－159	105	8.6	0	0.0	0	0.0	4	2.4	10	5.8	23	11.1	30	9.5	38	16.4	100	10.8
		160－169	40	3.3	0	0.0	0	0.0	1	0.6	1	0.6	5	2.4	10	3.2	23	9.9	39	4.2
		170－179	30	2.4	0	0.0	0	0.0	0	0.0	1	0.6	3	1.4	15	4.7	10	4.3	29	3.1
		180mmHg以上	17	1.4	0	0.0	0	0.0	0	0.0	0	0.0	2	1.0	8	2.5	7	3.0	16	1.7
		平均値 mmHg	132.0		110.7		120.1		123.2		126.1		133.3		137.4		142.9		135.7	
		標準偏差	18.2		9.6		11.3		12.5		13.5		16.1		18.4		19.0		18.1	
	最低血圧	総 数	1,226	100.0	47	100.0	84	100.0	169	100.0	171	100.0	207	100.0	316	100.0	232	100.0	923	100.0
		40mmHg未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		40－49	2	0.2	2	4.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		50－59	21	1.7	9	19.1	2	2.4	2	1.2	1	0.6	0	0.0	3	0.9	4	1.7	8	0.9
		60－69	138	11.3	16	34.0	21	25.0	25	14.8	11	6.4	11	5.3	19	6.0	35	15.1	76	8.2
		70－79	386	31.5	16	34.0	31	36.9	65	38.5	56	32.7	61	29.5	94	29.7	63	27.2	274	29.7
		80－89	406	33.1	3	6.4	24	28.6	50	29.6	64	37.4	66	31.9	113	35.8	86	37.1	326	35.3
		90－99	205	16.7	1	2.1	4	4.8	22	13.0	30	17.5	47	22.7	67	21.2	34	14.7	178	19.3
		100－109	58	4.7	0	0.0	2	2.4	4	2.4	7	4.1	20	9.7	17	5.4	8	3.4	52	5.6
		110－119	7	0.6	0	0.0	0	0.0	0	0.0	2	1.2	1	0.5	2	0.6	2	0.9	7	0.8
女性	最高血圧	120－129	3	0.2	0	0.0	0	0.0	1	0.6	0	0.0	1	0.5	1	0.3	0	0.0	2	0.2
		130－139	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		140mmHg以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		平均値 mmHg	81.4		66.7		76.0		79.6		82.9		85.2		83.4		80.2		82.9	
		標準偏差	11.0		9.4		9.7		10.5		9.7		11.0		10.3		10.6		10.6	
	最低血圧	総 数	1,888	100.0	54	100.0	118	100.0	340	100.0	317	100.0	364	100.0	397	100.0	298	100.0	1,374	100.0
		90mmHg未満	12	0.6	1	1.9	3	2.5	6	1.8	1	0.3	0	0.0	0	0.0	1	0.3	2	0.1
		90－99	86	4.6	8	14.8	12	10.2	39	11.5	14	4.4	11	3.0	2	0.5	0	0.0	27	2.0
		100－109	352	18.6	20	37.0	55	46.6	118	34.7	86	27.1	37	10.2	19	4.8	17	5.7	159	11.6
		110－119	405	21.5	21	38.9	35	29.7	97	28.5	88	27.8	75	20.6	59	14.9	30	10.1	252	18.3
		120－129	375	19.9	3	5.6	10	8.5	56	16.5	67	21.1	91	25.0	88	22.2	60	20.1	305	22.2
		130－139	267	14.1	1	1.9	3	2.5	15	4.4	34	10.7	66	18.1	81	20.4	67	22.5	247	18.0
		140－149	178	9.4	0	0.0	0	0.0	7	2.1	20	6.3	42	11.5	66	16.6	43	14.4	171	12.4
		150－159	116	6.1	0	0.0	0	0.0	2	0.6	3	0.9	23	6.3	39	9.8	49	16.4	114	8.3
		160－169	55	2.9	0	0.0	0	0.0	0	0.0	1	0.3	14	3.8	20	5.0	20	6.7	55	4.0
女性	最高血圧	170－179	22	1.2	0	0.0	0	0.0	0	0.0	1	0.3	5	1.4	11	2.8	5	1.7	22	1.6
		180mmHg以上	20	1.1	0	0.0	0	0.0	0	0.0	2	0.6	0	0.0	12	3.0	6	2.0	20	1.5
		平均値 mmHg	124.4		107.2		107.8		111.4		117.9		127.4		135.7		137.0		129.7	
		標準偏差	19.5		8.7		9.3		11.9		15.3		17.0		19.4		18.8		19.2	
	最低血圧	総 数	1,888	100.0	54	100.0	118	100.0	340	100.0	317	100.0	364	100.0	397	100.0	298	100.0	1,374	100.0
		40mmHg未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		40－49	8	0.4	3	5.6	2	1.7	2	0.6	0	0.0	0	0.0	0	0.0	1	0.3	1	0.1
		50－59	88	4.7	12	22.2	16	13.6	33	9.7	11	3.5	6	1.6	2	0.5	8	2.7	27	2.0
		60－69	423	22.4	24	44.4	54	45.8	117	34.4	81	25.6	50	13.7	51	12.8	46	15.4	227	16.5
		70－79	698	37.0	11	20.4	37	31.4	129	37.9	126	39.7	138	37.9	137	34.5	120	40.3	520	37.8
		80－89	471	24.9	4	7.4	8	6.8	50	14.7	77	24.3	110	30.2	135	34.0	87	29.2	409	29.8
		90－99	154	8.2	0	0.0	0	0.0	8	2.4	16	5.0	48	13.2	55	13.9	27	9.1	146	10.6
		100－109	35	1.9	0	0.0	1	0.8	1	0.3	4	1.3	11	3.0	11	2.8	7	2.3	33	2.4
		110－119	7	0.4	0	0.0	0	0.0	0	0.0	2	0.6	1	0.3	3	0.8	1	0.3	7	0.5
	女性	最高血圧	120－129	3	0.2	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.5	1	0.3	3
130－139			0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
140mmHg以上			1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	1	0.1
平均値 mmHg			76.0		64.5		67.1		70.5		75.4		79.2		81.0		77.7		78.5	
標準偏差			11.1		8.8		8.4		9.3		10.1		10.3		11.0		10.4		10.6	
最低血圧		総 数	1,869	100.0	54	100.0	112	100.0	328	100.0	316	100.0	364	100.0	397	100.0	298	100.0	1,373	100.0
		40mmHg未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		40－49	8	0.4	3	5.6	2	1.8	2	0.6	0	0.0	0	0.0	0	0.0	1	0.3	1	0.1
		50－59	84	4.5	12	22.2	15	13.4	30	9.1	11	3.5	6	1.6	2	0.5	8	2.7	27	2.0
		60－69	413	22.1	24	44.4	50	44.6	112	34.1	80	25.3	50	13.7	51	12.8	46	15.4	226	16.5
		70－79	694	37.1	11	20.4	36	32.1	126	38.4	126	39.9	138	37.9	137	34.5	120	40.3	520	37.9
		80－89	470	25.1	4	7.4	8	7.1	49	14.9	77	24.4	110	30.2	135	34.0	87	29.2	409	29.8
		90－99	154	8.2	0	0.0	0	0.0	8	2.4	16	5.1	48	13.2	55	13.9	27	9.1	146	10.6
		100－109	35	1.9	0	0.0	1	0.9	1	0.3	4	1.3	11	3.0	11	2.8	7	2.3	33	2.4
		110－119	7	0.4	0	0.0	0	0.0	0	0.0	2	0.6	1	0.3	3	0.8	1	0.3	7	0.5
女性	最高血圧	120－129	3	0.2	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.5	1	0.3	3	0.2
		130－139	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		140mmHg以上	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0		

第22表 血色素量の分布（性・年齢階級別）〔服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,570	100.0	75	100.0	161	100.0	174	100.0	257	100.0	458	100.0	445	100.0	412	100.0	261	100.0
	10.0g/dl未満	6	0.4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	6	1.3	2	0.5	4	1.5
	10.0-10.9	19	1.2	0	0.0	0	0.0	2	1.1	0	0.0	4	0.9	13	2.9	6	1.5	9	3.4
	11.0-11.9	35	2.2	0	0.0	0	0.0	0	0.0	1	0.4	4	0.9	30	6.7	7	1.7	25	9.6
	12.0-12.9	93	5.9	1	1.3	0	0.0	8	4.6	8	3.1	22	4.8	54	12.1	34	8.3	36	13.8
	13.0-13.9	293	18.7	0	0.0	11	6.8	14	8.0	35	13.6	102	22.3	131	29.4	105	25.5	79	30.3
	14.0-14.9	459	29.2	15	20.0	50	31.1	52	29.9	67	26.1	144	31.4	131	29.4	125	30.3	75	28.7
	15.0-15.9	410	26.1	34	45.3	54	33.5	58	33.3	93	36.2	120	26.2	51	11.5	83	20.1	25	9.6
	16.0-16.9	206	13.1	21	28.0	37	23.0	34	19.5	46	17.9	45	9.8	23	5.2	38	9.2	8	3.1
	17.0-17.9	37	2.4	4	5.3	7	4.3	5	2.9	5	1.9	12	2.6	4	0.9	10	2.4	0	0.0
	18.0g/dl以上	12	0.8	0	0.0	2	1.2	1	0.6	2	0.8	5	1.1	2	0.4	2	0.5	0	0.0
平均値(g/dl)		14.6		15.7		15.3		15.1		15.0		14.7		13.8		14.4		13.5	
標準偏差		1.4		0.8		1.0		1.2		1.1		1.3		1.5		1.4		1.4	
女性	総 数	2,220	100.0	103	100.0	315	100.0	302	100.0	385	100.0	558	100.0	557	100.0	534	100.0	321	100.0
	10.0g/dl未満	49	2.2	1	1.0	11	3.5	25	8.3	4	1.0	2	0.4	6	1.1	4	0.7	4	1.2
	10.0-10.9	63	2.8	0	0.0	14	4.4	18	6.0	7	1.8	4	0.7	20	3.6	8	1.5	15	4.7
	11.0-11.9	222	10.0	6	5.8	35	11.1	33	10.9	26	6.8	30	5.4	92	16.5	54	10.1	61	19.0
	12.0-12.9	689	31.0	24	23.3	95	30.2	94	31.1	115	29.9	158	28.3	203	36.4	162	30.3	124	38.6
	13.0-13.9	794	35.8	47	45.6	118	37.5	93	30.8	152	39.5	224	40.1	160	28.7	198	37.1	75	23.4
	14.0-14.9	334	15.0	22	21.4	39	12.4	36	11.9	62	16.1	112	20.1	63	11.3	87	16.3	37	11.5
	15.0-15.9	57	2.6	2	1.9	3	1.0	3	1.0	15	3.9	22	3.9	12	2.2	17	3.2	4	1.2
	16.0-16.9	9	0.4	1	1.0	0	0.0	0	0.0	4	1.0	4	0.7	0	0.0	2	0.4	0	0.0
	17.0-17.9	3	0.1	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	1	0.2	2	0.4	1	0.3
	18.0g/dl以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
平均値(g/dl)		13.0		13.3		12.8		12.5		13.2		13.3		12.8		13.1		12.7	
標準偏差		1.2		0.9		1.2		1.6		1.1		1.0		1.1		1.1		1.1	

注) 貧血治療のための薬服用者除外。

(22年)

第23表 赤血球数の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,584	100.0	75	100.0	161	100.0	174	100.0	257	100.0	461	100.0	456	100.0	418	100.0	268	100.0
	400万個/mm ³ 未満	121	7.6	0	0.0	0	0.0	1	0.6	4	1.6	32	6.9	84	18.4	43	10.3	59	22.0
	400-449	354	22.3	2	2.7	8	5.0	27	15.5	38	14.8	117	25.4	162	35.5	117	28.0	108	40.3
	450-499	661	41.7	14	18.7	62	38.5	66	37.9	140	54.5	210	45.6	169	37.1	189	45.2	85	31.7
	500-449	385	24.3	49	65.3	73	45.3	69	39.7	68	26.5	91	19.7	35	7.7	60	14.4	15	5.6
	550-599	59	3.7	10	13.3	18	11.2	11	6.3	6	2.3	10	2.2	4	0.9	7	1.7	1	0.4
	600-649	3	0.2	0	0.0	0	0.0	0	0.0	1	0.4	1	0.2	1	0.2	1	0.2	0	0.0
	650万個/mm ³ 以上	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	0	0.0
平均値(g/dl)		471.1		518.5		504.7		492.1		482.3		466.7		441.6		459.6		431.9	
標準偏差		49.4		30.2		35.1		40.3		36.2		42.6		53.3		49.5		47.3	
女性	総 数	2,261	100.0	105	100.0	318	100.0	307	100.0	391	100.0	563	100.0	577	100.0	545	100.0	333	100.0
	400万個/mm ³ 未満	320	14.2	3	2.9	23	7.2	36	11.7	39	10.0	59	10.5	160	27.7	94	17.2	106	31.8
	400-449	1,144	50.6	54	51.4	176	55.3	147	47.9	194	49.6	286	50.8	287	49.7	271	49.7	163	48.9
	450-499	726	32.1	43	41.0	112	35.2	116	37.8	141	36.1	189	33.6	125	21.7	164	30.1	61	18.3
	500-449	67	3.0	5	4.8	7	2.2	7	2.3	15	3.8	28	5.0	5	0.9	15	2.8	3	0.9
	550-599	3	0.1	0	0.0	0	0.0	1	0.3	2	0.5	0	0.0	0	0.0	0	0.0	0	0.0
	600-649	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.2	0	0.0
	650万個/mm ³ 以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
平均値(g/dl)		435.7		447.2		441.8		437.8		441.4		440.4		420.5		432.9		416.0	
標準偏差		35.2		30.9		29.3		32.9		35.2		34.3		36.4		35.4		37.7	

(22年)

第24表 ヘマトクリット値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,584	100.0	75	100.0	161	100.0	174	100.0	257	100.0	461	100.0	456	100.0	418	100.0	268	100.0
	25.0%未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	25.0-29.9	4	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	4	0.9	1	0.2	3	1.1
	30.0-34.9	28	1.8	0	0.0	0	0.0	2	1.1	0	0.0	3	0.7	23	5.0	9	2.2	15	5.6
	35.0-39.9	144	9.1	0	0.0	0	0.0	6	3.4	10	3.9	37	8.0	91	20.0	52	12.4	62	23.1
	40.0-44.9	673	42.5	14	18.7	53	32.9	59	33.9	100	38.9	220	47.7	227	49.8	202	48.3	138	51.5
	45.0-49.9	639	40.3	52	69.3	95	59.0	90	51.7	138	53.7	168	36.4	96	21.1	124	29.7	48	17.9
	50.0-54.9	90	5.7	9	12.0	12	7.5	17	9.8	8	3.1	31	6.7	13	2.9	28	6.7	2	0.7
	55.0%以上	6	0.4	0	0.0	1	0.6	0	0.0	1	0.4	2	0.4	2	0.4	2	0.5	0	0.0
	(再掲)39.8未満	160	10.1	0	0.0	0	0.0	7	4.0	8	3.1	34	7.4	111	24.3	56	13.4	77	28.7
	(再掲)51.9以上	33	2.1	4	5.3	3	1.9	6	3.4	3	1.2	11	2.4	6	1.3	8	1.9	1	0.4
平均値(g/dl)		44.4		47.0		46.3		45.8		45.5		44.6		42.2		43.8		41.4	
標準偏差		3.9		2.6		2.8		3.4		2.9		3.6		4.4		4.1		4.2	
女性	総 数	2,261	100.0	105	100.0	318	100.0	307	100.0	391	100.0	563	100.0	577	100.0	545	100.0	333	100.0
	25.0%未満	1	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	25.0-29.9	13	0.6	0	0.0	1	0.3	9	2.9	1	0.3	0	0.0	2	0.3	1	0.2	1	0.3
	30.0-34.9	113	5.0	3	2.9	25	7.9	30	9.8	11	2.8	8	1.4	36	6.2	15	2.8	27	8.1
	35.0-39.9	878	38.8	31	29.5	127	39.9	130	42.3	145	37.1	173	30.7	272	47.1	203	37.2	167	50.2
	40.0-44.9	1,113	49.2	69	65.7	157	49.4	126	41.0	199	50.9	324	57.5	238	41.2	286	52.5	119	35.7
	45.0-49.9	135	6.0	2	1.9	7	2.2	11	3.6	33	8.4	54	9.6	28	4.9	37	6.8	18	5.4
	50.0-54.9	7	0.3	0	0.0	1	0.3	0	0.0	2	0.5	3	0.5	1	0.2	2	0.4	1	0.3
	55.0%以上	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.2	0	0.0
	(再掲)33.4未満	69	3.1	1	1.0	10	3.1	29	9.4	6	1.5	5	0.9	18	3.1	8	1.5	14	4.2
	(再掲)45.0以上	143	6.3	2	1.9	8	2.5	11	3.6	35	9.0	58	10.3	29	5.0	40	7.3	19	5.7
平均値(g/dl)		40.3		40.7		39.9		39.1		40.8		41.3		39.8		40.7		39.5	
標準偏差		3.3		2.6		3.1		3.9		3.2		3.0		3.2		3.1		3.4	

(22年)

第25表 平均赤血球容積（MCV）の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,584	100.0	75	100.0	161	100.0	174	100.0	257	100.0	461	100.0	456	100.0	418	100.0	268	100.0
	70fl未満	3	0.2	0	0.0	0	0.0	1	0.6	0	0.0	0	0.0	2	0.4	1	0.2	1	0.4
	70-75未満	5	0.3	0	0.0	0	0.0	1	0.6	0	0.0	1	0.2	3	0.7	3	0.7	1	0.4
	75-80未満	6	0.4	1	1.3	1	0.6	0	0.0	1	0.4	1	0.2	2	0.4	1	0.2	2	0.7
	80-85未満	31	2.0	3	4.0	6	3.7	6	3.4	4	1.6	4	0.9	8	1.8	4	1.0	6	2.2
	85-90未満	225	14.2	27	36.0	48	29.8	35	20.1	28	10.9	46	10.0	41	9.0	42	10.0	20	7.5
	90-95未満	581	36.7	33	44.0	67	41.6	68	39.1	108	42.0	168	36.4	137	30.0	134	32.1	78	29.1
	95-100未満	499	31.5	11	14.7	32	19.9	46	26.4	88	34.2	160	34.7	162	35.5	162	38.8	94	35.1
	100-105未満	188	11.9	0	0.0	6	3.7	15	8.6	25	9.7	57	12.4	85	18.6	56	13.4	56	20.9
	105-110未満	33	2.1	0	0.0	0	0.0	1	0.6	2	0.8	19	4.1	11	2.4	11	2.6	7	2.6
	110-115未満	7	0.4	0	0.0	1	0.6	1	0.6	1	0.4	3	0.7	1	0.2	1	0.2	1	0.4
	115-120未満	4	0.3	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	3	0.7	2	0.5	2	0.7
	120fl以上	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	1	0.2	0	0.0
平均値(fl)		94.7		90.8		91.9		93.3		94.5		95.7		95.8		95.6		96.1	
標準偏差		5.5		3.9		4.6		5.4		4.4		5.4		6.1		5.6		6.0	
女性	総 数	2,261	100.0	105	100.0	318	100.0	307	100.0	391	100.0	563	100.0	577	100.0	545	100.0	333	100.0
	70fl未満	25	1.1	1	1.0	5	1.6	16	5.2	2	0.5	0	0.0	1	0.2	1	0.2	0	0.0
	70-75未満	15	0.7	0	0.0	6	1.9	5	1.6	4	1.0	0	0.0	0	0.0	0	0.0	0	0.0
	75-80未満	32	1.4	0	0.0	10	3.1	14	4.6	2	0.5	3	0.5	3	0.5	2	0.4	3	0.9
	80-85未満	78	3.4	8	7.6	21	6.6	31	10.1	7	1.8	6	1.1	5	0.9	6	1.1	2	0.6
	85-90未満	385	17.0	33	31.4	84	26.4	60	19.5	85	21.7	73	13.0	50	8.7	56	10.3	27	8.1
	90-95未満	1,001	44.3	43	41.0	138	43.4	117	38.1	172	44.0	286	50.8	245	42.5	270	49.5	131	39.3
	95-100未満	574	25.4	18	17.1	45	14.2	52	16.9	100	25.6	147	26.1	212	36.7	164	30.1	126	37.8
	100-105未満	131	5.8	1	1.0	8	2.5	9	2.9	16	4.1	43	7.6	54	9.4	41	7.5	38	11.4
	105-110未満	18	0.8	1	1.0	1	0.3	3	1.0	3	0.8	5	0.9	5	0.9	5	0.9	4	1.2
	110-115未満	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.3	0	0.0	2	0.6
	115-120未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	120fl以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
平均値(fl)		92.7		91.2		90.4		89.4		92.6		94.0		94.8		94.2		95.2	
標準偏差		5.8		4.7		6.2		8.1		5.2		4.2		4.5		4.3		4.6	

(22年)

第26表 平均赤血球血色素量（MCH）の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,584	100.0	75	100.0	161	100.0	174	100.0	257	100.0	461	100.0	456	100.0	418	100.0	268	100.0
	20pg未満	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	1	0.2	1	0.4
	20-25未満	12	0.8	1	1.3	1	0.6	2	1.1	0	0.0	2	0.4	6	1.3	5	1.2	3	1.1
	25-27未満	25	1.6	1	1.3	2	1.2	3	1.7	4	1.6	6	1.3	9	2.0	7	1.7	5	1.9
	27-29未満	99	6.3	9	12.0	16	9.9	16	9.2	14	5.4	17	3.7	27	5.9	14	3.3	16	6.0
	29-31未満	614	38.8	40	53.3	94	58.4	75	43.1	98	38.1	166	36.0	141	30.9	148	35.4	80	29.9
	31-33未満	614	38.8	22	29.3	42	26.1	66	37.9	107	41.6	184	39.9	193	42.3	176	42.1	112	41.8
	33-35未満	186	11.7	2	2.7	5	3.1	11	6.3	31	12.1	66	14.3	71	15.6	54	12.9	48	17.9
	35-40未満	31	2.0	0	0.0	1	0.6	1	0.6	3	1.2	19	4.1	7	1.5	13	3.1	3	1.1
	40pg以上	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0
女性	総 数	2,261	100.0	105	100.0	318	100.0	307	100.0	391	100.0	563	100.0	577	100.0	545	100.0	333	100.0
	20pg未満	21	0.9	1	1.0	4	1.3	13	4.2	2	0.5	0	0.0	1	0.2	1	0.2	0	0.0
	20-25未満	66	2.9	0	0.0	22	6.9	30	9.8	8	2.0	4	0.7	2	0.3	3	0.6	2	0.6
	25-27未満	72	3.2	4	3.8	14	4.4	27	8.8	9	2.3	6	1.1	12	2.1	7	1.3	8	2.4
	27-29未満	381	16.9	29	27.6	80	25.2	63	20.5	75	19.2	69	12.3	65	11.3	66	12.1	34	10.2
	29-31未満	1,160	51.3	51	48.6	151	47.5	115	37.5	198	50.6	342	60.7	303	52.5	320	58.7	173	52.0
	31-33未満	522	23.1	19	18.1	46	14.5	55	17.9	92	23.5	129	22.9	181	31.4	133	24.4	108	32.4
	33-35未満	37	1.6	1	1.0	1	0.3	4	1.3	6	1.5	12	2.1	13	2.3	14	2.6	8	2.4
	35-40未満	2	0.1	0	0.0	0	0.0	0	0.0	1	0.3	1	0.2	0	0.0	1	0.2	0	0.0
	40pg以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	平均値(pg)	29.8		29.6		29.0		28.5		29.9		30.2		30.4		30.3		30.4	
	標準偏差	2.2		1.7		2.5		3.4		2.0		1.4		1.6		1.5		1.6	

(22年)

第27表 平均赤血球血色素濃度（MCHC）の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,584	100.0	75	100.0	161	100.0	174	100.0	257	100.0	461	100.0	456	100.0	418	100.0	268	100.0
	22%未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	22-24未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	24-26未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	26-28未満	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	0	0.0
	28-30未満	17	1.1	1	1.3	1	0.6	1	0.6	1	0.4	2	0.4	11	2.4	5	1.2	7	2.6
	30-32未満	290	18.3	7	9.3	24	14.9	36	20.7	33	12.8	76	16.5	114	25.0	78	18.7	75	28.0
	32-34未満	1,067	67.4	50	66.7	101	62.7	111	63.8	185	72.0	325	70.5	295	64.7	292	69.9	167	62.3
	34%以上	209	13.2	17	22.7	35	21.7	26	14.9	38	14.8	58	12.6	35	7.7	42	10.0	19	7.1
	平均値(%)	32.9		33.3		33.1		32.9		33.1		32.9		32.6		32.8		32.5	
女性	総 数	2,261	100.0	105	100.0	318	100.0	307	100.0	391	100.0	563	100.0	577	100.0	545	100.0	333	100.0
	22%未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	22-24未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	24-26未満	1	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	26-28未満	20	0.9	0	0.0	4	1.3	13	4.2	3	0.8	0	0.0	0	0.0	0	0.0	0	0.0
	28-30未満	75	3.3	3	2.9	11	3.5	31	10.1	8	2.0	8	1.4	14	2.4	9	1.7	11	3.3
	30-32未満	810	35.8	26	24.8	115	36.2	94	30.6	122	31.2	203	36.1	250	43.3	210	38.5	144	43.2
	32-34未満	1,313	58.1	72	68.6	182	57.2	163	53.1	247	63.2	344	61.1	305	52.9	316	58.0	176	52.9
	34%以上	42	1.9	4	3.8	6	1.9	5	1.6	11	2.8	8	1.4	8	1.4	10	1.8	2	0.6
	平均値(%)	32.1		32.5		32.1		31.8		32.3		32.2		32.1		32.2		32.0	
	標準偏差	1.1		1.0		1.3		1.6		1.1		0.9		0.9		0.9		0.9	

(22年)

第 28 表 血清総たんぱく質値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75-79歳		(再掲) 80歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,597	100.0	75	100.0	162	100.0	175	100.0	259	100.0	462	100.0	464	100.0	424	100.0	140	100.0	131	100.0
	6.0g/dl未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	6.0-6.4	15	0.9	0	0.0	0	0.0	2	1.1	2	0.8	7	1.5	4	0.9	4	0.9	1	0.7	1	0.8
	6.5-6.9	154	9.6	1	1.3	12	7.4	17	9.7	17	6.6	50	10.8	57	12.3	50	11.8	18	12.9	18	13.7
	7.0-7.4	684	42.8	20	26.7	66	40.7	80	45.7	110	42.5	205	44.4	203	43.8	178	42.0	59	42.1	61	46.6
	7.5-7.9	575	36.0	39	52.0	59	36.4	60	34.3	109	42.1	153	33.1	155	33.4	142	33.5	54	38.6	37	28.2
	8.0-8.4	157	9.8	14	18.7	22	13.6	16	9.1	20	7.7	43	9.3	42	9.1	46	10.8	7	5.0	13	9.9
	8.5g/dll以上	12	0.8	1	1.3	3	1.9	0	0.0	1	0.4	4	0.9	3	0.6	4	0.9	1	0.7	1	0.8
	(再掲)6.5未満	15	0.9	0	0.0	0	0.0	2	1.1	2	0.8	7	1.5	4	0.9	4	0.9	1	0.7	1	0.8
平均値 (g/dl)	7.4		7.7		7.5		7.4		7.5		7.4		7.4		7.4		7.4		7.3		
標準偏差	0.4		0.3		0.4		0.4		0.4		0.4		0.4		0.4		0.4		0.4		
女性	総 数	2,268	100.0	105	100.0	321	100.0	308	100.0	392	100.0	564	100.0	578	100.0	546	100.0	172	100.0	161	100.0
	6.0g/dl未満	2	0.1	0	0.0	0	0.0	1	0.3	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0
	6.0-6.4	11	0.5	1	1.0	1	0.3	1	0.3	0	0.0	5	0.9	3	0.5	7	1.3	0	0.0	1	0.6
	6.5-6.9	227	10.0	8	7.6	29	9.0	35	11.4	31	7.9	46	8.2	78	13.5	56	10.3	24	14.0	28	17.4
	7.0-7.4	914	40.3	33	31.4	138	43.0	136	44.2	169	43.1	200	35.5	238	41.2	216	39.6	69	40.1	65	40.4
	7.5-7.9	880	38.8	46	43.8	127	39.6	109	35.4	152	38.8	245	43.4	201	34.8	208	38.1	61	35.5	53	32.9
	8.0-8.4	200	8.8	14	13.3	25	7.8	23	7.5	31	7.9	58	10.3	49	8.5	54	9.9	13	7.6	12	7.5
	8.5g/dll以上	34	1.5	3	2.9	1	0.3	3	1.0	9	2.3	9	1.6	9	1.6	5	0.9	5	2.9	2	1.2
	(再掲)6.5未満	13	0.6	1	1.0	1	0.3	2	0.6	0	0.0	6	1.1	3	0.5	7	1.3	0	0.0	1	0.6
平均値 (g/dl)	7.4		7.5		7.4		7.4		7.5		7.5		7.4		7.4		7.4		7.4		
標準偏差	0.4		0.5		0.4		0.4		0.4		0.4		0.4		0.4		0.4		0.4		

第30表 ヘモグロビンA1cの分布(性・年齢階級別)[インスリン注射または服薬者除外]

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	総 数	3,612	100.0	179	100.0	480	100.0	476	100.0	621	100.0	932	100.0	924	100.0	862	100.0	541	100.0
	3.9%以下	2	0.1	0	0.0	1	0.2	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0
	4.0-4.1	3	0.1	0	0.0	1	0.2	0	0.0	2	0.3	0	0.0	0	0.0	0	0.0	0	0.0
	4.2-4.3	9	0.2	2	1.1	5	1.0	0	0.0	1	0.2	1	0.1	0	0.0	0	0.0	0	0.0
	4.4-4.5	60	1.7	11	6.1	24	5.0	9	1.9	3	0.5	7	0.8	6	0.6	5	0.6	3	0.6
	4.6-4.7	187	5.2	31	17.3	49	10.2	36	7.6	22	3.5	26	2.8	23	2.5	23	2.7	12	2.2
	4.8-4.9	462	12.8	61	34.1	111	23.1	87	18.3	70	11.3	89	9.5	44	4.8	64	7.4	27	5.0
	5.0-5.1	724	20.0	40	22.3	138	28.8	121	25.4	132	21.3	158	17.0	135	14.6	129	15.0	84	15.5
	5.2	386	10.7	17	9.5	50	10.4	49	10.3	72	11.6	93	10.0	105	11.4	96	11.1	53	9.8
	5.3	376	10.4	8	4.5	35	7.3	39	8.2	71	11.4	110	11.8	113	12.2	103	11.9	66	12.2
	5.4	333	9.2	3	1.7	18	3.8	38	8.0	69	11.1	98	10.5	107	11.6	94	10.9	62	11.5
	5.5	260	7.2	4	2.2	21	4.4	30	6.3	49	7.9	68	7.3	88	9.5	84	9.7	44	8.1
	5.6	215	6.0	1	0.6	13	2.7	15	3.2	40	6.4	77	8.3	69	7.5	67	7.8	48	8.9
	5.7	141	3.9	1	0.6	3	0.6	17	3.6	26	4.2	44	4.7	50	5.4	51	5.9	24	4.4
	5.8	119	3.3	0	0.0	2	0.4	10	2.1	16	2.6	47	5.0	44	4.8	39	4.5	27	5.0
	5.9	70	1.9	0	0.0	3	0.6	2	0.4	7	1.1	22	2.4	36	3.9	27	3.1	23	4.3
	6.0	57	1.6	0	0.0	0	0.0	5	1.1	9	1.4	20	2.1	23	2.5	17	2.0	16	3.0
	6.1	33	0.9	0	0.0	2	0.4	3	0.6	2	0.3	13	1.4	13	1.4	12	1.4	10	1.8
	6.2-6.3	57	1.6	0	0.0	3	0.6	4	0.8	10	1.6	16	1.7	24	2.6	16	1.9	15	2.8
	6.4-6.5	30	0.8	0	0.0	0	0.0	0	0.0	5	0.8	11	1.2	14	1.5	11	1.3	7	1.3
	6.6-6.7	18	0.5	0	0.0	0	0.0	2	0.4	2	0.3	9	1.0	5	0.5	5	0.6	3	0.6
	6.8-6.9	13	0.4	0	0.0	0	0.0	1	0.2	2	0.3	3	0.3	7	0.8	3	0.3	6	1.1
	7.0-7.1	8	0.2	0	0.0	0	0.0	1	0.2	0	0.0	2	0.2	5	0.5	3	0.3	3	0.6
	7.2-7.3	8	0.2	0	0.0	0	0.0	1	0.2	1	0.2	2	0.2	4	0.4	3	0.3	2	0.4
	7.4-7.5	7	0.2	0	0.0	0	0.0	0	0.0	1	0.2	3	0.3	3	0.3	1	0.1	3	0.6
	7.6-7.7	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1	1	0.1	0	0.0
	7.8-7.9	3	0.1	0	0.0	1	0.2	0	0.0	1	0.2	1	0.1	0	0.0	0	0.0	0	0.0
	8.0-8.1	4	0.1	0	0.0	0	0.0	0	0.0	1	0.2	1	0.1	2	0.2	1	0.1	2	0.4
	8.2-8.3	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	2	0.2	0	0.0	1	0.1	0	0.0
	8.4-8.5	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.6-8.7	3	0.1	0	0.0	0	0.0	1	0.2	0	0.0	1	0.1	1	0.1	1	0.1	0	0.0
	8.8-8.9	1	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0
	9.0%以上	20	0.6	0	0.0	0	0.0	5	1.1	5	0.8	8	0.9	2	0.2	5	0.6	1	0.2
	(再掲)6.1以上	208	5.8	0	0.0	6	1.3	18	3.8	31	5.0	72	7.7	81	8.8	63	7.3	52	9.6
平均値(%) 標準偏差		5.3 0.6		4.9 0.3		5.0 0.3		5.2 0.8		5.3 0.6		5.4 0.6		5.5 0.6		5.4 0.6		5.5 0.6	
男 性	総 数	1,456	100.0	75	100.0	159	100.0	171	100.0	237	100.0	405	100.0	409	100.0	370	100.0	239	100.0
	3.9%以下	1	0.1	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0
	4.0-4.1	2	0.1	0	0.0	0	0.0	0	0.0	2	0.8	0	0.0	0	0.0	0	0.0	0	0.0
	4.2-4.3	4	0.3	0	0.0	2	1.3	0	0.0	1	0.4	1	0.2	0	0.0	0	0.0	0	0.0
	4.4-4.5	30	2.1	5	6.7	9	5.7	3	1.8	2	0.8	7	1.7	4	1.0	5	1.4	1	0.4
	4.6-4.7	91	6.3	15	20.0	22	13.8	17	9.9	7	3.0	17	4.2	13	3.2	14	3.8	6	2.5
	4.8-4.9	177	12.2	24	32.0	39	24.5	29	17.0	18	7.6	45	11.1	22	5.4	33	8.9	13	5.4
	5.0-5.1	295	20.3	18	24.0	46	28.9	46	26.9	53	22.4	75	18.5	57	13.9	52	14.1	39	16.3
	5.2	167	11.5	4	5.3	16	10.1	20	11.7	36	15.2	40	9.9	51	12.5	38	10.3	29	12.1
	5.3	140	9.6	7	9.3	9	5.7	13	7.6	22	9.3	42	10.4	47	11.5	35	9.5	31	13.0
	5.4	102	7.0	2	2.7	4	2.5	9	5.3	24	10.1	31	7.7	32	7.8	31	8.4	16	6.7
	5.5	94	6.5	0	0.0	5	3.1	10	5.8	17	7.2	24	5.9	38	9.3	34	9.2	20	8.4
	5.6	81	5.6	0	0.0	4	2.5	3	1.8	14	5.9	30	7.4	30	7.3	31	8.4	19	7.9
	5.7	54	3.7	0	0.0	0	0.0	8	4.7	8	3.4	18	4.4	20	4.9	24	6.5	9	3.8
	5.8	46	3.2	0	0.0	0	0.0	2	1.2	7	3.0	19	4.7	18	4.4	14	3.8	11	4.6
	5.9	34	2.3	0	0.0	0	0.0	0	0.0	3	1.3	8	2.0	23	5.6	14	3.8	13	5.4
	6.0	25	1.7	0	0.0	0	0.0	1	0.6	4	1.7	11	2.7	9	2.2	8	2.2	7	2.9
	6.1	18	1.2	0	0.0	2	1.3	2	1.2	0	0.0	8	2.0	6	1.5	7	1.9	3	1.3
	6.2-6.3	32	2.2	0	0.0	0	0.0	2	1.2	6	2.5	10	2.5	14	3.4	10	2.7	8	3.3
	6.4-6.5	15	1.0	0	0.0	0	0.0	0	0.0	5	2.1	4	1.0	6	1.5	5	1.4	2	0.8
	6.6-6.7	7	0.5	0	0.0	0	0.0	1	0.6	0	0.0	4	1.0	2	0.5	2	0.5	1	0.4
	6.8-6.9	7	0.5	0	0.0	0	0.0	0	0.0	1	0.4	1	0.2	5	1.2	1	0.3	5	2.1
	7.0-7.1	5	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	5	1.2	2	0.5	3	1.3
	7.2-7.3	6	0.4	0	0.0	0	0.0	1	0.6	1	0.4	1	0.2	3	0.7	3	0.8	1	0.4
	7.4-7.5	1	0.1	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0
	7.6-7.7	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.3	0	0.0
	7.8-7.9	3	0.2	0	0.0	1	0.6	0	0.0	1	0.4	1	0.2	0	0.0	0	0.0	0	0.0
	8.0-8.1	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	1	0.3	1	0.4
	8.2-8.3	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.3	0	0.0
	8.4-8.5	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.6-8.7	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0
	8.8-8.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	9.0%以上	14	1.0	0	0.0	0	0.0	4	2.3	3	1.3	5	1.2	2	0.5	4	1.1	1	0.4
	(再掲)6.1以上	113	7.8	0	0.0	3	1.9	10	5.8	18	7.6	37	9.1	45	11.0	37	10.0	25	10.5
平均値(%) 標準偏差		5.4 0.7		4.9 0.2		5.0 0.4		5.3 1.1		5.4 0.7		5.4 0.7		5.5 0.7		5.5 0.7		5.5 0.6	

(22年)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
女 性	総 数	2,156	100.0	104	100.0	321	100.0	305	100.0	384	100.0	527	100.0	515	100.0	492	100.0	302	100.0
	3.9%以下	1	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	4.0-4.1	1	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	4.2-4.3	5	0.2	2	1.9	3	0.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	4.4-4.5	30	1.4	6	5.8	15	4.7	6	2.0	1	0.3	0	0.0	2	0.4	0	0.0	2	0.7
	4.6-4.7	96	4.5	16	15.4	27	8.4	19	6.2	15	3.9	9	1.7	10	1.9	9	1.8	6	2.0
	4.8-4.9	285	13.2	37	35.6	72	22.4	58	19.0	52	13.5	44	8.3	22	4.3	31	6.3	14	4.6
	5.0-5.1	429	19.9	22	21.2	92	28.7	75	24.6	79	20.6	83	15.7	78	15.1	77	15.7	45	14.9
	5.2	219	10.2	13	12.5	34	10.6	29	9.5	36	9.4	53	10.1	54	10.5	58	11.8	24	7.9
	5.3	236	10.9	1	1.0	26	8.1	26	8.5	49	12.8	68	12.9	66	12.8	68	13.8	35	11.6
	5.4	231	10.7	1	1.0	14	4.4	29	9.5	45	11.7	67	12.7	75	14.6	63	12.8	46	15.2
	5.5	166	7.7	4	3.8	16	5.0	20	6.6	32	8.3	44	8.3	50	9.7	50	10.2	24	7.9
	5.6	134	6.2	1	1.0	9	2.8	12	3.9	26	6.8	47	8.9	39	7.6	36	7.3	29	9.6
	5.7	87	4.0	1	1.0	3	0.9	9	3.0	18	4.7	26	4.9	30	5.8	27	5.5	15	5.0
	5.8	73	3.4	0	0.0	2	0.6	8	2.6	9	2.3	28	5.3	26	5.0	25	5.1	16	5.3
	5.9	36	1.7	0	0.0	3	0.9	2	0.7	4	1.0	14	2.7	13	2.5	13	2.6	10	3.3
	6.0	32	1.5	0	0.0	0	0.0	4	1.3	5	1.3	9	1.7	14	2.7	9	1.8	9	3.0
	6.1	15	0.7	0	0.0	0	0.0	1	0.3	2	0.5	5	0.9	7	1.4	5	1.0	7	2.3
	6.2-6.3	25	1.2	0	0.0	3	0.9	2	0.7	4	1.0	6	1.1	10	1.9	6	1.2	7	2.3
	6.4-6.5	15	0.7	0	0.0	0	0.0	0	0.0	0	0.0	7	1.3	8	1.6	6	1.2	5	1.7
	6.6-6.7	11	0.5	0	0.0	0	0.0	1	0.3	2	0.5	5	0.9	3	0.6	3	0.6	2	0.7
	6.8-6.9	6	0.3	0	0.0	0	0.0	1	0.3	1	0.3	2	0.4	2	0.4	2	0.4	1	0.3
	7.0-7.1	3	0.1	0	0.0	0	0.0	1	0.3	0	0.0	2	0.4	0	0.0	1	0.2	0	0.0
	7.2-7.3	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	0	0.0	1	0.3
	7.4-7.5	6	0.3	0	0.0	0	0.0	0	0.0	0	0.0	3	0.6	3	0.6	1	0.2	3	1.0
	7.6-7.7	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	7.8-7.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.0-8.1	2	0.1	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	1	0.2	0	0.0	1	0.3
	8.2-8.3	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0
	8.4-8.5	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.6-8.7	2	0.1	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	1	0.2	1	0.2	0	0.0
	8.8-8.9	1	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0
	9.0%以上	6	0.3	0	0.0	0	0.0	1	0.3	2	0.5	3	0.6	0	0.0	1	0.2	0	0.0
	(再掲)6.1以上	95	4.4	0	0.0	3	0.9	8	2.6	13	3.4	35	6.6	36	7.0	26	5.3	27	8.9
平均値 (%)		5.3		4.9		5.1		5.2		5.3		5.4		5.4		5.4		5.5	
標準偏差		0.5		0.3		0.3		0.5		0.6		0.6		0.5		0.5		0.5	

注) インスリン注射または血糖を下げる薬服用者除外

(22年)

第31表 血糖値の分布（食後3時間以上、性・年齢階級別）〔インスリン注射または服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総 数	1,160	100.0	62	100.0	130	100.0	134	100.0	207	100.0	329	100.0	298	100.0	291	100.0	169	100.0
	40mg/dl未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	40-59	1	0.1	0	0.0	1	0.8	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	60-79	77	6.6	12	19.4	16	12.3	2	1.5	16	7.7	21	6.4	10	3.4	16	5.5	6	3.6
	80-99	684	59.0	42	67.7	98	75.4	110	82.1	145	70.0	171	52.0	118	39.6	144	49.5	58	34.3
	100-119	283	24.4	8	12.9	13	10.0	16	11.9	35	16.9	95	28.9	116	38.9	89	30.6	67	39.6
	120-140	70	6.0	0	0.0	2	1.5	3	2.2	7	3.4	26	7.9	32	10.7	30	10.3	21	12.4
	140-160	25	2.2	0	0.0	0	0.0	0	0.0	1	0.5	9	2.7	15	5.0	6	2.1	11	6.5
	160-180	4	0.3	0	0.0	0	0.0	0	0.0	0	0.0	2	0.6	2	0.7	1	0.3	2	1.2
	180-200	3	0.3	0	0.0	0	0.0	0	0.0	2	1.0	1	0.3	0	0.0	1	0.3	0	0.0
	200-220	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	1	0.6
	220-240	2	0.2	0	0.0	0	0.0	0	0.0	1	0.5	0	0.0	1	0.3	0	0.0	1	0.6
	240-260	3	0.3	0	0.0	0	0.0	1	0.7	0	0.0	2	0.6	0	0.0	1	0.3	0	0.0
	260mg/dl以上	7	0.6	0	0.0	0	0.0	2	1.5	0	0.0	2	0.6	3	1.0	3	1.0	2	1.2
	(再掲)110以上	206	17.8	3	4.8	7	5.4	8	6.0	20	9.7	65	19.8	103	34.6	79	27.1	64	37.9
	平均値(mg/dl) 標準偏差	99.4 26.3		88.8 10.5		89.6 11.1		96.4 33.4		94.8 17.2		101.5 24.8		108.3 32.8		103.5 27.5		111.4 36.9	
女 性	総 数	1,616	100.0	76	100.0	226	100.0	233	100.0	293	100.0	406	100.0	382	100.0	373	100.0	229	100.0
	40mg/dl未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	40-59	2	0.1	1	1.3	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0
	60-79	95	5.9	11	14.5	33	14.6	16	6.9	14	4.8	10	2.5	11	2.9	9	2.4	7	3.1
	80-99	1,004	62.1	59	77.6	164	72.6	181	77.7	208	71.0	205	50.5	187	49.0	193	51.7	106	46.3
	100-119	389	24.1	5	6.6	22	9.7	30	12.9	59	20.1	141	34.7	132	34.6	127	34.0	81	35.4
	120-140	83	5.1	0	0.0	6	2.7	5	2.1	7	2.4	33	8.1	32	8.4	30	8.0	19	8.3
	140-160	25	1.5	0	0.0	1	0.4	0	0.0	1	0.3	11	2.7	12	3.1	9	2.4	9	3.9
	160-180	11	0.7	0	0.0	0	0.0	1	0.4	1	0.3	3	0.7	6	1.6	3	0.8	5	2.2
	180-200	4	0.2	0	0.0	0	0.0	0	0.0	1	0.3	2	0.5	1	0.3	1	0.3	1	0.4
	200-220	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	1	0.4
	220-240	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	240-260	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	260mg/dl以上	2	0.1	0	0.0	0	0.0	0	0.0	1	0.3	1	0.2	0	0.0	1	0.3	0	0.0
	(再掲)110以上	251	15.5	1	1.3	11	4.9	14	6.0	25	8.5	101	24.9	99	25.9	90	24.1	63	27.5
	平均値(mg/dl) 標準偏差	97.6 17.6		87.0 9.1		89.8 12.1		92.2 11.0		95.8 17.7		102.5 20.2		103.6 17.8		102.3 19.8		105.0 19.7	

注) インスリン注射または血糖を下げる薬服用者除外

(22年)

第 32 表の 1 血清総コレステロール値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-79歳		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,597	100.0	75	100.0	162	100.0	175	100.0	259	100.0	462	100.0	464	100.0	1,229	100.0	424	100.0	271	100.0
	120mg/dl未満	9	0.6	0	0.0	2	1.2	0	0.0	1	0.4	3	0.6	3	0.6	6	0.5	4	0.9	1	0.4
	120-139	24	1.5	3	4.0	1	0.6	1	0.6	3	1.2	6	1.3	10	2.2	17	1.4	8	1.9	4	1.5
	140-159	140	8.8	14	18.7	16	9.9	2	1.1	16	6.2	28	6.1	64	13.8	91	7.4	33	7.8	45	16.6
	160-179	267	16.7	24	32.0	32	19.8	19	10.9	26	10.0	69	14.9	97	20.9	183	14.9	81	19.1	61	22.5
	180-199	357	22.4	12	16.0	29	17.9	38	21.7	48	18.5	114	24.7	116	25.0	280	22.8	102	24.1	68	25.1
	200-219	353	22.1	12	16.0	36	22.2	41	23.4	63	24.3	106	22.9	95	20.5	283	23.0	97	22.9	49	18.1
	220-239	245	15.3	6	8.0	24	14.8	38	21.7	56	21.6	75	16.2	46	9.9	200	16.3	58	13.7	26	9.6
	240-259	113	7.1	3	4.0	13	8.0	18	10.3	20	7.7	40	8.7	19	4.1	92	7.5	24	5.7	10	3.7
	260-279	62	3.9	0	0.0	6	3.7	12	6.9	18	6.9	17	3.7	9	1.9	54	4.4	11	2.6	6	2.2
	280-299	17	1.1	0	0.0	2	1.2	3	1.7	5	1.9	3	0.6	4	0.9	15	1.2	4	0.9	1	0.4
	300mg/dl以上	10	0.6	1	1.3	1	0.6	3	1.7	3	1.2	1	0.2	1	0.2	8	0.7	2	0.5	0	0.0
	(再掲)220以上	447	28.0	10	13.3	46	28.4	74	42.3	102	39.4	136	29.4	79	17.0	369	30.0	99	23.3	43	15.9
	(再掲)240以上	202	12.6	4	5.3	22	13.6	36	20.6	46	17.8	61	13.2	33	7.1	169	13.8	41	9.7	17	6.3
平均値(mg/dl)		200.7		182.9		200.1		214.8		210.3		203.0		190.9		203.2		197.4		188.9	
標準偏差		34.3		32.2		36.1		33.3		35.5		31.9		32.1		34.1		32.9		30.5	
女 性	総数	2,268	100.0	105	100.0	321	100.0	308	100.0	392	100.0	564	100.0	578	100.0	1,681	100.0	546	100.0	333	100.0
	120mg/dl未満	2	0.1	0	0.0	0	0.0	2	0.6	0	0.0	0	0.0	0	0.0	2	0.1	0	0.0	0	0.0
	120-139	29	1.3	7	6.7	9	2.8	3	1.0	2	0.5	1	0.2	7	1.2	13	0.8	4	0.7	4	1.2
	140-159	138	6.1	20	19.0	46	14.3	23	7.5	12	3.1	10	1.8	27	4.7	62	3.7	16	2.9	14	4.2
	160-179	301	13.3	32	30.5	80	24.9	47	15.3	28	7.1	35	6.2	79	13.7	167	9.9	48	8.8	48	14.4
	180-199	497	21.9	19	18.1	99	30.8	84	27.3	65	16.6	104	18.4	126	21.8	340	20.2	108	19.8	75	22.5
	200-219	528	23.3	14	13.3	45	14.0	69	22.4	101	25.8	139	24.6	160	27.7	423	25.2	143	26.2	98	29.4
	220-239	362	16.0	6	5.7	24	7.5	46	14.9	79	20.2	113	20.0	94	16.3	305	18.1	94	17.2	54	16.2
	240-259	221	9.7	5	4.8	11	3.4	21	6.8	49	12.5	87	15.4	48	8.3	193	11.5	77	14.1	22	6.6
	260-279	120	5.3	2	1.9	7	2.2	9	2.9	33	8.4	44	7.8	25	4.3	109	6.5	36	6.6	10	3.0
	280-299	46	2.0	0	0.0	0	0.0	2	0.6	17	4.3	18	3.2	9	1.6	43	2.6	13	2.4	6	1.8
	300mg/dl以上	24	1.1	0	0.0	0	0.0	2	0.6	6	1.5	13	2.3	3	0.5	24	1.4	7	1.3	2	0.6
	(再掲)220以上	773	34.1	13	12.4	42	13.1	80	26.0	184	46.9	275	48.8	179	31.0	674	40.1	227	41.6	94	28.2
	(再掲)240以上	411	18.1	7	6.7	18	5.6	34	11.0	105	26.8	162	28.7	85	14.7	369	22.0	133	24.4	40	12.0
平均値(mg/dl)		208.1		181.2		186.4		201.3		220.2		222.0		206.9		214.3		216.0		205.6	
標準偏差		35.7		31.8		29.5		33.1		34.8		35.5		31.6		35.1		34.0		30.9	

(22年)

第 32 表の 2 血清総コレステロール値の分布（性・年齢階級別）〔服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-79歳		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,382	100.0	75	100.0	161	100.0	163	100.0	224	100.0	388	100.0	371	100.0	1,041	100.0	343	100.0	217	100.0
	120mg/dl未満	9	0.7	0	0.0	2	1.2	0	0.0	1	0.4	3	0.8	3	0.8	6	0.6	4	1.2	1	0.5
	120-139	20	1.4	3	4.0	1	0.6	1	0.6	3	1.3	6	1.5	6	1.6	15	1.4	6	1.7	2	0.9
	140-159	107	7.7	14	18.7	16	9.9	2	1.2	11	4.9	20	5.2	44	11.9	63	6.1	22	6.4	31	14.3
	160-179	224	16.2	24	32.0	32	19.9	17	10.4	21	9.4	53	13.7	77	20.8	146	14.0	61	17.8	49	22.6
	180-199	304	22.0	12	16.0	29	18.0	35	21.5	42	18.8	92	23.7	94	25.3	233	22.4	82	23.9	55	25.3
	200-219	316	22.9	12	16.0	36	22.4	40	24.5	54	24.1	95	24.5	79	21.3	250	24.0	82	23.9	42	19.4
	220-239	223	16.1	6	8.0	24	14.9	37	22.7	50	22.3	67	17.3	39	10.5	180	17.3	51	14.9	22	10.1
	240-259	102	7.4	3	4.0	13	8.1	16	9.8	19	8.5	34	8.8	17	4.6	82	7.9	22	6.4	8	3.7
	260-279	54	3.9	0	0.0	5	3.1	10	6.1	17	7.6	14	3.6	8	2.2	47	4.5	8	2.3	6	2.8
	280-299	13	0.9	0	0.0	2	1.2	2	1.2	3	1.3	3	0.8	3	0.8	11	1.1	3	0.9	1	0.5
	300mg/dl以上	10	0.7	1	1.3	1	0.6	3	1.8	3	1.3	1	0.3	1	0.3	8	0.8	2	0.6	0	0.0
	(再掲)220以上	402	29.1	10	13.3	45	28.0	68	41.7	92	41.1	119	30.7	68	18.3	328	31.5	86	25.1	37	17.1
	(再掲)240以上	179	13.0	4	5.3	21	13.0	31	19.0	42	18.8	52	13.4	29	7.8	148	14.2	35	10.2	15	6.9
平均値(mg/dl)		202.0		182.9		199.7		214.3		211.9		204.4		192.8		204.9		199.2		191.0	
標準偏差		34.2		32.2		35.7		32.8		35.2		32.0		32.1		33.8		32.7		30.8	
女 性	総数	1,911	100.0	105	100.0	318	100.0	306	100.0	354	100.0	439	100.0	389	100.0	1,375	100.0	384	100.0	228	100.0
	120mg/dl未満	2	0.1	0	0.0	0	0.0	2	0.7	0	0.0	0	0.0	0	0.0	2	0.1	0	0.0	0	0.0
	120-139	24	1.3	7	6.7	9	2.8	3	1.0	2	0.6	1	0.2	2	0.5	8	0.6	3	0.8	0	0.0
	140-159	115	6.0	20	19.0	46	14.5	23	7.5	8	2.3	6	1.4	12	3.1	43	3.1	6	1.6	7	3.1
	160-179	248	13.0	32	30.5	79	24.8	47	15.4	22	6.2	21	4.8	47	12.1	125	9.1	30	7.8	29	12.7
	180-199	407	21.3	19	18.1	98	30.8	83	27.1	58	16.4	69	15.7	80	20.6	266	19.3	68	17.7	47	20.6
	200-219	424	22.2	14	13.3	44	13.8	69	22.5	89	25.1	99	22.6	109	28.0	333	24.2	90	23.4	72	31.6
	220-239	318	16.6	6	5.7	24	7.5	45	14.7	74	20.9	98	22.3	71	18.3	264	19.2	74	19.3	42	18.4
	240-259	195	10.2	5	4.8	11	3.5	21	6.9	46	13.0	75	17.1	37	9.5	169	12.3	62	16.1	17	7.5
	260-279	112	5.9	2	1.9	7	2.2	9	2.9	33	9.3	40	9.1	21	5.4	102	7.4	32	8.3	8	3.5
	280-299	45	2.4	0	0.0	0	0.0	2	0.7	17	4.8	18	4.1	8	2.1	42	3.1	13	3.4	5	2.2
	300mg/dl以上	21	1.1	0	0.0	0	0.0	2	0.7	5	1.4	12	2.7	2	0.5	21	1.5	6	1.6	1	0.4
	(再掲)220以上	691	36.2	13	12.4	42	13.2	79	25.8	175	49.4	243	55.4	139	35.7	598	43.5	187	48.7	73	32.0
	(再掲)240以上	373	19.5	7	6.7	18	5.7	34	11.1	101	28.5	145	33.0	68	17.5	334	24.3	113	29.4	31	13.6
平均値(mg/dl)		209.4		181.2		186.4		201.2		222.1		226.7		211.3		217.0		221.1		209.8	
標準偏差		36.5		31.8		29.5		33.1		34.3		35.9		31.2		35.5		34.8		29.5	

第33表 血清中性脂肪（トリグリセライド）値の分布（性・年齢階級別）〔服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,382	100.0	75	100.0	161	100.0	163	100.0	224	100.0	388	100.0	371	100.0	343	100.0
	20mg/dl未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	20-49	59	4.3	6	8.0	10	6.2	8	4.9	9	4.0	9	2.3	17	4.6	11	3.2
	50-79	230	16.6	23	30.7	29	18.0	22	13.5	40	17.9	47	12.1	69	18.6	47	13.7
	80-109	272	19.7	12	16.0	29	18.0	26	16.0	36	16.1	85	21.9	84	22.6	82	23.9
	110-139	230	16.6	13	17.3	17	10.6	24	14.7	34	15.2	67	17.3	75	20.2	63	18.4
	140-169	178	12.9	9	12.0	23	14.3	20	12.3	27	12.1	51	13.1	48	12.9	40	11.7
	170-199	121	8.8	6	8.0	10	6.2	17	10.4	26	11.6	31	8.0	31	8.4	31	9.0
	200mg/dl以上	292	21.1	6	8.0	43	26.7	46	28.2	52	23.2	98	25.3	47	12.7	69	20.1
	(再掲)150以上	525	38.0	17	22.7	67	41.6	77	47.2	97	43.3	158	40.7	109	29.4	127	37.0
平均値(mg/dl)		152.5		121.4		161.6		171.2		155.3		164.0		132.8		154.5	
標準偏差		103.2		92.1		123.0		115.3		93.5		108.0		85.2		108.6	
女 性	総数	1,911	100.0	105	100.0	318	100.0	306	100.0	354	100.0	439	100.0	389	100.0	384	100.0
	20mg/dl未満	2	0.1	2	1.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	20-49	228	11.9	30	28.6	63	19.8	51	16.7	29	8.2	27	6.2	28	7.2	28	7.3
	50-79	516	27.0	40	38.1	121	38.1	100	32.7	92	26.0	95	21.6	68	17.5	73	19.0
	80-109	420	22.0	15	14.3	60	18.9	64	20.9	89	25.1	96	21.9	96	24.7	77	20.1
	110-139	287	15.0	5	4.8	38	11.9	41	13.4	43	12.1	76	17.3	84	21.6	73	19.0
	140-169	169	8.8	2	1.9	18	5.7	21	6.9	35	9.9	51	11.6	42	10.8	43	11.2
	170-199	96	5.0	4	3.8	8	2.5	6	2.0	21	5.9	28	6.4	29	7.5	34	8.9
	200mg/dl以上	193	10.1	7	6.7	10	3.1	23	7.5	45	12.7	66	15.0	42	10.8	56	14.6
	(再掲)150以上	394	20.6	12	11.4	31	9.7	40	13.1	87	24.6	125	28.5	99	25.4	118	30.7
平均値(mg/dl)		114.1		82.7		86.4		103.6		124.4		132.5		123.4		133.3	
標準偏差		80.9		67.5		49.7		89.8		90.5		88.7		69.2		85.7	

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

(22年)

第34表 血清HDL- コレステロール値の分布（性・年齢階級別）〔服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,382	100.0	75	100.0	161	100.0	163	100.0	224	100.0	388	100.0	371	100.0	343	100.0
	30mg/dl未満	14	1.0	0	0.0	1	0.6	1	0.6	3	1.3	3	0.8	6	1.6	4	1.2
	30-39	140	10.1	5	6.7	16	9.9	14	8.6	21	9.4	48	12.4	36	9.7	37	10.8
	40-49	325	23.5	17	22.7	39	24.2	32	19.6	57	25.4	80	20.6	100	27.0	87	25.4
	50-59	366	26.5	24	32.0	50	31.1	48	29.4	56	25.0	95	24.5	93	25.1	93	27.1
	60-69	269	19.5	17	22.7	26	16.1	35	21.5	40	17.9	81	20.9	70	18.9	63	18.4
	70-79	144	10.4	8	10.7	17	10.6	13	8.0	18	8.0	48	12.4	40	10.8	34	9.9
	80-89	78	5.6	4	5.3	9	5.6	10	6.1	16	7.1	20	5.2	19	5.1	17	5.0
	90-99	28	2.0	0	0.0	3	1.9	6	3.7	6	2.7	6	1.5	7	1.9	5	1.5
	100mg/dl以上	18	1.3	0	0.0	0	0.0	4	2.5	7	3.1	7	1.8	0	0.0	3	0.9
平均値(mg/dl)		57.0		56.9		56.3		58.2		58.1		57.5		55.7		55.6	
標準偏差		15.4		12.1		14.3		16.0		17.1		15.8		14.5		14.6	
女 性	総数	1,911	100.0	105	100.0	318	100.0	306	100.0	354	100.0	439	100.0	389	100.0	384	100.0
	30mg/dl未満	2	0.1	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	1	0.3	1	0.3
	30-39	36	1.9	1	1.0	1	0.3	3	1.0	7	2.0	10	2.3	14	3.6	13	3.4
	40-49	197	10.3	9	8.6	25	7.9	28	9.2	35	9.9	43	9.8	57	14.7	46	12.0
	50-59	411	21.5	21	20.0	67	21.1	55	18.0	59	16.7	110	25.1	99	25.4	95	24.7
	60-69	490	25.6	32	30.5	84	26.4	81	26.5	86	24.3	114	26.0	93	23.9	94	24.5
	70-79	407	21.3	19	18.1	82	25.8	72	23.5	74	20.9	82	18.7	78	20.1	81	21.1
	80-89	233	12.2	16	15.2	38	11.9	44	14.4	56	15.8	47	10.7	32	8.2	32	8.3
	90-99	95	5.0	3	2.9	17	5.3	15	4.9	23	6.5	23	5.2	14	3.6	18	4.7
	100mg/dl以上	40	2.1	4	3.8	4	1.3	7	2.3	14	4.0	10	2.3	1	0.3	4	1.0
平均値(mg/dl)		66.8		67.4		68.0		68.5		69.4		66.1		62.7		64.3	
標準偏差		15.2		13.9		13.3		15.7		16.6		15.3		14.1		15.0	

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

(22年)

第 35 表の 1 血清LDL-コレステロール値 (Friedewaldの式) の分布 (性・年齢階級別) [服薬者除外]

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,339	100.0	72	100.0	156	100.0	155	100.0	218	100.0	373	100.0	365	100.0	332	100.0
	40mg/dl未満	5	0.4	0	0.0	0	0.0	0	0.0	1	0.5	3	0.8	1	0.3	1	0.5
	40-50未満	12	0.9	0	0.0	3	1.9	1	0.6	1	0.5	4	1.1	3	0.8	6	1.8
	50-60未満	20	1.5	3	4.2	2	1.3	2	1.3	1	0.5	6	1.6	6	1.6	4	1.2
	60-70未満	36	2.7	8	11.1	5	3.2	3	1.9	3	1.4	9	2.4	8	2.2	7	2.1
	70-80未満	67	5.0	6	8.3	16	10.3	2	1.3	9	4.1	17	4.6	17	4.7	13	3.9
	80-90未満	104	7.8	7	9.7	13	8.3	8	5.2	21	9.6	19	5.1	36	9.9	23	6.9
	90-100未満	168	12.5	10	13.9	22	14.1	21	13.5	15	6.9	42	11.3	58	15.9	44	13.3
	100-110未満	171	12.8	11	15.3	18	11.5	13	8.4	25	11.5	46	12.3	58	15.9	47	14.2
	110-120未満	181	13.5	7	9.7	18	11.5	22	14.2	22	10.1	56	15.0	56	15.3	54	16.3
	120-130未満	174	13.0	11	15.3	15	9.6	25	16.1	34	15.6	50	13.4	39	10.7	42	12.7
	130-140未満	137	10.2	2	2.8	16	10.3	15	9.7	27	12.4	46	12.3	31	8.5	34	10.2
	140-150未満	107	8.0	4	5.6	5	3.2	16	10.3	21	9.6	33	8.8	28	7.7	27	8.1
	150-160未満	55	4.1	1	1.4	11	7.1	9	5.8	10	4.6	17	4.6	7	1.9	11	3.3
	160-170未満	41	3.1	1	1.4	6	3.8	4	2.6	10	4.6	12	3.2	8	2.2	10	3.0
	170-180未満	28	2.1	1	1.4	3	1.9	7	4.5	7	3.2	8	2.1	2	0.5	5	1.5
	180-190未満	17	1.3	0	0.0	2	1.3	2	1.3	6	2.8	2	0.5	5	1.4	3	0.9
	190-200未満	7	0.5	0	0.0	1	0.6	1	0.6	1	0.5	2	0.5	2	0.5	0	0.0
	200-210未満	2	0.1	0	0.0	0	0.0	2	1.3	0	0.0	0	0.0	0	0.0	0	0.0
	210-220未満	4	0.3	0	0.0	0	0.0	1	0.6	2	0.9	1	0.3	0	0.0	1	0.3
	220mg/dl以上	3	0.2	0	0.0	0	0.0	1	0.6	2	0.9	0	0.0	0	0.0	0	0.0
	平均値 (mg/dl) 標準偏差	115.7 30.1		101.9 27.4		111.7 31.2		123.7 30.9		123.8 32.8		116.5 29.3		111.0 26.8		114.2 28.2	109.1 25.9
女 性	総数	1,887	100.0	105	100.0	318	100.0	302	100.0	345	100.0	430	100.0	387	100.0	379	100.0
	40mg/dl未満	3	0.2	0	0.0	2	0.6	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0
	40-50未満	7	0.4	3	2.9	1	0.3	2	0.7	0	0.0	1	0.2	0	0.0	0	0.0
	50-60未満	17	0.9	3	2.9	5	1.6	5	1.7	1	0.3	2	0.5	1	0.3	2	0.5
	60-70未満	45	2.4	11	10.5	19	6.0	7	2.3	3	0.9	2	0.5	3	0.8	3	0.8
	70-80未満	88	4.7	14	13.3	37	11.6	16	5.3	9	2.6	5	1.2	7	1.8	6	1.6
	80-90未満	148	7.8	16	15.2	52	16.4	29	9.6	14	4.1	13	3.0	24	6.2	16	4.2
	90-100未満	218	11.6	15	14.3	52	16.4	46	15.2	38	11.0	26	6.0	41	10.6	31	8.2
	100-110未満	241	12.8	9	8.6	52	16.4	50	16.6	42	12.2	43	10.0	45	11.6	35	9.2
	110-120未満	224	11.9	12	11.4	33	10.4	36	11.9	40	11.6	51	11.9	52	13.4	45	11.9
	120-130未満	237	12.6	10	9.5	21	6.6	34	11.3	50	14.5	54	12.6	68	17.6	52	13.7
	130-140未満	209	11.1	5	4.8	17	5.3	29	9.6	36	10.4	70	16.3	52	13.4	64	16.9
	140-150未満	129	6.8	2	1.9	10	3.1	20	6.6	19	5.5	46	10.7	32	8.3	39	10.3
	150-160未満	110	5.8	3	2.9	7	2.2	7	2.3	34	9.9	34	7.9	25	6.5	31	8.2
	160-170未満	74	3.9	1	1.0	5	1.6	11	3.6	22	6.4	23	5.3	12	3.1	14	3.7
	170-180未満	59	3.1	1	1.0	5	1.6	5	1.7	15	4.3	25	5.8	8	2.1	17	4.5
	180-190未満	31	1.6	0	0.0	0	0.0	1	0.3	9	2.6	14	3.3	7	1.8	9	2.4
	190-200未満	17	0.9	0	0.0	0	0.0	0	0.0	6	1.7	6	1.4	5	1.3	5	1.3
	200-210未満	11	0.6	0	0.0	0	0.0	1	0.3	4	1.2	5	1.2	1	0.3	2	0.5
	210-220未満	7	0.4	0	0.0	0	0.0	0	0.0	2	0.6	4	0.9	1	0.3	5	1.3
	220mg/dl以上	12	0.6	0	0.0	0	0.0	2	0.7	1	0.3	6	1.4	3	0.8	3	0.8
	平均値 (mg/dl) 標準偏差	120.2 32.4		97.2 27.0		101.1 25.8		112.5 30.6		128.8 31.0		135.0 33.0		124.3 27.9		130.9 30.8	123.0 26.5

(22年)

Friedewald の式 LDL コレステロール = 総コレステロール - HDL コレステロール - 中性脂肪 / 5
(中性脂肪 (トリグリセライド) が 400mg/dl 未満の場合。)

注) コレステロールを下げる薬または中性脂肪 (トリグリセライド) を下げる薬服用者除外

(参考) 第35表の2 血清LDL- コレステロール値(直接測定法)の分布(性・年齢階級別) [服薬者除外]

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,382	100.0	75	100.0	161	100.0	163	100.0	224	100.0	388	100.0	371	100.0	343	100.0
	40mg/dl未満	3	0.2	0	0.0	0	0.0	0	0.0	0	0.0	2	0.5	1	0.3	1	0.5
	40-49	8	0.6	0	0.0	1	0.6	1	0.6	3	1.3	0	0.0	3	0.8	2	0.5
	50-59	20	1.4	1	1.3	1	0.6	0	0.0	2	0.9	12	3.1	4	1.1	6	0.9
	60-69	34	2.5	5	6.7	3	1.9	1	0.6	3	1.3	6	1.5	16	4.3	9	2.6
	70-79	64	4.6	6	8.0	10	6.2	1	0.6	7	3.1	22	5.7	18	4.9	16	4.7
	80-89	96	6.9	5	6.7	14	8.7	11	6.7	9	4.0	22	5.7	35	9.4	22	6.4
	90-99	137	9.9	10	13.3	21	13.0	7	4.3	19	8.5	32	8.2	48	12.9	33	9.6
	100-109	187	13.5	14	18.7	14	8.7	20	12.3	19	8.5	57	14.7	63	17.0	61	17.8
	110-119	183	13.2	12	16.0	21	13.0	19	11.7	26	11.6	44	11.3	61	16.4	47	13.7
	120-129	170	12.3	6	8.0	11	6.8	25	15.3	39	17.4	49	12.6	40	10.8	41	12.0
	130-139	157	11.4	6	8.0	19	11.8	19	11.7	23	10.3	58	14.9	32	8.6	47	13.7
	140-149	131	9.5	4	5.3	16	9.9	23	14.1	24	10.7	40	10.3	24	6.5	24	7.0
	150-159	72	5.2	3	4.0	10	6.2	12	7.4	21	9.4	18	4.6	8	2.2	14	4.1
	160-169	50	3.6	1	1.3	10	6.2	8	4.9	9	4.0	13	3.4	9	2.4	10	2.9
	170-179	31	2.2	2	2.7	6	3.7	4	2.5	10	4.5	5	1.3	4	1.1	6	1.7
	180-189	22	1.6	0	0.0	3	1.9	6	3.7	4	1.8	5	1.3	4	1.1	3	0.9
	190-199	7	0.5	0	0.0	1	0.6	1	0.6	2	0.9	2	0.5	1	0.3	0	0.0
	200-209	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	1	0.3
	210-219	5	0.4	0	0.0	0	0.0	3	1.8	2	0.9	0	0.0	0	0.0	0	0.0
	220mg/dl以上	4	0.3	0	0.0	0	0.0	2	1.2	2	0.9	0	0.0	0	0.0	0	0.0
平均値(mg/dl) 標準偏差		118.4 30.0		107.6 26.8		119.9 30.8		130.4 31.0		126.7 31.5		117.5 29.1		110.5 26.8		115.1 27.4	108.4 26.0
女 性	総数	1,911	100.0	105	100.0	318.0	100.0	306	100.0	354	100.0	439	100.0	389	100.0	384	100.0
	40mg/dl未満	3	0.2	0	0.0	2	0.6	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0
	40-49	8	0.4	2	1.9	1	0.3	3	1.0	0	0.0	1	0.2	1	0.3	2	0.5
	50-59	17	0.9	2	1.9	7	2.2	3	1.0	1	0.3	3	0.7	1	0.3	3	0.8
	60-69	30	1.6	7	6.7	11	3.5	5	1.6	1	0.3	3	0.7	3	0.8	2	0.5
	70-79	96	5.0	19	18.1	30	9.4	18	5.9	8	2.3	7	1.6	14	3.6	10	2.6
	80-89	160	8.4	17	16.2	50	15.7	26	8.5	24	6.8	17	3.9	26	6.7	22	5.7
	90-99	203	10.6	11	10.5	59	18.6	37	12.1	28	7.9	28	6.4	40	10.3	25	6.5
	100-109	231	12.1	13	12.4	43	13.5	48	15.7	39	11.0	45	10.3	43	11.1	45	11.7
	110-119	250	13.1	9	8.6	37	11.6	46	15.0	49	13.8	42	9.6	67	17.2	49	12.8
	120-129	223	11.7	9	8.6	27	8.5	26	8.5	42	11.9	63	14.4	56	14.4	42	10.9
	130-139	196	10.3	5	4.8	13	4.1	32	10.5	40	11.3	65	14.8	41	10.5	53	13.8
	140-149	160	8.4	3	2.9	14	4.4	28	9.2	37	10.5	41	9.3	37	9.5	33	8.6
	150-159	123	6.4	2	1.9	10	3.1	12	3.9	25	7.1	45	10.3	29	7.5	46	12.0
	160-169	75	3.9	4	3.8	4	1.3	7	2.3	24	6.8	23	5.2	13	3.3	16	4.2
	170-179	55	2.9	2	1.9	5	1.6	9	2.9	16	4.5	19	4.3	4	1.0	10	2.6
	180-189	35	1.8	0	0.0	5	1.6	1	0.3	9	2.5	14	3.2	6	1.5	10	2.6
	190-199	17	0.9	0	0.0	0	0.0	1	0.3	4	1.1	9	2.1	3	0.8	5	1.3
	200-209	13	0.7	0	0.0	0	0.0	1	0.3	4	1.1	5	1.1	3	0.8	6	1.6
	210-219	5	0.3	0	0.0	0	0.0	0	0.0	3	0.8	2	0.5	0	0.0	1	0.3
	220mg/dl以上	11	0.6	0	0.0	0	0.0	2	0.7	0	0.0	7	1.6	2	0.5	4	1.0
平均値(mg/dl) 標準偏差		120.6 32.2		99.8 29.1		104.0 27.7		115.5 30.6		128.9 30.6		133.4 33.7		121.9 27.9		129.0 32.2	121.2 26.8

注) コレステロールを下げる薬または中性脂肪(トリグリセライド)を下げる薬服用者除外

(22年)

第 36 表 血清クレアチニン値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,597	100.0	75	100.0	162	100.0	175	100.0	259	100.0	462	100.0	464	100.0	424	100.0	271	100.0
	0.4mg/dl未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	0.4-0.49	8	0.5	0	0.0	1	0.6	1	0.6	0	0.0	5	1.1	1	0.2	3	0.7	1	0.4
	0.5-0.59	34	2.1	1	1.3	2	1.2	4	2.3	6	2.3	13	2.8	8	1.7	8	1.9	3	1.1
	0.6-0.69	195	12.2	10	13.3	8	4.9	21	12.0	40	15.4	67	14.5	49	10.6	57	13.4	26	9.6
	0.7-0.79	465	29.1	24	32.0	50	30.9	54	30.9	80	30.9	130	28.1	127	27.4	116	27.4	72	26.6
	0.8-0.89	447	28.0	24	32.0	76	46.9	52	29.7	66	25.5	122	26.4	107	23.1	115	27.1	54	19.9
	0.9-0.99	235	14.7	12	16.0	19	11.7	30	17.1	45	17.4	73	15.8	56	12.1	61	14.4	34	12.5
	1.0-1.19	151	9.5	4	5.3	6	3.7	11	6.3	21	8.1	38	8.2	71	15.3	46	10.8	48	17.7
	1.2-1.39	26	1.6	0	0.0	0	0.0	1	0.6	0	0.0	4	0.9	21	4.5	5	1.2	17	6.3
	1.4-1.59	11	0.7	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	9	1.9	5	1.2	5	1.8
	1.6-1.79	6	0.4	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	5	1.1	3	0.7	3	1.1
	1.8-1.99	6	0.4	0	0.0	0	0.0	1	0.6	0	0.0	2	0.4	3	0.6	0	0.0	3	1.1
	2.0mg/dl以上	13	0.8	0	0.0	0	0.0	0	0.0	1	0.4	5	1.1	7	1.5	5	1.2	5	1.8
	平均値(mg/dl) 標準偏差	0.9 0.3		0.8 0.1		0.8 0.1		0.8 0.1		0.8 0.2		0.9 0.5		0.9 0.3		0.9 0.4		0.9 0.3	
女 性	総数	2,268	100.0	105	100.0	321	100.0	308	100.0	392	100.0	564	100.0	578	100.0	546	100.0	333	100.0
	0.4mg/dl未満	13	0.6	2	1.9	1	0.3	1	0.3	6	1.5	2	0.4	1	0.2	0	0.0	1	0.3
	0.4-0.49	248	10.9	16	15.2	44	13.7	31	10.1	40	10.2	69	12.2	48	8.3	54	9.9	26	7.8
	0.5-0.59	870	38.4	44	41.9	153	47.7	119	38.6	165	42.1	216	38.3	173	29.9	194	35.5	92	27.6
	0.6-0.69	739	32.6	33	31.4	98	30.5	118	38.3	122	31.1	185	32.8	183	31.7	188	34.4	100	30.0
	0.7-0.79	277	12.2	9	8.6	21	6.5	31	10.1	48	12.2	69	12.2	99	17.1	77	14.1	61	18.3
	0.8-0.89	73	3.2	1	1.0	3	0.9	6	1.9	8	2.0	18	3.2	37	6.4	22	4.0	23	6.9
	0.9-0.99	22	1.0	0	0.0	1	0.3	1	0.3	2	0.5	0	0.0	18	3.1	3	0.5	15	4.5
	1.0-1.19	13	0.6	0	0.0	0	0.0	0	0.0	1	0.3	2	0.4	10	1.7	4	0.7	8	2.4
	1.2-1.39	4	0.2	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	3	0.5	2	0.4	1	0.3
	1.4-1.59	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0
	1.6-1.79	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.3
	1.8-1.99	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	2.0mg/dl以上	7	0.3	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	5	0.9	2	0.4	5	1.5
	平均値(mg/dl) 標準偏差	0.6 0.3		0.6 0.1		0.6 0.1		0.6 0.1		0.6 0.1		0.6 0.5		0.7 0.3		0.6 0.5		0.7 0.4	

(22年)

第 37 表 推算 GFR 値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,597	100.0	75	100.0	162	100.0	175	100.0	259	100.0	462	100.0	464	100.0	424	100.0	271	100.0
	15ml/min/1.73m ² 未満	3	0.2	0	0.0	0	0.0	0	0.0	0	0.0	3	0.6	0	0.0	2	0.5	0	0.0
	15-29	14	0.9	0	0.0	0	0.0	0	0.0	1	0.4	3	0.6	10	2.2	3	0.7	8	3.0
	30-59	206	12.9	0	0.0	0	0.0	3	1.7	16	6.2	58	12.6	129	27.8	75	17.7	92	33.9
	60-89	1,053	65.9	22	29.3	95	58.6	118	67.4	185	71.4	331	71.6	302	65.1	304	71.7	161	59.4
	90ml/min/1.73m ² 以上	321	20.1	53	70.7	67	41.4	54	30.9	57	22.0	67	14.5	23	5.0	40	9.4	10	3.7
	平均値 標準偏差	76.8 17.8		98.3 15.7		89.3 13.1		83.9 15.1		79.5 13.9		74.8 16.5		66.9 16.5		71.8 15.8		64.4 16.7	
女 性	総数	2,268	100.0	105	100.0	321	100.0	308	100.0	392	100.0	564	100.0	578	100.0	546	100.0	333	100.0
	15ml/min/1.73m ² 未満	4	0.2	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	2	0.3	2	0.4	2	0.6
	15-29	6	0.3	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	5	0.9	0	0.0	5	1.5
	30-59	226	10.0	0	0.0	2	0.6	4	1.3	19	4.8	53	9.4	148	25.6	80	14.7	101	30.3
	60-89	1,373	60.5	18	17.1	111	34.6	197	64.0	269	68.6	405	71.8	373	64.5	391	71.6	202	60.7
	90ml/min/1.73m ² 以上	659	29.1	87	82.9	208	64.8	107	34.7	104	26.5	103	18.3	50	8.7	73	13.4	23	6.9
	平均値 標準偏差	81.3 18.3		107.0 19.7		95.9 14.8		86.6 14.2		82.2 14.8		77.5 14.5		68.8 15.4		74.0 14.1		66.5 16.0	

(22年)

推算 GFR (ml/min/1.73m²) = 194 × 血清クレアチニン値 (mg/dl)^{-1.094} × 年齢 (歳)^{-0.287} (男性)
 = 194 × 血清クレアチニン値 (mg/dl)^{-1.094} × 年齢 (歳)^{-0.287} × 0.739 (女性)

注) CKD ガイドライン 2009 年版の推算式

第38表 血清鉄の分布（性・年齢階級別）〔妊婦・服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,583	100.0	75	100.0	162	100.0	175	100.0	259	100.0	459	100.0	453	100.0	418	100.0	264	100.0
	40 μ g/dl未満	53	3.3	2	2.7	4	2.5	4	2.3	5	1.9	14	3.1	24	5.3	14	3.3	17	6.4
	40-59	173	10.9	9	12.0	17	10.5	22	12.6	21	8.1	36	7.8	68	15.0	38	9.1	45	17.0
	60-79	344	21.7	11	14.7	33	20.4	35	20.0	55	21.2	93	20.3	117	25.8	91	21.8	73	27.7
	80-99	397	25.1	22	29.3	40	24.7	38	21.7	67	25.9	117	25.5	113	24.9	111	26.6	59	22.3
	100-119	293	18.5	15	20.0	30	18.5	43	24.6	44	17.0	93	20.3	68	15.0	80	19.1	39	14.8
	120-139	163	10.3	9	12.0	26	16.0	19	10.9	28	10.8	52	11.3	29	6.4	42	10.0	15	5.7
	140-159	72	4.5	1	1.3	7	4.3	5	2.9	21	8.1	23	5.0	15	3.3	16	3.8	10	3.8
	160-179	40	2.5	4	5.3	2	1.2	4	2.3	6	2.3	16	3.5	8	1.8	13	3.1	3	1.1
	180-199	27	1.7	0	0.0	3	1.9	3	1.7	8	3.1	6	1.3	7	1.5	8	1.9	2	0.8
	200 μ g/dl以上	21	1.3	2	2.7	0	0.0	2	1.1	4	1.5	9	2.0	4	0.9	5	1.2	1	0.4
	平均値(μ g/dl) 標準偏差	94.7 37.0		96.7 36.9		95.0 32.4		94.9 35.0		100.0 36.6		99.7 40.1		86.3 35.0		95.6 37.4		82.7 32.6	
女 性	総数	2,215	100.0	100	100.0	309	100.0	303	100.0	386	100.0	559	100.0	558	100.0	535	100.0	321	100.0
	40 μ g/dl未満	256	11.6	10	10.0	66	21.4	87	28.7	31	8.0	24	4.3	38	6.8	26	4.9	27	8.4
	40-59	500	22.6	16	16.0	56	18.1	57	18.8	77	19.9	135	24.2	159	28.5	150	28.0	90	28.0
	60-79	694	31.3	26	26.0	70	22.7	69	22.8	128	33.2	206	36.9	195	34.9	188	35.1	116	36.1
	80-99	418	18.9	13	13.0	57	18.4	37	12.2	79	20.5	126	22.5	106	19.0	109	20.4	59	18.4
	100-119	199	9.0	17	17.0	34	11.0	23	7.6	38	9.8	46	8.2	41	7.3	48	9.0	19	5.9
	120-139	87	3.9	8	8.0	12	3.9	18	5.9	22	5.7	17	3.0	10	1.8	6	1.1	6	1.9
	140-159	38	1.7	6	6.0	9	2.9	4	1.3	7	1.8	5	0.9	7	1.3	6	1.1	4	1.2
	160-179	14	0.6	2	2.0	3	1.0	5	1.7	2	0.5	0	0.0	2	0.4	2	0.4	0	0.0
	180-199	7	0.3	2	2.0	1	0.3	2	0.7	2	0.5	0	0.0	0	0.0	0	0.0	0	0.0
	200 μ g/dl以上	2	0.1	0	0.0	1	0.3	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	平均値(μ g/dl) 標準偏差	72.1 29.7		85.3 37.7		70.4 35.7		65.3 38.5		76.5 28.7		73.2 22.7		70.2 24.1		71.6 23.0		68.6 23.9	

注) 妊婦と貧血治療薬服用者除外

(22年)

第39表 総鉄結合能（TIBC）の分布（性・年齢階級別）〔妊婦・服薬者除外〕

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,583	100.0	75	100.0	162	100.0	175	100.0	259	100.0	459	100.0	453	100.0	418	100.0	264	100.0
	200 μ g/dl未満	1	0.1	0	0.0	1	0.6	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	200-249	29	1.8	0	0.0	1	0.6	1	0.6	1	0.4	4	0.9	22	4.9	13	3.1	12	4.5
	250-299	370	23.4	13	17.3	35	21.6	22	12.6	42	16.2	120	26.1	138	30.5	106	25.4	95	36.0
	300-349	720	45.5	47	62.7	77	47.5	91	52.0	129	49.8	194	42.3	182	40.2	182	43.5	95	36.0
	350-399	364	23.0	13	17.3	43	26.5	50	28.6	68	26.3	107	23.3	83	18.3	91	21.8	43	16.3
	400-449	84	5.3	1	1.3	3	1.9	8	4.6	16	6.2	29	6.3	27	6.0	22	5.3	18	6.8
	450-499	11	0.7	1	1.3	1	0.6	3	1.7	1	0.4	4	0.9	1	0.2	3	0.7	1	0.4
	500 μ g/dl以上	4	0.3	0	0.0	1	0.6	0	0.0	2	0.8	1	0.2	0	0.0	1	0.2	0	0.0
	平均値(μ g/dl) 標準偏差	328.9 44.6		327.1 36.3		331.2 44.2		337.8 38.2		336.5 43.1		330.4 45.8		319.3 46.2		326.2 45.8		317.3 49.0	
女 性	総数	2,215	100.0	100	100.0	309	100.0	303	100.0	386	100.0	559	100.0	558	100.0	535	100.0	321	100.0
	200 μ g/dl未満	3	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	2	0.4	2	0.4	1	0.3
	200-249	25	1.1	1	1.0	2	0.6	0	0.0	3	0.8	8	1.4	11	2.0	7	1.3	9	2.8
	250-299	374	16.9	7	7.0	35	11.3	29	9.6	67	17.4	109	19.5	127	22.8	119	22.2	69	21.5
	300-349	987	44.6	48	48.0	104	33.7	97	32.0	187	48.4	295	52.8	256	45.9	261	48.8	148	46.1
	350-399	563	25.4	31	31.0	105	34.0	88	29.0	96	24.9	117	20.9	126	22.6	113	21.1	75	23.4
	400-449	203	9.2	11	11.0	44	14.2	74	24.4	25	6.5	20	3.6	29	5.2	24	4.5	15	4.7
	450-499	55	2.5	2	2.0	17	5.5	15	5.0	6	1.6	8	1.4	7	1.3	9	1.7	4	1.2
	500 μ g/dl以上	5	0.2	0	0.0	2	0.6	0	0.0	2	0.5	1	0.2	0	0.0	0	0.0	0	0.0
	平均値(μ g/dl) 標準偏差	340.4 48.0		350.9 42.0		359.2 52.1		366.7 54.1		337.2 44.2		328.9 40.5		327.8 43.2		327.9 42.8		327.6 43.4	

注) 妊婦と貧血治療薬服用者除外

(22年)

第 40 表 AST (GOT) の分布 (性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,597	100.0	75	100.0	162	100.0	175	100.0	259	100.0	462	100.0	464	100.0	424	100.0	271	100.0
	5IU/L未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	5-9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	10-19	360	22.5	26	34.7	41	25.3	51	29.1	68	26.3	77	16.7	97	20.9	84	19.8	50	18.5
	20-29	859	53.8	34	45.3	88	54.3	85	48.6	144	55.6	272	58.9	236	50.9	230	54.2	140	51.7
	30-39	263	16.5	11	14.7	15	9.3	29	16.6	36	13.9	75	16.2	97	20.9	76	17.9	64	23.6
	40-49	66	4.1	4	5.3	7	4.3	7	4.0	8	3.1	16	3.5	24	5.2	17	4.0	12	4.4
	50IU/L以上	49	3.1	0	0.0	11	6.8	3	1.7	3	1.2	22	4.8	10	2.2	17	4.0	5	1.8
	平均値 (IU/L) 標準偏差	26.3 14.0		23.7 7.6		26.4 12.8		25.1 9.8		24.4 7.6		27.8 21.1		26.6 9.8		26.9 11.6		26.6 8.5	
女 性	総数	2,268	100.0	105	100.0	321	100.0	308	100.0	392	100.0	564	100.0	578	100.0	546	100.0	333	100.0
	5IU/L未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	5-9	1	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	10-19	903	39.8	79	75.2	230	71.7	207	67.2	126	32.1	141	25.0	120	20.8	133	24.4	70	21.0
	20-29	1,138	50.2	23	21.9	78	24.3	88	28.6	229	58.4	342	60.6	378	65.4	333	61.0	221	66.4
	30-39	162	7.1	2	1.9	9	2.8	7	2.3	27	6.9	55	9.8	62	10.7	58	10.6	33	9.9
	40-49	31	1.4	0	0.0	1	0.3	1	0.3	6	1.5	16	2.8	7	1.2	10	1.8	4	1.2
	50IU/L以上	33	1.5	1	1.0	3	0.9	4	1.3	4	1.0	10	1.8	11	1.9	12	2.2	5	1.5
	平均値 (IU/L) 標準偏差	22.3 8.7		18.0 8.0		18.8 8.2		18.9 7.2		22.8 7.5		24.4 9.5		24.7 8.4		24.5 9.6		24.5 7.7	

(22年)

第 41 表 ALT (GPT) の分布 (性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,597	100.0	75	100.0	162	100.0	175	100.0	259	100.0	462	100.0	464	100.0	424	100.0	271	100.0
	5IU/L未満	1	0.1	0	0.0	0	0.0	1	0.6	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	5-9	27	1.7	0	0.0	2	1.2	3	1.7	1	0.4	5	1.1	16	3.4	10	2.4	8	3.0
	10-19	608	38.1	29	38.7	49	30.2	49	28.0	73	28.2	173	37.4	235	50.6	177	41.7	147	54.2
	20-29	508	31.8	20	26.7	45	27.8	51	29.1	97	37.5	162	35.1	133	28.7	145	34.2	78	28.8
	30-39	227	14.2	12	16.0	17	10.5	31	17.7	52	20.1	67	14.5	48	10.3	52	12.3	25	9.2
	40-49	97	6.1	3	4.0	15	9.3	19	10.9	21	8.1	22	4.8	17	3.7	15	3.5	7	2.6
	50IU/L以上	129	8.1	11	14.7	34	21.0	21	12.0	15	5.8	33	7.1	15	3.2	25	5.9	6	2.2
	平均値 (IU/L) 標準偏差	27.0 17.8		30.2 19.0		35.7 30.0		32.6 22.4		27.4 12.7		26.5 15.9		21.6 11.0		24.6 14.5		20.5 9.6	
女 性	総数	2,268	100.0	105	100.0	321	100.0	308	100.0	392	100.0	564	100.0	578	100.0	546	100.0	333	100.0
	5IU/L未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	5-9	126	5.6	18	17.1	40	12.5	27	8.8	7	1.8	8	1.4	26	4.5	11	2.0	21	6.3
	10-19	1,408	62.1	69	65.7	224	69.8	224	72.7	212	54.1	303	53.7	376	65.1	318	58.2	228	68.5
	20-29	513	22.6	11	10.5	40	12.5	42	13.6	118	30.1	165	29.3	137	23.7	162	29.7	69	20.7
	30-39	130	5.7	6	5.7	8	2.5	7	2.3	35	8.9	53	9.4	21	3.6	30	5.5	9	2.7
	40-49	31	1.4	0	0.0	4	1.2	2	0.6	5	1.3	14	2.5	6	1.0	7	1.3	1	0.3
	50IU/L以上	60	2.6	1	1.0	5	1.6	6	1.9	15	3.8	21	3.7	12	2.1	18	3.3	5	1.5
	平均値 (IU/L) 標準偏差	19.2 13.1		15.2 11.6		16.6 15.9		16.5 11.0		21.3 11.9		22.3 15.2		18.6 10.1		20.8 14.3		17.3 8.3	

(22年)

第42表 γ -GTの分布(性・年齢階級別)

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総数	1,597	100.0	75	100.0	162	100.0	175	100.0	259	100.0	462	100.0	464	100.0	424	100.0
	10IU/L未満	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	0	0.0
	10-19	311	19.5	24	32.0	39	24.1	24	13.7	34	13.1	66	14.3	124	26.7	83	19.6
	20-29	433	27.1	27	36.0	47	29.0	38	21.7	64	24.7	115	24.9	142	30.6	127	30.0
	30-39	265	16.6	11	14.7	22	13.6	23	13.1	41	15.8	89	19.3	79	17.0	74	17.5
	40-49	127	8.0	4	5.3	13	8.0	19	10.9	16	6.2	38	8.2	37	8.0	40	9.4
	50-59	96	6.0	5	6.7	6	3.7	13	7.4	23	8.9	29	6.3	20	4.3	19	4.5
	60-69	81	5.1	0	0.0	6	3.7	14	8.0	22	8.5	28	6.1	11	2.4	15	3.5
	70-79	61	3.8	2	2.7	7	4.3	13	7.4	13	5.0	20	4.3	6	1.3	15	3.5
	80IU/L以上	221	13.8	2	2.7	22	13.6	31	17.7	46	17.8	77	16.7	43	9.3	51	12.0
女性	総数	2,268	100.0	105	100.0	321	100.0	308	100.0	392	100.0	564	100.0	578	100.0	546	100.0
	10IU/L未満	81	3.6	16	15.2	25	7.8	12	3.9	7	1.8	4	0.7	17	2.9	6	1.1
	10-19	1,278	56.3	67	63.8	223	69.5	193	62.7	183	46.7	273	48.4	339	58.7	281	51.5
	20-29	514	22.7	15	14.3	41	12.8	60	19.5	93	23.7	162	28.7	143	24.7	160	29.3
	30-39	148	6.5	3	2.9	16	5.0	14	4.5	36	9.2	40	7.1	39	6.7	34	6.2
	40-49	76	3.4	1	1.0	8	2.5	7	2.3	19	4.8	29	5.1	12	2.1	18	3.3
	50-59	56	2.5	0	0.0	3	0.9	10	3.2	19	4.8	12	2.1	12	2.1	15	2.7
	60-69	24	1.1	1	1.0	2	0.6	3	1.0	5	1.3	9	1.6	4	0.7	8	1.5
	70-79	18	0.8	1	1.0	1	0.3	1	0.3	7	1.8	5	0.9	3	0.5	4	0.7
	80IU/L以上	73	3.2	1	1.0	2	0.6	8	2.6	23	5.9	30	5.3	9	1.6	20	3.7
	平均値(IU/L)	53.9		30.5		50.4		58.7		59.5		67.2		40.8		48.9	
	標準偏差	94.6		23.9		64.1		56.2		91.9		142.5		52.3		67.9	

(22年)

第43表 血清尿酸値の分布(性・年齢階級別)

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総数	1,597	100.0	75	100.0	162	100.0	175	100.0	259	100.0	462	100.0	464	100.0	424	100.0
	2.0mg/dl未満	3	0.2	0	0.0	0	0.0	0	0.0	1	0.4	1	0.2	1	0.2	1	0.0
	2.0-2.9	20	1.3	0	0.0	1	0.6	2	1.1	2	0.8	5	1.1	10	2.2	8	1.9
	3.0-3.9	83	5.2	2	2.7	5	3.1	5	2.9	12	4.6	33	7.1	26	5.6	25	5.9
	4.0-4.9	284	17.8	8	10.7	25	15.4	31	17.7	43	16.6	83	18.0	94	20.3	80	18.9
	5.0-5.9	479	30.0	26	34.7	50	30.9	44	25.1	81	31.3	140	30.3	138	29.7	123	29.0
	6.0-6.9	440	27.6	22	29.3	47	29.0	46	26.3	82	31.7	123	26.6	120	25.9	109	25.7
	7.0-7.9	200	12.5	11	14.7	26	16.0	32	18.3	29	11.2	49	10.6	53	11.4	54	12.7
	8.0-8.9	62	3.9	4	5.3	7	4.3	9	5.1	7	2.7	21	4.5	14	3.0	20	4.7
	9.0-9.9	19	1.2	2	2.7	1	0.6	6	3.4	2	0.8	3	0.6	5	1.1	3	0.7
	10mg/dl以上	7	0.4	0	0.0	0	0.0	0	0.0	0	0.0	4	0.9	3	0.6	1	0.2
	平均値(μ g/dl)	5.8		6.1		6.0		6.1		5.8		5.8		5.7		5.8	
女性	総数	2,268	100.0	105	100.0	321	100.0	308	100.0	392	100.0	564	100.0	578	100.0	546	100.0
	2.0mg/dl未満	13	0.6	0	0.0	2	0.6	4	1.3	0	0.0	4	0.7	3	0.5	4	0.7
	2.0-2.9	148	6.5	8	7.6	32	10.0	36	11.7	20	5.1	24	4.3	28	4.8	28	5.1
	3.0-3.9	654	28.8	33	31.4	122	38.0	113	36.7	109	27.8	140	24.8	137	23.7	140	25.6
	4.0-4.9	870	38.4	43	41.0	121	37.7	121	39.3	158	40.3	217	38.5	210	36.3	206	37.7
	5.0-5.9	423	18.7	19	18.1	33	10.3	29	9.4	79	20.2	131	23.2	132	22.8	115	21.1
	6.0-6.9	120	5.3	2	1.9	8	2.5	3	1.0	20	5.1	38	6.7	49	8.5	37	6.8
	7.0-7.9	32	1.4	0	0.0	3	0.9	2	0.6	4	1.0	7	1.2	16	2.8	10	1.8
	8.0-8.9	7	0.3	0	0.0	0	0.0	0	0.0	2	0.5	3	0.5	2	0.3	5	0.9
	9.0-9.9	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2
	10mg/dl以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	平均値(μ g/dl)	4.4		4.2		4.0		4.0		4.4		4.5		4.6		4.5	
	標準偏差	1.1		0.9		0.9		0.9		1.0		1.1		1.1		1.1	

(22年)

第44表 メタボリックシンドローム（内臓脂肪症候群）の状況（性・年齢階級別）

	総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	3,829	100.0	175	100.0	469	100.0	483	100.0	649	100.0	1,022	100.0	1,031	100.0	2,588	100.0	963	100.0	597	100.0
	707	18.5	0	0.0	11	2.3	43	8.9	110	16.9	252	24.7	291	28.2	527	20.4	259	26.9	169	28.3
	494	12.9	7	4.0	42	9.0	44	9.1	95	14.6	152	14.9	154	14.9	352	13.6	128	13.3	93	15.6
	2,628	68.6	168	96.0	416	88.7	396	82.0	444	68.4	618	60.5	586	56.8	1,709	66.0	576	59.8	335	56.1
男 性	1,586	100.0	75	100.0	161	100.0	175	100.0	257	100.0	460	100.0	458	100.0	1,082	100.0	420	100.0	268	100.0
	443	27.9	0	0.0	11	6.8	34	19.4	79	30.7	152	33.0	167	36.5	334	30.9	154	36.7	98	36.6
	331	20.9	7	9.3	33	20.5	27	15.4	59	23.0	109	23.7	96	21.0	236	21.8	89	21.2	55	20.5
	812	51.2	68	90.7	117	72.7	114	65.1	119	46.3	199	43.3	195	42.6	512	47.3	177	42.1	115	42.9
女 性	2,243	100.0	100	100.0	308	100.0	308	100.0	392	100.0	562	100.0	573	100.0	1,506	100.0	543	100.0	329	100.0
	264	11.8	0	0.0	0	0.0	9	2.9	31	7.9	100	17.8	124	21.6	193	12.8	105	19.3	71	21.6
	163	7.3	0	0.0	9	2.9	17	5.5	36	9.2	43	7.7	58	10.1	116	7.7	39	7.2	38	11.6
	1,816	81.0	100	100.0	299	97.1	282	91.6	325	82.9	419	74.6	391	68.2	1,197	79.5	399	73.5	220	66.9

(22年)

注) 妊婦除外，ヘモグロビン A1c ≥ 5.5% の場合。項目の基準は p.18 参照

第45表の1 腹囲区分別、血中脂質、血圧、血糖のいずれかのリスクを有する割合（性・年齢階級別）

		腹 囲	総 数		リ ス ク 保 有 数							
					0		1つ		2つ		3つ	
			人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	<85cm(男)/90cm(女)	2,475	100.0	908	36.7	939	37.9	505	20.4	123	5.0
		≥85cm(男)/90cm(女)	1,354	100.0	153	11.3	494	36.5	479	35.4	228	16.8
	20-29歳	<85cm(男)/90cm(女)	150	100.0	129	86.0	18	12.0	3	2.0	0	0.0
		≥85cm(男)/90cm(女)	25	100.0	18	72.0	7	28.0	0	0.0	0	0.0
	30-39歳	<85cm(男)/90cm(女)	382	100.0	287	75.1	87	22.8	8	2.1	0	0.0
		≥85cm(男)/90cm(女)	87	100.0	34	39.1	42	48.3	10	11.5	1	1.1
	40-49歳	<85cm(男)/90cm(女)	364	100.0	198	54.4	149	40.9	16	4.4	1	0.3
		≥85cm(男)/90cm(女)	119	100.0	32	26.9	44	37.0	36	30.3	7	5.9
	50-59歳	<85cm(男)/90cm(女)	423	100.0	145	34.3	199	47.0	72	17.0	7	1.7
		≥85cm(男)/90cm(女)	226	100.0	21	9.3	95	42.0	76	33.6	34	15.0
	60-69歳	<85cm(男)/90cm(女)	585	100.0	99	16.9	258	44.1	187	32.0	41	7.0
		≥85cm(男)/90cm(女)	437	100.0	33	7.6	152	34.8	166	38.0	86	19.7
	70歳以上	<85cm(男)/90cm(女)	571	100.0	50	8.8	228	39.9	219	38.4	74	13.0
		≥85cm(男)/90cm(女)	460	100.0	15	3.3	154	33.5	191	41.5	100	21.7
	(再掲) 40-74歳	<85cm(男)/90cm(女)	1,616	100.0	469	29.0	699	43.3	369	22.8	79	4.9
		≥85cm(男)/90cm(女)	972	100.0	93	9.6	352	36.2	355	36.5	172	17.7
	(再掲) 65-74歳	<85cm(男)/90cm(女)	553	100.0	75	13.6	229	41.4	191	34.5	58	10.5
		≥85cm(男)/90cm(女)	410	100.0	23	5.6	128	31.2	166	40.5	93	22.7
	(再掲) 75歳以上	<85cm(男)/90cm(女)	327	100.0	23	7.0	135	41.3	125	38.2	44	13.5
		≥85cm(男)/90cm(女)	270	100.0	8	3.0	93	34.4	114	42.2	55	20.4
男性	総 数	<85cm	715	100.0	202	28.3	307	42.9	170	23.8	36	5.0
		≥85cm	871	100.0	97	11.1	331	38.0	304	34.9	139	16.0
	20-29歳	<85cm	54	100.0	42	77.8	10	18.5	2	3.7	0	0.0
		≥85cm	21	100.0	14	66.7	7	33.3	0	0.0	0	0.0
	30-39歳	<85cm	99	100.0	56	56.6	41	41.4	2	2.0	0	0.0
		≥85cm	62	100.0	18	29.0	33	53.2	10	16.1	1	1.6
	40-49歳	<85cm	93	100.0	43	46.2	43	46.2	6	6.5	1	1.1
		≥85cm	82	100.0	21	25.6	27	32.9	28	34.1	6	7.3
	50-59歳	<85cm	105	100.0	27	25.7	57	54.3	18	17.1	3	2.9
		≥85cm	152	100.0	14	9.2	59	38.8	50	32.9	29	19.1
	60-69歳	<85cm	177	100.0	23	13.0	80	45.2	61	34.5	13	7.3
		≥85cm	283	100.0	22	7.8	109	38.5	102	36.0	50	17.7
	70歳以上	<85cm	187	100.0	11	5.9	76	40.6	81	43.3	19	10.2
		≥85cm	271	100.0	8	3.0	96	35.4	114	42.1	53	19.6
	(再掲) 40-74歳	<85cm	451	100.0	96	21.3	209	46.3	121	26.8	25	5.5
		≥85cm	631	100.0	61	9.7	236	37.4	226	35.8	108	17.1
	(再掲) 65-74歳	<85cm	164	100.0	16	9.8	68	41.5	64	39.0	16	9.8
		≥85cm	256	100.0	13	5.1	89	34.8	102	39.8	52	20.3
	(再掲) 75歳以上	<85cm	111	100.0	8	7.2	47	42.3	45	40.5	11	9.9
		≥85cm	157	100.0	4	2.5	55	35.0	68	43.3	30	19.1
女性	総 数	<90cm	1,760	100.0	706	40.1	632	35.9	335	19.0	87	4.9
		≥90cm	483	100.0	56	11.6	163	33.7	175	36.2	89	18.4
	20-29歳	<90cm	96	100.0	87	90.6	8	8.3	1	1.0	0	0.0
		≥90cm	4	100.0	4	100.0	0	0.0	0	0.0	0	0.0
	30-39歳	<90cm	283	100.0	231	81.6	46	16.3	6	2.1	0	0.0
		≥90cm	25	100.0	16	64.0	9	36.0	0	0.0	0	0.0
	40-49歳	<90cm	271	100.0	155	57.2	106	39.1	10	3.7	0	0.0
		≥90cm	37	100.0	11	29.7	17	45.9	8	21.6	1	2.7
	50-59歳	<90cm	318	100.0	118	37.1	142	44.7	54	17.0	4	1.3
		≥90cm	74	100.0	7	9.5	36	48.6	26	35.1	5	6.8
	60-69歳	<90cm	408	100.0	76	18.6	178	43.6	126	30.9	28	6.9
		≥90cm	154	100.0	11	7.1	43	27.9	64	41.6	36	23.4
	70歳以上	<90cm	384	100.0	39	10.2	152	39.6	138	35.9	55	14.3
		≥90cm	189	100.0	7	3.7	58	30.7	77	40.7	47	24.9
	(再掲) 40-74歳	<90cm	1,165	100.0	373	32.0	490	42.1	248	21.3	54	4.6
		≥90cm	341	100.0	32	9.4	116	34.0	129	37.8	64	18.8
	(再掲) 65-74歳	<90cm	389	100.0	59	15.2	161	41.4	127	32.6	42	10.8
		≥90cm	154	100.0	10	6.5	39	25.3	64	41.6	41	26.6
	(再掲) 75歳以上	<90cm	216	100.0	15	6.9	88	40.7	80	37.0	33	15.3
		≥90cm	113	100.0	4	3.5	38	33.6	46	40.7	25	22.1

注) 妊婦除外，ヘモグロビン A1c ≥ 5.5% の場合。リスクの項目は基準は p.18 参照

(22年)

第 45 表の 2 腹囲区分別，血中脂質，血圧，血糖のリスク保有状況〔すべての組み合わせ①血中脂質②血圧③血糖〕（性・年齢階級別）

		腹 囲	総 数		リ ス ク 保 有 数															
					①有②有③有		①有②有③無		①無②有③有		①有②無③有		①有②無③無		①無②有③無		①無②無③有		①無②無③無	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	総 数	<85cm(男)/90cm(女)	2,475	100.0	123	5.0	131	5.3	347	14.0	27	1.1	59	2.4	688	27.8	192	7.8	908	36.7
		≥85cm(男)/90cm(女)	1,354	100.0	228	16.8	143	10.6	314	23.2	22	1.6	25	1.8	414	30.6	55	4.1	153	11.3
	20-29歳	<85cm(男)/90cm(女)	150	100.0	0	0.0	2	1.3	0	0.0	1	0.7	3	2.0	9	6.0	6	4.0	129	86.0
		≥85cm(男)/90cm(女)	25	100.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	7	28.0	0	0.0	18	72.0
	30-39歳	<85cm(男)/90cm(女)	382	100.0	0	0.0	2	0.5	6	1.6	0	0.0	10	2.6	48	12.6	29	7.6	287	75.1
		≥85cm(男)/90cm(女)	87	100.0	1	1.1	5	5.7	5	5.7	0	0.0	4	4.6	30	34.5	8	9.2	34	39.1
	40-49歳	<85cm(男)/90cm(女)	364	100.0	1	0.3	3	0.8	13	3.6	0	0.0	5	1.4	89	24.5	55	15.1	198	54.4
		≥85cm(男)/90cm(女)	119	100.0	7	5.9	13	10.9	22	18.5	1	0.8	3	2.5	36	30.3	5	4.2	32	26.9
	50-59歳	<85cm(男)/90cm(女)	423	100.0	7	1.7	17	4.0	49	11.6	6	1.4	11	2.6	141	33.3	47	11.1	145	34.3
		≥85cm(男)/90cm(女)	226	100.0	34	15.0	24	10.6	52	23.0	0	0.0	5	2.2	79	35.0	11	4.9	21	9.3
	60-69歳	<85cm(男)/90cm(女)	585	100.0	41	7.0	44	7.5	130	22.2	13	2.2	14	2.4	213	36.4	31	5.3	99	16.9
		≥85cm(男)/90cm(女)	437	100.0	86	19.7	43	9.8	115	26.3	8	1.8	8	1.8	125	28.6	19	4.3	33	7.6
	70歳以上	<85cm(男)/90cm(女)	571	100.0	74	13.0	63	11.0	149	26.1	7	1.2	16	2.8	188	32.9	24	4.2	50	8.8
		≥85cm(男)/90cm(女)	460	100.0	100	21.7	58	12.6	120	26.1	13	2.8	5	1.1	137	29.8	12	2.6	15	3.3
	(再掲) 40-74歳	<85cm(男)/90cm(女)	1,616	100.0	79	4.9	93	5.8	253	15.7	23	1.4	37	2.3	519	32.1	143	8.8	469	29.0
		≥85cm(男)/90cm(女)	972	100.0	172	17.7	107	11.0	232	23.9	16	1.6	17	1.7	292	30.0	43	4.4	93	9.6
	(再掲) 65-74歳	<85cm(男)/90cm(女)	553	100.0	58	10.5	50	9.0	128	23.1	13	2.4	15	2.7	186	33.6	28	5.1	75	13.6
		≥85cm(男)/90cm(女)	410	100.0	93	22.7	51	12.4	103	25.1	12	2.9	3	0.7	109	26.6	16	3.9	23	5.6
	(再掲) 75歳以上	<85cm(男)/90cm(女)	327	100.0	44	13.5	34	10.4	88	26.9	3	0.9	9	2.8	112	34.3	14	4.3	23	7.0
		≥85cm(男)/90cm(女)	270	100.0	55	20.4	31	11.5	77	28.5	6	2.2	4	1.5	85	31.5	4	1.5	8	3.0
男 性	総 数	<85cm	715	100.0	36	5.0	40	5.6	123	17.2	7	1.0	18	2.5	258	36.1	31	4.3	202	28.3
		≥85cm	871	100.0	139	16.0	96	11.0	197	22.6	11	1.3	19	2.2	278	31.9	34	3.9	97	11.1
	20-29歳	<85cm	54	100.0	0	0.0	2	3.7	0	0.0	0	0.0	3	5.6	7	13.0	0	0.0	42	77.8
		≥85cm	21	100.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	7	33.3	0	0.0	14	66.7
	30-39歳	<85cm	99	100.0	0	0.0	2	2.0	0	0.0	0	0.0	6	6.1	32	32.3	3	3.0	56	56.6
		≥85cm	62	100.0	1	1.6	5	8.1	5	8.1	0	0.0	4	6.5	24	38.7	5	8.1	18	29.0
	40-49歳	<85cm	93	100.0	1	1.1	2	2.2	4	4.3	0	0.0	3	3.2	31	33.3	9	9.7	43	46.2
		≥85cm	82	100.0	6	7.3	13	15.9	15	18.3	0	0.0	2	2.4	22	26.8	3	3.7	21	25.6
	50-59歳	<85cm	105	100.0	3	2.9	4	3.8	13	12.4	1	1.0	1	1.0	50	47.6	6	5.7	27	25.7
		≥85cm	152	100.0	29	19.1	17	11.2	33	21.7	0	0.0	4	2.6	49	32.2	6	3.9	14	9.2
	60-69歳	<85cm	177	100.0	13	7.3	14	7.9	43	24.3	4	2.3	2	1.1	72	40.7	6	3.4	23	13.0
		≥85cm	283	100.0	50	17.7	29	10.2	68	24.0	5	1.8	6	2.1	89	31.4	14	4.9	22	7.8
	70歳以上	<85cm	187	100.0	19	10.2	16	8.6	63	33.7	2	1.1	3	1.6	66	35.3	7	3.7	11	5.9
		≥85cm	271	100.0	53	19.6	32	11.8	76	28.0	6	2.2	3	1.1	87	32.1	6	2.2	8	3.0
	(再掲) 40-74歳	<85cm	451	100.0	25	5.5	27	6.0	87	19.3	7	1.6	6	1.3	179	39.7	24	5.3	96	21.3
		≥85cm	631	100.0	108	17.1	73	11.6	145	23.0	8	1.3	13	2.1	196	31.1	27	4.3	61	9.7
	(再掲) 65-74歳	<85cm	164	100.0	16	9.8	12	7.3	47	28.7	5	3.0	0	0.0	62	37.8	6	3.7	16	9.8
		≥85cm	256	100.0	52	20.3	28	10.9	69	27.0	5	2.0	2	0.8	75	29.3	12	4.7	13	5.1
	(再掲) 75歳以上	<85cm	111	100.0	11	9.9	9	8.1	36	32.4	0	0.0	3	2.7	40	36.0	4	3.6	8	7.2
		≥85cm	157	100.0	30	19.1	18	11.5	47	29.9	3	1.9	2	1.3	51	32.5	2	1.3	4	2.5
女 性	総 数	<90cm	1,760	100.0	87	4.9	91	5.2	224	12.7	20	1.1	41	2.3	430	24.4	161	9.1	706	40.1
		≥90cm	483	100.0	89	18.4	47	9.7	117	24.2	11	2.3	6	1.2	136	28.2	21	4.3	56	11.6
	20-29歳	<90cm	96	100.0	0	0.0	0	0.0	0	0.0	1	1.0	0	0.0	2	2.1	6	6.3	87	90.6
		≥90cm	4	100.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	4	100.0
	30-39歳	<90cm	283	100.0	0	0.0	0	0.0	6	2.1	0	0.0	4	1.4	16	5.7	26	9.2	231	81.6
		≥90cm	25	100.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	6	24.0	3	12.0	16	64.0
	40-49歳	<90cm	271	100.0	0	0.0	1	0.4	9	3.3	0	0.0	2	0.7	58	21.4	46	17.0	155	57.2
		≥90cm	37	100.0	1	2.7	0	0.0	7	18.9	1	2.7	1	2.7	14	37.8	2	5.4	11	29.7
	50-59歳	<90cm	318	100.0	4	1.3	13	4.1	36	11.3	5	1.6	10	3.1	91	28.6	41	12.9	118	37.1
		≥90cm	74	100.0	5	6.8	7	9.5	19	25.7	0	0.0	1	1.4	30	40.5	5	6.8	7	9.5
	60-69歳	<90cm	408	100.0	28	6.9	30	7.4	87	21.3	9	2.2	12	2.9	141	34.6	25	6.1	76	18.6
		≥90cm	154	100.0	36	23.4	14	9.1	47	30.5	3	1.9	2	1.3	36	23.4	5	3.2	11	7.1
	70歳以上	<90cm	384	100.0	55	14.3	47	12.2	86	22.4	5	1.3	13	3.4	122	31.8	17	4.4	39	10.2
		≥90cm	189	100.0	47	24.9	26	13.8	44	23.3	7	3.7	2	1.1	50	26.5	6	3.2	7	3.7
	(再掲) 40-74歳	<90cm	1,165	100.0	54	4.6	66	5.7	166	14.2	16	1.4	31	2.7	340	29.2	119	10.2	373	32.0
		≥90cm	341	100.0	64	18.8	34	10.0	87	25.5	8	2.3	4	1.2	96	28.2	16	4.7	32	9.4
	(再掲) 65-74歳	<90cm	389	100.0	42	10.8	38	9.8	81	20.8	8	2.1	15	3.9	124	31.9	22	5.7	59	15.2
		≥90cm	154	100.0	41	26.6	23	14.9	34	22.1	7	4.5	1	0.6	34	22.1	4	2.6	10	6.5
	(再掲) 75歳以上	<90cm	216	100.0	33	15.3	25	11.6	52	24.1	3	1.4	6	2.8	72	33.3	10	4.6	15	6.9
		≥90cm	113	100.0	25	22.1	13	11.5	30	26.5	3	2.7	2	1.8	34	30.1	2	1.8	4	3.5

注) 妊婦除外，ヘモグロビン A1c ≥ 5.5% の場合。リスクの項目は基準は p.18 参照

(22年)

第 46 表 「糖尿病が強く疑われる人」及び「糖尿病の可能性を否定できない人」の状況（性・年齢階級別）

	総 数	20～29歳		30～39歳		40～49歳		50～59歳		60～69歳		70歳以上		(再掲) 40～74歳		(再掲) 65～74歳		(再掲) 75歳以上	
		人数		人数		人数		人数		人数		人数		人数		人数		人数	
		%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%	%
総 数	総 数	3,857	100.0	180	100.0	482	100.0	483	100.0	649	100.0	1,026	100.0	1,037	100.0	2,594	100.0	968	100.0
	糖尿病が強く疑われる人	472	12.2	1	0.6	8	1.7	25	5.2	62	9.6	178	17.3	198	19.1	350	13.5	180	18.6
	(再掲)糖尿病が強く疑われる人のうち服薬者*	238	50.4	1	100.0	2	25.0	7	28.0	28	45.2	94	52.8	106	53.5	180	51.4	104	57.8
	糖尿病の可能性が否定できない人	590	15.3	2	1.1	21	4.4	49	10.1	97	14.9	201	19.6	220	21.2	429	16.5	193	19.9
	上記以外	2,795	72.5	177	98.3	453	94.0	409	84.7	490	75.5	647	63.1	619	59.7	1,815	70.0	595	61.5
男 性	総 数	1,589	100.0	75	100.0	161	100.0	175	100.0	257	100.0	461	100.0	460	100.0	1,085	100.0	422	100.0
	糖尿病が強く疑われる人	264	16.6	0	0.0	5	3.1	14	8.0	40	15.6	102	22.1	103	22.4	202	18.6	100	23.7
	(再掲)糖尿病が強く疑われる人のうち服薬者*	133	50.4	0	0.0	2	40.0	4	28.6	20	50.0	56	54.9	51	49.5	102	50.5	52	52.0
	糖尿病の可能性が否定できない人	231	14.5	0	0.0	4	2.5	14	8.0	35	13.6	80	17.4	98	21.3	168	15.5	84	19.9
	上記以外	1,094	68.8	75	100.0	152	94.4	147	84.0	182	70.8	279	60.5	259	56.3	715	65.9	238	56.4
女 性	総 数	2,268	100.0	105	100.0	321	100.0	308	100.0	392	100.0	565	100.0	577	100.0	1,509	100.0	546	100.0
	糖尿病が強く疑われる人	208	9.2	1	1.0	3	0.9	11	3.6	22	5.6	76	13.5	95	16.5	148	9.8	80	14.7
	(再掲)糖尿病が強く疑われる人のうち服薬者*	105	50.5	1	100.0	0	0.0	3	27.3	8	36.4	38	50.0	55	57.9	78	52.7	52	65.0
	糖尿病の可能性が否定できない人	359	15.8	2	1.9	17	5.3	35	11.4	62	15.8	121	21.4	122	21.1	261	17.3	109	20.0
	上記以外	1,701	75.0	102	97.1	301	93.8	262	85.1	308	78.6	368	65.1	360	62.4	1,100	72.9	357	65.4

(22年)

注) ヘモグロビン A1c $\geq$ 5.6% の場合 判定基準は p.18 参照
* 「服薬者」とは、質問票で「インスリン注射または血糖値を下げる薬」の使用有と回答した者

第 47 表 高血圧症有病者の状況（性・年齢階級別）

		年齢(歳)														(再掲)			
		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		40-74歳		65-74歳	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	総 数	4,119	100.0	202	100.0	511	100.0	509	100.0	696	100.0	1,076	100.0	1,125	100.0	2,743	100.0	1,022	100.0
	高血圧症有病者	1,999	48.5	11	5.4	46	9.0	103	20.2	312	44.8	667	62.0	860	76.4	1,421	51.8	692	67.7
	(再掲)高血圧症有病者のうち服薬者*	1,106	55.3	0	0.0	2	4.3	21	20.4	125	40.1	363	54.4	595	69.2	731	51.4	418	60.4
	正常高値血圧者	556	13.5	14	6.9	59	11.5	88	17.3	114	16.4	167	15.5	114	10.1	419	15.3	133	13.0
	上記以外	1,564	38.0	177	87.6	406	79.5	318	62.5	270	38.8	242	22.5	151	13.4	903	32.9	197	19.3
男 性	総 数	1,697	100.0	84	100.0	170	100.0	185	100.0	277	100.0	486	100.0	495	100.0	1,152	100.0	447	100.0
	高血圧症有病者	977	57.6	10	11.9	33	19.4	62	33.5	160	57.8	313	64.4	399	80.6	705	61.2	330	73.8
	(再掲)高血圧症有病者のうち服薬者*	518	53.0	0	0.0	1	3.0	14	22.6	70	43.8	170	54.3	263	65.9	363	51.5	200	60.6
	正常高値血圧者	261	15.4	11	13.1	39	22.9	37	20.0	47	17.0	81	16.7	46	9.3	179	15.5	53	11.9
	上記以外	459	27.0	63	75.0	98	57.6	86	46.5	70	25.3	92	18.9	50	10.1	268	23.3	64	14.3
女 性	総 数	2,422	100.0	118	100.0	341	100.0	324	100.0	419	100.0	590	100.0	630	100.0	1,591	100.0	575	100.0
	高血圧症有病者	1,022	42.2	1	0.8	13	3.8	41	12.7	152	36.3	354	60.0	461	73.2	716	45.0	362	63.0
	(再掲)高血圧症有病者のうち服薬者*	588	57.5	0	0.0	1	7.7	7	17.1	55	36.2	193	54.5	332	72.0	368	51.4	218	60.2
	正常高値血圧者	295	12.2	3	2.5	20	5.9	51	15.7	67	16.0	86	14.6	68	10.8	240	15.1	80	13.9
	上記以外	1,105	45.6	114	96.6	308	90.3	232	71.6	200	47.7	150	25.4	101	16.0	635	39.9	133	23.1

* 「服薬者」とは、質問票で「血圧を下げる薬」の使用有と回答した者

「高血圧症有病者」の判定
収縮期血圧 140mmHg 以上または拡張期血圧 90mmHg 以上、もしくは血圧を下げる薬を服用している者

(22年)

第48表 「脂質異常症が疑われる人」の状況（採血時間によらず，性・年齢階級別）

		総 数		20～29歳		30～39歳		40～49歳		50～59歳		60～69歳		70歳以上		(再掲) 40～74歳		(再掲) 65～74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	総 数	3,863	100.0	180	100.0	483	100.0	483	100.0	651	100.0	1,025	100.0	1,041	100.0	2,597	100.0	970	100.0	603	100.0
	脂質異常症が疑われる人	762	19.7	6	3.3	22	4.6	33	6.8	104	16.0	259	25.3	338	32.5	547	21.1	298	30.7	187	31.0
	(再掲)脂質異常症が疑われる人のうち服薬者*	571	74.9	0	0.0	4	18.2	14	42.4	73	70.2	199	76.8	281	83.1	409	74.8	243	81.5	158	84.5
	上記以外	3,101	80.3	174	96.7	461	95.4	450	93.2	547	84.0	766	74.7	703	67.5	2,050	78.9	672	69.3	416	69.0
男 性	総 数	1,596	100.0	75	100.0	162	100.0	175	100.0	259	100.0	461	100.0	464	100.0	1,088	100.0	424	100.0	271	100.0
	脂質異常症が疑われる人	368	23.1	5	6.7	18	11.1	27	15.4	59	22.8	124	26.9	135	29.1	269	24.7	122	28.8	76	28.0
	(再掲)脂質異常症が疑われる人のうち服薬者*	215	58.4	0	0.0	1	5.6	12	44.4	35	59.3	74	59.7	93	68.9	160	59.5	81	66.4	54	71.1
	上記以外	1,228	76.9	70	93.3	144	88.9	148	84.6	200	77.2	337	73.1	329	70.9	819	75.3	302	71.2	195	72.0
女 性	総 数	2,267	100.0	105	100.0	321	100.0	308	100.0	392	100.0	564	100.0	577	100.0	1,509	100.0	546	100.0	332	100.0
	脂質異常症が疑われる人	394	17.4	1	1.0	4	1.2	6	1.9	45	11.5	135	23.9	203	35.2	278	18.4	176	32.2	111	33.4
	(再掲)脂質異常症が疑われる人のうち服薬者*	356	90.4	0	0.0	3	75.0	2	33.3	38	84.4	125	92.6	188	92.6	249	89.6	162	92.0	104	93.7
	上記以外	1,873	82.6	104	99.0	317	98.8	302	98.1	347	88.5	429	76.1	374	64.8	1,231	81.6	370	67.8	221	66.6

* 「服薬者」とは，質問票で「コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬」の使用有と回答した者

(22年)

「脂質異常症が疑われる人」の判定
国民健康・栄養調査の血液検査では，空腹時採血が困難であるため，脂質異常症の診断基準項目である中性脂肪による判定は行わず，下記の通りとした。
・「脂質異常症が疑われる人」
HDL コレステロールが40mg/dl未満，もしくはコレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬を服用している者。

第 49 表 運動習慣の状況（20 歳以上、性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 20-64歳	(再掲) 65歳以上	(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	4,621	100.0	284	100.0	594	100.0	596	100.0	780	100.0	1,156	100.0	1,211	100.0	2,814	100.0	1,807	100.0
	習慣無(健康上の理由)	321	6.9	5	1.8	14	2.4	19	3.2	31	4.0	72	6.2	180	14.9	96	3.4	225	12.5
	習慣無(その他)	2,858	61.8	226	79.6	463	77.9	477	80.0	524	67.2	618	53.5	550	45.4	2,034	72.3	824	45.6
	運動の習慣有	1,442	31.2	53	18.7	117	19.7	100	16.8	225	28.8	466	40.3	481	39.7	684	24.3	758	41.9
男性	総 数	1,963	100.0	126	100.0	218	100.0	237	100.0	324	100.0	531	100.0	527	100.0	1,173	100.0	790	100.0
	習慣無(健康上の理由)	97	4.9	1	0.8	1	0.5	5	2.1	12	3.7	28	5.3	50	9.5	31	2.6	66	8.4
	習慣無(その他)	1,182	60.2	89	70.6	163	74.8	186	78.5	227	70.1	277	52.2	240	45.5	834	71.1	348	44.1
	運動の習慣有	684	34.8	36	28.6	54	24.8	46	19.4	85	26.2	226	42.6	237	45.0	308	26.3	376	47.6
女性	総 数	2,658	100.0	158	100.0	376	100.0	359	100.0	456	100.0	625	100.0	684	100.0	1,641	100.0	1,017	100.0
	習慣無(健康上の理由)	224	8.4	4	2.5	13	3.5	14	3.9	19	4.2	44	7.0	130	19.0	65	4.0	159	15.6
	習慣無(その他)	1,676	63.1	137	86.7	300	79.8	291	81.1	297	65.1	341	54.6	310	45.3	1,200	73.1	476	46.8
	運動の習慣有	758	28.5	17	10.8	63	16.8	54	15.0	140	30.7	240	38.4	244	35.7	376	22.9	382	37.6

(22年)

第 50 表 1 週間の運動日数（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	4,621	100.0	284	100.0	594	100.0	596	100.0	780	100.0	1,156	100.0	1,211	100.0	1,081	100.0
	運動無し	3,179	68.8	231	81.3	477	80.3	496	83.2	555	71.2	690	59.7	730	60.3	568	52.5
	2日/週	308	6.7	26	9.2	43	7.2	27	4.5	59	7.6	81	7.0	72	5.9	82	7.6
	3日/週	212	4.6	10	3.5	18	3.0	19	3.2	33	4.2	64	5.5	68	5.6	68	6.3
	4日/週	144	3.1	4	1.4	8	1.3	9	1.5	20	2.6	49	4.2	54	4.5	64	5.9
	5日/週	180	3.9	6	2.1	20	3.4	16	2.7	33	4.2	59	5.1	46	3.8	56	5.2
	6日/週	79	1.7	2	0.7	9	1.5	3	0.5	16	2.1	30	2.6	19	1.6	20	1.9
	7日/週	519	11.2	5	1.8	19	3.2	26	4.4	64	8.2	183	15.8	222	18.3	223	20.6
男性	総 数	1,963	100.0	126	100.0	218	100.0	237	100.0	324	100.0	531	100.0	527	100.0	480	100.0
	運動無し	1,279	65.2	90	71.4	164	75.2	191	80.6	239	73.8	305	57.4	290	55.0	229	47.7
	2日/週	146	7.4	23	18.3	22	10.1	15	6.3	22	6.8	32	6.0	32	6.1	30	6.3
	3日/週	94	4.8	5	4.0	9	4.1	5	2.1	10	3.1	33	6.2	32	6.1	35	7.3
	4日/週	65	3.3	1	0.8	3	1.4	5	2.1	7	2.2	24	4.5	25	4.7	33	6.9
	5日/週	79	4.0	2	1.6	7	3.2	10	4.2	13	4.0	25	4.7	22	4.2	23	4.8
	6日/週	42	2.1	1	0.8	7	3.2	2	0.8	7	2.2	17	3.2	8	1.5	11	2.3
	7日/週	258	13.1	4	3.2	6	2.8	9	3.8	26	8.0	95	17.9	118	22.4	119	24.8
女性	総 数	2,658	100.0	158	100.0	376	100.0	359	100.0	456	100.0	625	100.0	684	100.0	601	100.0
	運動無し	1,900	71.5	141	89.2	313	83.2	305	85.0	316	69.3	385	61.6	440	64.3	339	56.4
	2日/週	162	6.1	3	1.9	21	5.6	12	3.3	37	8.1	49	7.8	40	5.8	52	8.7
	3日/週	118	4.4	5	3.2	9	2.4	14	3.9	23	5.0	31	5.0	36	5.3	33	5.5
	4日/週	79	3.0	3	1.9	5	1.3	4	1.1	13	2.9	25	4.0	29	4.2	31	5.2
	5日/週	101	3.8	4	2.5	13	3.5	6	1.7	20	4.4	34	5.4	24	3.5	33	5.5
	6日/週	37	1.4	1	0.6	2	0.5	1	0.3	9	2.0	13	2.1	11	1.6	9	1.5
	7日/週	261	9.8	1	0.6	13	3.5	17	4.7	38	8.3	88	14.1	104	15.2	104	17.3

(22年)

第 51 表 運動を行う日の平均運動時間（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	4,621	100.0	284	100.0	594	100.0	596	100.0	780	100.0	1,156	100.0	1,211	100.0	1,081	100.0
	運動無し	3,179	68.8	231	81.3	477	80.3	496	83.2	555	71.2	690	59.7	730	60.3	568	52.5
	30分以上60分未満	597	12.9	17	6.0	35	5.9	35	5.9	102	13.1	196	17.0	212	17.5	214	19.8
	60分以上120分未満	535	11.6	15	5.3	48	8.1	41	6.9	79	10.1	171	14.8	181	14.9	191	17.7
	120分以上180分未満	202	4.4	15	5.3	20	3.4	16	2.7	33	4.2	63	5.4	55	4.5	65	6.0
	180分以上	108	2.3	6	2.1	14	2.4	8	1.3	11	1.4	36	3.1	33	2.7	43	4.0
男性	総 数	1,963	100.0	126	100.0	218	100.0	237	100.0	324	100.0	531	100.0	527	100.0	480	100.0
	運動無し	1,279	65.2	90	71.4	164	75.2	191	80.6	239	73.8	305	57.4	290	55.0	229	47.7
	30分以上60分未満	238	12.1	9	7.1	13	6.0	13	5.5	28	8.6	78	14.7	97	18.4	88	18.3
	60分以上120分未満	267	13.6	10	7.9	22	10.1	20	8.4	38	11.7	90	16.9	87	16.5	100	20.8
	120分以上180分未満	110	5.6	12	9.5	9	4.1	7	3.0	15	4.6	36	6.8	31	5.9	35	7.3
	180分以上	69	3.5	5	4.0	10	4.6	6	2.5	4	1.2	22	4.1	22	4.2	28	5.8
女性	総 数	2,658	100.0	158	100.0	376	100.0	359	100.0	456	100.0	625	100.0	684	100.0	601	100.0
	運動無し	1,900	71.5	141	89.2	313	83.2	305	85.0	316	69.3	385	61.6	440	64.3	339	56.4
	30分以上60分未満	359	13.5	8	5.1	22	5.9	22	6.1	74	16.2	118	18.9	115	16.8	126	21.0
	60分以上120分未満	268	10.1	5	3.2	26	6.9	21	5.8	41	9.0	81	13.0	94	13.7	91	15.1
	120分以上180分未満	92	3.5	3	1.9	11	2.9	9	2.5	18	3.9	27	4.3	24	3.5	30	5.0
	180分以上	39	1.5	1	0.6	4	1.1	2	0.6	7	1.5	14	2.2	11	1.6	15	2.5

(22年)

第52表 運動強度の状況（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	4,621	100.0	284	100.0	594	100.0	596	100.0	780	100.0	1,156	100.0	1,211	100.0	1,081	100.0	726	100.0
	運動無し	3,179	68.8	231	81.3	477	80.3	496	83.2	555	71.2	690	59.7	730	60.3	568	52.5	481	66.3
	高強度	102	2.2	17	6.0	19	3.2	17	2.9	21	2.7	18	1.6	10	0.8	18	1.7	2	0.3
	中強度	468	10.1	23	8.1	55	9.3	41	6.9	86	11.0	154	13.3	109	9.0	139	12.9	48	6.6
	低強度	872	18.9	13	4.6	43	7.2	42	7.0	118	15.1	294	25.4	362	29.9	356	32.9	195	26.9
男性	総 数	1,963	100.0	126	100.0	218	100.0	237	100.0	324	100.0	531	100.0	527	100.0	480	100.0	310	100.0
	運動無し	1,279	65.2	90	71.4	164	75.2	191	80.6	239	73.8	305	57.4	290	55.0	229	47.7	185	59.7
	高強度	73	3.7	16	12.7	15	6.9	10	4.2	14	4.3	12	2.3	6	1.1	12	2.5	1	0.3
	中強度	233	11.9	14	11.1	24	11.0	20	8.4	34	10.5	79	14.9	62	11.8	72	15.0	32	10.3
	低強度	378	19.3	6	4.8	15	6.9	16	6.8	37	11.4	135	25.4	169	32.1	167	34.8	92	29.7
女性	総 数	2,658	100.0	158	100.0	376	100.0	359	100.0	456	100.0	625	100.0	684	100.0	601	100.0	416	100.0
	運動無し	1,900	71.5	141	89.2	313	83.2	305	85.0	316	69.3	385	61.6	440	64.3	339	56.4	296	71.2
	高強度	29	1.1	1	0.6	4	1.1	7	1.9	7	1.5	6	1.0	4	0.6	6	1.0	1	0.2
	中強度	235	8.8	9	5.7	31	8.2	21	5.8	52	11.4	75	12.0	47	6.9	67	11.1	16	3.8
	低強度	494	18.6	7	4.4	28	7.4	26	7.2	81	17.8	159	25.4	193	28.2	189	31.4	103	24.8

(22年)

第53表の1 歩行数の分布（性・年齢階級別）

		総 数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 20歳以上		(再掲) 20-64歳		(再掲) 65歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	7,141	100.0	340	100.0	588	100.0	1,029	100.0	1,036	100.0	1,140	100.0	1,525	100.0	1,483	100.0	6,801	100.0	4,580	100.0	2,221	100.0	1,322	100.0	899	100.0
	2,000歩未満	792	11.1	12	3.5	35	6.0	51	5.0	39	3.8	49	4.3	138	9.0	468	31.6	780	11.5	239	5.2	541	24.4	180	13.6	361	40.2
	2,000-3,999	1,320	18.5	56	16.5	88	15.0	150	14.6	161	15.5	183	16.1	305	20.0	377	25.4	1,264	18.6	729	15.9	535	24.1	290	21.9	245	27.3
	4,000-5,999	1,397	19.6	62	18.2	108	18.4	226	22.0	207	20.0	228	20.0	323	21.2	243	16.4	1,335	19.6	944	20.6	391	17.6	251	19.0	140	15.6
	6,000-7,999	1,292	18.1	74	21.8	120	20.4	211	20.5	231	22.3	227	19.9	261	17.1	168	11.3	1,218	17.9	932	20.3	286	12.9	206	15.6	80	8.9
	8,000-9,999	1,047	14.7	52	15.3	87	14.8	176	17.1	188	18.1	204	17.9	219	14.4	121	8.2	995	14.6	773	16.9	222	10.0	176	13.3	46	5.1
	10,000歩以上	1,293	18.1	84	24.7	150	25.5	215	20.9	210	20.3	249	21.8	279	18.3	106	7.1	1,209	17.8	963	21.0	246	11.1	219	16.6	27	3.0
	9,200歩(男)/8,300歩(女)以上	1,909	26.7	111	32.6	199	33.8	327	31.8	309	29.8	371	32.5	421	27.6	171	11.5	1,798	26.4	1,417	30.9	381	17.2	329	24.9	52	5.8
	平均値(歩/日)	6,636		7,665		7,672		7,410		7,404		7,412		6,630		4,324		6,584		7,328		5,051		6,164		3,416	
	標準偏差	4,225		4,201		4,482		4,217		4,026		3,987		4,007		3,785		4,220		4,127		3,991		4,016		3,341	
男性	総 数	3,290	100.0	170	100.0	274	100.0	474	100.0	488	100.0	520	100.0	705	100.0	659	100.0	3,120	100.0	2,126	100.0	994	100.0	608	100.0	386	100.0
	2,000歩未満	338	10.3	5	2.9	12	4.4	21	4.4	21	4.3	28	5.4	58	8.2	193	29.3	333	10.7	107	5.0	226	22.7	83	13.7	143	37.0
	2,000-3,999	553	16.8	31	18.2	36	13.1	58	12.2	72	14.8	81	15.6	128	18.2	147	22.3	522	16.7	311	14.6	211	21.2	114	18.8	97	25.1
	4,000-5,999	595	18.1	29	17.1	50	18.2	86	18.1	83	17.0	88	16.9	153	21.7	106	16.1	566	18.1	393	18.5	173	17.4	111	18.3	62	16.1
	6,000-7,999	582	17.7	37	21.8	48	17.5	100	21.1	102	20.9	97	18.7	112	15.9	86	13.1	545	17.5	415	19.5	130	13.1	89	14.6	41	10.6
	8,000-9,999	479	14.6	23	13.5	43	15.7	74	15.6	93	19.1	98	18.8	92	13.0	56	8.5	456	14.6	361	17.0	95	9.6	71	11.7	24	6.2
	10,000歩以上	743	22.6	45	26.5	85	31.0	135	28.5	117	24.0	128	24.6	162	23.0	71	10.8	698	22.4	539	25.4	159	16.0	140	23.0	19	4.9
	(再掲)9,200歩以上	892	27.1	52	30.6	97	35.4	161	34.0	143	29.3	158	30.4	196	27.8	85	12.9	840	26.9	653	30.7	187	18.8	162	26.6	25	6.5
	平均値(歩/日)	7,174		7,872		8,322		8,278		7,873		7,684		7,092		4,890		7,136		7,841		5,628		6,703		3,935	
	標準偏差	4,685		4,545		4,833		4,901		4,576		4,368		4,379		4,362		4,690		4,590		4,546		4,482		4,115	
女性	総 数	3,851	100.0	170	100.0	314	100.0	555	100.0	548	100.0	620	100.0	820	100.0	824	100.0	3,681	100.0	2,454	100.0	1,227	100.0	714	100.0	513	100.0
	2,000歩未満	454	11.8	7	4.1	23	7.3	30	5.4	18	3.3	21	3.4	80	9.8	275	33.4	447	12.1	132	5.4	315	25.7	97	13.6	218	42.5
	2,000-3,999	767	19.9	25	14.7	52	16.6	92	16.6	89	16.2	102	16.5	177	21.6	230	27.9	742	20.2	418	17.0	324	26.4	176	24.6	148	28.8
	4,000-5,999	802	20.8	33	19.4	58	18.5	140	25.2	124	22.6	140	22.6	170	20.7	137	16.6	769	20.9	551	22.5	218	17.8	140	19.6	78	15.2
	6,000-7,999	710	18.4	37	21.8	72	22.9	111	20.0	129	23.5	130	21.0	149	18.2	82	10.0	673	18.3	517	21.1	156	12.7	117	16.4	39	7.6
	8,000-9,999	568	14.7	29	17.1	44	14.0	102	18.4	95	17.3	106	17.1	127	15.5	65	7.9	539	14.6	412	16.8	127	10.4	105	14.7	22	4.3
	10,000歩以上	550	14.3	39	22.9	65	20.7	80	14.4	93	17.0	121	19.5	117	14.3	35	4.2	511	13.9	424	17.3	87	7.1	79	11.1	8	1.6
	(再掲)8,300歩以上	1,017	26.4	59	34.7	102	32.5	166	29.9	166	30.3	213	34.4	225	27.4	86	10.4	958	26.0	764	31.1	194	15.8	167	23.4	27	5.3
	平均値(歩/日)	6,176		7,458		7,104		6,669		6,986		7,184		6,234		3,872		6,117		6,883		4,584		5,705		3,025	
	標準偏差	3,728		3,829		4,075		3,361		3,414		3,624		3,612		3,183		3,713		3,622		3,409		3,510		2,547	

(22年)

第 53 表の 2 運動習慣の状況別，歩行数の分布（性・年齢階級別）

			総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	総 数	4,294	100.0	248	100.0	565	100.0	558	100.0	735	100.0	1,102	100.0	1,086	100.0	1,015	100.0	640	100.0
		2,000歩未満	478	11.1	17	6.9	25	4.4	16	2.9	30	4.1	87	7.9	303	27.9	125	12.3	230	35.9
		2,000-3,999	821	19.1	39	15.7	91	16.1	85	15.2	120	16.3	213	19.3	273	25.1	207	20.4	177	27.7
		4,000-5,999	903	21.0	51	20.6	131	23.2	125	22.4	157	21.4	248	22.5	191	17.6	201	19.8	110	17.2
		6,000-7,999	789	18.4	53	21.4	116	20.5	126	22.6	153	20.8	201	18.2	140	12.9	173	17.0	66	10.3
		8,000-9,999	592	13.8	36	14.5	96	17.0	101	18.1	113	15.4	155	14.1	91	8.4	132	13.0	34	5.3
		10,000歩以上	711	16.6	52	21.0	106	18.8	105	18.8	162	22.0	198	18.0	88	8.1	177	17.4	23	3.6
		9,200歩(男)/8,300歩(女)以上	1,085	25.3	74	29.8	172	30.4	161	28.9	233	31.7	302	27.4	143	13.2	272	26.8	40	6.3
	習慣無 (健康上の理由)	総 数	264	100.0	3	100.0	14	100.0	16	100.0	26	100.0	65	100.0	140	100.0	73	100.0	108	100.0
		2,000歩未満	103	39.0	1	33.3	1	7.1	0	0.0	4	15.4	16	24.6	81	57.9	25	34.2	66	61.1
		2,000-3,999	79	29.9	0	0.0	4	28.6	3	18.8	7	26.9	29	44.6	36	25.7	26	35.6	28	25.9
		4,000-5,999	31	11.7	1	33.3	3	21.4	4	25.0	3	11.5	9	13.8	11	7.9	9	12.3	8	7.4
		6,000-7,999	24	9.1	1	33.3	3	21.4	4	25.0	4	15.4	5	7.7	7	5.0	4	5.5	6	5.6
		8,000-9,999	15	5.7	0	0.0	1	7.1	3	18.8	5	19.2	2	3.1	4	2.9	6	8.2	0	0.0
		10,000歩以上	12	4.5	0	0.0	2	14.3	2	12.5	3	11.5	4	6.2	1	0.7	3	4.1	0	0.0
		9,200歩(男)/8,300歩(女)以上	22	8.3	0	0.0	3	21.4	4	25.0	6	23.1	5	7.7	4	2.9	7	9.6	0	0.0
	習慣無 (その他)	総 数	2,677	100.0	201	100.0	438	100.0	445	100.0	496	100.0	587	100.0	510	100.0	458	100.0	314	100.0
		2,000歩未満	304	11.4	14	7.0	20	4.6	12	2.7	22	4.4	62	10.6	174	34.1	82	17.9	129	41.1
		2,000-3,999	547	20.4	33	16.4	78	17.8	71	16.0	90	18.1	133	22.7	142	27.8	113	24.7	87	27.7
		4,000-5,999	620	23.2	45	22.4	107	24.4	108	24.3	121	24.4	140	23.9	99	19.4	100	21.8	58	18.5
		6,000-7,999	487	18.2	38	18.9	89	20.3	97	21.8	107	21.6	109	18.6	47	9.2	72	15.7	21	6.7
		8,000-9,999	350	13.1	31	15.4	76	17.4	81	18.2	66	13.3	71	12.1	25	4.9	42	9.2	13	4.1
		10,000歩以上	369	13.8	40	19.9	68	15.5	76	17.1	90	18.1	72	12.3	23	4.5	49	10.7	6	1.9
		9,200歩(男)/8,300歩(女)以上	586	21.9	60	29.9	122	27.9	119	26.7	129	26.0	118	20.1	38	7.5	81	17.7	13	4.1
	運動の 習慣有	総 数	1,353	100.0	44	100.0	113	100.0	97	100.0	213	100.0	450	100.0	436	100.0	484	100.0	218	100.0
		2,000歩未満	71	5.2	2	4.5	4	3.5	4	4.1	4	1.9	9	2.0	48	11.0	18	3.7	35	16.1
		2,000-3,999	195	14.4	6	13.6	9	8.0	11	11.3	23	10.8	51	11.3	95	21.8	68	14.0	62	28.4
		4,000-5,999	252	18.6	5	11.4	21	18.6	13	13.4	33	15.5	99	22.0	81	18.6	92	19.0	44	20.2
		6,000-7,999	278	20.5	14	31.8	24	21.2	25	25.8	42	19.7	87	19.3	86	19.7	97	20.0	39	17.9
		8,000-9,999	227	16.8	5	11.4	19	16.8	17	17.5	42	19.7	82	18.2	62	14.2	84	17.4	21	9.6
		10,000歩以上	330	24.4	12	27.3	36	31.9	27	27.8	69	32.4	122	27.1	64	14.7	125	25.8	17	7.8
		9,200歩(男)/8,300歩(女)以上	477	35.3	14	31.8	47	41.6	38	39.2	98	46.0	179	39.8	101	23.2	184	38.0	27	12.4
男性	総 数	総 数	1,794	100.0	105	100.0	203	100.0	220	100.0	297	100.0	496	100.0	473	100.0	447	100.0	273	100.0
		2,000歩未満	198	11.0	6	5.7	10	4.9	7	3.2	15	5.1	39	7.9	121	25.6	58	13.0	88	32.2
		2,000-3,999	309	17.2	13	12.4	24	11.8	36	16.4	47	15.8	87	17.5	102	21.6	75	16.8	69	25.3
		4,000-5,999	359	20.0	20	19.0	43	21.2	40	18.2	60	20.2	113	22.8	83	17.5	87	19.5	48	17.6
		6,000-7,999	326	18.2	21	20.0	42	20.7	50	22.7	61	20.5	80	16.1	72	15.2	71	15.9	34	12.5
		8,000-9,999	226	12.6	15	14.3	29	14.3	39	17.7	44	14.8	61	12.3	38	8.0	43	9.6	18	6.6
		10,000歩以上	376	21.0	30	28.6	55	27.1	48	21.8	70	23.6	116	23.4	57	12.1	113	25.3	16	5.9
		(再掲)9,200歩以上	441	24.6	34	32.4	64	31.5	57	25.9	78	26.3	139	28.0	69	14.6	132	29.5	20	7.3
	習慣無 (健康上の理由)	総 数	77	100.0	0	0.0	1	100.0	5	100.0	10	100.0	24	100.0	37	100.0	21	100.0	30	100.0
		2,000歩未満	27	35.1	0	0.0	0	0.0	0	0.0	1	10.0	6	25.0	20	54.1	7	33.3	18	60.0
		2,000-3,999	21	27.3	0	0.0	0	0.0	1	20.0	1	10.0	11	45.8	8	21.6	5	23.8	7	23.3
		4,000-5,999	7	9.1	0	0.0	0	0.0	1	20.0	0	0.0	2	8.3	4	10.8	4	19.0	1	3.3
		6,000-7,999	11	14.3	0	0.0	1	100.0	2	40.0	3	30.0	1	4.2	4	10.8	1	4.8	4	13.3
		8,000-9,999	6	7.8	0	0.0	0	0.0	1	20.0	4	40.0	1	4.2	0	0.0	1	4.8	0	0.0
		10,000歩以上	5	6.5	0	0.0	0	0.0	0	0.0	1	10.0	3	12.5	1	2.7	3	14.3	0	0.0
		(再掲)9,200歩以上	7	9.1	0	0.0	0	0.0	0	0.0	3	30.0	3	12.5	1	2.7	3	14.3	0	0.0
	習慣無 (その他)	総 数	1,083	100.0	75	100.0	151	100.0	170	100.0	209	100.0	259	100.0	219	100.0	190	100.0	131	100.0
		2,000歩未満	130	12.0	5	6.7	7	4.6	4	2.4	10	4.8	29	11.2	75	34.2	41	21.6	52	39.7
		2,000-3,999	210	19.4	9	12.0	21	13.9	29	17.1	36	17.2	60	23.2	55	25.1	46	24.2	35	26.7
		4,000-5,999	230	21.2	16	21.3	34	22.5	32	18.8	50	23.9	58	22.4	40	18.3	36	18.9	25	19.1
		6,000-7,999	196	18.1	13	17.3	30	19.9	38	22.4	42	20.1	44	17.0	29	13.2	32	16.8	11	8.4
		8,000-9,999	125	11.5	12	16.0	20	13.2	33	19.4	29	13.9	24	9.3	7	3.2	7	3.7	5	3.8
		10,000歩以上	192	17.7	20	26.7	39	25.8	34	20.0	42	20.1	44	17.0	13	5.9	28	14.7	3	2.3
		(再掲)9,200歩以上	223	20.6	24	32.0	45	29.8	42	24.7	47	22.5	50	19.3	15	6.8	30	15.8	4	3.1
	運動の 習慣有	総 数	634	100.0	30	100.0	51	100.0	45	100.0	78	100.0	213	100.0	217	100.0	236	100.0	112	100.0
		2,000歩未満	41	6.5	1	3.3	3	5.9	3	6.7	4	5.1	4	1.9	26	12.0	10	4.2	18	16.1
		2,000-3,999	78	12.3	4	13.3	3	5.9	6	13.3	10	12.8	16	7.5	39	18.0	24	10.2	27	24.1
		4,000-5,999	122	19.2	4	13.3	9	17.6	7	15.6	10	12.8	53	24.9	39	18.0	47	19.9	22	19.6
		6,000-7,999	119	18.8	8	26.7	11	21.6	10	22.2	16	20.5	35	16.4	39	18.0	38	16.1	19	17.0
		8,000-9,999	95	15.0	3	10.0	9	17.6	5	11.1	11	14.1	36	16.9	31	14.3	35	14.8	13	11.6
		10,000歩以上	179	28.2	10	33.3	16	31.4	14	31.1	27	34.6	69	32.4	43	19.8	82	34.7	13	11.6
		(再掲)9,200歩以上	211	33.3	10	33.3	19	37.3	15	33.3	28	35.9	86	40.4	53	24.4	99	41.9	16	14.3

			総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
女性	総 数	総 数	2,500	100.0	143	100.0	362	100.0	338	100.0	438	100.0	606	100.0	613	100.0	568	100.0	367	100.0
		2,000歩未満	280	11.2	11	7.7	15	4.1	9	2.7	15	3.4	48	7.9	182	29.7	67	11.8	142	38.7
		2,000-3,999	512	20.5	26	18.2	67	18.5	49	14.5	73	16.7	126	20.8	171	27.9	132	23.2	108	29.4
		4,000-5,999	544	21.8	31	21.7	88	24.3	85	25.1	97	22.1	135	22.3	108	17.6	114	20.1	62	16.9
		6,000-7,999	463	18.5	32	22.4	74	20.4	76	22.5	92	21.0	121	20.0	68	11.1	102	18.0	32	8.7
		8,000-9,999	366	14.6	21	14.7	67	18.5	62	18.3	69	15.8	94	15.5	53	8.6	89	15.7	16	4.4
		10,000歩以上	335	13.4	22	15.4	51	14.1	57	16.9	92	21.0	82	13.5	31	5.1	64	11.3	7	1.9
		(再掲)8,300歩以上	644	25.8	40	28.0	108	29.8	104	30.8	155	35.4	163	26.9	74	12.1	140	24.6	20	5.4
	習慣無 (健康上の理由)	総 数	187	100.0	3	100.0	13	100.0	11	100.0	16	100.0	41	100.0	103	100.0	52	100.0	78	100.0
		2,000歩未満	76	40.6	1	33.3	1	7.7	0	0.0	3	18.8	10	24.4	61	59.2	18	34.6	48	61.5
		2,000-3,999	58	31.0	0	0.0	4	30.8	2	18.2	6	37.5	18	43.9	28	27.2	21	40.4	21	26.9
		4,000-5,999	24	12.8	1	33.3	3	23.1	3	27.3	3	18.8	7	17.1	7	6.8	5	9.6	7	9.0
		6,000-7,999	13	7.0	1	33.3	2	15.4	2	18.2	1	6.3	4	9.8	3	2.9	3	5.8	2	2.6
		8,000-9,999	9	4.8	0	0.0	1	7.7	2	18.2	1	6.3	1	2.4	4	3.9	5	9.6	0	0.0
		10,000歩以上	7	3.7	0	0.0	2	15.4	2	18.2	2	12.5	1	2.4	0	0.0	0	0.0	0	0.0
		(再掲)8,300歩以上	15	8.0	0	0.0	3	23.1	4	36.4	3	18.8	2	4.9	3	2.9	4	7.7	0	0.0
	習慣無 (その他)	総 数	1,594	100.0	126	100.0	287	100.0	275	100.0	287	100.0	328	100.0	291	100.0	268	100.0	183	100.0
		2,000歩未満	174	10.9	9	7.1	13	4.5	8	2.9	12	4.2	33	10.1	99	34.0	41	15.3	77	42.1
		2,000-3,999	337	21.1	24	19.0	57	19.9	42	15.3	54	18.8	73	22.3	87	29.9	67	25.0	52	28.4
		4,000-5,999	390	24.5	29	23.0	73	25.4	76	27.6	71	24.7	82	25.0	59	20.3	64	23.9	33	18.0
		6,000-7,999	291	18.3	25	19.8	59	20.6	59	21.5	65	22.6	65	19.8	18	6.2	40	14.9	10	5.5
		8,000-9,999	225	14.1	19	15.1	56	19.5	48	17.5	37	12.9	47	14.3	18	6.2	35	13.1	8	4.4
		10,000歩以上	177	11.1	20	15.9	29	10.1	42	15.3	48	16.7	28	8.5	10	3.4	21	7.8	3	1.6
		(再掲)8,300歩以上	363	22.8	36	28.6	77	26.8	77	28.0	82	28.6	68	20.7	23	7.9	51	19.0	9	4.9
	運動の 習慣有	総 数	719	100.0	14	100.0	62	100.0	52	100.0	135	100.0	237	100.0	219	100.0	248	100.0	106	100.0
		2,000歩未満	30	4.2	1	7.1	1	1.6	1	1.9	0	0.0	5	2.1	22	10.0	8	3.2	17	16.0
		2,000-3,999	117	16.3	2	14.3	6	9.7	5	9.6	13	9.6	35	14.8	56	25.6	44	17.7	35	33.0
		4,000-5,999	130	18.1	1	7.1	12	19.4	6	11.5	23	17.0	46	19.4	42	19.2	45	18.1	22	20.8
		6,000-7,999	159	22.1	6	42.9	13	21.0	15	28.8	26	19.3	52	21.9	47	21.5	59	23.8	20	18.9
		8,000-9,999	132	18.4	2	14.3	10	16.1	12	23.1	31	23.0	46	19.4	31	14.2	49	19.8	8	7.5
		10,000歩以上	151	21.0	2	14.3	20	32.3	13	25.0	42	31.1	53	22.4	21	9.6	43	17.3	4	3.8
		(再掲)8,300歩以上	266	37.0	4	28.6	28	45.2	23	44.2	70	51.9	93	39.2	48	21.9	85	34.3	11	10.4

(22年)

