

第2部 身体状況調査の結果

第11表 身長・体重の平均値及び標準偏差（性・年齢階級別）

年齢(歳)	男 性						女 性					
	身 長(cm)			体 重(kg)			身 長(cm)			体 重(kg)		
	人 数	平均値	標準偏差	人 数	平均値	標準偏差	人 数	平均値	標準偏差	人 数	平均値	標準偏差
総 数	3,639	161.0	18.7	3,630	60.0	17.6	4,361	150.1	14.2	4,336	49.6	12.8
1	22	80.1	3.2	25	10.9	1.3	35	77.2	4.8	35	10.0	1.2
2	43	88.0	4.6	43	12.6	1.4	20	87.9	3.9	20	12.4	1.6
3	35	98.2	4.2	35	15.2	1.5	39	94.8	3.6	39	14.0	1.6
4	40	103.9	4.6	40	16.8	2.0	39	103.5	4.2	39	16.6	1.8
5	30	108.6	4.4	30	17.8	2.0	29	109.7	5.9	29	18.2	2.4
6	36	117.0	5.1	36	21.4	3.7	44	115.7	5.2	44	20.5	3.4
7	40	121.9	4.4	40	23.9	3.3	33	121.9	5.5	33	22.9	4.4
8	40	126.5	5.2	40	25.3	4.0	46	125.4	6.0	46	25.7	4.9
9	49	133.6	5.9	49	30.8	6.2	41	133.5	6.0	41	30.7	6.2
10	40	138.1	6.2	40	33.9	7.2	50	139.4	6.7	50	33.8	7.5
11	37	144.0	6.8	37	37.8	11.4	53	146.7	6.9	53	39.5	9.6
12	32	150.7	8.9	32	41.4	7.0	36	153.0	4.6	36	43.1	6.5
13	53	158.0	7.3	53	46.8	8.4	54	153.9	4.7	54	46.0	8.0
14	39	164.4	6.7	39	51.0	8.3	28	157.9	5.4	29	49.6	7.4
15	32	168.4	5.6	32	58.4	11.7	33	158.6	5.4	33	49.4	7.3
16	38	170.0	5.2	36	59.8	10.5	39	158.5	5.1	39	51.7	7.8
17	44	171.8	5.4	43	61.5	8.6	26	157.6	4.4	26	51.4	7.3
18	33	171.7	5.4	33	63.3	9.6	27	158.2	4.4	27	52.3	6.2
19	24	172.0	5.2	24	65.1	10.4	37	157.7	4.9	36	51.4	5.8
20	16	173.5	5.8	16	64.0	18.0	38	157.0	6.2	38	48.5	6.7
21	24	170.1	5.8	24	60.9	10.9	16	159.0	5.2	16	50.8	6.8
22	25	172.3	5.4	25	65.3	11.2	24	158.3	5.5	24	50.0	5.7
23	21	171.4	5.6	21	63.2	11.6	35	158.6	5.7	35	51.6	7.9
24	28	169.9	5.1	28	66.9	14.3	29	158.8	5.5	28	50.1	5.5
25	23	171.9	6.9	23	62.8	9.4	38	156.0	4.7	37	50.0	8.1
26-29	106	169.8	6.0	106	65.6	12.7	116	157.1	5.5	114	51.7	8.2
30-39	406	171.4	5.9	406	70.8	12.6	514	158.1	5.1	495	53.9	10.4
40-49	474	170.9	5.7	469	70.7	10.9	547	157.7	5.0	545	55.6	10.7
50-59	524	168.1	5.9	520	67.0	10.1	609	154.6	5.6	606	53.9	8.7
60-69	635	164.8	5.6	635	64.3	9.1	776	152.2	5.2	776	53.4	8.5
70歳以上	650	160.3	6.8	650	59.7	9.1	910	147.2	6.2	913	49.7	8.7
(再掲)												
20歳以上	2,932	166.8	7.3	2,923	65.7	11.2	3,652	153.4	6.9	3,627	52.7	9.4
20-29	243	170.7	5.9	243	64.7	12.6	296	157.5	5.5	292	50.7	7.5
60-64	322	165.0	5.6	322	64.4	9.8	406	153.2	5.2	405	54.2	8.8
65-69	313	164.6	5.6	313	64.1	8.4	370	151.1	5.0	371	52.5	8.2
70-74	261	162.1	5.6	261	62.5	8.6	321	149.3	5.7	321	51.4	8.3
75-79	214	160.2	6.0	214	59.5	8.3	278	147.3	5.5	277	49.8	8.6
80歳以上	175	157.6	8.3	175	55.6	9.3	311	145.1	6.5	315	47.8	8.9

注) 体重は妊婦除外

第12表 BMIの平均値及び標準偏差（15歳以上，性・年齢階級別）

年齢(歳)	男 性			女 性		
	人 数	平均値	標準偏差	人 数	平均値	標準偏差
総 数	3,090	23.44	3.43	3,783	22.33	3.68
15-19	168	21.03	3.15	161	20.47	2.68
20-29	243	22.18	3.97	292	20.42	2.81
30-39	405	24.06	3.94	495	21.60	3.95
40-49	469	24.19	3.45	545	22.36	4.06
50-59	520	23.70	3.13	606	22.55	3.51
60-69	635	23.63	3.00	775	23.05	3.53
70歳以上	650	23.22	3.15	909	22.90	3.54
(再掲)						
20歳以上	2,922	23.58	3.40	3,622	22.41	3.70
20-69	2,272	23.68	3.46	2,713	22.25	3.74
40-69	1,624	23.81	3.19	1,926	22.70	3.69
65-69	313	23.62	2.72	370	22.99	3.48
70-74	261	23.78	3.00	321	23.07	3.44
75-79	214	23.18	2.96	277	22.91	3.54
80-84	108	22.90	3.44	184	22.87	3.65
80歳以上	175	22.41	3.44	311	22.72	3.65
85歳以上	67	21.63	3.31	127	22.52	3.67

注) 妊婦除外

第13表 BMIの分布（15歳以上，性・年齢階級別）

	総 数		男 性		女 性	
	人 数	%	人 数	%	人 数	%
総 数	6,873	100.0	3,090	100.0	3,783	100.0
15未満	8	0.1	2	0.1	6	0.2
15-16未満	32	0.5	4	0.1	28	0.7
16-17	99	1.4	32	1.0	67	1.8
17-18	236	3.4	70	2.3	166	4.4
18-19	487	7.1	130	4.2	357	9.4
19-20	663	9.6	201	6.5	462	12.2
20-21	775	11.3	309	10.0	466	12.3
21-22	797	11.6	354	11.5	443	11.7
22-23	812	11.8	399	12.9	413	10.9
23-24	700	10.2	379	12.3	321	8.5
24-25	595	8.7	304	9.8	291	7.7
25-26	494	7.2	280	9.1	214	5.7
26-27	362	5.3	205	6.6	157	4.2
27-28	251	3.7	139	4.5	112	3.0
28-29	178	2.6	89	2.9	89	2.4
29-30	126	1.8	64	2.1	62	1.6
30-31	73	1.1	44	1.4	29	0.8
31-32	65	0.9	32	1.0	33	0.9
32-33	38	0.6	17	0.6	21	0.6
33-34	23	0.3	10	0.3	13	0.3
34-35	17	0.2	11	0.4	6	0.2
35以上	42	0.6	15	0.5	27	0.7

注) 妊婦除外

第 14 表 BMI の状況（15 歳以上、性・年齢階級別）

		総 数		低体重(やせ)		普 通		肥 満		(再掲) 25以上30未満		(再掲) 30以上	
				18.5未満		18.5以上25未満		25以上					
		人 数	%	人 数	%	人 数	%	人 数	%	人 数	%	人 数	%
総数	総 数	6,873	100.0	592	8.6	4,612	67.1	1,669	24.3	1,411	20.5	258	3.8
	15-19歳	329	100.0	66	20.1	239	72.6	24	7.3	21	6.4	3	0.9
	20-29歳	535	100.0	95	17.8	374	69.9	66	12.3	49	9.2	17	3.2
	30-39歳	900	100.0	86	9.6	600	66.7	214	23.8	161	17.9	53	5.9
	40-49歳	1,014	100.0	67	6.6	668	65.9	279	27.5	226	22.3	53	5.2
	50-59歳	1,126	100.0	64	5.7	772	68.6	290	25.8	245	21.8	45	4.0
	60-69歳	1,410	100.0	74	5.2	951	67.4	385	27.3	336	23.8	49	3.5
	70歳以上	1,559	100.0	140	9.0	1,008	64.7	411	26.4	373	23.9	38	2.4
	(再掲)												
	20歳以上	6,544	100.0	526	8.0	4,373	66.8	1,645	25.1	1,390	21.2	255	3.9
	20-69歳	4,985	100.0	386	7.7	3,365	67.5	1,234	24.8	1,017	20.4	217	4.4
	40-69歳	3,550	100.0	205	5.8	2,391	67.4	954	26.9	807	22.7	147	4.1
	65-69歳	683	100.0	32	4.7	467	68.4	184	26.9	162	23.7	22	3.2
	70-74歳	582	100.0	37	6.4	385	66.2	160	27.5	142	24.4	18	3.1
	75-79歳	491	100.0	41	8.4	325	66.2	125	25.5	115	23.4	10	2.0
	80-84歳	292	100.0	31	10.6	180	61.6	81	27.7	75	25.7	6	2.1
	80歳以上	486	100.0	62	12.8	298	61.3	126	25.9	116	23.9	10	2.1
	85歳以上	194	100.0	31	16.0	118	60.8	45	23.2	41	21.1	4	2
男性	総 数	3,090	100.0	161	5.2	2,023	65.5	906	29.3	777	25.1	129	4.2
	15-19歳	168	100.0	32	19.0	121	72.0	15	8.9	13	7.7	2	1.2
	20-29歳	243	100.0	30	12.3	168	69.1	45	18.5	32	13.2	13	5.3
	30-39歳	405	100.0	15	3.7	249	61.5	141	34.8	105	25.9	36	8.9
	40-49歳	469	100.0	10	2.1	289	61.6	170	36.2	140	29.9	30	6.4
	50-59歳	520	100.0	14	2.7	333	64.0	173	33.3	154	29.6	19	3.7
	60-69歳	635	100.0	21	3.3	422	66.5	192	30.2	177	27.9	15	2.4
	70歳以上	650	100.0	39	6.0	441	67.8	170	26.2	156	24.0	14	2.2
	(再掲)												
	20歳以上	2,922	100.0	129	4.4	1,902	65.1	891	30.5	764	26.1	127	4.3
	20-69歳	2,272	100.0	90	4.0	1,461	64.3	721	31.7	608	26.8	113	5.0
	40-69歳	1,624	100.0	45	2.8	1,044	64.3	535	32.9	471	29.0	64	3.9
	65-69歳	313	100.0	7	2.2	217	69.3	89	28.4	83	26.5	6	1.9
	70-74歳	261	100.0	6	2.3	180	69.0	75	28.7	67	25.7	8	3.1
	75-79歳	214	100.0	13	6.1	145	67.8	56	26.2	53	24.8	3	1.4
	80-84歳	108	100.0	9	8.3	72	66.7	27	25.0	24	22.2	3	2.8
	80歳以上	175	100.0	20	11.4	116	66.3	39	22.3	36	20.6	3	1.7
	85歳以上	67	100.0	11	16.4	44	65.7	12	17.9	12	17.9	0	0
女性	総 数	3,783	100.0	431	11.4	2,589	68.4	763	20.2	634	16.8	129	3.4
	15-19歳	161	100.0	34	21.1	118	73.3	9	5.6	8	5.0	1	0.6
	20-29歳	292	100.0	65	22.3	206	70.5	21	7.2	17	5.8	4	1.4
	30-39歳	495	100.0	71	14.3	351	70.9	73	14.7	56	11.3	17	3.4
	40-49歳	545	100.0	57	10.5	379	69.5	109	20.0	86	15.8	23	4.2
	50-59歳	606	100.0	50	8.3	439	72.4	117	19.3	91	15.0	26	4.3
	60-69歳	775	100.0	53	6.8	529	68.3	193	24.9	159	20.5	34	4.4
	70歳以上	909	100.0	101	11.1	567	62.4	241	26.5	217	23.9	24	2.6
	(再掲)												
	20歳以上	3,622	100.0	397	11.0	2,471	68.2	754	20.8	626	17.3	128	3.5
	20-69歳	2,713	100.0	296	10.9	1,904	70.2	513	18.9	409	15.1	104	3.8
	40-69歳	1,926	100.0	160	8.3	1,347	69.9	419	21.8	336	17.4	83	4.3
	65-69歳	370	100.0	25	6.8	250	67.6	95	25.7	79	21.4	16	4.3
	70-74歳	321	100.0	31	9.7	205	63.9	85	26.5	75	23.4	10	3.1
	75-79歳	277	100.0	28	10.1	180	65.0	69	24.9	62	22.4	7	2.5
	80-84歳	184	100.0	22	12.0	108	58.7	54	29.3	51	27.7	3	1.6
	80歳以上	311	100.0	42	13.5	182	58.5	87	28.0	80	25.7	7	2.3
	85歳以上	127	100.0	20	15.7	74	58.3	33	26.0	29	22.8	4	3.1

注) 妊婦除外。BMI の判定は p. 17 参照

第15表 学校保健統計調査方式による肥満区分（6－14歳，性・年齢階級別）

		総 数		6－8歳		9－11歳		12－14歳	
		人 数	%	人 数	%	人 数	%	人 数	%
総数	総 数	751	100.0	239	100.0	270	100.0	242	100.0
	－30%未満	1	0.1	0	0.0	0	0.0	1	0.4
	－30%以上～－20%未満	25	3.3	1	0.4	12	4.4	12	5.0
	－20%以上～－10%未満	161	21.4	46	19.2	55	20.4	60	24.8
	－10%以上～ 0%未満	273	36.4	102	42.7	87	32.2	84	34.7
	0%以上～+10%未満	152	20.2	52	21.8	54	20.0	46	19.0
	+10%以上～+20%未満	74	9.9	24	10.0	29	10.7	21	8.7
	+20%以上～+30%未満	30	4.0	6	2.5	15	5.6	9	3.7
	+30%以上～+40%未満	22	2.9	6	2.5	10	3.7	6	2.5
	+40%以上～+50%未満	4	0.5	0	0.0	3	1.1	1	0.4
	+50%以上	9	1.2	2	0.8	5	1.9	2	0.8
男性	総 数	366	100.0	116	100.0	126	100.0	124	100.0
	－30%未満	0	0.0	0	0.0	0	0.0	0	0.0
	－30%以上～－20%未満	11	3.0	0	0.0	6	4.8	5	4.0
	－20%以上～－10%未満	85	23.2	25	21.6	26	20.6	34	27.4
	－10%以上～ 0%未満	131	35.8	46	39.7	42	33.3	43	34.7
	0%以上～+10%未満	72	19.7	27	23.3	23	18.3	22	17.7
	+10%以上～+20%未満	33	9.0	12	10.3	10	7.9	11	8.9
	+20%以上～+30%未満	16	4.4	3	2.6	9	7.1	4	3.2
	+30%以上～+40%未満	10	2.7	2	1.7	5	4.0	3	2.4
	+40%以上～+50%未満	3	0.8	0	0.0	2	1.6	1	0.8
	+50%以上	5	1.4	1	0.9	3	2.4	1	0.8
女性	総 数	385	100.0	123	100.0	144	100.0	118	100.0
	－30%未満	1	0.3	0	0.0	0	0.0	1	0.8
	－30%以上～－20%未満	14	3.6	1	0.8	6	4.2	7	5.9
	－20%以上～－10%未満	76	19.7	21	17.1	29	20.1	26	22.0
	－10%以上～ 0%未満	142	36.9	56	45.5	45	31.3	41	34.7
	0%以上～+10%未満	80	20.8	25	20.3	31	21.5	24	20.3
	+10%以上～+20%未満	41	10.6	12	9.8	19	13.2	10	8.5
	+20%以上～+30%未満	14	3.6	3	2.4	6	4.2	5	4.2
	+30%以上～+40%未満	12	3.1	4	3.3	5	3.5	3	2.5
	+40%以上～+50%未満	1	0.3	0	0.0	1	0.7	0	0.0
	+50%以上	4	1.0	1	0.8	2	1.4	1	0.8

注) 学校保健統計調査方式による区分は、p.17 参照

第16表 腹囲の分布（6歳以上、性・年齢階級別）

総数		6-8歳		9-11歳		12-14歳		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 20歳以上		(再掲) 40-74歳	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総数	7,163	100.0	226	100.0	253	100.0	214	100.0	292	100.0	479	100.0	973	100.0	1,067	100.0	1,359	100.0	1,455	100.0	6,178	100.0	3,954	100.0
	50cm未満	23	0.3	20	8.8	2	0.8	1	0.5	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	50－55cm未満	135	1.9	95	42.0	32	12.6	2	0.9	2	0.7	0	0.0	1	0.1	0	0.0	0	0.0	1	0.1	4	0.1	1	0.0
	55－60	178	2.5	77	34.1	63	24.9	15	7.0	2	0.7	6	1.3	1	0.1	3	0.3	3	0.2	3	0.2	21	0.3	7	0.2
	60－65	358	5.0	20	8.8	84	33.2	67	31.3	20	6.8	37	7.7	25	2.6	20	1.9	19	1.4	27	1.9	167	2.7	74	1.9
	65－70	631	8.8	9	4.0	32	12.6	61	28.5	87	29.8	100	20.9	82	8.4	46	4.3	58	4.3	58	4.0	442	7.2	204	5.2
	70－75	968	13.5	0	0.0	17	6.7	31	14.5	95	32.5	127	26.5	148	15.2	143	13.4	117	8.6	129	8.9	825	13.4	451	11.4
	75－80	1,084	15.1	4	1.8	13	5.1	20	9.3	45	15.4	83	17.3	170	17.5	200	18.7	195	14.3	195	13.4	1,002	16.2	640	16.2
	80－85	1,213	16.9	1	0.4	2	0.8	8	3.7	19	6.5	61	12.7	172	17.7	221	20.7	295	21.7	302	20.8	1,183	19.1	797	20.2
	85－90	1,129	15.8	0	0.0	4	1.6	8	3.7	11	3.8	50	10.4	160	16.4	198	18.6	297	21.9	313	21.5	1,106	17.9	779	19.7
	90－95	736	10.3	0	0.0	1	0.4	1	0.5	4	1.4	17	3.5	97	10.0	130	12.2	200	14.7	228	15.7	730	11.8	518	13.1
	95－100	397	5.5	0	0.0	2	0.8	0	0.0	3	1.0	12	2.5	41	7.3	60	5.6	100	7.4	114	7.8	392	6.3	286	7.2
男性	100－105	185	2.6	0	0.0	1	0.4	0	0.0	2	0.7	7	1.5	21	2.2	29	2.7	50	3.7	57	3.9	182	2.9	115	2.9
	105－110	70	1.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	10	1.0	10	0.9	14	1.0	19	1.3	70	1.1	45	1.1
	110－115	32	0.4	0	0.0	0	0.0	0	0.0	1	0.3	1	0.2	4	0.5	5	0.5	7	0.5	6	0.4	31	0.5	23	0.6
	115－120	11	0.2	0	0.0	0	0.0	0	0.0	2	0.4	2	0.2	2	0.2	2	0.2	1	0.1	2	0.1	11	0.2	6	0.2
	120cm以上	13	0.2	0	0.0	0	0.0	1	0.3	1	0.2	2	0.2	5	0.5	0	0.0	3	0.2	1	0.1	12	0.2	8	0.2
	総数	3,233	100.0	112	100.0	118	100.0	106	100.0	151	100.0	223	100.0	448	100.0	486	100.0	607	100.0	609	100.0	2,746	100.0	1,788	100.0
	50cm未満	7	0.2	6	5.4	1	0.8	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	50－55cm未満	61	1.9	47	42.0	13	11.0	1	0.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	55－60	83	2.6	40	35.7	37	31.4	5	4.7	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.0	1	0.1
	60－65	104	3.2	11	9.8	36	30.5	35	33.0	7	4.6	5	2.2	3	0.8	1	0.2	2	0.3	2	0.3	15	0.5	5	0.3
	65－70	177	5.5	6	5.4	17	14.4	30	28.3	34	22.5	32	14.3	14	3.8	4	0.8	11	1.8	17	2.8	90	3.3	32	1.8
	70－75	306	9.5	0	0.0	5	4.2	18	17.0	54	35.8	56	25.1	44	11.8	30	6.2	32	5.3	42	6.9	229	8.3	99	5.5
	75－80	429	13.3	2	1.8	4	3.4	9	8.5	29	19.2	48	21.5	62	16.6	79	16.3	61	10.0	72	11.8	385	14.0	230	12.9
	80－85	632	19.5	0	0.0	0	0.0	3	2.8	11	7.3	35	15.7	81	21.7	112	23.0	149	24.5	145	23.8	618	22.5	415	23.2
	85－90	637	19.7	0	0.0	2	1.7	5	4.7	8	5.3	15	6.7	70	18.8	114	23.5	155	25.5	163	26.8	622	22.7	445	24.9
女性	90－95	420	13.0	0	0.0	1	0.8	0	0.0	4	2.6	12	5.4	41	11.0	81	16.7	116	19.1	95	15.6	415	15.1	309	17.3
	95－100	220	6.8	0	0.0	0	0.0	0	0.0	2	1.3	9	4.0	25	6.7	35	7.2	54	8.9	46	7.6	217	7.9	161	9.0
	100－105	99	3.1	0	0.0	1	0.8	0	0.0	0	0.0	6	2.7	18	4.8	19	3.9	17	2.8	22	3.6	98	3.6	57	3.2
	105－110	36	1.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.4	11	2.9	8	1.6	5	0.8	4	0.7	36	1.3	23	1.3
	110－115	11	0.3	0	0.0	0	0.0	0	0.0	1	0.7	1	0.4	2	0.4	2	0.4	1	0.2	1	0.2	10	0.4	5	0.3
	115－120	5	0.2	0	0.0	0	0.0	0	0.0	2	0.9	1	0.3	1	0.3	1	0.2	0	0.0	0	0.0	5	0.2	2	0.1
	120cm以上	6	0.2	0	0.0	0	0.0	1	0.7	1	0.4	0	0.0	1	0.2	0	0.0	3	0.5	0	0.0	5	0.2	4	0.2
	総数	3,930	100.0	114	100.0	135	100.0	108	100.0	141	100.0	256	100.0	525	100.0	581	100.0	752	100.0	846	100.0	3,432	100.0	2,166	100.0
	50cm未満	16	0.4	14	12.3	1	0.7	1	0.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	50－55cm未満	74	1.9	48	42.1	19	14.1	1	0.9	2	1.4	0	0.0	1	0.2	0	0.0	0	0.0	1	0.1	4	0.1	1	0.0
	55－60	95	2.4	37	32.5	26	19.3	10	9.3	2	1.4	6	2.3	5	1.1	3	0.5	2	0.3	3	0.4	20	0.6	6	0.3
	60－65	254	6.5	9	7.9	48	35.6	32	29.6	13	9.2	32	12.5	23	4.4	19	3.3	17	2.3	25	3.0	152	4.4	69	3.2
	65－70	454	11.6	3	2.6	15	11.1	31	28.7	53	37.6	68	26.6	70	13.3	42	7.2	47	6.3	41	4.8	352	10.3	172	7.9
	70－75	662	16.8	0	0.0	12	8.9	13	12.0	41	29.1	71	27.7	123	23.4	113	19.4	85	11.3	87	10.3	596	17.4	352	16.3
	75－80	655	16.7	2	1.8	9	6.7	11	10.2	16	11.3	35	13.7	97	20.6	121	20.8	134	17.8	123	14.5	617	18.0	410	18.9
	80－85	581	14.8	1	0.9	2	1.5	5	4.6	8	5.7	26	10.2	76	14.5	109	18.8	146	19.4	157	18.6	565	16.5	382	17.6
	85－90	492	12.5	0	0.0	2	1.5	3	2.8	3	2.1	9	3.5	55	10.5	84	14.5	142	18.9	150	17.7	484	14.1	334	15.4
	90－95	316	8.0	0	0.0	0	0.0	1	0.9	0	0.0	5	2.0	17	3.6	27	5.1	49	8.4	133	15.7	315	9.2	209	9.6
注）妊婦除外	95－100	177	4.5	0	0.0	1	0.7	0	0.0	1	0.7	3	1.2	10	2.1	25	4.3	46	6.1	68	8.0	175	5.1	125	5.8
	100－105	86	2.2	0	0.0	0	0.0	0	0.0	2	1.4	1	0.4	5	1.0	10	1.7	33	4.4	35	4.1	84	2.4	58	2.7
	105－110	34	0.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	3	0.6	2	0.3	9	1.2	15	1.8	34	1.0	22	1.0
	110－115	21	0.5	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	6	1.1	3	0.5	6	0.8	5	0.6	21	0.6	18	0.8
	115－120	6	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	1	0.2	1	0.1	2	0.2	6	0.2	4	0.2
	120cm以上	7	0.2	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	4	0.8	0	0.0	0	0.0	1	0.1	7	0.2	4	0.2

第 17 表 腹囲の区分（20 歳以上，性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	総 数	6,178	100.0	479	100.0	845	100.0	973	100.0	1,067	100.0	1,359	100.0	1,455	100.0	3,954	100.0
	<85cm(男)/<90cm(女) ≥85cm(男)/≥90cm(女)	4,128 2,050	66.8 33.2	423 56	88.3 11.7	640 205	75.7 24.3	654 319	67.2 32.8	717 350	67.2 32.8	829 530	61.0 39.0	865 590	59.5 40.5	2,508 1,446	63.4 36.6
男 性	総 数	2,746	100.0	223	100.0	373	100.0	448	100.0	486	100.0	607	100.0	609	100.0	1,788	100.0
	<85cm ≥85cm	1,338 1,408	48.7 51.3	176 47	78.9 21.1	204 169	54.7 45.3	198 250	44.2 55.8	226 260	46.5 53.5	256 351	42.2 57.8	278 331	45.6 54.4	782 1,006	43.7 56.3
女 性	総 数	3,432	100.0	256	100.0	472	100.0	525	100.0	581	100.0	752	100.0	846	100.0	2,166	100.0
	<90cm ≥90cm	2,790 642	81.3 18.7	247 9	96.5 3.5	436 36	92.4 7.6	456 69	86.9 13.1	491 90	84.5 15.5	573 179	76.2 23.8	587 259	69.4 30.6	1,726 440	79.7 20.3

注) 妊婦除外

第 18 表 BMI・腹囲の区分（20 歳以上，性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	6,147	100.0	477	100.0	839	100.0	963	100.0	1,064	100.0	1,354	100.0	1,450	100.0	3,936	100.0
	BMI<25,腹囲<85cm(男)/90cm(女)	3,804	61.9	410	86.0	588	70.1	594	61.7	663	62.3	762	56.3	787	54.3	2,300	58.4
	BMIのみ超(BMI≥25)	305	5.0	11	2.3	48	5.7	53	5.5	53	5.0	64	4.7	76	5.2	197	5.0
	腹囲のみ超(腹囲≥85cm(男)/90cm(女))	788	12.8	8	1.7	53	6.3	99	10.3	127	11.9	223	16.5	278	19.2	571	14.5
男性	BMI≥25,腹囲≥85cm(男)/90cm(女)	1,250	20.3	48	10.1	150	17.9	217	22.5	221	20.8	305	22.5	309	21.3	868	22.1
	総 数	2,727	100.0	221	100.0	369	100.0	443	100.0	483	100.0	604	100.0	607	100.0	1,777	100.0
	BMI<25,腹囲<85cm	1,276	46.8	173	78.3	189	51.2	187	42.2	214	44.3	249	41.2	264	43.5	744	41.9
	BMIのみ超(BMI≥25)	52	1.9	1	0.5	13	3.5	8	1.8	11	2.3	5	0.8	14	2.3	32	1.8
女性	腹囲のみ超(腹囲≥85cm)	609	22.3	7	3.2	49	13.3	91	20.5	109	22.6	172	28.5	181	29.8	454	25.5
	BMI≥25,腹囲≥85cm	790	29.0	40	18.1	118	32.0	157	35.4	149	30.8	178	29.5	148	24.4	547	30.8
	総 数	3,420	100.0	256	100.0	470	100.0	520	100.0	581	100.0	750	100.0	843	100.0	2,159	100.0
	BMI<25,腹囲<90cm	2,528	73.9	237	92.6	399	84.9	407	78.3	449	77.3	513	68.4	523	62.0	1,556	72.1
	BMIのみ超(BMI≥25)	253	7.4	10	3.9	35	7.4	45	8.7	42	7.2	59	7.9	62	7.4	165	7.6
	腹囲のみ超(腹囲≥90cm)	179	5.2	1	0.4	4	0.9	8	1.5	18	3.1	51	6.8	97	11.5	117	5.4
	BMI≥25,腹囲≥90cm	460	13.5	8	3.1	32	6.8	60	11.5	72	12.4	127	16.9	161	19.1	321	14.9

注) 妊婦除外

【参考】 内臓脂肪型肥満の診断基準：
● BMI 25 以上で，男性のウエスト周囲径 85cm 以上，女性のウエスト周囲径 90cm 以上を上半身肥満の疑いとする。
● 上半身肥満の疑いと判定され，腹部CT法による内臓脂肪面積 100cm²以上（男女とも）を内臓脂肪型肥満と診断する。（日本肥満学会肥満症診断基準検討委員会，2000 年）
※国民健康・栄養調査の「腹囲」は，「立位のへその高さ」で計測したが，計測位置は「ウエスト周囲径」と同じである。

第19表 薬の服用状況（性・年齢階級別）

		総数		20－29歳		30－39歳		40－49歳		50－59歳		60－69歳		70歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	4,872	100.0	298	100.0	601	100.0	708	100.0	841	100.0	1,149	100.0	1,275	100.0
	使用なし	3,270	67.1	293	98.3	591	98.3	643	90.8	649	77.2	634	55.2	460	36.1
	使用あり(一つ以上)	1,602	32.9	5	1.7	10	1.7	65	9.2	192	22.8	515	44.8	815	63.9
	血圧を下げる薬	1,246	25.6	2	0.7	3	0.5	42	5.9	133	15.8	391	34.0	675	52.9
	脈の乱れを治す薬	188	3.9	0	0.0	0	0.0	6	0.8	9	1.1	44	3.8	129	10.1
	インスリン注射又は血糖を下げる薬	289	5.9	1	0.3	0	0.0	16	2.3	45	5.4	91	7.9	136	10.7
	コレステロールを下げる薬	573	11.8	1	0.3	2	0.3	12	1.7	60	7.1	216	18.8	282	22.1
	中性脂肪(トリグリセライド)を下げる薬	157	3.2	1	0.3	1	0.2	8	1.1	15	1.8	56	4.9	76	6.0
	貧血治療のための薬(鉄剤)	76	1.6	1	0.3	5	0.8	12	1.7	11	1.3	19	1.7	28	2.2
男性	総 数	2,019	100.0	121	100.0	239	100.0	288	100.0	346	100.0	498	100.0	527	100.0
	使用なし	1,337	66.2	119	98.3	236	98.7	253	87.8	249	72.0	271	54.4	209	39.7
	使用あり(一つ以上)	682	33.8	2	1.7	3	1.3	35	12.2	97	28.0	227	45.6	318	60.3
	血圧を下げる薬	545	27.0	1	0.8	0	0.0	27	9.4	70	20.2	190	38.2	257	48.8
	脈の乱れを治す薬	90	4.5	0	0.0	0	0.0	2	0.7	6	1.7	25	5.0	57	10.8
	インスリン注射又は血糖を下げる薬	160	7.9	0	0.0	0	0.0	10	3.5	27	7.8	52	10.4	71	13.5
	コレステロールを下げる薬	170	8.4	1	0.8	2	0.8	6	2.1	20	5.8	62	12.4	79	15.0
	中性脂肪(トリグリセライド)を下げる薬	74	3.7	0	0.0	1	0.4	6	2.1	9	2.6	29	5.8	29	5.5
	貧血治療のための薬(鉄剤)	22	1.1	0	0.0	0	0.0	1	0.3	2	0.6	8	1.6	11	2.1
女性	総 数	2,853	100.0	177	100.0	362	100.0	420	100.0	495	100.0	651	100.0	748	100.0
	使用なし	1,933	67.8	174	98.3	355	98.1	390	92.9	400	80.8	363	55.8	251	33.6
	使用あり(一つ以上)	920	32.2	3	1.7	7	1.9	30	7.1	95	19.2	288	44.2	497	66.4
	血圧を下げる薬	701	24.6	1	0.6	3	0.8	15	3.6	63	12.7	201	30.9	418	55.9
	脈の乱れを治す薬	98	3.4	0	0.0	0	0.0	4	1.0	3	0.6	19	2.9	72	9.6
	インスリン注射又は血糖を下げる薬	129	4.5	1	0.6	0	0.0	6	1.4	18	3.6	39	6.0	65	8.7
	コレステロールを下げる薬	403	14.1	0	0.0	0	0.0	6	1.4	40	8.1	154	23.7	203	27.1
	中性脂肪(トリグリセライド)を下げる薬	83	2.9	1	0.6	0	0.0	2	0.5	6	1.2	27	4.1	47	6.3
	貧血治療のための薬(鉄剤)	54	1.9	1	0.6	5	1.4	11	2.6	9	1.8	11	1.7	17	2.3

注) 妊婦除外。

* 複数回答

第20表 血圧の状況（性・年齢階級別）

		総数		至適血圧		正常血圧		正常高値血圧		Ⅰ度高血圧		Ⅱ度高血圧		Ⅲ度高血圧		(再掲) 収縮期高血圧	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	3,425	100.0	1,201	35.1	674	19.7	628	18.3	649	18.9	208	6.1	65	1.9	400	11.7
	15-19歳	129	100.0	105	81.4	18	14.0	6	4.7	0	0.0	0	0.0	0	0.0	0	0.0
	20-29歳	241	100.0	171	71.0	47	19.5	19	7.9	3	1.2	0	0.0	1	0.4	1	0.0
	30-39歳	528	100.0	318	60.2	104	19.7	57	10.8	35	6.6	11	2.1	3	0.6	12	2.3
	40-49歳	596	100.0	257	43.1	122	20.5	106	17.8	83	13.9	22	3.7	6	1.0	17	2.9
	50-59歳	650	100.0	160	24.6	138	21.2	148	22.8	145	22.3	45	6.9	14	2.2	67	10.3
	60-69歳	717	100.0	124	17.3	146	20.4	175	24.4	184	25.7	68	9.5	20	2.8	114	15.9
	70歳以上	564	100.0	66	11.7	99	17.6	117	20.7	199	35.3	62	11.0	21	3.7	189	33.5
	(再掲) 65-74歳	591	100.0	78	13.2	110	18.6	142	24.0	185	31.3	56	9.5	20	3.4	133	22.5
	(再掲) 75歳以上	305	100.0	32	10.5	53	17.4	67	22.0	106	34.8	41	13.4	6	2.0	113	37.0
男性	総 数	1,363	100.0	316	23.2	272	20.0	287	21.1	332	24.4	114	8.4	42	3.1	183	13.4
	15-19歳	67	100.0	46	68.7	16	23.9	5	7.5	0	0.0	0	0.0	0	0.0	0	0.0
	20-29歳	96	100.0	51	53.1	27	28.1	15	15.6	2	2.1	0	0.0	1	1.0	1	1.0
	30-39歳	202	100.0	73	36.1	52	25.7	44	21.8	24	11.9	6	3.0	3	1.5	8	4.0
	40-49歳	222	100.0	63	28.4	50	22.5	42	18.9	44	19.8	18	8.1	5	2.3	5	2.3
	50-59歳	241	100.0	37	15.4	42	17.4	60	24.9	67	27.8	24	10.0	11	4.6	25	10.4
	60-69歳	288	100.0	29	10.1	48	16.7	78	27.1	84	29.2	37	12.8	12	4.2	46	16.0
	70歳以上	247	100.0	17	6.9	37	15.0	43	17.4	111	44.9	29	11.7	10	4.0	98	39.7
	(再掲) 65-74歳	251	100.0	20	8.0	34	13.5	65	25.9	88	35.1	32	12.7	12	4.8	62	24.7
	(再掲) 75歳以上	137	100.0	10	7.3	25	18.2	23	16.8	60	43.8	17	12.4	2	1.5	56	40.9
女性	総 数	2,062	100.0	885	42.9	402	19.5	341	16.5	317	15.4	94	4.6	23	1.1	217	10.5
	15-19歳	62	100.0	59	95.2	2	3.2	1	1.6	0	0.0	0	0.0	0	0.0	0	0.0
	20-29歳	145	100.0	120	82.8	20	13.8	4	2.8	1	0.7	0	0.0	0	0.0	0	0.0
	30-39歳	326	100.0	245	75.2	52	16.0	13	4.0	11	3.4	5	1.5	0	0.0	4	1.2
	40-49歳	374	100.0	194	51.9	72	19.3	64	17.1	39	10.4	4	1.1	1	0.3	12	3.2
	50-59歳	409	100.0	123	30.1	96	23.5	88	21.5	78	19.1	21	5.1	3	0.7	42	10.3
	60-69歳	429	100.0	95	22.1	98	22.8	97	22.6	100	23.3	31	7.2	8	1.9	68	15.9
	70歳以上	317	100.0	49	15.5	62	19.6	74	23.3	88	27.8	33	10.4	11	3.5	91	28.7
	(再掲) 65-74歳	340	100.0	58	17.1	76	22.4	77	22.6	97	28.5	24	7.1	8	2.4	71	20.9
	(再掲) 75歳以上	168	100.0	22	13.1	28	16.7	44	26.2	46	27.4	24	14.3	4	2.4	57	33.9

注) 妊婦，血圧を下げる薬服用者除外。日本高血圧学会による血圧の分類は p .16 を参照。

数値は2回の測定値の平均値を用いた。なお，1回しか測定できなかった者については，その値を採用した。

* 総数に対する割合

第 21 表 収縮期（最高）・拡張期（最低）血圧の分布（性・年齢階級別）

			総 数		15－19歳		20－29歳		30－39歳		40－49歳		50－59歳		60－69歳		70歳以上	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	収縮期（最高）血圧	総 数	1,363	100.0	67	100.0	96	100.0	202	100.0	222	100.0	241	100.0	288	100.0	247	100.0
		90mmHg未満	3	0.2	1	1.5	0	0.0	1	0.5	0	0.0	0	0.0	1	0.3	0	0.0
		90－99	19	1.4	3	4.5	5	5.2	5	2.5	1	0.5	3	1.2	0	0.0	2	0.8
		100－109	111	8.1	18	26.9	18	18.8	17	8.4	29	13.1	13	5.4	11	3.8	5	2.0
		110－119	233	17.1	26	38.8	31	32.3	64	31.7	46	20.7	28	11.6	27	9.4	11	4.5
		120－129	304	22.3	14	20.9	28	29.2	60	29.7	62	27.9	49	20.3	53	18.4	38	15.4
		130－139	280	20.5	5	7.5	11	11.5	35	17.3	45	20.3	63	26.1	77	26.7	44	17.8
		140－149	192	14.1	0	0.0	2	2.1	10	5.0	23	10.4	41	17.0	51	17.7	65	26.3
		150－159	122	9.0	0	0.0	0	0.0	8	4.0	9	4.1	25	10.4	31	10.8	49	19.8
		160－169	50	3.7	0	0.0	0	0.0	1	0.5	6	2.7	10	4.1	18	6.3	15	6.1
		170－179	28	2.1	0	0.0	1	1.0	0	0.0	1	0.5	5	2.1	11	3.8	10	4.0
		180mmHg以上	21	1.5	0	0.0	0	0.0	1	0.5	0	0.0	4	1.7	8	2.8	8	3.2
		平均値 mmHg 標準偏差	131.7 18.8		113.8 10.0		118.1 12.5		122.8 13.3		126.2 15.0		134.9 17.8		138.4 19.5		143 18.1	
	拡張期（最低）血圧	総 数	1,363	100.0	67	100.0	96	100.0	202	100.0	222	100.0	241	100.0	288	100.0	247	100.0
		40mmHg未満	1	0.1	1	1.5	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		40－49	3	0.2	3	4.5	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		50－59	33	2.4	11	16.4	5	5.2	4	2.0	1	0.5	2	0.8	5	1.7	5	2.0
		60－69	207	15.2	32	47.8	32	33.3	36	17.8	22	9.9	18	7.5	28	9.7	39	15.8
		70－79	390	28.6	14	20.9	39	40.6	78	38.6	63	28.4	62	25.7	64	22.2	70	28.3
		80－89	424	31.1	6	9.0	18	18.8	59	29.2	74	33.3	82	34.0	104	36.1	81	32.8
		90－99	208	15.3	0	0.0	1	1.0	16	7.9	42	18.9	50	20.7	60	20.8	39	15.8
		100－109	66	4.8	0	0.0	0	0.0	6	3.0	15	6.8	19	7.9	19	6.6	7	2.8
		110－119	23	1.7	0	0.0	0	0.0	3	1.5	3	1.4	7	2.9	6	2.1	4	1.6
女性	収縮期（最高）血圧	120－129	6	0.4	0	0.0	1	1.0	0	0.0	2	0.9	0	0.0	1	0.3	2	0.8
		130－139	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.4	1	0.3	0	0.0
		140mmHg以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		平均値 mmHg 標準偏差	80.9 12.5		65.4 9.8		73.4 10.0		77.8 10.8		83.5 11.6		85.0 11.9		84.0 12.0		80.6 12.0	
	拡張期（最低）血圧	総 数	2,061	100.0	62	100.0	145	100.0	326	100.0	374	100.0	409	100.0	429	100.0	316	100.0
		90mmHg未満	11	0.5	1	1.6	2	1.4	4	1.2	3	0.8	0	0.0	0	0.0	1	0.3
		90－99	154	7.5	15	24.2	24	16.6	49	15.0	37	9.9	13	3.2	10	2.3	6	1.9
		100－109	379	18.4	30	48.4	63	43.4	111	34.0	75	20.1	48	11.7	35	8.2	17	5.4
		110－119	409	19.8	13	21.0	34	23.4	92	28.2	99	26.5	79	19.3	63	14.7	29	9.2
		120－129	398	19.3	2	3.2	19	13.1	49	15.0	77	20.6	98	24.0	90	21.0	63	19.9
		130－139	329	16.0	1	1.6	3	2.1	10	3.1	56	15.0	83	20.3	101	23.5	75	23.7
		140－149	196	9.5	0	0.0	0	0.0	9	2.8	17	4.5	55	13.4	60	14.0	55	17.4
		150－159	92	4.5	0	0.0	0	0.0	0	0.0	7	1.9	17	4.2	40	9.3	28	8.9
		160－169	51	2.5	0	0.0	0	0.0	2	0.6	2	0.5	9	2.2	17	4.0	21	6.6
女性	収縮期（最高）血圧	170－179	22	1.1	0	0.0	0	0.0	0	0.0	0	0.0	4	1.0	8	1.9	10	3.2
		180mmHg以上	20	1.0	0	0.0	0	0.0	0	0.0	1	0.3	3	0.7	5	1.2	11	3.5
		平均値 mmHg 標準偏差	123.3 19.4		105.1 9.1		107.7 9.8		110.5 12.7		117.7 15.2		127.0 17.3		132.1 18.4		137.2 20.4	
	拡張期（最低）血圧	総 数	2,061	100.0	62	100.0	145	100.0	326	100.0	374	100.0	409	100.0	429	100.0	316	100.0
		40mmHg未満	1	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0
		40－49	5	0.2	1	1.6	0	0.0	3	0.9	1	0.3	0	0.0	0	0.0	0	0.0
		50－59	118	5.7	18	29.0	24	16.6	25	7.7	26	7.0	4	1.0	8	1.9	13	4.1
		60－69	509	24.7	31	50.0	66	45.5	134	41.1	92	24.6	60	14.7	70	16.3	56	17.7
		70－79	725	35.2	12	19.4	47	32.4	119	36.5	138	36.9	142	34.7	156	36.4	111	35.1
		80－89	486	23.6	0	0.0	7	4.8	32	9.8	85	22.7	143	35.0	124	28.9	95	30.1
		90－99	173	8.4	0	0.0	1	0.7	7	2.1	29	7.8	49	12.0	54	12.6	33	10.4
		100－109	39	1.9	0	0.0	0	0.0	5	1.5	3	0.8	10	2.4	13	3.0	8	2.5
		110－119	3	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	3	0.7	0	0.0
女性	収縮期（最高）血圧	120－129	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
		130－139	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0
		140mmHg以上	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0
		平均値 mmHg 標準偏差	75.5 11.1		63.3 7.1		67.4 7.7		70.3 9.7		74.5 10.4		79.6 10.2		79.2 11.1		77.5 10.6	
	拡張期（最低）血圧	総 数	2,061	100.0	62	100.0	145	100.0	326	100.0	374	100.0	409	100.0	429	100.0	316	100.0
		40mmHg未満	1	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0
		40－49	5	0.2	1	1.6	0	0.0	3	0.9	1	0.3	0	0.0	0	0.0	0	0.0
		50－59	118	5.7	18	29.0	24	16.6	25	7.7	26	7.0	4	1.0	8	1.9	13	4.1
		60－69	509	24.7	31	50.0	66	45.5	134	41.1	92	24.6	60	14.7	70	16.3	56	17.7
		70－79	725	35.2	12	19.4	47	32.4	119	36.5	138	36.9	142	34.7	156	36.4	111	35.1
		80－89	486	23.6	0	0.0	7	4.8	32	9.8	85	22.7	143	35.0	124	28.9	95	30.1
		90－99	173	8.4	0	0.0	1	0.7	7	2.1	29	7.8	49	12.0	54	12.6	33	10.4
		100－109	39	1.9	0	0.0	0	0.0	5	1.5	3	0.8	10	2.4	13	3.0	8	2.5
		110－119	3	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	3	0.7	0	0.0

注）妊婦，血圧を下げる薬服用者除外。

数値は 2 回の測定値の平均値を用いた。なお，1 回しか測定できなかった者については，その値を採用した。

第22表 血色素量の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,708	100.0	90	100.0	196	100.0	238	100.0	293	100.0	440	100.0	451	100.0	416	100.0	263	100.0
	10.0g/dl未満	6	0.4	0	0.0	0	0.0	0	0.0	1	0.3	2	0.5	3	0.7	1	0.2	3	1.1
	10.0-10.9	9	0.5	0	0.0	0	0.0	0	0.0	2	0.7	2	0.5	5	1.1	3	0.7	3	1.1
	11.0-11.9	37	2.2	0	0.0	0	0.0	0	0.0	4	1.4	12	2.7	21	4.7	13	3.1	18	6.8
	12.0-12.9	97	5.7	0	0.0	1	0.5	5	2.1	5	1.7	26	5.9	60	13.3	32	7.7	40	15.2
	13.0-13.9	232	13.6	6	6.7	14	7.1	10	4.2	32	10.9	76	17.3	94	20.8	85	20.4	55	20.9
	14.0-14.9	501	29.3	18	20.0	45	23.0	59	24.8	93	31.7	147	33.4	139	30.8	144	34.6	79	30.0
	15.0-15.9	537	31.4	43	47.8	89	45.4	108	45.4	91	31.1	112	25.5	94	20.8	95	22.8	49	18.6
	16.0-16.9	222	13.0	15	16.7	34	17.3	42	17.6	52	17.7	51	11.6	28	6.2	35	8.4	13	4.9
	17.0-17.9	57	3.3	6	6.7	13	6.6	13	5.5	9	3.1	9	2.0	7	1.6	6	1.4	3	1.1
	18.0g/dl以上	10	0.6	2	2.2	0	0.0	1	0.4	4	1.4	3	0.7	0	0.0	2	0.5	0	0.0
平均値(g/dl)		14.8		15.5		15.4		15.3		15.0		14.6		14.1		14.4		13.9	
標準偏差		1.3		1.0		0.9		1.0		1.2		1.3		1.4		1.3		1.4	
女性	総 数	2,484	100.0	138	100.0	331	100.0	364	100.0	439	100.0	568	100.0	644	100.0	528	100.0	398	100.0
	10.0g/dl未満	57	2.3	3	2.2	13	3.9	25	6.9	4	0.9	3	0.5	9	1.4	3	0.6	8	2.0
	10.0-10.9	76	3.1	0	0.0	21	6.3	16	4.4	8	1.8	2	0.4	29	4.5	10	1.9	20	5.0
	11.0-11.9	306	12.3	22	15.9	37	11.2	44	12.1	32	7.3	48	8.5	123	19.1	60	11.4	87	21.9
	12.0-12.9	754	30.4	49	35.5	114	34.4	109	29.9	126	28.7	159	28.0	197	30.6	152	28.8	129	32.4
	13.0-13.9	841	33.9	40	29.0	118	35.6	111	30.5	164	37.4	228	40.1	180	28.0	193	36.6	97	24.4
	14.0-14.9	379	15.3	21	15.2	24	7.3	51	14.0	85	19.4	102	18.0	96	14.9	96	18.2	52	13.1
	15.0-15.9	65	2.6	3	2.2	4	1.2	6	1.6	17	3.9	25	4.4	10	1.6	14	2.7	5	1.3
	16.0-16.9	5	0.2	0	0.0	0	0.0	1	0.3	3	0.7	1	0.2	0	0.0	0	0.0	0	0.0
	17.0-17.9	1	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	18.0g/dl以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
平均値(g/dl)		12.9		12.8		12.6		12.6		13.2		13.3		12.8		13.1		12.6	
標準偏差		1.2		1.3		1.3		1.6		1.1		1.0		1.2		1.1		1.2	

注) 貧血治療のための薬服用者除外。

第23表 赤血球数の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,725	100.0	90	100.0	196	100.0	239	100.0	295	100.0	445	100.0	460	100.0	423	100.0	268	100.0
	400万個/mm ³ 未満	115	6.7	0	0.0	1	0.5	2	0.8	11	3.7	27	6.1	74	16.1	38	9.0	54	20.1
	400-449	323	18.7	3	3.3	10	5.1	18	7.5	44	14.9	103	23.1	145	31.5	114	27.0	88	32.8
	450-499	746	43.2	26	28.9	71	36.2	104	43.5	147	49.8	215	48.3	183	39.8	199	47.0	98	36.6
	500-449	482	27.9	49	54.4	98	50.0	105	43.9	87	29.5	89	20.0	54	11.7	67	15.8	26	9.7
	550-599	52	3.0	10	11.1	16	8.2	9	3.8	5	1.7	9	2.0	3	0.7	5	1.2	1	0.4
	600-649	7	0.4	2	2.2	0	0.0	1	0.4	1	0.3	2	0.4	1	0.2	0	0.0	1	0.4
	650万個/mm ³ 以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
平均値(g/dl)		474.9		514.4		504.0		495.0		479.5		467.9		448.2		459.9		442.5	
標準偏差		47.2		33.2		33.6		35.1		41.8		44.3		48.6		44.9		49.1	
女性	総 数	2,534	100.0	139	100.0	336	100.0	375	100.0	446	100.0	579	100.0	659	100.0	538	100.0	408	100.0
	400万個/mm ³ 未満	399	15.7	11	7.9	33	9.8	45	12.0	40	9.0	73	12.6	197	29.9	95	17.7	138	33.8
	400-449	1,233	48.7	63	45.3	173	51.5	187	49.9	214	48.0	289	49.9	307	46.6	267	49.6	190	46.6
	450-499	807	31.8	57	41.0	119	35.4	123	32.8	170	38.1	196	33.9	142	21.5	159	29.6	74	18.1
	500-449	91	3.6	8	5.8	10	3.0	18	4.8	21	4.7	21	3.6	13	2.0	17	3.2	6	1.5
	550-599	3	0.1	0	0.0	0	0.0	2	0.5	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0
	600-649	1	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	650万個/mm ³ 以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
平均値(g/dl)		435.4		447.5		439.7		438.9		444.3		438.6		419.7		432.0		415.1	
標準偏差		37.6		33.9		33.0		35.5		34.7		34.8		41.0		38.4		39.8	

第24表 ヘマトクリット値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,725	100.0	90	100.0	196	100.0	239	100.0	295	100.0	445	100.0	460	100.0	423	100.0
	25%未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	25-29	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.0
	30-34	23	1.3	0	0.0	0	0.0	0	0.0	4	1.4	6	1.3	13	2.8	6	4.1
	35-39	115	6.7	1	1.1	1	0.5	3	1.3	8	2.7	32	7.2	70	15.2	35	20.5
	40-44	617	35.8	23	25.6	44	22.4	56	23.4	97	32.9	180	40.4	217	47.2	198	44.8
	45-49	809	46.9	50	55.6	125	63.8	156	65.3	145	49.2	191	42.9	142	30.9	158	29.1
	50-54	155	9.0	16	17.8	25	12.8	23	9.6	39	13.2	35	7.9	17	3.7	25	1.5
	55%以上	5	0.3	0	0.0	1	0.5	1	0.4	2	0.7	1	0.2	0	0.0	0	0.0
	(再掲)39.8未満	133	7.7	1	1.1	0	0.0	3	1.3	12	4.1	37	8.3	80	17.4	40	23.5
	(再掲)51.9以上	43	2.5	2	2.2	6	3.1	5	2.1	13	4.4	12	2.7	5	1.1	7	1.1
平均値(g/dl)		45.2		46.9		46.9		46.6		45.9		44.9		43.3		44.3	
標準偏差		3.8		2.9		2.7		2.8		3.7		3.7		4.1		3.8	
女性	総 数	2,534	100.0	139	100.0	336	100.0	375	100.0	446	100.0	579	100.0	659	100.0	538	100.0
	25%未満	3	0.1	1	0.7	0	0.0	1	0.3	1	0.2	0	0.0	0	0.0	0	0.0
	25-29	20	0.8	0	0.0	5	1.5	8	2.1	1	0.2	1	0.2	5	0.8	3	0.6
	30-34	110	4.3	5	3.6	22	6.5	28	7.5	8	1.8	7	1.2	40	6.1	14	2.6
	35-39	884	34.9	57	41.0	138	41.1	133	35.5	123	27.6	161	27.8	272	41.3	171	31.8
	40-44	1,308	51.6	64	46.0	160	47.6	186	49.6	268	60.1	344	59.4	286	43.4	296	55.0
	45-49	203	8.0	12	8.6	11	3.3	18	4.8	44	9.9	63	10.9	55	8.3	51	9.5
	50-54	6	0.2	0	0.0	0	0.0	1	0.3	1	0.2	3	0.5	1	0.2	3	0.6
	55%以上	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	(再掲)33.4未満	76	3.0	3	2.2	17	5.1	26	6.9	2	0.4	4	0.7	24	3.6	9	1.7
	(再掲)45.0以上	209	8.2	12	8.6	11	3.3	19	5.1	45	10.1	66	11.4	56	8.5	54	10.0
平均値(g/dl)		40.5		40.3		39.7		39.9		41.4		41.4		40.0		41.0	
標準偏差		3.5		3.5		3.4		4.0		3.2		3.1		3.6		3.3	

第25表 平均赤血球容積（MCV）の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,725	100.0	90	100.0	196	100.0	239	100.0	295	100.0	445	100.0	460	100.0	423	100.0
	70fl未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	70-74	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	75-79	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	0	0.0
	80-84	21	1.2	3	3.3	2	1.0	6	2.5	1	0.3	4	0.9	5	1.1	5	1.2
	85-89	200	11.6	32	35.6	50	25.5	29	12.1	27	9.2	35	7.9	27	5.9	29	6.9
	90-94	614	35.6	39	43.3	85	43.4	106	44.4	104	35.3	157	35.3	123	26.7	135	31.9
	95-99	603	35.0	15	16.7	47	24.0	78	32.6	114	38.6	159	35.7	190	41.3	158	37.4
	100-104	221	12.8	1	1.1	8	4.1	17	7.1	38	12.9	67	15.1	90	19.6	70	16.5
	105-109	44	2.6	0	0.0	3	1.5	2	0.8	6	2.0	18	4.0	15	3.3	17	4.0
	110-114	13	0.8	0	0.0	0	0.0	0	0.0	4	1.4	2	0.4	7	1.5	6	1.4
	115-119	3	0.2	0	0.0	1	0.5	1	0.4	0	0.0	1	0.2	0	0.0	1	0.2
	120fl以上	4	0.2	0	0.0	0	0.0	0	0.0	1	0.3	2	0.4	1	0.2	2	0.5
平均値 (fl)		95.4		91.3		93.1		94.3		95.8		96.3		96.8		96.5	
標準偏差		5.2		3.8		4.8		4.5		4.8		5.3		5.2		5.4	
女性	総 数	2,534	100.0	139	100.0	336	100.0	375	100.0	446	100.0	579	100.0	659	100.0	538	100.0
	70fl未満	20	0.8	2	1.4	3	0.9	11	2.9	2	0.4	2	0.3	0	0.0	2	0.4
	70-74	22	0.9	2	1.4	8	2.4	9	2.4	2	0.4	1	0.2	0	0.0	0	0.0
	75-79	42	1.7	7	5.0	13	3.9	16	4.3	5	1.1	0	0.0	1	0.2	0	0.0
	80-84	73	2.9	7	5.0	26	7.7	19	5.1	11	2.5	3	0.5	7	1.1	5	0.9
	85-89	355	14.0	45	32.4	77	22.9	61	16.3	58	13.0	58	10.0	56	8.5	45	8.4
	90-94	1,068	42.1	54	38.8	137	40.8	160	42.7	218	48.9	247	42.7	252	38.2	222	41.3
	95-99	713	28.1	17	12.2	55	16.4	78	20.8	113	25.3	204	35.2	246	37.3	198	36.8
	100-104	204	8.1	4	2.9	13	3.9	15	4.0	33	7.4	57	9.8	82	12.4	54	10.0
	105-109	28	1.1	1	0.7	3	0.9	3	0.8	4	0.9	6	1.0	11	1.7	9	1.7
	110-114	7	0.3	0	0.0	1	0.3	2	0.5	0	0.0	1	0.2	3	0.5	2	0.4
	115-119	1	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0
	120fl以上	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2
平均値 (fl)		93.3		90.1		90.6		91.0		93.2		94.7		95.5		95.1	
標準偏差		6.1		6.3		6.7		7.8		5.1		94.7		4.8		5.1	

第26表 平均赤血球血色素量（MCH）の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,725	100.0	90	100.0	196	100.0	239	100.0	295	100.0	445	100.0	460	100.0	423	100.0	268	100.0
	20pg未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	20-24	4	0.2	0	0.0	1	0.5	0	0.0	0	0.0	1	0.2	2	0.4	1	0.2	1	0.4
	25-26	13	0.8	1	1.1	0	0.0	2	0.8	2	0.7	4	0.9	4	0.9	4	0.9	2	0.7
	27-28	111	6.4	14	15.6	24	12.2	13	5.4	11	3.7	24	5.4	25	5.4	22	5.2	17	6.3
	29-30	691	40.1	58	64.4	101	51.5	112	46.9	118	40.0	172	38.7	130	28.3	144	34.0	75	28.0
	31-32	688	39.9	17	18.9	61	31.1	96	40.2	123	41.7	178	40.0	213	46.3	183	43.3	122	45.5
	33-34	185	10.7	0	0.0	7	3.6	15	6.3	34	11.5	54	12.1	75	16.3	57	13.5	47	17.5
	35-39	31	1.8	0	0.0	2	1.0	1	0.4	7	2.4	11	2.5	10	2.2	11	2.6	4	1.5
	40pg以上	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	1	0.2	0	0.0
	平均値(pg) 標準偏差	31.2 1.7		30.1 1.1		30.5 1.5		31.0 1.5		31.4 1.6		31.3 1.8		31.5 1.9		31.5 1.9		31.5 1.7	
女性	総 数	2,534	100.0	139	100.0	336	100.0	375	100.0	446	100.0	579	100.0	659	100.0	538	100.0	408	100.0
	20pg未満	26	1.0	2	1.4	6	1.8	14	3.7	2	0.4	2	0.3	0	0.0	2	0.4	0	0.0
	20-24	80	3.2	9	6.5	26	7.7	30	8.0	11	2.5	1	0.2	3	0.5	1	0.2	2	0.5
	25-26	90	3.6	8	5.8	32	9.5	22	5.9	11	2.5	4	0.7	13	2.0	6	1.1	9	2.2
	27-28	414	16.3	39	28.1	81	24.1	67	17.9	73	16.4	77	13.3	77	11.7	63	11.7	54	13.2
	29-30	1,341	52.9	71	51.1	147	43.8	169	45.1	253	56.7	339	58.5	362	54.9	306	56.9	213	52.2
	31-32	533	21.0	10	7.2	40	11.9	66	17.6	90	20.2	146	25.2	181	27.5	148	27.5	116	28.4
	33-34	41	1.6	0	0.0	4	1.2	4	1.1	5	1.1	9	1.6	19	2.9	10	1.9	12	2.9
	35-39	8	0.3	0	0.0	0	0.0	3	0.8	1	0.2	1	0.2	3	0.5	1	0.2	2	0.5
	40pg以上	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	0	0.0
	平均値(pg) 標準偏差	29.7 2.3		28.7 2.5		28.7 2.6		28.8 3.3		29.8 1.9		30.2 1.5		30.4 1.6		30.3 1.6		30.4 1.6	

第27表 平均赤血球血色素濃度（MCHC）の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,725	100.0	90	100.0	196	100.0	239	100.0	295	100.0	445	100.0	460	100.0	423	100.0	268	100.0
	22%未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	22-23	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	24-25	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	26-27	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0
	28-29	12	0.7	0	0.0	2	1.0	2	0.8	2	0.7	3	0.7	3	0.7	4	0.9	0	0.0
	30-31	370	21.4	16	17.8	34	17.3	42	17.6	55	18.6	103	23.1	120	26.1	90	21.3	80	29.9
	32-33	1,194	69.2	57	63.3	142	72.4	168	70.3	215	72.9	310	69.7	302	65.7	298	70.4	169	63.1
	34%以上	148	8.6	17	18.9	18	9.2	27	11.3	23	7.8	28	6.3	35	7.6	31	7.3	19	7.1
	平均値(%) 標準偏差	32.7 1.0		33.0 1.1		32.8 1.0		32.9 1.0		32.8 1.0		32.6 1.0		32.6 1.0		32.6 1.0		32.5 1.0	
女性	総 数	2,534	100.0	139	100.0	336	100.0	375	100.0	446	100.0	579	100.0	659	100.0	538	100.0	408	100.0
	22%未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	22-23	2	0.1	1	0.7	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	24-25	4	0.2	0	0.0	0	0.0	3	0.8	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0
	26-27	22	0.9	0	0.0	7	2.1	11	2.9	3	0.7	10	2.2	0	0.0	1	0.2	0	0.0
	28-29	152	6.0	9	6.5	27	8.0	38	10.1	24	5.4	23	4.0	31	4.7	29	5.4	17	4.2
	30-31	1,105	43.6	57	41.0	147	43.8	145	38.7	182	40.8	251	43.4	323	49.0	252	46.8	201	49.3
	32-33	1,215	47.9	68	48.9	149	44.3	170	45.3	232	52.0	298	51.5	298	45.2	246	45.7	189	46.3
	34%以上	34	1.3	4	2.9	6	1.8	7	1.9	4	0.9	6	1.0	7	1.1	10	1.9	1	0.2
	平均値(%) 標準偏差	31.8 1.2		31.9 1.4		31.7 1.4		31.6 1.6		31.9 1.2		32.0 1.0		31.8 1.0		31.9 1.1		31.8 1.0	

第28表 血清フェリチン値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,717	100.0	90	100.0	198	100.0	238	100.0	294	100.0	440	100.0	457	100.0	418	100.0	267	100.0
	1.1ng/ml未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	1.1-5未満	6	0.3	0	0.0	1	0.5	1	0.4	1	0.3	1	0.2	2	0.4	1	0.2	1	0.4
	5-9	22	1.3	0	0.0	1	0.5	1	0.4	3	1.0	8	1.8	9	2.0	10	2.4	5	1.9
	10-14	27	1.6	0	0.0	1	0.5	3	1.3	4	1.4	9	2.0	10	2.2	6	1.4	7	2.6
	15-19	23	1.3	1	1.1	0	0.0	2	0.8	5	1.7	6	1.4	9	2.0	4	1.0	7	2.6
	20-24	42	2.4	0	0.0	2	1.0	3	1.3	5	1.7	13	3.0	19	4.2	11	2.6	12	4.5
	25-29	33	1.9	3	3.3	5	2.5	4	1.7	4	1.4	3	0.7	14	3.1	4	1.0	12	4.5
	30-34	30	1.7	1	1.1	3	1.5	4	1.7	6	2.0	7	1.6	9	2.0	7	1.7	6	2.2
	35-39	37	2.2	0	0.0	2	1.0	6	2.5	6	2.0	9	2.0	14	3.1	11	2.6	8	3.0
	40-44	38	2.2	6	6.7	5	2.5	1	0.4	4	1.4	8	1.8	14	3.1	12	2.9	7	2.6
	45-49	42	2.4	3	3.3	1	0.5	0	0.0	4	1.4	12	2.7	22	4.8	15	3.6	11	4.1
	50-59	81	4.7	3	3.3	7	3.5	12	5.0	13	4.4	23	5.2	23	5.0	22	5.3	14	5.2
	60-69	97	5.6	6	6.7	5	2.5	12	5.0	17	5.8	27	6.1	30	6.6	24	5.7	18	6.7
	70-79	91	5.3	5	5.6	9	4.5	15	6.3	13	4.4	28	6.4	21	4.6	20	4.8	15	5.6
	80-89	102	5.9	4	4.4	12	6.1	19	8.0	16	5.4	30	6.8	21	4.6	19	4.5	17	6.4
	90-99	103	6.0	4	4.4	16	8.1	13	5.5	22	7.5	21	4.8	27	5.9	22	5.3	15	5.6
	100-149	358	20.9	36	40.0	57	28.8	51	21.4	50	17.0	82	18.6	82	17.9	76	18.2	44	16.5
	150-199	242	14.1	6	6.7	31	15.7	42	17.6	51	17.3	54	12.3	58	12.7	65	15.6	26	9.7
	200ng/ml以上	343	20.0	12	13.3	40	20.2	49	20.6	70	23.8	99	22.5	73	16.0	89	21.3	42	15.7
	(再掲)27未満	132	7.7	3	3.3	6	3.0	10	4.2	18	6.1	40	9.1	55	12.0	35	8.4	37	13.9
	(再掲)320以上	114	6.6	0	0.0	10	5.1	14	5.9	22	7.5	45	10.2	23	5.0	36	8.6	12	4.5
	平均値 (%)	139.6		117.3		144.3		148.9		152.9		145.0		123.3		142.7		116.5	
	標準偏差	123.4		63.4		115.3		117.4		123.8		123.4		136.3		123.4		140.6	
女性	総 数	2,488	100.0	137	100.0	321	100.0	364	100.0	443	100.0	571	100.0	652	100.0	533	100.0	402	100.0
	1.1ng/ml未満	1	0.0	1	0.7	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	1.1-5未満	171	6.9	19	13.9	55	17.1	67	18.4	24	5.4	4	0.7	2	0.3	2	0.4	2	0.5
	5-9	206	8.3	19	13.9	48	15.0	60	16.5	23	5.2	15	2.6	41	6.3	18	3.4	30	7.5
	10-14	190	7.6	23	16.8	52	16.2	49	13.5	16	3.6	13	2.3	37	5.7	20	3.8	25	6.2
	15-19	136	5.5	18	13.1	25	7.8	33	9.1	12	2.7	14	2.5	34	5.2	20	3.8	23	5.7
	20-24	126	5.1	5	3.6	40	12.5	24	6.6	21	4.7	12	2.1	24	3.7	16	3.0	17	4.2
	25-29	143	5.7	7	5.1	28	8.7	26	7.1	18	4.1	25	4.4	39	6.0	26	4.9	26	6.5
	30-34	126	5.1	11	8.0	16	5.0	22	6.0	22	5.0	23	4.0	32	4.9	23	4.3	21	5.2
	35-39	128	5.1	9	6.6	13	4.0	15	4.1	23	5.2	28	4.9	40	6.1	31	5.8	23	5.7
	40-44	106	4.3	3	2.2	10	3.1	11	3.0	26	5.9	27	4.7	29	4.4	27	5.1	19	4.7
	45-49	104	4.2	5	3.6	8	2.5	6	1.6	33	7.4	31	5.4	21	3.2	26	4.9	13	3.2
	50-59	199	8.0	6	4.4	8	2.5	12	3.3	48	10.8	71	12.4	54	8.3	47	8.8	36	9.0
	60-69	168	6.8	4	2.9	10	3.1	13	3.6	34	7.7	42	7.4	65	10.0	45	8.4	40	10.0
	70-79	134	5.4	2	1.5	2	0.6	6	1.6	30	6.8	38	6.7	56	8.6	37	6.9	32	8.0
	80-89	113	4.5	1	0.7	1	0.3	2	0.5	23	5.2	47	8.2	39	6.0	40	7.5	21	5.2
	90-99	86	3.5	0	0.0	1	0.3	7	1.9	15	3.4	37	6.5	26	4.0	28	5.3	16	4.0
	100-149	218	8.8	4	2.9	1	0.3	8	2.2	57	12.9	89	15.6	59	9.0	77	14.4	30	7.5
	150-199	80	3.2	0	0.0	1	0.3	2	0.5	9	2.0	35	6.1	33	5.1	28	5.3	18	4.5
	200ng/ml以上	53	2.1	0	0.0	2	0.6	1	0.3	9	2.0	20	3.5	21	3.2	22	4.1	10	2.5
	(再掲)3.4未満	89	3.6	9	6.5	30	9.0	39	10.7	9	2.0	2	0.4	0	0.0	1	0.2	0	0.0
	(再掲)89以上	445	17.8	4	2.9	5	1.5	18	4.9	92	20.8	184	32.2	142	21.8	157	29.5	77	19.2
	平均値 (%)	55.0		25.0		22.5		25.7		64.8		78.5		66.5		75.3		62.3	
	標準偏差	60.5		23.7		26.3		28.7		84.8		58.0		58.1		59.7		58.5	

注) 妊婦，貧血治療のための薬服用者除外。

第29表 血清総たんぱく質値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75-79歳		(再掲) 80歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,736	100.0	90	100.0	198	100.0	239	100.0	297	100.0	445	100.0	467	100.0	425	100.0	153	100.0	120	100.0
	6.0g/dl未満	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0
	6.0-6.4	21	1.2	0	0.0	1	0.5	0	0.0	3	1.0	6	1.3	11	2.4	6	1.4	4	2.6	3	2.5
	6.5-6.9	278	16.0	4	4.4	20	10.1	33	13.8	43	14.5	69	15.5	109	23.3	78	18.4	31	20.3	33	27.5
	7.0-7.4	803	46.3	43	47.8	92	46.5	125	52.3	139	46.8	216	48.5	188	40.3	207	48.7	56	36.6	46	38.3
	7.5-7.9	514	29.6	36	40.0	66	33.3	68	28.5	89	30.0	127	28.5	128	27.4	106	24.9	51	33.3	32	26.7
	8.0-8.4	107	6.2	7	7.8	17	8.6	10	4.2	22	7.4	24	5.4	27	5.8	26	6.1	10	6.5	5	4.2
	8.5g/dl以上	12	0.7	0	0.0	2	1.0	3	1.3	1	0.3	2	0.4	4	0.9	2	0.5	1	0.7	1	0.8
	(再掲)6.5未満	22	1.3	0	0.0	1	0.5	0	0.0	3	1.0	7	1.6	11	2.4	6	1.4	4	2.6	3	2.5
	平均値 (g/dl) 標準偏差	7.3 0.4		7.5 0.3		7.4 0.4		7.3 0.4		7.4 0.4		7.3 0.4		7.3 0.5		7.3 0.4		7.3 0.5		7.2 0.5	
女性	総 数	2,553	100.0	140	100.0	337	100.0	376	100.0	450	100.0	583	100.0	667	100.0	544	100.0	218	100.0	194	100.0
	6.0g/dl未満	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.1	1	0.2	0	0.0	1	0.5
	6.0-6.4	29	1.1	0	0.0	6	1.8	4	1.1	3	0.7	2	0.3	14	2.1	5	0.9	4	1.8	6	3.1
	6.5-6.9	338	13.2	13	9.3	34	10.1	60	16.0	47	10.4	66	11.3	118	17.7	79	14.5	39	17.9	32	16.5
	7.0-7.4	1,170	45.8	53	37.9	159	47.2	181	48.1	202	44.9	270	46.3	305	45.7	248	45.6	102	46.8	86	44.3
	7.5-7.9	825	32.3	58	41.4	118	35.0	107	28.5	163	36.2	204	35.0	175	26.2	168	30.9	54	24.8	55	28.4
	8.0-8.4	168	6.6	14	10.0	17	5.0	23	6.1	28	6.2	38	6.5	48	7.2	39	7.2	18	8.3	12	6.2
	8.5g/dl以上	21	0.8	2	1.4	3	0.9	1	0.3	7	1.6	2	0.3	6	0.9	4	0.7	1	0.5	2	1.0
	(再掲)6.5未満	31	1.2	0	0.0	6	1.8	4	1.1	3	0.7	3	0.5	15	2.2	6	1.1	4	1.8	7	3.6
	平均値 (g/dl) 標準偏差	7.4 0.4		7.5 0.5		7.4 0.4		7.3 0.4		7.4 0.4		7.4 0.4		7.3 0.4		7.4 0.4		7.3 0.4		7.3 0.5	

第30表 血清アルブミン値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75-79歳		(再掲) 80歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男性	総 数	1,736	100.0	90	100.0	198	100.0	239	100.0	297	100.0	445	100.0	467	100.0	425	100.0	153	100.0	120	100.0
	3.0g/dl未満	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0
	3.0-3.2未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	3.2-3.4	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.2	0	0.0	0	0.0
	3.4-3.6	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	0	0.0	1	0.7	1	0.8
	3.6-3.8	20	1.2	0	0.0	0	0.0	1	0.4	4	1.3	2	0.4	13	2.8	2	0.5	3	2	9	7.5
	3.8-4.0	54	3.1	0	0.0	0	0.0	1	0.4	2	0.7	10	2.2	41	8.8	15	3.5	16	10.5	14	11.7
	4.0-4.2	189	10.9	0	0.0	0	0.0	10	4.2	16	5.4	57	12.8	106	22.7	73	17.2	33	21.6	34	28.3
	4.2-4.4	364	21.0	3	3.3	13	6.6	25	10.5	61	20.5	124	27.9	138	29.6	140	32.9	36	23.5	35	29.2
	4.4-4.6	495	28.5	17	18.9	43	21.7	79	33.1	96	32.3	141	31.7	119	25.5	126	29.6	47	30.7	19	15.8
	4.6-4.8	369	21.3	29	32.2	76	38.4	74	31.0	73	24.6	77	17.3	40	8.6	53	12.5	15	9.8	6	5.0
	4.8-5.0	175	10.1	26	28.9	48	24.2	35	14.6	39	13.1	24	5.4	3	0.6	8	1.9	1	0.7	1	0.8
	5.0-5.2	55	3.2	14	15.6	15	7.6	12	5.0	5	1.7	4	0.9	5	1.1	6	1.4	1	0.7	1	0.8
	5.2g/dl以上	11	0.6	1	1.1	3	1.5	2	0.8	1	0.3	4	0.9	0	0.0	1	0.2	0	0.0	0	0.0
	(再掲)3.5以下	4	0.2	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	2	0.4	1	0.2	1	0.7	1	0.8
	平均値 (g/dl) 標準偏差	4.4 0.3		4.7 0.2		4.7 0.2		4.6 0.2		4.5 0.2		4.4 0.3		4.2 0.3		4.3 0.2		4.3 0.3		4.2 0.3	
女性	総 数	2,553	100.0	140	100.0	337	100.0	376	100.0	450	100.0	583	100.0	667	100.0	544	100.0	218	100.0	194	100.0
	3.0g/dl未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	3.0-3.2未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	3.2-3.4	3	0.1	0	0.0	1	0.3	0	0.0	0	0.0	1	0.2	1	0.1	0	0.0	0	0.0	1	0.5
	3.4-3.6	6	0.2	0	0.0	2	0.6	0	0.0	1	0.2	0	0.0	3	0.4	0	0.0	1	0.5	2	1.0
	3.6-3.8	29	1.1	1	0.7	5	1.5	2	0.5	1	0.2	6	1.0	14	2.1	7	1.3	2	0.9	8	4.1
	3.8-4.0	63	2.5	1	0.7	4	1.2	2	0.5	4	0.9	14	2.4	38	5.7	16	2.9	12	5.5	19	9.8
	4.0-4.2	292	11.4	1	0.7	24	7.1	43	11.4	30	6.7	63	10.8	131	19.6	85	15.6	33	15.1	46	23.7
	4.2-4.4	698	27.3	28	20	72	21.4	113	30.1	102	22.7	158	27.1	225	33.7	169	31.1	80	36.7	57	29.4
	4.4-4.6	833	32.6	42	30	126	37.4	141	37.5	178	39.6	180	30.9	166	24.9	160	29.4	59	27.1	39	20.1
	4.6-4.8	469	18.4	43	30.7	76	22.6	61	16.2	96	21.3	120	20.6	73	10.9	81	14.9	25	11.5	19	9.8
	4.8-5.0	134	5.2	19	13.6	24	7.1	12	3.2	30	6.7	34	5.8	15	2.2	23	4.2	6	2.8	3	1.5
	5.0-5.2	23	0.9	5	3.6	3	0.9	2	0.5	6	1.3	6	1.0	1	0.1	3	0.6	0	0.0	0	0.0
	5.2g/dl以上	3	0.1	0	0.0	0	0.0	0	0.0	2	0.4	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0
	(再掲)3.5以下	9	0.4	0	0.0	3	0.9	0	0.0	1	0.2	1	0.2	4	0.6	0	0.0	1	0.5	3	1.5
	平均値 (g/dl) 標準偏差	4.4 0.3		4.5 0.2		4.4 0.3		4.4 0.2		4.4 0.2		4.4 0.3		4.3 0.3		4.3 0.2		4.3 0.2		4.2 0.3	

第31表 ヘモグロビンA1cの分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	総 数	4,028	100.0	229	100.0	533	100.0	600	100.0	706	100.0	952	100.0	1,008	100.0	885	100.0	603	100.0
	3.9%以下	3	0.1	0	0.0	0	0.0	0	0.0	0	0.0	2	0.2	1	0.1	1	0.1	1	0.2
	4.0-4.1	4	0.1	0	0.0	1	0.2	2	0.3	0	0.0	1	0.1	0	0.0	1	0.1	0	0.0
	4.2-4.3	21	0.5	6	2.6	7	1.3	5	0.8	1	0.1	0	0.0	2	0.2	0	0.0	2	0.3
	4.4-4.5	59	1.5	6	2.6	18	3.4	13	2.2	9	1.3	3	0.3	10	1.0	5	0.6	5	0.8
	4.6-4.7	195	4.8	39	17.0	50	9.4	41	6.8	26	3.7	22	2.3	17	1.7	16	1.8	11	1.8
	4.8-4.9	481	11.9	63	27.5	115	21.6	89	14.8	68	9.6	70	7.4	76	7.5	63	7.1	45	7.5
	5.0-5.1	758	18.8	57	24.9	133	25.0	142	23.7	132	18.7	143	15.0	151	15.0	138	15.6	85	14.1
	5.2	464	11.5	25	10.9	57	10.7	80	13.3	93	13.2	113	11.9	96	9.5	99	11.2	51	8.5
	5.3	401	10.0	15	6.6	47	8.8	66	11.0	79	11.2	101	10.6	93	9.2	82	9.3	65	10.8
	5.4	393	9.8	9	3.9	41	7.7	51	8.5	77	10.9	98	10.3	117	11.6	95	10.7	68	11.3
	5.5	298	7.4	5	2.2	19	3.6	28	4.7	63	8.9	83	8.7	100	9.9	82	9.3	64	10.6
	5.6	253	6.3	2	0.9	22	4.1	28	4.7	47	6.7	70	7.4	84	8.3	65	7.3	51	8.5
	5.7	162	4.0	0	0.0	4	0.8	17	2.8	22	3.1	56	5.9	63	6.3	53	6.0	37	6.1
	5.8	130	3.2	1	0.4	6	1.1	10	1.7	23	3.3	40	4.2	50	5.0	47	5.3	26	4.3
	5.9	80	2.0	0	0.0	2	0.4	5	0.8	17	2.4	25	2.6	31	3.1	19	2.1	21	3.5
	6.0	61	1.5	0	0.0	2	0.4	4	0.7	7	1.0	22	2.3	26	2.6	19	2.1	18	3.0
	6.1	46	1.1	1	0.4	0	0.0	3	0.5	9	1.3	18	1.9	15	1.5	22	2.5	7	1.2
	6.2-6.3	65	1.6	0	0.0	2	0.4	6	1.0	5	0.7	27	2.8	25	2.5	26	2.9	17	2.8
	6.4-6.5	46	1.1	0	0.0	1	0.2	2	0.3	6	0.8	16	1.7	21	2.1	16	1.8	12	2.0
	6.6-6.7	22	0.5	0	0.0	1	0.2	0	0.0	1	0.1	10	1.1	10	1.0	8	0.9	7	1.2
	6.8-6.9	16	0.4	0	0.0	0	0.0	0	0.0	5	0.7	6	0.6	5	0.5	6	0.7	3	0.5
	7.0-7.1	11	0.3	0	0.0	1	0.2	1	0.2	1	0.1	6	0.6	2	0.2	5	0.6	0	0.0
	7.2-7.3	8	0.2	0	0.0	0	0.0	0	0.0	1	0.1	4	0.4	3	0.3	2	0.2	3	0.5
	7.4-7.5	8	0.2	0	0.0	1	0.2	1	0.2	4	0.6	0	0.0	2	0.2	1	0.1	1	0.2
	7.6-7.7	2	0.0	0	0.0	0	0.0	0	0.0	0	0.0	2	0.2	0	0.0	1	0.1	0	0.0
	7.8-7.9	4	0.1	0	0.0	0	0.0	1	0.2	0	0.0	3	0.3	0	0.0	1	0.1	0	0.0
	8.0-8.1	2	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	1	0.1	1	0.1	0	0.0
	8.2-8.3	4	0.1	0	0.0	0	0.0	0	0.0	1	0.1	1	0.1	2	0.2	1	0.1	1	0.2
	8.4-8.5	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1	0	0.0	1	0.1	0	0.0
	8.6-8.7	3	0.1	0	0.0	0	0.0	1	0.2	1	0.1	1	0.1	0	0.0	0	0.0	0	0.0
	8.8-8.9	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1	0	0.0	1	0.1	0	0.0
	9.0%以上	26	0.6	0	0.0	3	0.6	3	0.5	8	1.1	7	0.7	5	0.5	8	0.9	2	0.3
	(再掲)6.1以上	265	6.6	1	0.4	9	1.7	19	3.2	42	5.9	103	10.8	91	9.0	100	11.3	53	8.8
	平均値 (%) 標準偏差	5.3 0.7		5.0 0.3		5.1 0.7		5.2 0.6		5.4 0.7		5.5 0.7		5.5 0.6		5.5 0.7		5.5 0.5	
男 性	総 数	1,595	100.0	90	100.0	197	100.0	230	100.0	275	100.0	403	100.0	400	100.0	379	100.0	231	100.0
	3.9%以下	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	1	0.4
	4.0-4.1	2	0.1	0	0.0	1	0.5	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	4.2-4.3	11	0.7	4	4.4	2	1.0	3	1.3	1	0.4	0	0.0	1	0.3	0	0.0	1	0.4
	4.4-4.5	35	2.2	5	5.6	9	4.6	8	3.5	5	1.8	1	0.2	7	1.8	3	0.8	4	1.7
	4.6-4.7	81	5.1	14	15.6	20	10.2	15	6.5	7	2.5	14	3.5	11	2.8	13	3.4	7	3.0
	4.8-4.9	197	12.4	30	33.3	38	19.3	33	14.3	28	10.2	33	8.2	35	8.8	33	8.7	18	7.8
	5.0-5.1	285	17.9	18	20.0	51	25.9	48	20.9	48	17.5	70	17.4	50	12.5	62	16.4	24	10.4
	5.2	171	10.7	6	6.7	21	10.7	32	13.9	35	12.7	39	9.7	38	9.5	37	9.8	23	10.0
	5.3	153	9.6	6	6.7	21	10.7	28	12.2	24	8.7	36	8.9	38	9.5	32	8.4	25	10.8
	5.4	149	9.3	2	2.2	14	7.1	20	8.7	33	12.0	37	9.2	43	10.8	37	9.8	25	10.8
	5.5	108	6.8	2	2.2	8	4.1	12	5.2	27	9.8	27	6.7	32	8.0	28	7.4	19	8.2
	5.6	95	6.0	1	1.1	5	2.5	7	3.0	17	6.2	43	10.7	22	5.5	30	7.9	12	5.2
	5.7	63	3.9	0	0.0	1	0.5	9	3.9	9	3.3	20	5.0	24	6.0	18	4.7	14	6.1
	5.8	46	2.9	1	1.1	2	1.0	3	1.3	8	2.9	14	3.5	18	4.5	17	4.5	7	3.0
	5.9	36	2.3	0	0.0	0	0.0	4	1.7	8	2.9	9	2.2	15	3.8	7	1.8	11	4.8
	6.0	28	1.8	0	0.0	1	0.5	0	0.0	2	0.7	9	2.2	16	4.0	11	2.9	9	3.9
	6.1	19	1.2	1	1.1	0	0.0	1	0.4	3	1.1	4	1.0	10	2.5	8	2.1	5	2.2
	6.2-6.3	31	1.9	0	0.0	0	0.0	1	0.4	2	0.7	14	3.5	14	3.5	14	3.7	10	4.3
	6.4-6.5	26	1.6	0	0.0	1	0.5	1	0.4	4	1.5	10	2.5	10	2.5	10	2.6	5	2.2
	6.6-6.7	16	1.0	0	0.0	1	0.5	0	0.0	0	0.0	7	1.7	8	2.0	6	1.6	7	3.0
	6.8-6.9	8	0.5	0	0.0	0	0.0	0	0.0	5	1.8	2	0.5	1	0.3	3	0.8	0	0.0
	7.0-7.1	5	0.3	0	0.0	1	0.5	0	0.0	1	0.4	3	0.7	0	0.0	2	0.5	0	0.0
	7.2-7.3	6	0.4	0	0.0	0	0.0	0	0.0	1	0.4	3	0.7	2	0.5	2	0.5	2	0.9
	7.4-7.5	4	0.3	0	0.0	0	0.0	1	0.4	2	0.7	0	0.0	1	0.3	1	0.3	0	0.0
	7.6-7.7	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.3	0	0.0
	7.8-7.9	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0
	8.0-8.1	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	1	0.3	0	0.0
	8.2-8.3	2	0.1	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.3	0	0.0	1	0.4
	8.4-8.5	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.3	0	0.0
	8.6-8.7	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0
	8.8-8.9	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.3	0	0.0
	9.0%以上	11	0.7	0	0.0	0	0.0	3	1.3	4	1.5	3	0.7	1	0.3	1	0.3	1	0.4
	(再掲)6.1以上	134	8.4	1	1.1	3	1.5	7	3.0	23	8.4	51	12.7	49	12.3	51	13.5	31	13.4
	平均値 (%) 標準偏差	5.4 0.7		4.9 0.3		5.1 0.4		5.2 0.8		5.4 0.9		5.5 0.8		5.5 0.6		5.5 0.7		5.5 0.6	

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
女 性	総 数	2,433	100.0	139	100.0	336	100.0	370	100.0	431	100.0	549	100.0	608	100.0	506	100.0	372	100.0
	3.9%以下	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	0	0.0	1	0.2	0	0.0
	4.0-4.1	2	0.1	0	0.0	0	0.0	1	0.3	0	0.0	1	0.2	0	0.0	1	0.2	0	0.0
	4.2-4.3	10	0.4	2	1.4	5	1.5	2	0.5	0	0.0	0	0.0	1	0.2	0	0.0	1	0.3
	4.4-4.5	24	1.0	1	0.7	9	2.7	5	1.4	4	0.9	2	0.4	3	0.5	2	0.4	1	0.3
	4.6-4.7	114	4.7	25	18.0	30	8.9	26	7.0	19	4.4	8	1.5	6	1.0	3	0.6	4	1.1
	4.8-4.9	284	11.7	33	23.7	77	22.9	56	15.1	40	9.3	37	6.7	41	6.7	30	5.9	27	7.3
	5.0-5.1	473	19.4	39	28.1	82	24.4	94	25.4	84	19.5	73	13.3	101	16.6	76	15.0	61	16.4
	5.2	293	12.0	19	13.7	36	10.7	48	13.0	58	13.5	74	13.5	58	9.5	62	12.3	28	7.5
	5.3	248	10.2	9	6.5	26	7.7	38	10.3	55	12.8	65	11.8	55	9.0	50	9.9	40	10.8
	5.4	244	10.0	7	5.0	27	8.0	31	8.4	44	10.2	61	11.1	74	12.2	58	11.5	43	11.6
	5.5	190	7.8	3	2.2	11	3.3	16	4.3	36	8.4	56	10.2	68	11.2	54	10.7	45	12.1
	5.6	158	6.5	1	0.7	17	5.1	21	5.7	30	7.0	27	4.9	62	10.2	35	6.9	39	10.5
	5.7	99	4.1	0	0.0	3	0.9	8	2.2	13	3.0	36	6.6	39	6.4	35	6.9	23	6.2
	5.8	84	3.5	0	0.0	4	1.2	7	1.9	15	3.5	26	4.7	32	5.3	30	5.9	19	5.1
	5.9	44	1.8	0	0.0	2	0.6	1	0.3	9	2.1	16	2.9	16	2.6	12	2.4	10	2.7
	6.0	33	1.4	0	0.0	1	0.3	4	1.1	5	1.2	13	2.4	10	1.6	8	1.6	9	2.4
	6.1	27	1.1	0	0.0	0	0.0	2	0.5	6	1.4	14	2.6	5	0.8	14	2.8	2	0.5
	6.2-6.3	34	1.4	0	0.0	2	0.6	5	1.4	3	0.7	13	2.4	11	1.8	12	2.4	7	1.9
	6.4-6.5	20	0.8	0	0.0	0	0.0	1	0.3	2	0.5	6	1.1	11	1.8	6	1.2	7	1.9
	6.6-6.7	6	0.2	0	0.0	0	0.0	0	0.0	1	0.2	3	0.5	2	0.3	2	0.4	0	0.0
	6.8-6.9	8	0.3	0	0.0	0	0.0	0	0.0	0	0.0	4	0.7	4	0.7	3	0.6	3	0.8
	7.0-7.1	6	0.2	0	0.0	0	0.0	1	0.3	0	0.0	3	0.5	2	0.3	3	0.6	0	0.0
	7.2-7.3	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	0	0.0	1	0.3
	7.4-7.5	4	0.2	0	0.0	1	0.3	0	0.0	2	0.5	0	0.0	1	0.2	0	0.0	1	0.3
	7.6-7.7	1	0.0	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	0	0.0
	7.8-7.9	3	0.1	0	0.0	0	0.0	1	0.3	0	0.0	2	0.4	0	0.0	1	0.2	0	0.0
	8.0-8.1	1	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.2-8.3	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	1	0.2	0	0.0
	8.4-8.5	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	8.6-8.7	2	0.1	0	0.0	0	0.0	1	0.3	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0
	8.8-8.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	9.0%以上	15	0.6	0	0.0	3	0.9	0	0.0	4	0.9	4	0.7	4	0.7	7	1.4	1	0.3
	(再掲)6.1以上	131	5.4	0	0.0	6	1.8	12	3.2	19	4.4	52	9.5	42	6.9	49	9.7	22	5.9
平均値 (%)		5.3		5.0		5.2		5.2		5.3		5.5		5.5		5.5		5.4	
標準偏差		0.6		0.3		0.8		0.4		0.6		0.6		0.6		0.7		0.5	

注) インスリン注射または血糖を下げる薬服用者除外

第 32 表 血糖値の分布（食後 3 時間以上、性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総 数	1,238	100.0	69	100.0	152	100.0	185	100.0	214	100.0	315	100.0	303	100.0	282	100.0	176	100.0
	40mg/dl未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	40-59	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	60-79	69	5.6	4	5.8	12	7.9	9	4.9	13	6.1	18	5.7	13	4.3	14	5.0	8	4.5
	80-99	738	59.6	56	81.2	115	75.7	134	72.4	138	64.5	166	52.7	129	42.6	140	49.6	70	39.8
	100-119	309	25.0	6	8.7	22	14.5	36	19.5	45	21.0	88	27.9	112	37.0	93	33.0	64	36.4
	120-140	76	6.1	2	2.9	2	1.3	3	1.6	10	4.7	27	8.6	32	10.6	25	8.9	21	11.9
	140-160	22	1.8	1	1.4	1	0.7	1	0.5	3	1.4	5	1.6	11	3.6	2	0.7	10	5.7
	160-180	5	0.4	0	0.0	0	0.0	0	0.0	1	0.5	3	1.0	1	0.3	2	0.7	1	0.6
	180-200	8	0.6	0	0.0	0	0.0	0	0.0	1	0.5	3	1.0	4	1.3	3	1.1	2	1.1
	200-220	1	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0
	220-240	2	0.2	0	0.0	0	0.0	1	0.5	0	0.0	1	0.3	0	0.0	1	0.4	0	0.0
	240-260	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	260mg/dl以上	8	0.6	0	0.0	0	0.0	1	0.5	3	1.4	3	1.0	1	0.3	2	0.7	0	0.0
	(再掲)110以上	206	16.6	4	5.8	6	3.9	15	8.1	28	13.1	69	21.9	84	27.7	64	22.7	54	30.7
	平均値 (mg/dl)	100.0		92.5		91.2		95.6		100.6		103.8		104.4		102.9		105.4	
	標準偏差	26.9		12.9		10.3		20.4		36.2		33.0		21.8		25.4		20.0	
女 性	総 数	1,875	100.0	107	100.0	255	100.0	280	100.0	329	100.0	420	100.0	484	100.0	389	100.0	301	100.0
	40mg/dl未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	40-59	1	0.1	1	0.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	60-79	96	5.1	10	9.3	29	11.4	19	6.8	14	4.3	13	3.1	11	2.3	8	2.1	9	3.0
	80-99	1,105	58.9	81	75.7	189	74.1	207	73.9	216	65.7	219	52.1	193	39.9	175	45.0	118	39.2
	100-119	529	28.2	15	14.0	29	11.4	48	17.1	86	26.1	149	35.5	202	41.7	151	38.8	128	42.5
	120-140	93	5.0	0	0.0	5	2.0	4	1.4	6	1.8	28	6.7	50	10.3	37	9.5	29	9.6
	140-160	25	1.3	0	0.0	0	0.0	0	0.0	2	0.6	6	1.4	17	3.5	9	2.3	11	3.7
	160-180	6	0.3	0	0.0	0	0.0	0	0.0	1	0.3	1	0.2	4	0.8	3	0.8	2	0.7
	180-200	5	0.3	0	0.0	0	0.0	0	0.0	1	0.3	2	0.5	2	0.4	2	0.5	1	0.3
	200-220	3	0.2	0	0.0	0	0.0	0	0.0	1	0.3	1	0.2	1	0.2	1	0.3	1	0.3
	220-240	1	0.1	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	240-260	3	0.2	0	0.0	0	0.0	1	0.4	1	0.3	0	0.0	1	0.2	0	0.0	1	0.3
	260mg/dl以上	8	0.4	0	0.0	2	0.8	1	0.4	1	0.3	1	0.2	3	0.6	3	0.8	1	0.3
	(再掲)110以上	296	15.8	3	2.8	14	5.5	16	5.7	30	9.1	73	17.4	160	33.1	93	23.9	104	34.6
	平均値 (mg/dl)	99.1		89.6		93.9		93.0		97.2		100.9		107.3		105.2		106.9	
	標準偏差	25.0		9.4		39.8		18.1		20.2		18.3		26.2		24.2		25.7	

注) インスリン注射または血糖を下げる薬服用者除外

第33表 血清総コレステロール値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,565	100.0	90	100.0	195	100.0	231	100.0	274	100.0	381	100.0	394	100.0	352	100.0	232	100.0
	120mg/dl未満	15	1.0	0	0.0	1	0.5	2	0.9	3	1.1	2	0.5	7	1.8	4	1.1	5	2.2
	120-139	29	1.9	5	5.6	1	0.5	2	0.9	3	1.1	7	1.8	11	2.8	11	3.1	5	2.2
	140-159	154	9.8	26	28.9	21	10.8	13	5.6	15	5.5	27	7.1	52	13.2	34	9.7	31	13.4
	160-179	267	17.1	25	27.8	27	13.8	39	16.9	28	10.2	70	18.4	78	19.8	72	20.5	51	22.0
	180-199	361	23.1	19	21.1	42	21.5	43	18.6	73	26.6	82	21.5	102	25.9	91	25.9	54	23.3
	200-219	349	22.3	10	11.1	54	27.7	51	22.1	73	26.6	76	19.9	85	21.6	67	19.0	52	22.4
	220-239	223	14.2	2	2.2	31	15.9	45	19.5	38	13.9	69	18.1	38	9.6	39	11.1	22	9.5
	240-259	102	6.5	1	1.1	13	6.7	21	9.1	25	9.1	27	7.1	15	3.8	20	5.7	8	3.4
	260-279	44	2.8	0	0.0	2	1.0	11	4.8	13	4.7	14	3.7	4	1.0	9	2.6	3	1.3
	280-299	15	1.0	1	1.1	1	0.5	3	1.3	3	1.1	5	1.3	2	0.5	3	0.9	1	0.4
	300mg/dl以上	6	0.4	1	1.1	2	1.0	1	0.4	0	0.0	2	0.5	0	0.0	2	0.6	0	0.0
	(再掲)220以上	390	24.9	5	5.6	49	25.1	81	35.1	79	28.8	117	30.7	59	15.0	73	20.7	34	14.7
	(再掲)240以上	167	10.7	3	3.3	18	9.2	36	15.6	41	15.0	48	12.6	21	5.3	34	9.7	12	5.2
	平均値(mg/dl) 標準偏差	198.3 33.8		177.4 32.6		200.3 32.2		206.2 33.9		205.8 32.7		202.1 34.0		188.6 31.2		193.4 33.6		188.5 31.7	
女 性	総数	2,186	100.0	139	100.0	337	100.0	369	100.0	416	100.0	443	100.0	482	100.0	402	100.0	293	100.0
	120mg/dl未満	3	0.1	0	0.0	1	0.3	0	0.0	0	0.0	1	0.2	1	0.2	0	0.0	1	0.3
	120-139	37	1.7	9	6.5	11	3.3	3	0.8	3	0.7	2	0.5	9	1.9	2	0.5	7	2.4
	140-159	162	7.4	36	25.9	52	15.4	32	8.7	4	1.0	8	1.8	30	6.2	14	3.5	23	7.8
	160-179	327	15.0	38	27.3	92	27.3	71	19.2	26	6.3	30	6.8	70	14.5	32	8.0	52	17.7
	180-199	455	20.8	35	25.2	84	24.9	82	22.2	75	18.0	68	15.3	111	23.0	76	18.9	69	23.5
	200-219	467	21.4	15	10.8	59	17.5	81	22.0	107	25.7	97	21.9	108	22.4	93	23.1	65	22.2
	220-239	356	16.3	4	2.9	20	5.9	54	14.6	91	21.9	107	24.2	80	16.6	93	23.1	36	12.3
	240-259	214	9.8	1	0.7	8	2.4	31	8.4	61	14.7	64	14.4	49	10.2	58	14.4	27	9.2
	260-279	103	4.7	1	0.7	6	1.8	12	3.3	31	7.5	40	9.0	13	2.7	20	5.0	8	2.7
	280-299	42	1.9	0	0.0	3	0.9	1	0.3	13	3.1	16	3.6	9	1.9	9	2.2	5	1.7
	300mg/dl以上	20	0.9	0	0.0	1	0.3	2	0.5	5	1.2	10	2.3	2	0.4	5	1.2	0	0.0
	(再掲)220以上	735	33.6	6	4.3	38	11.3	100	27.1	201	48.3	237	53.5	153	31.7	185	46.0	76	25.9
	(再掲)240以上	379	17.3	2	1.4	18	5.3	46	12.5	110	26.4	130	29.3	73	15.1	92	22.9	40	13.7
	平均値(mg/dl) 標準偏差	206.1 36.2		174.6 25.4		185.5 30.8		200.7 31.7		220.9 32.0		224.1 36.5		204.4 33.6		216.7 33.3		199.8 33.7	

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

第34表 血清中性脂肪（トリグリセライド）値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,565	100.0	90	100.0	195	100.0	231	100.0	274	100.0	381	100.0	394	100.0	352	100.0	232	100.0
	20mg/dl未満	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	20-49	76	4.9	10	11.1	16	8.2	7	3.0	8	2.9	15	3.9	20	5.1	15	4.3	14	6.0
	50-79	294	18.8	33	36.7	31	15.9	36	15.6	33	12.0	63	16.5	98	24.9	63	17.9	61	26.3
	80-109	298	19.0	16	17.8	24	12.3	51	22.1	63	23.0	65	17.1	79	20.1	63	17.9	48	20.7
	110-139	242	15.5	11	12.2	29	14.9	33	14.3	49	17.9	63	16.5	57	14.5	53	15.1	34	14.7
	140-169	183	11.7	7	7.8	26	13.3	27	11.7	30	10.9	42	11.0	51	12.9	41	11.6	34	14.7
	170-199	112	7.2	3	3.3	16	8.2	15	6.5	18	6.6	32	8.4	28	7.1	33	9.4	14	6.0
	200mg/dl以上	360	23.0	10	11.1	53	27.2	62	26.8	73	26.6	101	26.5	61	15.5	84	23.9	27	11.6
	(再掲)150以上	586	37.4	17	18.9	85	43.6	93	40.3	107	39.1	162	42.5	122	31.0	144	40.9	63	27.2
	平均値(mg/dl) 標準偏差	157.7 126.7		112.5 85.0		170.4 130.2		173.5 173.2		175.6 140.8		166.1 120.7		132.0 82.6		154.5 102.4		123.7 73.3	
女 性	総数	2,186	100.0	139	100.0	337	100.0	369	100.0	416	100.0	443	100.0	482	100.0	402	100.0	293	100.0
	20mg/dl未満	1	0.0	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	20-49	244	11.2	47	33.8	68	20.2	64	17.3	22	5.3	21	4.7	22	4.6	20	5.0	16	5.5
	50-79	580	26.5	56	40.3	125	37.1	99	26.8	100	24.0	87	19.6	113	23.4	80	19.9	70	23.9
	80-109	450	20.6	14	10.1	58	17.2	70	19.0	94	22.6	95	21.4	119	24.7	90	22.4	74	25.3
	110-139	328	15.0	11	7.9	33	9.8	50	13.6	72	17.3	74	16.7	88	18.3	74	18.4	55	18.8
	140-169	215	9.8	8	5.8	16	4.7	35	9.5	53	12.7	49	11.1	54	11.2	42	10.4	33	11.3
	170-199	114	5.2	0	0.0	11	3.3	16	4.3	22	5.3	31	7.0	34	7.1	30	7.5	18	6.1
	200mg/dl以上	254	11.6	3	2.2	25	7.4	35	9.5	53	12.7	86	19.4	52	10.8	66	16.4	27	9.2
	(再掲)150以上	495	22.6	6	4.3	46	13.6	71	19.2	102	24.5	149	33.6	121	25.1	122	30.3	69	23.5
	平均値(mg/dl) 標準偏差	117.8 84.3		71.3 38.5		95.3 76.1		110.1 81.2		126.6 87.4		141.5 100.9		123.2 72.1		133.3 80.6		118.9 66.3	

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

第35表 血清HDL-コレステロール値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,565	100.0	90	100.0	195	100.0	231	100.0	274	100.0	381	100.0	394	100.0	352	100.0	232	100.0
	30mg/dl未満	26	1.7	1	1.1	0	0.0	4	1.7	5	1.8	8	2.1	8	2.0	9	2.6	5	2.2
	30-39	174	11.1	4	4.4	23	11.8	19	8.2	42	15.3	39	10.2	47	11.9	46	13.1	27	11.6
	40-49	416	26.6	27	30.0	50	25.6	57	24.7	61	22.3	104	27.3	117	29.7	92	26.1	79	34.1
	50-59	402	25.7	24	26.7	56	28.7	73	31.6	52	19.0	105	27.6	92	23.4	97	27.6	48	20.7
	60-69	285	18.2	19	21.1	31	15.9	44	19.0	51	18.6	70	18.4	70	17.8	65	18.5	34	14.7
	70-79	152	9.7	9	10.0	25	12.8	20	8.7	33	12.0	31	8.1	34	8.6	25	7.1	21	9.1
	80-89	58	3.7	5	5.6	6	3.1	4	1.7	14	5.1	13	3.4	16	4.1	10	2.8	12	5.2
	90-99	32	2.0	1	1.1	2	1.0	6	2.6	6	2.2	10	2.6	7	1.8	7	2.0	4	1.7
	100mg/dl以上	20	1.3	0	0.0	2	1.0	4	1.7	10	3.6	1	0.3	3	0.8	1	0.3	2	0.9
	平均値(mg/dl) 標準偏差	55.5 15.4		56.7 13.1		55.4 14.5		55.8 15.2		57.6 18.7		54.7 14.4		54.3 15.0		53.6 14.3		54.1 15.4	
女 性	総数	2,186	100.0	139	100.0	337	100.0	369	100.0	416	100.0	443	100.0	482	100.0	402	100.0	293	100.0
	30mg/dl未満	5	0.2	0	0.0	1	0.3	0	0.0	1	0.2	1	0.2	2	0.4	0	0.0	2	0.7
	30-39	64	2.9	2	1.4	4	1.2	9	2.4	8	1.9	13	2.9	28	5.8	21	5.2	16	5.5
	40-49	281	12.9	7	5.0	29	8.6	34	9.2	42	10.1	72	16.3	97	20.1	79	19.7	56	19.1
	50-59	528	24.2	35	25.2	64	19.0	88	23.8	90	21.6	118	26.6	133	27.6	95	23.6	89	30.4
	60-69	538	24.6	42	30.2	104	30.9	85	23.0	98	23.6	102	23.0	107	22.2	99	24.6	65	22.2
	70-79	424	19.4	37	26.6	76	22.6	84	22.8	81	19.5	77	17.4	69	14.3	59	14.7	44	15.0
	80-89	232	10.6	15	10.8	43	12.8	40	10.8	66	15.9	38	8.6	30	6.2	35	8.7	12	4.1
	90-99	77	3.5	1	0.7	11	3.3	20	5.4	18	4.3	13	2.9	14	2.9	10	2.5	7	2.4
	100mg/dl以上	37	1.7	0	0.0	5	1.5	9	2.4	12	2.9	9	2.0	2	0.4	4	1.0	2	0.7
	平均値(mg/dl) 標準偏差	64.4 14.9		65.4 11.2		66.8 13.8		66.7 15.1		67.5 15.8		62.8 14.9		59.4 14.5		60.8 14.6		59.0 14.0	

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

第 36 表 血清 LDL- コレステロール値（Friedewald の式）の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,496	100.0	88	100.0	181	100.0	222	100.0	256	100.0	362	100.0	387	100.0	341	100.0
	40mg/dl未満	10	0.7	0	0.0	1	0.6	2	0.9	2	0.8	2	0.6	3	0.8	3	0.9
	40-49	8	0.5	0	0.0	0	0.0	0	0.0	4	1.6	2	0.6	2	0.5	3	0.9
	50-59	22	1.5	1	1.1	2	1.1	3	1.4	1	0.4	3	0.8	12	3.1	8	2.3
	60-69	57	3.8	9	10.2	8	4.4	7	3.2	5	2.0	13	3.6	15	3.9	17	5.0
	70-79	85	5.7	14	15.9	11	6.1	10	4.5	10	3.9	17	4.7	23	5.9	16	4.7
	80-89	131	8.8	13	14.8	16	8.8	17	7.7	19	7.4	26	7.2	40	10.3	33	9.7
	90-99	182	12.2	17	19.3	22	12.2	21	9.5	28	10.9	41	11.3	53	13.7	43	12.6
	100-109	224	15.0	8	9.1	23	12.7	29	13.1	46	18.0	59	16.3	59	15.2	55	16.1
	110-119	180	12.0	15	17.0	21	11.6	24	10.8	23	9.0	53	14.6	44	11.4	55	16.1
	120-129	189	12.6	6	6.8	25	13.8	28	12.6	40	15.6	38	10.5	52	13.4	29	8.5
	130-139	143	9.6	1	1.1	19	10.5	27	12.2	21	8.2	35	9.7	40	10.3	31	9.1
	140-149	111	7.4	2	2.3	15	8.3	21	9.5	24	9.4	32	8.8	17	4.4	18	5.3
	150-159	67	4.5	0	0.0	10	5.5	11	5.0	15	5.9	16	4.4	15	3.9	13	3.8
	160-169	40	2.7	0	0.0	3	1.7	13	5.9	6	2.3	10	2.8	8	2.1	8	2.3
	170-179	22	1.5	1	1.1	2	1.1	5	2.3	7	2.7	4	1.1	3	0.8	3	0.9
	180-189	8	0.5	0	0.0	1	0.6	0	0.0	3	1.2	4	1.1	0	0.0	2	0.6
	190-199	8	0.5	1	1.1	0	0.0	2	0.9	1	0.4	4	1.1	0	0.0	2	0.6
	200-209	3	0.2	0	0.0	0	0.0	2	0.9	0	0.0	1	0.3	0	0.0	0	0.0
	210-219	2	0.1	0	0.0	0	0.0	0	0.0	1	0.4	1	0.3	0	0.0	1	0.3
	220mg/dl以上	4	0.3	0	0.0	2	1.1	0	0.0	0	0.0	1	0.3	1	0.3	1	0.3
平均値(mg/dl) 標準偏差		113.1 29.7		97.4 24.1		114.1 30.0		118.8 30.6		116.7 29.8		115.6 30.0		108.3 28.0		109.6 30.0	110.0 27.2
女 性	総数	2,158	100.0	139	100.0	334	100.0	363	100.0	411	100.0	432	100.0	479	100.0	395	100.0
	40mg/dl未満	3	0.1	0	0.0	3	0.9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	40-49	10	0.5	1	0.7	0	0.0	3	0.8	2	0.5	2	0.5	2	0.4	1	0.3
	50-59	26	1.2	6	4.3	4	1.2	6	1.7	6	1.5	0	0.0	4	0.8	1	0.3
	60-69	50	2.3	8	5.8	21	6.3	10	2.8	1	0.2	1	0.2	9	1.9	2	0.5
	70-79	119	5.5	21	15.1	37	11.1	28	7.7	7	1.7	5	1.2	21	4.4	7	1.8
	80-89	209	9.7	28	20.1	64	19.2	35	9.6	17	4.1	24	5.6	41	8.6	20	5.1
	90-99	214	9.9	25	18.0	52	15.6	45	12.4	19	4.6	22	5.1	51	10.6	29	7.3
	100-109	273	12.7	16	11.5	56	16.8	52	14.3	53	12.9	38	8.8	58	12.1	39	9.9
	110-119	260	12.0	17	12.2	28	8.4	49	13.5	54	13.1	59	13.7	53	11.1	52	13.2
	120-129	257	11.9	8	5.8	31	9.3	41	11.3	67	16.3	52	12.0	58	12.1	54	13.7
	130-139	220	10.2	4	2.9	14	4.2	29	8.0	54	13.1	59	13.7	60	12.5	53	13.4
	140-149	174	8.1	3	2.2	8	2.4	27	7.4	45	10.9	51	11.8	40	8.4	44	11.1
	150-159	131	6.1	1	0.7	7	2.1	17	4.7	30	7.3	41	9.5	35	7.3	37	9.4
	160-169	73	3.4	0	0.0	3	0.9	10	2.8	22	5.4	21	4.9	17	3.5	15	3.8
	170-179	64	3.0	0	0.0	4	1.2	5	1.4	17	4.1	21	4.9	17	3.5	23	5.8
	180-189	29	1.3	1	0.7	1	0.3	2	0.6	10	2.4	11	2.5	4	0.8	6	1.5
	190-199	20	0.9	0	0.0	1	0.3	2	0.6	4	1.0	10	2.3	3	0.6	4	1.0
	200-209	15	0.7	0	0.0	0	0.0	1	0.3	1	0.2	9	2.1	4	0.8	4	1.0
	210-219	2	0.1	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.2	1	0.3
	220mg/dl以上	9	0.4	0	0.0	0	0.0	1	0.3	1	0.2	6	1.4	1	0.2	3	0.8
平均値(mg/dl) 標準偏差		118.7 31.8		94.9 22.6		100.0 25.6		112.5 29.0		128.5 28.6		134.4 32.4		120.9 30.4		130.3 29.8	117.2 30.6

Friedewald の式 LDL コレステロール＝総コレステロール－HDL コレステロール－中性脂肪 /5
 （中性脂肪（トリグリセライド）が 400mg/dl 未満の場合。）

注）コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬服用者除外

(参考) 第37表 血清LDL- コレステロール値(直接測定法)の分布(性・年齢階級別)

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,565	100.0	90	100.0	195	100.0	231	100.0	274	100.0	381	100.0	394	100.0	352	100.0
	40mg/dl未満	5	0.3	0	0.0	1	0.5	1	0.4	1	0.4	1	0.3	1	0.3	1	0.4
	40-49	7	0.4	0	0.0	0	0.0	1	0.4	5	1.8	1	0.3	0	0.0	1	0.0
	50-59	24	1.5	1	1.1	0	0.0	3	1.3	1	0.4	4	1.0	15	3.8	11	3.1
	60-69	48	3.1	5	5.6	1	0.5	4	1.7	7	2.6	12	3.1	19	4.8	18	5.1
	70-79	76	4.9	11	12.2	8	4.1	8	3.5	6	2.2	20	5.2	23	5.8	17	4.8
	80-89	124	7.9	16	17.8	18	9.2	15	6.5	16	5.8	24	6.3	35	8.9	28	8.0
	90-99	185	11.8	14	15.6	26	13.3	21	9.1	32	11.7	40	10.5	52	13.2	38	10.8
	100-109	196	12.5	10	11.1	17	8.7	23	10.0	36	13.1	52	13.6	58	14.7	55	15.6
	110-119	227	14.5	10	11.1	32	16.4	26	11.3	42	15.3	56	14.7	61	15.5	61	17.3
	120-129	177	11.3	11	12.2	19	9.7	26	11.3	32	11.7	46	12.1	43	10.9	38	10.8
	130-139	170	10.9	4	4.4	30	15.4	32	13.9	28	10.2	41	10.8	35	8.9	30	8.5
	140-149	114	7.3	4	4.4	15	7.7	18	7.8	23	8.4	31	8.1	23	5.8	22	6.3
	150-159	104	6.6	1	1.1	16	8.2	26	11.3	21	7.7	24	6.3	16	4.1	14	4.0
	160-169	46	2.9	0	0.0	5	2.6	11	4.8	11	4.0	10	2.6	9	2.3	7	2.0
	170-179	29	1.9	0	0.0	3	1.5	9	3.9	4	1.5	11	2.9	2	0.5	5	1.4
	180-189	13	0.8	0	0.0	1	0.5	4	1.7	3	1.1	4	1.0	1	0.3	3	0.9
	190-199	8	0.5	2	2.2	1	0.5	0	0.0	4	1.5	1	0.3	0	0.0	0	0.0
	200-209	3	0.2	0	0.0	0	0.0	1	0.4	1	0.4	1	0.3	0	0.0	1	0.3
	210-219	6	0.4	0	0.0	0	0.0	2	0.9	1	0.4	2	0.5	1	0.3	2	0.6
	220mg/dl以上	3	0.2	1	1.1	2	1.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
平均値(mg/dl) 標準偏差		116.1 29.8		104.2 30.2		119.4 28.4		123.9 31.3		119.6 30.4		117.7 29.4		108.8 27.3		111.4 29.2	109.7 26.4
女 性	総数	2,186	100.0	139	100.0	337.0	100.0	369	100.0	416	100.0	443	100.0	482	100.0	402	100.0
	40mg/dl未満	2	0.1	0	0.0	1	0.3	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0
	40-49	4	0.2	0	0.0	1	0.3	1	0.3	1	0.2	1	0.2	0	0.0	0	0.0
	50-59	26	1.2	5	3.6	2	0.6	6	1.6	5	1.2	0	0.0	8	1.7	4	1.4
	60-69	46	2.1	5	3.6	11	3.3	10	2.7	4	1.0	1	0.2	15	3.1	2	0.5
	70-79	117	5.4	20	14.4	38	11.3	25	6.8	9	2.2	2	0.5	23	4.8	6	1.5
	80-89	176	8.1	23	16.5	57	16.9	30	8.1	12	2.9	20	4.5	34	7.1	19	4.7
	90-99	217	9.9	26	18.7	53	15.7	40	10.8	19	4.6	32	7.2	47	9.8	29	7.2
	100-109	271	12.4	18	12.9	51	15.1	50	13.6	39	9.4	49	11.1	64	13.3	45	11.2
	110-119	280	12.8	20	14.4	36	10.7	48	13.0	68	16.3	49	11.1	59	12.2	58	14.4
	120-129	260	11.9	10	7.2	34	10.1	42	11.4	56	13.5	62	14.0	56	11.6	49	12.2
	130-139	224	10.2	6	4.3	18	5.3	36	9.8	56	13.5	49	11.1	59	12.2	50	12.4
	140-149	189	8.6	2	1.4	17	5.0	28	7.6	48	11.5	53	12.0	41	8.5	44	10.9
	150-159	136	6.2	2	1.4	5	1.5	22	6.0	38	9.1	38	8.6	31	6.4	38	9.5
	160-169	102	4.7	1	0.7	3	0.9	16	4.3	23	5.5	34	7.7	25	5.2	23	5.7
	170-179	55	2.5	0	0.0	4	1.2	9	2.4	19	4.6	16	3.6	7	1.5	15	3.7
	180-189	30	1.4	0	0.0	3	0.9	1	0.3	9	2.2	11	2.5	6	1.2	9	2.2
	190-199	28	1.3	1	0.7	2	0.6	2	0.5	7	1.7	13	2.9	3	0.6	5	1.2
	200-209	12	0.5	0	0.0	1	0.3	1	0.3	2	0.5	6	1.4	2	0.4	2	0.5
	210-219	2	0.1	0	0.0	0	0.0	1	0.3	0	0.0	1	0.2	0	0.0	1	0.2
	220mg/dl以上	9	0.4	0	0.0	0	0.0	1	0.3	1	0.2	5	1.1	2	0.4	3	0.7
平均値(mg/dl) 標準偏差		120.1 31.4		98.1 23.4		104.4 26.8		116.4 30.2		129.9 28.7		133.7 32.1		119.1 29.9		129.2 29.7	115.0 29.7

注) コレステロールを下げる薬または中性脂肪(トリグリセライド)を下げる薬服用者除外

第 38 表 血清クレアチニン値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,736	100.0	90	100.0	198	100.0	239	100.0	297	100.0	445	100.0	467	100.0	425	100.0	273	100.0
	0.4mg/dl未満	1	0.1	0	0.0	1	0.5	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	0.4-0.49	6	0.3	0	0.0	1	0.5	1	0.4	2	0.7	2	0.4	0	0.0	1	0.2	0	0.0
	0.5-0.59	58	3.3	1	1.1	4	2.0	10	4.2	13	4.4	15	3.4	15	3.2	13	3.1	9	3.3
	0.6-0.69	275	15.8	9	10.0	30	15.2	33	13.8	53	17.8	76	17.1	74	15.8	71	16.7	39	14.3
	0.7-0.79	559	32.2	35	38.9	78	39.4	97	40.6	99	33.3	139	31.2	111	23.8	116	27.3	64	23.4
	0.8-0.89	443	25.5	33	36.7	53	26.8	66	27.6	72	24.2	113	25.4	106	22.7	107	25.2	64	23.4
	0.9-0.99	223	12.8	11	12.2	28	14.1	24	10.0	34	11.4	55	12.4	71	15.2	64	15.1	38	13.9
	1.0-1.19	129	7.4	1	1.1	3	1.5	8	3.3	19	6.4	34	7.6	64	13.7	42	9.9	41	15.0
	1.2-1.39	27	1.6	0	0.0	0	0.0	0	0.0	4	1.3	3	0.7	20	4.3	7	1.6	15	5.5
	1.4-1.59	8	0.5	0	0.0	0	0.0	0	0.0	1	0.3	2	0.4	5	1.1	2	0.5	3	1.1
	1.6-1.79	3	0.2	0	0.0	0	0.0	0	0.0	0	0.0	3	0.7	0	0.0	0	0.0	0	0.0
	1.8-1.99	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	2.0mg/dl以上	4	0.2	0	0.0	0	0.0	0	0.0	0	0.0	3	0.7	1	0.2	2	0.5	0	0.0
	平均値(mg/dl) 標準偏差	0.8 0.4		0.8 0.1		0.8 0.1		0.8 0.1		0.8 0.1		0.8 0.7		0.9 0.3		0.8 0.3		0.9 0.2	
女 性	総数	2,553	100.0	140	100.0	337	100.0	376	100.0	450	100.0	583	100.0	667	100.0	544	100.0	412	100.0
	0.4mg/dl未満	18	0.7	2	1.4	5	1.5	1	0.3	5	1.1	4	0.7	1	0.1	1	0.2	1	0.2
	0.4-0.49	296	11.6	14	10.0	50	14.8	52	13.8	58	12.9	54	9.3	68	10.2	61	11.2	38	9.2
	0.5-0.59	1,016	39.8	70	50.0	162	48.1	158	42.0	185	41.1	254	43.6	187	28.0	214	39.3	103	25.0
	0.6-0.69	812	31.8	45	32.1	103	30.6	132	35.1	146	32.4	174	29.8	212	31.8	174	32.0	127	30.8
	0.7-0.79	282	11.0	8	5.7	16	4.7	29	7.7	44	9.8	74	12.7	111	16.6	61	11.2	81	19.7
	0.8-0.89	91	3.6	1	0.7	1	0.3	4	1.1	12	2.7	16	2.7	57	8.5	25	4.6	37	9.0
	0.9-0.99	15	0.6	0	0.0	0	0.0	0	0.0	0	0.0	3	0.5	12	1.8	3	0.6	10	2.4
	1.0-1.19	12	0.5	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	11	1.6	3	0.6	9	2.2
	1.2-1.39	3	0.1	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	3	0.4	0	0.0	3	0.7
	1.4-1.59	4	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	3	0.4	0	0.0	3	0.7
	1.6-1.79	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	1.8-1.99	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	2.0mg/dl以上	4	0.2	0	0.0	0	0.0	0	0.0	0	0.0	2	0.3	2	0.3	2	0.4	0	0.0
	平均値(mg/dl) 標準偏差	0.6 0.2		0.6 0.1		0.6 0.1		0.6 0.1		0.6 0.1		0.6 0.4		0.7 0.3		0.6 0.3		0.7 0.2	

第 39 表 推算 GFR 値の分布（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男 性	総数	1,736	100.0	90	100.0	198	100.0	239	100.0	297	100.0	445	100.0	467	100.0	425	100.0	273	100.0
	15ml/min/1.73m ² 未満	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	1	0.2	1	0.2	0	0.0
	15-29	2	0.1	0	0.0	0	0.0	0	0.0	0	0.0	2	0.4	0	0.0	1	0.2	0	0.0
	30-59	193	11.1	0	0.0	0	0.0	2	0.8	20	6.7	47	10.6	124	26.6	65	15.3	84	30.8
	60-89	1,074	61.9	19	21.1	83	41.9	145	60.7	200	67.3	331	74.4	296	63.4	306	72.0	166	60.8
	90ml/min/1.73m ² 以上	465	26.8	71	78.9	115	58.1	92	38.5	77	25.9	64	14.4	46	9.9	52	12.2	23	8.4
	平均値 (mg/dl) 標準偏差	80.1 17.7		99.7 12.0		93.3 16.1		87.8 14.3		81.6 15.4		76.3 15.4		69.5 15.4		73.6 14.6		68.0 15.2	
女 性	総数	2,553	100.0	140	100.0	337	100.0	376	100.0	450	100.0	583	100.0	667	100.0	544	100.0	412	100.0
	15ml/min/1.73m ² 未満	4	0.2	0	0.0	0	0.0	0	0.0	0	0.0	2	0.3	2	0.3	2	0.4	0	0.0
	15-29	5	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.2	4	0.6	0	0.0	4	1.0
	30-59	232	9.1	0	0.0	0	0.0	2	0.5	13	2.9	42	7.2	175	26.2	65	11.9	131	31.8
	60-89	1,529	59.9	21	15.0	116	34.4	218	58.0	310	68.9	438	75.1	426	63.9	393	72.2	249	60.4
	90ml/min/1.73m ² 以上	783	30.7	119	85.0	221	65.6	156	41.5	127	28.2	100	17.2	60	9.0	84	15.4	28	6.8
	平均値 (mg/dl) 標準偏差	82.4 18.4		105.9 17.1		98.1 15.6		88.8 14.1		83.2 14.6		77.9 14.0		69.3 15.3		75.7 14.1		67.1 15.2	

推算 GFR (ml/min/1.73m²) = 194 × 血清クレアチニン値 (mg/dl)^{-1.094} × 年齢 (歳)^{-0.287} (男性)
 = 194 × 血清クレアチニン値 (mg/dl)^{-1.094} × 年齢 (歳)^{-0.287} × 0.739 (女性)

注) CKD ガイドライン 2009 年版の推算式

第40表 メタボリックシンドローム（内臓脂肪症候群）の状況（性・年齢階級別）

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
		4,232	100.0	225	100.0	517	100.0	611	100.0	741	100.0	1,024	100.0	1,114	100.0	2,822	100.0	964	100.0
総数	総数	747	17.7	2	0.9	26	5.0	57	9.3	123	16.6	245	23.9	294	26.4	544	19.3	246	25.5
	メタボリックシンドローム(内臓脂肪症候群)が強く疑われる者	534	12.6	9	4.0	51	9.9	89	14.6	92	12.4	149	14.6	144	12.9	402	14.2	158	16.4
	メタボリックシンドローム(内臓脂肪症候群)の予備群と考えられる者	2,951	69.7	214	95.1	440	85.1	465	76.1	526	71.0	630	61.5	676	60.7	1,876	66.5	560	58.1
	上記以外																		
男性	総数	1,722	100.0	88	100.0	197	100.0	238	100.0	295	100.0	445	100.0	459	100.0	1,170	100.0	423	100.0
	メタボリックシンドローム(内臓脂肪症候群)が強く疑われる者	474	27.5	1	1.1	21	10.7	39	16.4	90	30.5	159	35.7	164	35.7	356	30.4	149	35.2
	メタボリックシンドローム(内臓脂肪症候群)の予備群と考えられる者	360	20.9	8	9.1	43	21.8	65	27.3	61	20.7	99	22.2	84	18.3	270	23.1	107	25.3
	上記以外	888	51.6	79	89.8	133	67.5	134	56.3	144	48.8	187	42.0	211	46.0	544	46.5	167	39.5
女性	総数	2,510	100.0	137	100.0	320	100.0	373	100.0	446	100.0	579	100.0	655	100.0	1,652	100.0	541	100.0
	メタボリックシンドローム(内臓脂肪症候群)が強く疑われる者	273	10.9	1	0.7	5	1.6	18	4.8	33	7.4	86	14.9	130	19.8	188	11.4	97	17.9
	メタボリックシンドローム(内臓脂肪症候群)の予備群と考えられる者	174	6.9	1	0.7	8	2.5	24	6.4	31	7.0	50	8.6	60	9.2	132	8.0	51	9.4
	上記以外	2,063	82.2	135	98.5	307	95.9	331	88.7	382	85.7	443	76.5	465	71.0	1,332	80.6	393	72.6

注) 妊婦除外, ヘモグロビン Alc ≧ 5.5% の場合。項目の基準は p.18 参照

第 41 表の 1 腹囲区分別，血中脂質，血圧，血糖のいずれかのリスクを有する割合（性・年齢階級別）

		腹 囲	総 数		リ ス ク 保 有 数							
					0		1 つ		2 つ		3 つ	
			人数	%	人数	%	人数	%	人数	%	人数	%
総 数	総 数	<85cm(男)/90cm(女)	2,796	100.0	1,089	38.9	991	35.4	557	19.9	159	5.7
		≥85cm(男)/90cm(女)	1,436	100.0	155	10.8	534	37.2	523	36.4	224	15.6
	20-29歳	<85cm(男)/90cm(女)	202	100.0	176	87.1	24	11.9	2	1.0	0	0.0
		≥85cm(男)/90cm(女)	23	100.0	12	52.2	9	39.1	2	8.7	0	0.0
	30-39歳	<85cm(男)/90cm(女)	399	100.0	316	79.2	77	19.3	6	1.5	0	0.0
		≥85cm(男)/90cm(女)	118	100.0	41	34.7	51	43.2	22	18.6	4	3.4
	40-49歳	<85cm(男)/90cm(女)	420	100.0	258	61.4	128	30.5	31	7.4	3	0.7
		≥85cm(男)/90cm(女)	191	100.0	45	23.6	89	46.6	43	22.5	14	7.3
	50-59歳	<85cm(男)/90cm(女)	508	100.0	167	32.9	238	46.9	91	17.9	12	2.4
		≥85cm(男)/90cm(女)	233	100.0	18	7.7	92	39.5	96	41.2	27	11.6
	60-69歳	<85cm(男)/90cm(女)	607	100.0	105	17.3	262	43.2	187	30.8	53	8.7
		≥85cm(男)/90cm(女)	417	100.0	23	5.5	149	35.7	172	41.2	73	17.5
	70歳以上	<85cm(男)/90cm(女)	660	100.0	67	10.2	262	39.7	240	36.4	91	13.8
		≥85cm(男)/90cm(女)	454	100.0	16	3.5	144	31.7	188	41.4	106	23.3
	(再掲) 40-74歳	<85cm(男)/90cm(女)	1,782	100.0	564	31.6	720	40.4	399	22.4	99	5.6
		≥85cm(男)/90cm(女)	1,040	100.0	94	9.0	402	38.7	385	37.0	159	15.3
	(再掲) 65-74歳	<85cm(男)/90cm(女)	542	100.0	68	12.5	209	38.6	201	37.1	64	11.8
		≥85cm(男)/90cm(女)	422	100.0	18	4.3	158	37.4	155	36.7	91	21.6
	(再掲) 75歳以上	<85cm(男)/90cm(女)	413	100.0	33	8.0	170	41.2	150	36.3	60	14.5
		≥85cm(男)/90cm(女)	255	100.0	8	3.1	72	28.2	114	44.7	61	23.9
男 性	総 数	<85cm	783	100.0	247	31.5	321	41.0	177	22.6	38	4.9
		≥85cm	939	100.0	105	11.2	360	38.3	340	36.2	134	14.3
	20-29歳	<85cm	71	100.0	54	76.1	17	23.9	0	0.0	0	0.0
		≥85cm	17	100.0	8	47.1	8	47.1	1	5.9	0	0.0
	30-39歳	<85cm	102	100.0	70	68.6	30	29.4	2	2.0	0	0.0
		≥85cm	95	100.0	31	32.6	43	45.3	17	17.9	4	4.2
	40-49歳	<85cm	103	100.0	55	53.4	37	35.9	11	10.7	0	0.0
		≥85cm	135	100.0	31	23.0	65	48.1	30	22.2	9	6.7
	50-59歳	<85cm	132	100.0	32	24.2	65	49.2	31	23.5	4	3.0
		≥85cm	163	100.0	12	7.4	61	37.4	69	42.3	21	12.9
	60-69歳	<85cm	172	100.0	18	10.5	86	50.0	54	31.4	14	8.1
		≥85cm	273	100.0	15	5.5	99	36.3	114	41.8	45	16.5
	70歳以上	<85cm	203	100.0	18	8.9	86	42.4	79	38.9	20	9.9
		≥85cm	256	100.0	8	3.1	84	32.8	109	42.6	55	21.5
	(再掲) 40-74歳	<85cm	484	100.0	109	22.5	223	46.1	126	26.0	26	5.4
		≥85cm	686	100.0	60	8.7	270	39.4	257	37.5	99	14.4
	(再掲) 65-74歳	<85cm	158	100.0	7	4.4	75	47.5	58	36.7	18	11.4
		≥85cm	265	100.0	9	3.4	107	40.4	94	35.5	55	20.8
	(再掲) 75歳以上	<85cm	126	100.0	14	11.1	51	40.5	49	38.9	12	9.5
		≥85cm	141	100.0	6	4.3	39	27.7	65	46.1	31	22.0
女 性	総 数	<90cm	2,013	100.0	842	41.8	670	33.3	380	18.9	121	6.0
		≥90cm	497	100.0	50	10.1	174	35.0	183	36.8	90	18.1
	20-29歳	<90cm	131	100.0	122	93.1	7	5.3	2	1.5	0	0.0
		≥90cm	6	100.0	4	66.7	1	16.7	1	16.7	0	0.0
	30-39歳	<90cm	297	100.0	246	82.8	47	15.8	4	1.3	0	0.0
		≥90cm	23	100.0	10	43.5	8	34.8	5	21.7	0	0.0
	40-49歳	<90cm	317	100.0	203	64.0	91	28.7	20	6.3	3	0.9
		≥90cm	56	100.0	14	25.0	24	42.9	13	23.2	5	8.9
	50-59歳	<90cm	376	100.0	135	35.9	173	46.0	60	16.0	8	2.1
		≥90cm	70	100.0	6	8.6	31	44.3	27	38.6	6	8.6
	60-69歳	<90cm	435	100.0	87	20.0	176	40.5	133	30.6	39	9.0
		≥90cm	144	100.0	8	5.6	50	34.7	58	40.3	28	19.4
	70歳以上	<90cm	457	100.0	49	10.7	176	38.5	161	35.2	71	15.5
		≥90cm	198	100.0	8	4.0	60	30.3	79	39.9	51	25.8
	(再掲) 40-74歳	<90cm	1,298	100.0	455	35.1	497	38.3	273	21.0	73	5.6
		≥90cm	354	100.0	34	9.6	132	37.3	128	36.2	60	16.9
	(再掲) 65-74歳	<90cm	384	100.0	61	15.9	134	34.9	143	37.2	46	12.0
		≥90cm	157	100.0	9	5.7	51	32.5	61	38.9	36	22.9
	(再掲) 75歳以上	<90cm	287	100.0	19	6.6	119	41.5	101	35.2	48	16.7
		≥90cm	114	100.0	2	1.8	33	28.9	49	43.0	30	26.3

注）妊婦除外，ヘモグロビン A1c ≥ 5.5% の場合。リスクの項目は基準は p.18 参照

第 41 表の 2 腹囲区分別, 血中脂質, 血圧, 血糖のリスク保有状況〔すべての組み合わせ①血中脂質②血圧③血糖〕(性・年齢階級別)

		腹 囲	総 数		リ ス ク 保 有 数															
					①有②有③有		①有②有③無		①無②有③有		①有②無③有		①有②無③無		①無②有③無		①無②無③有		①無②無③無	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	総 数	<85cm(男)/90cm(女)	2,796	100.0	159	5.7	134	4.8	390	13.9	33	1.2	54	1.9	720	25.8	217	7.8	1,089	38.9
		≥85cm(男)/90cm(女)	1,436	100.0	224	15.6	134	9.3	360	25.1	29	2.0	30	2.1	441	30.7	63	4.4	155	10.8
	20-29歳	<85cm(男)/90cm(女)	202	100.0	0	0.0	0	0.0	1	0.5	1	0.5	2	1.0	18	8.9	4	2.0	176	87.1
		≥85cm(男)/90cm(女)	23	100.0	0	0.0	0	0.0	1	4.3	1	4.3	3	13.0	5	21.7	1	4.3	12	52.2
	30-39歳	<85cm(男)/90cm(女)	399	100.0	0	0.0	1	0.3	5	1.3	0	0.0	3	0.8	42	10.5	32	8.0	316	79.2
		≥85cm(男)/90cm(女)	118	100.0	4	3.4	11	9.3	8	6.8	3	2.5	9	7.6	35	29.7	7	5.9	41	34.7
	40-49歳	<85cm(男)/90cm(女)	420	100.0	3	0.7	5	1.2	24	5.7	2	0.5	4	1.0	89	21.2	35	8.3	258	61.4
		≥85cm(男)/90cm(女)	191	100.0	14	7.3	12	6.3	28	14.7	3	1.6	4	2.1	70	36.6	15	7.9	45	23.6
	50-59歳	<85cm(男)/90cm(女)	508	100.0	12	2.4	22	4.3	64	12.6	5	1.0	9	1.8	166	32.7	63	12.4	167	32.9
		≥85cm(男)/90cm(女)	233	100.0	27	11.6	23	9.9	67	28.8	6	2.6	7	3.0	72	30.9	13	5.6	18	7.7
	60-69歳	<85cm(男)/90cm(女)	607	100.0	53	8.7	49	8.1	122	20.1	16	2.6	24	4.0	182	30.0	56	9.2	105	17.3
		≥85cm(男)/90cm(女)	417	100.0	73	17.5	37	8.9	127	30.5	8	1.9	5	1.2	127	30.5	17	4.1	23	5.5
	70歳以上	<85cm(男)/90cm(女)	660	100.0	91	13.8	57	8.6	174	26.4	9	1.4	12	1.8	223	33.8	27	4.1	67	10.2
		≥85cm(男)/90cm(女)	454	100.0	106	23.3	51	11.2	129	28.4	8	1.8	2	0.4	132	29.1	10	2.2	16	3.5
	(再掲) 40-74歳	<85cm(男)/90cm(女)	1,782	100.0	99	5.6	96	5.4	273	15.3	30	1.7	42	2.4	512	28.7	166	9.3	564	31.6
		≥85cm(男)/90cm(女)	1,040	100.0	159	15.3	92	8.8	273	26.3	20	1.9	18	1.7	335	32.2	49	4.7	94	9.0
	(再掲) 65-74歳	<85cm(男)/90cm(女)	542	100.0	64	11.8	52	9.6	137	25.3	12	2.2	16	3.0	159	29.3	34	6.3	68	12.5
		≥85cm(男)/90cm(女)	422	100.0	91	21.6	41	9.7	105	24.9	9	2.1	6	1.4	139	32.9	13	3.1	18	4.3
	(再掲) 75歳以上	<85cm(男)/90cm(女)	413	100.0	60	14.5	37	9.0	111	26.9	2	0.5	7	1.7	148	35.8	15	3.6	33	8.0
		≥85cm(男)/90cm(女)	255	100.0	61	23.9	31	12.2	78	30.6	5	2.0	0	0.0	66	25.9	6	2.4	8	3.1
男 性	総 数	<85cm	783	100.0	38	4.9	37	4.7	132	16.9	8	1.0	16	2.0	255	32.6	50	6.4	247	31.5
		≥85cm	939	100.0	134	14.3	91	9.7	231	24.6	18	1.9	24	2.6	306	32.6	30	3.2	105	11.2
	20-29歳	<85cm	71	100.0	0	0.0	0	0.0	0	0.0	0	0.0	1	1.4	14	19.7	2	2.8	54	76.1
		≥85cm	17	100.0	0	0.0	0	0.0	1	5.9	0	0.0	3	17.6	4	23.5	1	5.9	8	47.1
	30-39歳	<85cm	102	100.0	0	0.0	1	1.0	1	1.0	0	0.0	2	2.0	25	24.5	3	2.9	70	68.6
		≥85cm	95	100.0	4	4.2	10	10.5	5	5.3	2	2.1	7	7.4	31	32.6	5	5.3	31	32.6
	40-49歳	<85cm	103	100.0	0	0.0	2	1.9	8	7.8	1	1.0	3	2.9	27	26.2	7	6.8	55	53.4
		≥85cm	135	100.0	9	6.7	10	7.4	18	13.3	2	1.5	4	3.0	55	40.7	6	4.4	31	23.0
	50-59歳	<85cm	132	100.0	4	3.0	8	6.1	22	16.7	1	0.8	4	3.0	46	34.8	15	11.4	32	24.2
		≥85cm	163	100.0	21	12.9	21	12.9	43	26.4	5	3.1	5	3.1	52	31.9	4	2.5	12	7.4
	60-69歳	<85cm	172	100.0	14	8.1	11	6.4	38	22.1	5	2.9	3	1.7	68	39.5	15	8.7	18	10.5
		≥85cm	273	100.0	45	16.5	25	9.2	84	30.8	5	1.8	3	1.1	87	31.9	9	3.3	15	5.5
	70歳以上	<85cm	203	100.0	20	9.9	15	7.4	63	31.0	1	0.5	3	1.5	75	36.9	8	3.9	18	8.9
		≥85cm	256	100.0	55	21.5	25	9.8	80	31.3	4	1.6	2	0.8	77	30.1	5	2.0	8	3.1
	(再掲) 40-74歳	<85cm	484	100.0	26	5.4	24	5.0	94	19.4	8	1.7	12	2.5	170	35.1	41	8.5	109	22.5
		≥85cm	686	100.0	99	14.4	69	10.1	174	25.4	14	2.0	14	2.0	235	34.3	21	3.1	60	8.7
	(再掲) 65-74歳	<85cm	158	100.0	18	11.4	9	5.7	46	29.1	3	1.9	4	2.5	63	39.9	8	5.1	7	4.4
		≥85cm	265	100.0	55	20.8	27	10.2	60	22.6	7	2.6	5	1.9	93	35.1	9	3.4	9	3.4
	(再掲) 75歳以上	<85cm	126	100.0	12	9.5	12	9.5	37	29.4	0	0.0	1	0.8	46	36.5	4	3.2	14	11.1
		≥85cm	141	100.0	31	22.0	12	8.5	51	36.2	2	1.4	0	0.0	36	25.5	3	2.1	6	4.3
女 性	総 数	<90cm	2,013	100.0	121	6.0	97	4.8	258	12.8	25	1.2	38	1.9	465	23.1	167	8.3	842	41.8
		≥90cm	497	100.0	90	18.1	43	8.7	129	26.0	11	2.2	6	1.2	135	27.2	33	6.6	50	10.1
	20-29歳	<90cm	131	100.0	0	0.0	0	0.0	1	0.8	1	0.8	1	0.8	4	3.1	2	1.5	122	93.1
		≥90cm	6	100.0	0	0.0	0	0.0	0	0.0	1	16.7	0	0.0	1	16.7	0	0.0	4	66.7
	30-39歳	<90cm	297	100.0	0	0.0	0	0.0	4	1.3	0	0.0	1	0.3	17	5.7	29	9.8	246	82.8
		≥90cm	23	100.0	0	0.0	1	4.3	3	13.0	1	4.3	2	8.7	4	17.4	2	8.7	10	43.5
	40-49歳	<90cm	317	100.0	3	0.9	3	0.9	16	5.0	1	0.3	1	0.3	62	19.6	28	8.8	203	64.0
		≥90cm	56	100.0	5	8.9	2	3.6	10	17.9	1	1.8	0	0.0	15	26.8	9	16.1	14	25.0
	50-59歳	<90cm	376	100.0	8	2.1	14	3.7	42	11.2	4	1.1	5	1.3	120	31.9	48	12.8	135	35.9
		≥90cm	70	100.0	6	8.6	2	2.9	24	34.3	1	1.4	2	2.9	20	28.6	9	12.9	6	8.6
	60-69歳	<90cm	435	100.0	39	9.0	38	8.7	84	19.3	11	2.5	21	4.8	114	26.2	41	9.4	87	20.0
		≥90cm	144	100.0	28	19.4	12	8.3	43	29.9	3	2.1	2	1.4	40	27.8	8	5.6	8	5.6
	70歳以上	<90cm	457	100.0	71	15.5	42	9.2	111	24.3	8	1.8	9	2.0	148	32.4	19	4.2	49	10.7
		≥90cm	198	100.0	51	25.8	26	13.1	49	24.7	4	2.0	0	0.0	55	27.8	5	2.5	8	4.0
	(再掲) 40-74歳	<90cm	1,298	100.0	73	5.6	72	5.5	179	13.8	22	1.7	30	2.3	342	26.3	125	9.6	455	35.1
		≥90cm	354	100.0	60	16.9	23	6.5	99	28.0	6	1.7	4	1.1	100	28.2	28	7.9	34	9.6
	(再掲) 65-74歳	<90cm	384	100.0	46	12.0	43	11.2	91	23.7	9	2.3	12	3.1	96	25.0	26	6.8	61	15.9
		≥90cm	157	100.0	36	22.9	14	8.9	45	28.7	2	1.3	1	0.6	46	29.3	4	2.5	9	5.7
	(再掲) 75歳以上	<90cm	287	100.0	48	16.7	25	8.7	74	25.8	2	0.7	6	2.1	102	35.5	11	3.8	19	6.6
		≥90cm	114	100.0	30	26.3	19	16.7	27	23.7	3	2.6	0	0.0	30	26.3	3	2.6	2	1.8

注) 妊婦除外, ヘモグロビン A1c ≥ 5.5% の場合。リスクの項目は基準は p.18 参照

第 42 表 「糖尿病が強く疑われる人」及び「糖尿病の可能性を否定できない人」の状況（性・年齢階級別）

	総 数	20～29歳		30～39歳		40～49歳		50～59歳		60～69歳		70歳以上		(再掲) 40～74歳		(再掲) 65～74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	糖尿病が強く疑われる人	4,273	100.0	230	100.0	533	100.0	614	100.0	743	100.0	1,026	100.0	1,127	100.0	2,830	100.0	966	100.0
		513	12.0	2	0.9	9	1.7	33	5.4	79	10.6	176	17.2	214	19.0	372	13.1	182	18.8
		230	44.8	1	50.0	0	0.0	14	42.4	36	45.6	70	39.8	109	50.9	161	43.3	77	42.3
		680	15.9	3	1.3	36	6.8	64	10.4	116	15.6	213	20.8	248	22.0	491	17.3	201	20.8
		3,080	72.1	225	97.8	488	91.6	517	84.2	548	73.8	637	62.1	665	59.0	1,967	69.5	583	60.4
男 性	糖尿病が強く疑われる人	1,730	100.0	90	100.0	197	100.0	239	100.0	296	100.0	445	100.0	463	100.0	1,173	100.0	424	100.0
		275	15.9	1	1.1	3	1.5	16	6.7	44	14.9	94	21.1	117	25.3	198	16.9	97	22.9
		130	47.3	0	0.0	0	0.0	9	56.3	20	45.5	41	43.6	60	51.3	93	47.0	43	44.3
		265	15.3	2	2.2	9	4.6	23	9.6	44	14.9	95	21.3	92	19.9	202	17.2	82	19.3
		1,190	68.8	87	96.7	185	93.9	200	83.7	208	70.3	256	57.5	254	54.9	773	65.9	245	57.8
女 性	糖尿病が強く疑われる人	2,543	100.0	140	100.0	336	100.0	375	100.0	447	100.0	581	100.0	664	100.0	1,657	100.0	542	100.0
		238	9.4	1	0.7	6	1.8	17	4.5	35	7.8	82	14.1	97	14.6	174	10.5	85	15.7
		100	42.0	1	100.0	0	0.0	5	29.4	16	45.7	29	35.4	49	50.5	68	39.1	34	40.0
		415	16.3	1	0.7	27	8.0	41	10.9	72	16.1	118	20.3	156	23.5	289	17.4	119	22.0
		1,890	74.3	138	98.6	303	90.2	317	84.5	340	76.1	381	65.6	411	61.9	1,194	72.1	338	62.4

注) ヘモグロビン A1c $\geq$ 5.6% の場合 判定基準は p.18 参照

* 「服薬者」とは、質問票で「インスリン注射または血糖値を下げる薬」の使用有と回答した者

第43表 高血圧症有病者の状況（性・年齢階級別）

		年齢(歳)																			
		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
				人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	総 数	4,449	100.0	244	100.0	540	100.0	632	100.0	769	100.0	1,082	100.0	1,182	100.0	2,952	100.0	1,013	100.0	713	100.0
	高血圧症有病者	2,070	46.5	6	2.5	50	9.3	150	23.7	323	42.0	638	59.0	903	76.4	1,451	49.2	685	67.6	563	79.0
	(再掲) 高血圧症有病者のうち服薬者*	1,153	55.7	2	33.3	2	4.0	39	26.0	119	36.8	366	57.4	625	69.2	735	50.7	424	61.9	414	73.5
	正常高値血圧者	620	13.9	20	8.2	57	10.6	105	16.6	148	19.2	174	16.1	116	9.8	477	16.2	141	13.9	66	9.3
	上記以外	1,759	39.5	218	89.3	433	80.2	377	59.7	298	38.8	270	25.0	163	13.8	1,024	34.7	187	18.5	84	11.8
男 性	総 数	1,799	100.0	96	100.0	202	100.0	244	100.0	303	100.0	466	100.0	488	100.0	1,215	100.0	442	100.0	286	100.0
	高血圧症有病者	996	55.4	4	4.2	33	16.3	92	37.7	164	54.1	312	67.0	391	80.1	731	60.2	324	73.3	228	79.7
	(再掲) 高血圧症有病者のうち服薬者*	509	51.1	1	25.0	0	0.0	25	27.2	62	37.8	179	57.4	242	61.9	358	49.0	192	59.3	150	65.8
	正常高値血圧者	280	15.6	15	15.6	44	21.8	41	16.8	60	19.8	77	16.5	43	8.8	198	16.3	64	14.5	23	8.0
	上記以外	523	29.1	77	80.2	125	61.9	111	45.5	79	26.1	77	16.5	54	11.1	286	23.5	54	12.2	35	12.2
女 性	総 数	2,650	100.0	148	100.0	338	100.0	388	100.0	466	100.0	616	100.0	694	100.0	1,737	100.0	571	100.0	427	100.0
	高血圧症有病者	1,074	40.5	2	1.4	17	5.0	58	14.9	159	34.1	326	52.9	512	73.8	720	41.5	361	63.2	335	78.5
	(再掲) 高血圧症有病者のうち服薬者*	644	60.0	1	50.0	2	11.8	14	24.1	57	35.8	187	57.4	383	74.8	377	52.4	232	64.3	264	78.8
	正常高値血圧者	340	12.8	5	3.4	13	3.8	64	16.5	88	18.9	97	15.7	73	10.5	279	16.1	77	13.5	43	10.1
	上記以外	1,236	46.6	141	95.3	308	91.1	266	68.6	219	47.0	193	31.3	109	15.7	738	42.5	133	23.3	49	11.5

注) 判定の基準は p.16 参照
* 「服薬者」とは、質問票で「血圧を下げる薬」の使用有と回答した者

第 44 表 「脂質異常症が疑われる人」の状況（採血時間によらず，性・年齢階級別）

			年齢(歳)																			
			総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳		(再掲) 65-74歳		(再掲) 75歳以上	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	総 数		4,279	100.0	229	100.0	532	100.0	613	100.0	746	100.0	1,028	100.0	1,131	100.0	2,836	100.0	969	100.0	682	100.0
	脂質異常症が疑われる人		806	18.8	8	3.5	31	5.8	47	7.7	113	15.1	265	25.8	342	30.2	558	19.7	291	30.0	209	30.6
	(再掲)脂質異常症が疑われる人のうち服薬者*		538	66.7	1	12.5	3	9.7	15	31.9	57	50.4	204	77.0	258	75.4	374	67.0	215	73.9	160	76.6
	上記以外		3,473	81.2	221	96.5	501	94.2	566	92.3	633	84.9	763	74.2	789	69.8	2,278	80.3	678	70.0	473	69.4
男 性	総 数		1,731	100.0	89	100.0	198	100.0	238	100.0	296	100.0	445	100.0	465	100.0	1,173	100.0	425	100.0	271	100.0
	脂質異常症が疑われる人		370	21.4	5	5.6	26	13.1	31	13.0	70	23.6	111	24.9	127	27.3	267	22.8	128	30.1	72	26.6
	(再掲)脂質異常症が疑われる人のうち服薬者*		171	46.2	0	0.0	3	11.5	8	25.8	23	32.9	64	57.7	73	57.5	127	47.6	73	57.0	41	56.9
	上記以外		1,361	78.6	84	94.4	172	86.9	207	87.0	226	76.4	334	75.1	338	72.7	906	77.2	297	69.9	199	73.4
女 性	総 数		2,548	100.0	140	100.0	334	100.0	375	100.0	450	100.0	583	100.0	666	100.0	1,663	100.0	544	100.0	411	100.0
	脂質異常症が疑われる人		436	17.1	3	2.1	5	1.5	16	4.3	43	9.6	154	26.4	215	32.3	291	17.5	163	30.0	137	33.3
	(再掲)脂質異常症が疑われる人のうち服薬者*		367	84.2	1	33.3	0	0.0	7	43.8	34	79.1	140	90.9	185	86.0	247	84.9	142	87.1	119	86.9
	上記以外		2,112	82.9	137	97.9	329	98.5	359	95.7	407	90.4	429	73.6	451	67.7	1,372	82.5	381	70.0	274	66.7

* 「服薬者」とは，質問票で「コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬」の使用有と回答した者

「脂質異常症が疑われる人」の判定
国民健康・栄養調査の血液検査では，空腹時採血が困難であるため，脂質異常症の診断基準項目である中性脂肪による判定は行わず，下記の通りとした。
・「脂質異常症が疑われる人」
HDL コレステロールが 40mg/dl 未満 もしくはコレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬を服用している者。

第45表 運動習慣の状況（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	4,871	100.0	295	100.0	614	100.0	713	100.0	831	100.0	1,143	100.0	1,275	100.0	1,058	100.0
	習慣無(健康上の理由)	379	7.8	3	1.0	13	2.1	18	2.5	24	2.9	78	6.8	243	19.1	105	9.9
	習慣無(その他)	3,072	63.1	240	81.4	490	79.8	546	76.6	614	73.9	590	51.6	592	46.4	498	47.1
	運動の習慣有	1,420	29.2	52	17.6	111	18.1	149	20.9	193	23.2	475	41.6	440	34.5	455	43.0
男性	総 数	2,010	100.0	118	100.0	236	100.0	291	100.0	343	100.0	496	100.0	526	100.0	462	100.0
	習慣無(健康上の理由)	131	6.5	1	0.8	4	1.7	8	2.7	7	2.0	25	5.0	86	16.3	35	7.6
	習慣無(その他)	1,231	61.2	87	73.7	181	76.7	214	73.5	256	74.6	263	53.0	230	43.7	207	44.8
	運動の習慣有	648	32.2	30	25.4	51	21.6	69	23.7	80	23.3	208	41.9	210	39.9	220	47.6
女性	総 数	2,861	100.0	177	100.0	378	100.0	422	100.0	488	100.0	647	100.0	749	100.0	596	100.0
	習慣無(健康上の理由)	248	8.7	2	1.1	9	2.4	10	2.4	17	3.5	53	8.2	157	21.0	70	11.7
	習慣無(その他)	1,841	64.3	153	86.4	309	81.7	332	78.7	358	73.4	327	50.5	362	48.3	291	48.8
	運動の習慣有	772	27.0	22	12.4	60	15.9	80	19.0	113	23.2	267	41.3	230	30.7	235	39.4

第46表 1週間の運動日数（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	4,871	100.0	295	100.0	614	100.0	713	100.0	831	100.0	1,143	100.0	1,275	100.0	1,058	100.0
	運動無し	3,451	70.8	243	82.4	503	81.9	564	79.1	638	76.8	668	58.4	835	65.5	603	57.0
	2日/週	298	6.1	17	5.8	35	5.7	38	5.3	44	5.3	99	8.7	65	5.1	76	7.2
	3日/週	233	4.8	10	3.4	14	2.3	26	3.6	32	3.9	81	7.1	70	5.5	77	7.3
	4日/週	134	2.8	1	0.3	11	1.8	12	1.7	20	2.4	46	4.0	44	3.5	47	4.4
	5日/週	170	3.5	17	5.8	22	3.6	20	2.8	24	2.9	43	3.8	44	3.5	52	4.9
	6日/週	79	1.6	0	0.0	7	1.1	12	1.7	10	1.2	27	2.4	23	1.8	23	2.2
	7日/週	506	10.4	7	2.4	22	3.6	41	5.8	63	7.6	179	15.7	194	15.2	180	17.0
男性	総 数	2,010	100.0	118	100.0	236	100.0	291	100.0	343	100.0	496	100.0	526	100.0	462	100.0
	運動無し	1,362	67.8	88	74.6	185	78.4	222	76.3	263	76.7	288	58.1	316	60.1	242	52.4
	2日/週	132	6.6	11	9.3	14	5.9	22	7.6	22	6.4	33	6.7	30	5.7	30	6.5
	3日/週	105	5.2	7	5.9	8	3.4	12	4.1	11	3.2	33	6.7	34	6.5	30	6.5
	4日/週	59	2.9	1	0.8	6	2.5	3	1.0	8	2.3	18	3.6	23	4.4	25	5.4
	5日/週	71	3.5	10	8.5	9	3.8	6	2.1	11	3.2	19	3.8	16	3.0	24	5.2
	6日/週	46	2.3	0	0.0	3	1.3	8	2.7	6	1.7	16	3.2	13	2.5	14	3.0
	7日/週	235	11.7	1	0.8	11	4.7	18	6.2	22	6.4	89	17.9	94	17.9	97	21.0
女性	総 数	2,861	100.0	177	100.0	378	100.0	422	100.0	488	100.0	647	100.0	749	100.0	596	100.0
	運動無し	2,089	73.0	155	87.6	318	84.1	342	81.0	375	76.8	380	58.7	519	69.3	361	60.6
	2日/週	166	5.8	6	3.4	21	5.6	16	3.8	22	4.5	66	10.2	35	4.7	46	7.7
	3日/週	128	4.5	3	1.7	6	1.6	14	3.3	21	4.3	48	7.4	36	4.8	47	7.9
	4日/週	75	2.6	0	0.0	5	1.3	9	2.1	12	2.5	28	4.3	21	2.8	22	3.7
	5日/週	99	3.5	7	4.0	13	3.4	14	3.3	13	2.7	24	3.7	28	3.7	28	4.7
	6日/週	33	1.2	0	0.0	4	1.1	4	0.9	4	0.8	11	1.7	10	1.3	9	1.5
	7日/週	271	9.5	6	3.4	11	2.9	23	5.5	41	8.4	90	13.9	100	13.4	83	13.9

第47表 運動を行う日の平均運動時間（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳	(再掲) 75歳以上
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	4,871	100.0	295	100.0	614	100.0	713	100.0	831	100.0	1,143	100.0	1,275	100.0	1,058	100.0
	運動無し	3,451	70.8	243	82.4	503	81.9	564	79.1	638	76.8	668	58.4	835	65.5	603	57.0
	30分以上60分未満	619	12.7	17	5.8	39	6.4	73	10.2	88	10.6	203	17.8	199	15.6	211	19.9
	60分以上120分未満	539	11.1	17	5.8	44	7.2	49	6.9	69	8.3	196	17.1	164	12.9	168	15.9
	120分以上180分未満	183	3.8	12	4.1	20	3.3	20	2.8	25	3.0	50	4.4	56	4.4	52	4.9
	180分以上	79	1.6	6	2.0	8	1.3	7	1.0	11	1.3	26	2.3	21	1.6	24	2.3
男性	総 数	2,010	100.0	118	100.0	236	100.0	291	100.0	343	100.0	496	100.0	526	100.0	462	100.0
	運動無し	1,362	67.8	88	74.6	185	78.4	222	76.3	263	76.7	288	58.1	316	60.1	242	52.4
	30分以上60分未満	239	11.9	5	4.2	14	5.9	30	10.3	34	9.9	84	16.9	72	13.7	88	19.0
	60分以上120分未満	270	13.4	10	8.5	22	9.3	27	9.3	26	7.6	90	18.1	95	18.1	95	20.6
	120分以上180分未満	87	4.3	10	8.5	8	3.4	5	1.7	13	3.8	21	4.2	30	5.7	23	5.0
	180分以上	52	2.6	5	4.2	7	3.0	7	2.4	7	2.0	13	2.6	13	2.5	14	3.0
女性	総 数	2,861	100.0	177	100.0	378	100.0	422	100.0	488	100.0	647	100.0	749	100.0	596	100.0
	運動無し	2,089	73.0	155	87.6	318	84.1	342	81.0	375	76.8	380	58.7	519	69.3	361	60.6
	30分以上60分未満	380	13.3	12	6.8	25	6.6	43	10.2	54	11.1	119	18.4	127	17.0	123	20.6
	60分以上120分未満	269	9.4	7	4.0	22	5.8	22	5.2	43	8.8	106	16.4	69	9.2	73	12.2
	120分以上180分未満	96	3.4	2	1.1	12	3.2	15	3.6	12	2.5	29	4.5	26	3.5	29	4.9
	180分以上	27	0.9	1	0.6	1	0.3	0	0.0	4	0.8	13	2.0	8	1.1	10	1.7

第 48 表 運動強度の状況（性・年齢階級別）

		総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	4,871	100.0	295	100.0	614	100.0	713	100.0	831	100.0	1,143	100.0	1,275	100.0	1,058	100.0	792	100.0
	運動無し	3,451	70.8	243	82.4	503	81.9	564	79.1	638	76.8	668	58.4	835	65.5	603	57.0	553	69.8
	高強度	105	2.2	17	5.8	23	3.7	22	3.1	12	1.4	19	1.7	12	0.9	14	1.3	5	0.6
	中強度	447	9.2	15	5.1	47	7.7	60	8.4	69	8.3	158	13.8	98	7.7	130	12.3	47	5.9
	低強度	868	17.8	20	6.8	41	6.7	67	9.4	112	13.5	298	26.1	330	25.9	311	29.4	187	23.6
男性	総 数	2,010	100.0	118	100.0	236	100.0	291	100.0	343	100.0	496	100.0	526	100.0	462	100.0	319	100.0
	運動無し	1,362	67.8	88	74.6	185	78.4	222	76.3	263	76.7	288	58.1	316	60.1	242	52.4	206	64.6
	高強度	65	3.2	13	11.0	14	5.9	16	5.5	4	1.2	10	2.0	8	1.5	9	1.9	4	1.3
	中強度	211	10.5	11	9.3	25	10.6	26	8.9	31	9.0	65	13.1	53	10.1	66	14.3	26	8.2
	低強度	372	18.5	6	5.1	12	5.1	27	9.3	45	13.1	133	26.8	149	28.3	145	31.4	83	26.0
女性	総 数	2,861	100.0	177	100.0	378	100.0	422	100.0	488	100.0	647	100.0	749	100.0	596	100.0	473	100.0
	運動無し	2,089	73.0	155	87.6	318	84.1	342	81.0	375	76.8	380	58.7	519	69.3	361	60.6	347	73.4
	高強度	40	1.4	4	2.3	9	2.4	6	1.4	8	1.6	9	1.4	4	0.5	5	0.8	1	0.2
	中強度	236	8.2	4	2.3	22	5.8	34	8.1	38	7.8	93	14.4	45	6.0	64	10.7	21	4.4
	低強度	496	17.3	14	7.9	29	7.7	40	9.5	67	13.7	165	25.5	181	24.2	166	27.9	104	22.0

第 49 表 歩行数の平均値及び標準偏差（性・年齢階級別）

		人数	平均値(歩)	標準偏差
総数	総 数	7,320	6,800	4,506
	15-19歳	363	7,929	4,759
	20-29	624	8,586	6,390
	30-39	1,016	7,632	4,291
	40-49	1,093	7,769	4,130
	50-59	1,210	7,521	4,033
	60-69	1,461	6,642	4,053
	70歳以上	1,553	4,178	3,450
	(再掲)20歳以上	6,957	6,741	4,485
	(再掲)65-74歳	1,274	5,951	3,764
	(再掲)75歳以上	976	3,393	3,086
男性	総 数	3,323	7,243	4,908
	15-19歳	187	7,722	4,875
	20-29	277	9,107	7,663
	30-39	459	8,537	4,974
	40-49	506	7,919	4,280
	50-59	573	7,772	4,280
	60-69	671	6,949	4,297
	70歳以上	650	4,707	3,840
	(再掲)20歳以上	3,136	7,214	4,910
	(再掲)65-74歳	588	6,379	4,106
	(再掲)75歳以上	384	3,765	3,418
女性	総 数	3,997	6,431	4,106
	15-19歳	176	8,149	4,635
	20-29	347	8,170	5,128
	30-39	557	6,886	3,465
	40-49	587	7,639	3,996
	50-59	637	7,296	3,786
	60-69	790	6,381	3,817
	70歳以上	903	3,797	3,086
	(再掲)20歳以上	3,821	6,352	4,063
	(再掲)65-74歳	686	5,584	3,404
	(再掲)75歳以上	592	3,151	2,828

第50表の1 歩行数の分布（性・年齢階級別）

		総 数		15-19歳		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 20歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総 数	総 数	7,320	100.0	363	100.0	624	100.0	1,016	100.0	1,093	100.0	1,210	100.0	1,461	100.0	1,553	100.0	6,957	100.0	1,274	100.0	976	100.0
	2,000歩未満	795	10.9	20	5.5	25	4.0	39	3.8	29	2.7	56	4.6	130	8.9	496	31.9	775	11.1	157	12.3	406	41.6
	2,000-4,000歩未満	1,284	17.5	39	10.7	72	11.5	150	14.8	148	13.5	167	13.8	307	21.0	401	25.8	1,245	17.9	305	23.9	253	25.9
	4,000-6,000	1,447	19.8	79	21.8	123	19.7	211	20.8	232	21.2	252	20.8	292	20.0	258	16.6	1,368	19.7	251	19.7	147	15.1
	6,000-8,000	1,319	18.0	80	22.0	114	18.3	207	20.4	237	21.7	252	20.8	252	17.2	177	11.4	1,239	17.8	231	18.1	75	7.7
	8,000-10,000	1,044	14.3	49	13.5	100	16.0	165	16.2	175	16.0	214	17.7	228	15.6	113	7.3	995	14.3	154	12.1	57	5.8
	10,000歩以上	1,431	19.5	96	26.4	190	30.4	244	24.0	272	24.9	269	22.2	252	17.2	108	7.0	1,335	19.2	176	13.8	38	3.9
	(再掲)9,200歩(男)/8,300歩(女)以上	2,041	27.9	123	33.9	251	40.2	350	34.4	369	33.8	394	32.6	381	26.1	173	11.1	1,918	27.6	264	20.7	68	7.0
男 性	総 数	3,323	100.0	187	100.0	277	100.0	459	100.0	506	100.0	573	100.0	671	100.0	650	100.0	3,136	100.0	588	100.0	384	100.0
	2,000歩未満	331	10.0	15	8.0	15	5.4	12	2.6	19	3.8	28	4.9	55	8.2	187	28.8	316	10.1	70	11.9	150	39.1
	2,000-4,000歩未満	539	16.2	21	11.2	31	11.2	55	12.0	69	13.6	75	13.1	129	19.2	159	24.5	518	16.5	122	20.7	98	25.5
	4,000-6,000	619	18.6	37	19.8	50	18.1	85	18.5	96	19.0	108	18.8	138	20.6	105	16.2	582	18.6	122	20.7	57	14.8
	6,000-8,000	570	17.2	42	22.5	44	15.9	90	19.6	97	19.2	117	20.4	108	16.1	72	11.1	528	16.8	92	15.6	29	7.6
	8,000-10,000	507	15.3	25	13.4	44	15.9	73	15.9	87	17.2	108	18.8	110	16.4	60	9.2	482	15.4	75	12.8	29	7.6
	10,000歩以上	757	22.8	47	25.1	93	33.6	144	31.4	138	27.3	137	23.9	131	19.5	67	10.3	710	22.6	107	18.2	21	5.5
	(再掲)9,200歩以上	923	27.8	53	28.3	106	38.3	170	37.0	167	33.0	172	30.0	168	25.0	87	13.4	870	27.7	135	23.0	26	6.8
女 性	総 数	3,997	100.0	176	100.0	347	100.0	557	100.0	587	100.0	637	100.0	790	100.0	903	100.0	3,821	100.0	686	100.0	592	100.0
	2,000歩未満	464	11.6	5	2.8	10	2.9	27	4.8	10	1.7	28	4.4	75	9.5	309	34.2	459	12.0	87	12.7	256	43.2
	2,000-4,000歩未満	745	18.6	18	10.2	41	11.8	95	17.1	79	13.5	92	14.4	178	22.5	242	26.8	727	19.0	183	26.7	155	26.2
	4,000-6,000	828	20.7	42	23.9	73	21.0	126	22.6	136	23.2	144	22.6	154	19.5	153	16.9	786	20.6	129	18.8	90	15.2
	6,000-8,000	749	18.7	38	21.6	70	20.2	117	21.0	140	23.9	135	21.2	144	18.2	105	11.6	711	18.6	139	20.3	46	7.8
	8,000-10,000	537	13.4	24	13.6	56	16.1	92	16.5	88	15.0	106	16.6	118	14.9	53	5.9	513	13.4	79	11.5	28	4.7
	10,000歩以上	674	16.9	49	27.8	97	28.0	100	18.0	134	22.8	132	20.7	121	15.3	41	4.5	625	16.4	69	10.1	17	2.9
	(再掲)8,300歩以上	1,118	28.0	70	39.8	145	41.8	180	32.3	202	34.4	222	34.9	213	27.0	86	9.5	1,048	27.4	129	18.8	42	7.1

第 50 表の 2 運動習慣の状況別，歩行数の分布（20 歳以上，性・年齢階級別）

			総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総 数	総 数	4,568	100.0	268	100.0	585	100.0	675	100.0	781	100.0	1,089	100.0	1,170	100.0	1,998	100.0	1,430	100.0
		2,000歩未満	489	10.7	8	3.0	16	2.7	15	2.2	36	4.6	85	7.8	329	28.1	232	11.6	524	36.6
		2,000-4,000歩未満	854	18.7	35	13.1	74	12.6	94	13.9	108	13.8	239	21.9	304	26.0	488	24.4	378	26.4
		4,000-6,000	929	20.3	55	20.5	134	22.9	139	20.6	166	21.3	230	21.1	205	17.5	388	19.4	238	16.6
		6,000-8,000	827	18.1	47	17.5	130	22.2	151	22.4	163	20.9	180	16.5	156	13.3	382	19.1	136	9.5
		8,000-10,000	655	14.3	46	17.2	102	17.4	116	17.2	136	17.4	167	15.3	88	7.5	238	11.9	96	6.7
		10,000歩以上	814	17.8	77	28.7	129	22.1	160	23.7	172	22.0	188	17.3	88	7.5	270	13.5	58	4.1
		9,200歩(男)/8,300歩(女)以上	1,218	26.7	110	41.0	195	33.3	225	33.3	256	32.8	292	26.8	140	12.0	414	20.7	108	7.6
	習慣無 (健康上の理由)	総 数	331	100.0	3	100.0	12	100.0	18	100.0	20	100.0	71	100.0	207	100.0	192	100.0	302	100.0
		2,000歩未満	149	45.0	0	0.0	1	8.3	1	5.6	4	20.0	14	19.7	129	62.3	68	35.4	206	68.2
		2,000-4,000歩未満	89	26.9	2	66.7	3	25.0	5	27.8	6	30.0	24	33.8	49	23.7	58	30.2	66	21.9
		4,000-6,000	47	14.2	0	0.0	4	33.3	7	38.9	5	25.0	17	23.9	14	6.8	26	13.5	18	6.0
		6,000-8,000	27	8.2	1	33.3	4	33.3	1	5.6	2	10.0	9	12.7	10	4.8	26	13.5	10	3.3
		8,000-10,000	9	2.7	0	0.0	0	0.0	1	5.6	1	5.0	4	5.6	3	1.4	10	5.2	2	0.7
		10,000歩以上	10	3.0	0	0.0	0	0.0	3	16.7	2	10.0	3	4.2	2	1.0	4	2.1	0	0.0
		9,200歩(男)/8,300歩(女)以上	15	4.5	0	0.0	0	0.0	4	22.2	2	10.0	6	8.5	3	1.4	10	5.2	2	0.7
	習慣無 (その他)	総 数	2,877	100.0	218	100.0	468	100.0	515	100.0	578	100.0	556	100.0	542	100.0	928	100.0	670	100.0
		2,000歩未満	262	9.1	8	3.7	13	2.8	12	2.3	26	4.5	51	9.2	152	28.0	120	12.9	238	35.5
		2,000-4,000歩未満	571	19.8	28	12.8	62	13.2	74	14.4	90	15.6	146	26.3	171	31.5	276	29.7	214	31.9
		4,000-6,000	621	21.6	45	20.6	105	22.4	117	22.7	126	21.8	136	24.5	92	17.0	190	20.5	112	16.7
		6,000-8,000	554	19.3	40	18.3	105	22.4	112	21.7	135	23.4	91	16.4	71	13.1	178	19.2	58	8.7
		8,000-10,000	412	14.3	38	17.4	84	17.9	91	17.7	99	17.1	70	12.6	30	5.5	88	9.5	28	4.2
		10,000歩以上	457	15.9	59	27.1	99	21.2	109	21.2	102	17.6	62	11.2	26	4.8	76	8.2	20	3.0
		9,200歩(男)/8,300歩(女)以上	711	24.7	87	39.9	153	32.7	160	31.1	166	28.7	102	18.3	43	7.9	122	13.1	34	5.1
	運動の 習慣有	総 数	1,360	100.0	47	100.0	105	100.0	142	100.0	183	100.0	462	100.0	421	100.0	878	100.0	458	100.0
		2,000歩未満	78	5.7	0	0.0	2	1.9	2	1.4	6	3.3	20	4.3	48	11.4	44	5.0	80	17.5
		2,000-4,000歩未満	194	14.3	5	10.6	9	8.6	15	10.6	12	6.6	69	14.9	84	20.0	154	17.5	98	21.4
		4,000-6,000	261	19.2	10	21.3	25	23.8	15	10.6	35	19.1	77	16.7	99	23.5	172	19.6	108	23.6
		6,000-8,000	246	18.1	6	12.8	21	20.0	38	26.8	26	14.2	80	17.3	75	17.8	178	20.3	68	14.8
		8,000-10,000	234	17.2	8	17.0	18	17.1	24	16.9	36	19.7	93	20.1	55	13.1	140	15.9	66	14.4
		10,000歩以上	347	25.5	18	38.3	30	28.6	48	33.8	68	37.2	123	26.6	60	14.3	190	21.6	38	8.3
		9,200歩(男)/8,300歩(女)以上	492	36.2	23	48.9	42	40.0	61	43.0	88	48.1	184	39.8	94	22.3	282	32.1	72	15.7
男性	総 数	総 数	1,861	100.0	105	100.0	222	100.0	269	100.0	313	100.0	472	100.0	480	100.0	866	100.0	572	100.0
		2,000歩未満	182	9.8	4	3.8	0	0.0	7	2.6	16	5.1	36	7.6	119	24.8	84	9.7	200	35.0
		2,000-4,000歩未満	332	17.8	14	13.3	23	10.4	40	14.9	39	12.5	96	20.3	120	25.0	192	22.2	146	25.5
		4,000-6,000	351	18.9	18	17.1	50	22.5	47	17.5	54	17.3	104	22.0	78	16.3	178	20.6	90	15.7
		6,000-8,000	316	17.0	18	17.1	45	20.3	54	20.1	69	22.0	69	14.6	61	12.7	146	16.9	52	9.1
		8,000-10,000	288	15.5	14	13.3	39	17.6	56	20.8	56	17.9	77	16.3	46	9.6	110	12.7	50	8.7
		10,000歩以上	392	21.1	37	35.2	65	29.3	65	24.2	79	25.2	90	19.1	56	11.7	156	18.0	34	5.9
		(再掲)9,200歩以上	494	26.5	40	38.1	79	35.6	86	32.0	95	30.4	122	25.8	72	15.0	200	23.1	44	7.7
	習慣無 (健康上の理由)	総 数	114	100.0	1	100.0	4	100.0	8	100.0	7	100.0	21	100.0	73	100.0	58	100.0	112	100.0
		2,000歩未満	50	43.9	0	0.0	0	0.0	0	0.0	1	14.3	5	23.8	44	60.3	16	27.6	78	69.6
		2,000-4,000歩未満	30	26.3	1	100.0	0	0.0	3	37.5	2	28.6	7	33.3	17	23.3	20	34.5	24	21.4
		4,000-6,000	12	10.5	0	0.0	2	50.0	2	25.0	1	14.3	3	14.3	4	5.5	8	13.8	2	1.8
		6,000-8,000	11	9.6	0	0.0	2	50.0	1	12.5	0	0.0	2	9.5	6	8.2	8	13.8	8	7.1
		8,000-10,000	4	3.5	0	0.0	0	0.0	0	0.0	1	14.3	2	9.5	1	1.4	4	6.9	0	0.0
		10,000歩以上	7	6.1	0	0.0	0	0.0	2	25.0	2	28.6	2	9.5	1	1.4	2	3.4	0	0.0
		(再掲)9,200歩以上	8	7.0	0	0.0	0	0.0	2	25.0	2	28.6	3	14.3	1	1.4	4	6.9	0	0.0
	習慣無 (その他)	総 数	1,133	100.0	77	100.0	170	100.0	199	100.0	233	100.0	249	100.0	205	100.0	384	100.0	240	100.0
		2,000歩未満	102	9.0	4	5.2	0	0.0	5	2.5	13	5.6	26	10.4	54	26.3	54	14.1	86	35.8
		2,000-4,000歩未満	208	18.4	9	11.7	20	11.8	31	15.6	31	13.3	56	22.5	61	29.8	98	25.5	72	30.0
		4,000-6,000	233	20.6	13	16.9	37	21.8	40	20.1	42	18.0	65	26.1	36	17.6	96	25.0	40	16.7
		6,000-8,000	196	17.3	14	18.2	30	17.6	40	20.1	57	24.5	34	13.7	21	10.2	54	14.1	16	6.7
		8,000-10,000	178	15.7	11	14.3	31	18.2	43	21.6	41	17.6	36	14.5	16	7.8	44	11.5	16	6.7
		10,000歩以上	216	19.1	26	33.8	52	30.6	40	20.1	49	21.0	32	12.9	17	8.3	38	9.9	10	4.2
		(再掲)9,200歩以上	276	24.4	28	36.4	62	36.5	58	29.1	63	27.0	43	17.3	22	10.7	48	12.5	14	5.8
	運動の 習慣有	総 数	614	100.0	27	100.0	48	100.0	62	100.0	73	100.0	202	100.0	202	100.0	424	100.0	220	100.0
		2,000歩未満	30	4.9	0	0.0	0	0.0	2	3.2	2	2.7	5	2.5	21	10.4	14	3.3	36	16.4
		2,000-4,000歩未満	94	15.3	4	14.8	3	6.3	6	9.7	6	8.2	33	16.3	42	20.8	74	17.5	50	22.7
		4,000-6,000	106	17.3	5	18.5	11	22.9	5	8.1	11	15.1	36	17.8	38	18.8	74	17.5	48	21.8
		6,000-8,000	109	17.8	4	14.8	13	27.1	13	21.0	12	16.4	33	16.3	34	16.8	84	19.8	28	12.7
		8,000-10,000	106	17.3	3	11.1	8	16.7	13	21.0	14	19.2	39	19.3	29	14.4	62	14.6	34	15.5
		10,000歩以上	169	27.5	11	40.7	13	27.1	23	37.1	28	38.4	56	27.7	38	18.8	116	27.4	24	10.9
		(再掲)9,200歩以上	210	34.2	12	44.4	17	35.4	26	41.9	30	41.1	76	37.6	49	24.3	148	34.9	30	13.6

			総 数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 65-74歳		(再掲) 75歳以上	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
女性	総 数	総 数	2,707	100.0	163	100.0	363	100.0	406	100.0	468	100.0	617	100.0	690	100.0	1,132	100.0	858	100.0
		2,000歩未満	307	11.3	4	2.5	16	4.4	8	2.0	20	4.3	49	7.9	210	30.4	148	13.1	324	37.8
		2,000-4,000歩未満	522	19.3	21	12.9	51	14.0	54	13.3	69	14.7	143	23.2	184	26.7	296	26.1	232	27.0
		4,000-6,000	578	21.4	37	22.7	84	23.1	92	22.7	112	23.9	126	20.4	127	18.4	210	18.6	148	17.2
		6,000-8,000	511	18.9	29	17.8	85	23.4	97	23.9	94	20.1	111	18.0	95	13.8	236	20.8	84	9.8
		8,000-10,000	367	13.6	32	19.6	63	17.4	60	14.8	80	17.1	90	14.6	42	6.1	128	11.3	46	5.4
		10,000歩以上	422	15.6	40	24.5	64	17.6	95	23.4	93	19.9	98	15.9	32	4.6	114	10.1	24	2.8
		(再掲)8,300歩以上	724	26.7	70	42.9	116	32.0	139	34.2	161	34.4	170	27.6	68	9.9	214	18.9	64	7.5
	習慣無 (健康上の理由)	総 数	217	100.0	2	100.0	8	100.0	10	100.0	13	100.0	50	100.0	134	100.0	134	100.0	190	100.0
		2,000歩未満	99	45.6	0	0.0	1	12.5	1	10.0	3	23.1	9	18.0	85	63.4	52	38.8	128	67.4
		2,000-4,000歩未満	59	27.2	1	50.0	3	37.5	2	20.0	4	30.8	17	34.0	32	23.9	38	28.4	42	22.1
		4,000-6,000	35	16.1	0	0.0	2	25.0	5	50.0	4	30.8	14	28.0	10	7.5	18	13.4	16	8.4
		6,000-8,000	16	7.4	1	50.0	2	25.0	0	0.0	2	15.4	7	14.0	4	3.0	18	13.4	2	1.1
		8,000-10,000	5	2.3	0	0.0	0	0.0	1	10.0	0	0.0	2	4.0	2	1.5	6	4.5	2	1.1
		10,000歩以上	3	1.4	0	0.0	0	0.0	1	10.0	0	0.0	1	2.0	1	0.7	2	1.5	0	0.0
		(再掲)8,300歩以上	7	3.2	0	0.0	0	0.0	2	20.0	0	0.0	3	6.0	2	1.5	6	4.5	2	1.1
	習慣無 (その他)	総 数	1,744	100.0	141	100.0	298	100.0	316	100.0	345	100.0	307	100.0	337	100.0	544	100.0	430	100.0
		2,000歩未満	160	9.2	4	2.8	13	4.4	7	2.2	13	3.8	25	8.1	98	29.1	66	12.1	152	35.3
		2,000-4,000歩未満	363	20.8	19	13.5	42	14.1	43	13.6	59	17.1	90	29.3	110	32.6	178	32.7	142	33.0
		4,000-6,000	388	22.2	32	22.7	68	22.8	77	24.4	84	24.3	71	23.1	56	16.6	94	17.3	72	16.7
		6,000-8,000	358	20.5	26	18.4	75	25.2	72	22.8	78	22.6	57	18.6	50	14.8	124	22.8	42	9.8
		8,000-10,000	234	13.4	27	19.1	53	17.8	48	15.2	58	16.8	34	11.1	14	4.2	44	8.1	12	2.8
		10,000歩以上	241	13.8	33	23.4	47	15.8	69	21.8	53	15.4	30	9.8	9	2.7	38	7.0	10	2.3
		(再掲)8,300歩以上	435	24.9	59	41.8	91	30.5	102	32.3	103	29.9	59	19.2	21	6.2	74	13.6	20	4.7
	運動の 習慣有	総 数	746	100.0	20	100.0	57	100.0	80	100.0	110	100.0	260	100.0	219	100.0	454	100.0	238	100.0
		2,000歩未満	48	6.4	0	0.0	2	3.5	0	0.0	4	3.6	15	5.8	27	12.3	30	6.6	44	18.5
		2,000-4,000歩未満	100	13.4	1	5.0	6	10.5	9	11.3	6	5.5	36	13.8	42	19.2	80	17.6	48	20.2
		4,000-6,000	155	20.8	5	25.0	14	24.6	10	12.5	24	21.8	41	15.8	61	27.9	98	21.6	60	25.2
		6,000-8,000	137	18.4	2	10.0	8	14.0	25	31.3	14	12.7	47	18.1	41	18.7	94	20.7	40	16.8
		8,000-10,000	128	17.2	5	25.0	10	17.5	11	13.8	22	20.0	54	20.8	26	11.9	78	17.2	32	13.4
		10,000歩以上	178	23.9	7	35.0	17	29.8	25	31.3	40	36.4	67	25.8	22	10.0	74	16.3	14	5.9
		(再掲)8,300歩以上	282	37.8	11	55.0	25	43.9	35	43.8	58	52.7	108	41.5	45	20.5	134	29.5	42	17.6

