

**Stockholm Convention
on Persistent Organic
Pollutants**

Persistent Organic Pollutants Review Committee
Seventeenth meeting
Geneva, 24–28 January 2022

**Report of the Persistent Organic Pollutants Review Committee
on the work of its seventeenth meeting****Addendum****Risk profile for Dechlorane Plus**

At its seventeenth meeting, by its decision POPRC-17/2, the Persistent Organic Pollutants Review Committee adopted a risk profile for Dechlorane Plus on the basis of the draft contained in the note by the Secretariat (UNEP/POPS/POPRC.17/3), as revised during the meeting. The text of the risk profile as adopted is set out in the annex to the present addendum. It has not been formally edited.

Annex*

Dechlorane Plus

Risk profile

January 2022

* The studies and other information referred to in this risk profile do not necessarily reflect the views of the Secretariat, the United Nations Environment Programme (UNEP) or the United Nations. The designations employed and the presentation of the material within such studies and references do not imply the expression of any opinion whatsoever on the part of the Secretariat, UNEP or the United Nations concerning geopolitical situations or the legal status of any country, territory, area or city or their authorities.
The annex has not been formally edited.

Table of contents

Executive summary	4
1. Introduction	6
1.1 Chemical Identity.....	6
1.2 Conclusion of the POPs Review Committee regarding Annex D information.....	7
1.3 Data sources.....	8
1.4 Status of the chemical under international conventions.....	8
2. Summary information relevant to the risk profile.....	8
2.1 Sources	8
2.1.1 Production, trade, stockpiles	8
2.1.2 Uses.....	9
2.1.3 Releases and emissions to the environment	10
2.2. Environmental fate.....	11
2.2.1 Persistence	11
2.2.2 Bioaccumulation and toxicokinetics	13
2.2.3 Potential for long-range transport	15
2.3 Exposure	18
2.3.1 Environmental levels and trends	18
2.3.2 Human exposure	24
2.4 Hazard assessment for endpoint of concern.....	26
2.4.1 Toxicity to aquatic organisms	26
2.4.2 Toxicity in terrestrial organisms	29
2.4.3 Human toxicity.....	30
2.4.4 Human epidemiology.....	32
2.4.5 Other concerns	33
3. Synthesis of information.....	33
4. Concluding statement.....	34
References	35

Executive summary

1. At its fifteenth meeting, the Persistent Organic Pollutants Review Committee concluded that Dechlorane Plus (DP, CAS No. 13560-89-9) and its *syn*-isomer (CAS No. 135821-03-3) and *anti*-isomer (CAS No. 135821-74-8) fulfilled the screening criteria in Annex D of the Stockholm Convention on Persistent Organic Pollutants (Decision POPRC-15/2), and decided to establish an intersessional working group to review the proposal further and to prepare a draft risk profile in accordance with Annex E to the Convention. Parties and observers were invited to submit information specified in Annex E, before 2 December 2019. This risk profile presents and considers data on Dechlorane plus and its *syn*- and *anti*-isomers, but for reasons of simplicity denoted as "Dechlorane Plus".

2. The POPs Review Committee considered the draft risk profile at its sixteenth meeting and adopted decision POPRC-16/1, by which it decided to defer its decision on the draft risk profile (UNEP/POPS/POPRC.16/INF/19) to its seventeenth meeting. In decision POPRC-16/1, the Committee noted that, while information on persistence, bioaccumulation and the potential for long-range environmental transport was conclusive, the Committee had been unable to reach agreement that the information on adverse effects was sufficient to reach a conclusion on the risk profile. The Committee also decided to establish an intersessional working group to review and update the draft risk profile; and invited Parties and observers to submit to the Secretariat additional information relating to the adverse effects of DP before 1 March 2021.

3. The "Dechlorane Plus"TM technical mixture is a commercially available polychlorinated flame retardant that has been in use since the 1960s. It is used as an additive flame retardant in electrical wire and cable coatings, plastic roofing materials, connectors in TV and computer monitors, and as a non-plasticizing flame retardant in polymeric systems, such as nylon and polypropylene plastic. It is currently marketed as a replacement substance for deca-BDE.

4. The technical DP mixture contains two stereoisomers, *syn*-DP and *anti*-DP, that are present in ratios of about 1:3 or 25 % *syn*-DP and 75 % *anti*-DP. Commercially available DP mixtures may also contain DP monoadducts, mono-dechlorinated DP and other substances as impurities. DP and its isomers are not known to be unintentionally produced.

5. DP production started in the 1960s at the manufacturing plant of Hooker Chemicals and Plastics Corporation (now Occidental Chemical Company (OxyChem)) in Niagara Falls, New York. DP has been considered as a high production volume chemical in the United States with an estimated annual production of 450-4500 tonnes from 1986. Production in China started in 2003 and the annual production is reported to be 300-1000 tonnes. The Chinese manufacturer, Anpon Electrochemical Company Ltd., was recently acquired by ADAMA, which may now be the sole global manufacturer (ADAMA, 2019).

6. DP is released to the environment during production, processing and use, as well as from waste disposal and recycling activities. Releases from use include industrial and professional uses as well as releases from consumer products. DP entering the environment is expected to predominantly reside in soil and/or sediment, depending on the compartment of release, but DP remaining in air and in water has the potential for dispersion and may undergo environmental transport by particle binding. DP has been detected globally in many locations, ranging from production and recycling sites to urban, rural and remote areas. It has been detected in air, water, sediment, ice, soil, wastewater, sludge, biosolids, landfill leachate, indoor and outdoor dust, wildlife and humans.

7. On a global scale, the highest DP concentrations were detected close to known production sites or electronic waste (e-waste) treatment facilities. Monitoring studies document the long-range transport of DP to remote regions via the atmosphere, ocean currents and possibly also via migratory birds. Modelling studies suggest DP has transport and persistence properties similar to listed POPs. DP has been detected in different environmental matrices and biota in the Arctic, Antarctic, Tibetan Plateau, a mountain region in Asia and in the remote Xilingol Prairie in Inner Mongolia, China. The atmospheric half-life of DP is in one study calculated to be ~14 hours, i.e. below the criterion of two days set in Annex D (d) (iii) of the Convention. Another study report estimated half-lives in air of 0.468 and 160.12 days for atmospheric oxidation and ozone reaction, respectively, using the AOPWIN 2010 model (12-hr day). However, actual half-lives may be longer as a result of sorption to particles, which is presumed to be the primary mode of transport for DP due to its low vapour pressure and a high log octanol-air partitioning coefficient. Available monitoring data confirms this assumption and shows that DP is found predominantly in the particulate phase both in air and water. Long-range transport of DP is thus likely mediated by particle binding. Sorption to particles may slow down reaction rates, increase the actual half-lives in air and facilitate the long-range transport of DP.

8. DP is persistent in water, soil and sediment, and is also considered to be persistent in air under natural conditions as a result of its binding to air particles and limited photodegradation. It has very low water solubility and a high octanol-water partition coefficient. It binds to organic matter in soil and sediments and is therefore probably not easily bioavailable for microorganisms. Furthermore, DP has no functional groups that are susceptible for hydrolysis and is not expected to hydrolyse. Modelling predicts that aerobic biodegradation of DP would be very slow and similar to analogue chemicals (mirex, chlordane, heptachlor, dieldrin/endrin, aldrin and endosulfan) that are already listed under the Stockholm Convention. DP is detected in deep soil and sediment cores, indicating its persistence. Monitoring data show limited degradation in soil and sediments over time and support the conclusion that DP is

persistent. This is confirmed in a recent laboratory study which shows that the half-life of DP in soil is greater than 180 days.

9. Bioconcentration factors above 5000 have recently been reported for the DP isomers in a fish study. Depuration half-lives corresponding to a bioconcentration factor above 5000 L/kg have also been reported for DP, as well as biomagnification and trophic magnification factors above 1 in a variety of food chains and locations, including an Antarctic food chain. DP has been identified as a very persistent and very bioaccumulative substance (vPvB) in the European Union.

10. A Canadian screening assessment concludes that there is risk of harm to the environment from DP, and that DP meets the criteria under paragraph 64(a) of the Canadian Environmental Protection Act, 1999 (CEPA) as it is entering or may enter the environment in a quantity or concentration or under conditions that have or may have an immediate or long-term harmful effect on the environment or its biological diversity.

11. Exposure of the general population to DP takes place by consumption of food and drinking water, inhalation of indoor and ambient air, as well as respiratory and oral uptake of dust containing DP. DP has been detected in human blood and breast milk in many regions of the world. Moreover, the human foetus may be exposed to DP via the umbilical cord blood, and breast milk may be an important source of exposure for infants. The highest DP levels have been observed in occupationally exposed workers and residents living near production facilities and e-waste recycling sites in China.

12. Available animal studies suggest low concern for acute toxicity via oral, inhalative or dermal routes of exposure in humans and other organisms, and chronic toxicity data on DP are limited. Toxicity studies with DP report effects such as oxidative damage, indications for neurodevelopmental toxicity and potential for endocrine disruption. Oxidative stress has been observed in marine macroalgae, fish, marine bivalves, earthworms, birds, and mice. In a marine macroalgae exposed to DP, photosynthesis was reduced at low concentrations. In a carp embryo study, DP increased mortality rate and morphological deformities, reduced hatching rate, delayed hatching time and decreased body length. In addition, neurodevelopmental toxicity was observed in fish embryos/larvae. DP has been shown to activate peroxisome proliferator activator receptor-gamma and inhibit insulin signalling in adipocytes in vitro. Furthermore, at low doses, DP promoted adipose tissue dysfunction, alter insulin secretion in mice. In mice receiving high fat diet glucose intolerance was observed at low dose DP exposure. A few human epidemiology studies found associations between DP and the sex- and thyroid hormone pathways. DP has also been reported to cross the blood-brain barrier and to be maternally transferred to offspring in several species. A study in zebrafish embryos indicate synergistic effects on neurotoxicity parameters and elevated bioaccumulation of both compounds as a synergistic effect of co-exposure to DP and the 3-methyl phenanthrene.

13. DP is widely detected in the global environment, including in remote regions. It is transported to locations far from production sites and places of use. Available scientific data show that DP is persistent and bioaccumulative. Although data are limited and chronic toxicity studies are lacking, there are studies showing that DP has adverse effects on the environment and that it can be potentially toxic to mammals and humans. Based on evidence for persistence, bioaccumulation and adverse effects of DP observed in some organisms and its widespread occurrence in the global environment including at remote regions and mindful of Article 1 referring to the precautionary approach as set forth in Principle 15 of the Rio Declaration on Environment and Development and Article 8 para 7 (a) stating that a lack of full scientific certainty should not prevent the proposal from proceeding, it is concluded that DP is likely, as a result of its long-range environmental transport, to lead to significant adverse human health and/or environmental effects such that global action is warranted.

1. Introduction

14. In May 2019, Norway submitted a proposal to include the chlorinated flame retardant (FR) Dechlorane Plus (CAS No. 13560-89-9) and its *syn*- (CAS No. 135821-03-3) and *anti*- (CAS No. 135821-74-8) isomers in Annexes A, B and/or C to the Stockholm Convention on Persistent Organic Pollutants (POPs). The proposal (UNEP/POPS/POPRC.15/3) was submitted in accordance with Article 8 of the Convention and reviewed by the Persistent Organic Pollutants Review Committee (POPRC) at its fifteenth meeting in October 2019.

15. The draft risk profile was discussed by the POPs Review Committee at its sixteenth meeting in January 2021. In considering the sub-item, the Committee had before it a note by the Secretariat on the draft risk profile prepared by the intersessional working group (UNEP/POPS/POPRC.16/2) and notes by the Secretariat containing additional information on the draft risk profile (UNEP/POPS/POPRC.16/INF/14) and a compilation of comments and responses relating to the draft risk profile (UNEP/POPS/POPRC.16/INF/4).

1.1 Chemical Identity

16. The commercial mixture “Dechlorane Plus”™ is an additive FR that has been in use since the 1960s (Shen et al., 2011). For Dechlorane Plus the acronyms DP or DDC-CO are commonly used. In this risk profile, DP is used. When environmental levels are reported the abbreviation, DP, represents the sum of the *syn*- and *anti*-DP isomers.

17. The technical mixture Dechlorane Plus™ (CAS No. 13560-89-9) is a commercially available formulation that contains two stereoisomers, *syn*-DP (CAS No. 135821-03-3) and *anti*-DP (CAS No. 135821-74-8) in a ratio of about 1:3 or 25 % *syn*-DP and 75 % *anti*-DP (Sverko et al., 2011). According to the North American manufacturer OxyChem, their commercial product contains approximately 35% *syn*-DP and 65% *anti*-DP (OxyChem, 2013). The fraction of *syn*-DP (f_{syn}) values for commercial DP from the Anpon Electrochemical Co and OxyChem manufacturers are in the range 0.20-0.41 (Wang et al., 2010a).

18. The structural formula of DP and its two isomers is shown in Figure 1 below. The chemical identity, and the modelled and experimental physico-chemical properties of DP and its two isomers are listed in Tables 1 and 2, below.

Figure 1. Structural formula of DP and its two isomers




Table 1. The chemical identity of Dechlorane Plus, with its *syn*- and *anti*-isomers.

CAS number:	13560-89-9 (Dechlorane Plus™), 135821-03-3 (<i>syn</i> -DP), 135821-74-8 (<i>anti</i> -DP)
IUPAC name:	1,6,7,8,9,14,15,16,17,17,18,18-dodecachloropentacyclo-[12.2.1.1 ^{6,9} .0 ^{2,13} .0 ^{5,10}] octadeca-7,15-diene (Dechlorane Plus™), (1S,2S,5R,6R,9S,10S,13R,14R)-1,6,7,8,9,14,15,16,17,17,18,18-dodecachloropentacyclo[12.2.1.1 ^{6,9} .0 ^{2,13} .0 ^{5,10}]octadeca-7,15-diene (<i>syn</i> -DP), (1R,2R,5R,6R,9S,10S,13S,14S)-1,6,7,8,9,14,15,16,17,17,18,18-dodecachloropentacyclo[12.2.1.1 ^{6,9} .0 ^{2,13} .0 ^{5,10}]octadeca-7,15-diene (<i>anti</i> -DP)
EC number:	236-948-9
EC name:	1,6,7,8,9,14,15,16,17,17,18,18-Dodecachloropentacyclo[12.2.1.16,9,02,13,05,10] octadeca-7,15-diene
Molecular formula:	C ₁₈ H ₁₂ Cl ₁₂
Molecular weight:	653.73 g/mol
Synonyms:	Bis(hexachlorocyclopentadieno)cyclooctane; 1,2,3,4,7,8,9,10,13,13,14,14-Dodecachloro-1,4,4a,5,6,6a,7,10,10a,11,12,12a-dodecahydro-1,4:7,10-dimethanodibenzo[<i>a,e</i>]cyclooctene; 1,4:7,10-Dimethanodibenzo[<i>a,e</i>]cyclooctene 1,2,3,4,7,8,9,10,13,13,14,14-dodecachloro-1,4,4a,5,6,6a,7,10,10a,11,12,12a-dodecahydro-; Dodecachlorodimethanodibenzocyclooctane; Dodecachlorododecahydrodimethanodibenzocyclooctane; Dodecachlorododecahydrodimethanodibenzocyclooctene; DDCDiMeDiBzcOb

Trade names	Dechlorane Plus; Dechlorane Plus 25 (Dech Plus); Dechlorane Plus 35 (Dech Plus-2); Dechlorane Plus 515; Dechlorane 605; Dechlorane Plus 1000; Dechlorane Plus 2520; Dechlorane A; DP; Escapeflam DK-15 (China); PyroVex SG (grade 515, 25 and 35)
-------------	---

Table 2. Overview of selected physicochemical properties of Dechlorane Plus, with its *syn*- and *anti*-isomers

Property	Value	Reference
Physical state at 20 °C and at 101.3 kPa	Solid white powder	ECHA, 2017b
Melting/freezing point	340-382 °C 350 °C	ECHA, 2017b OxyChem, 2004b
Vapour pressure	0.006 mm Hg \pm 0.8 Pa (at 200 °C)	OxyChem, 2004b
Water solubility*	<1.67 ng/L (20 – 25 °C) 0.044 – 249 μ g/L (insoluble)	ECHA, 2017b OxyChem, 2004b
n-Octanol/water partition coefficient, K_{ow} (log value)	9.3	OxyChem, 2004b
n-Octanol-air partition coefficient K_{oa} (log value)	12.26	OxyChem, 2004b
Sediment/water partition coefficient K_p (log value)	6.65	OxyChem, 2004b
Air-water partition coefficient K_{aw} (log value)	The following log K_{AW} values are estimated at 25 °C -3.2 (from measured water solubility and estimated vapour pressure) 0.44 (from measured water solubility and vapour pressure) -2.8 (from EPIWIN **predicted water solubility using log K_{ow} of 9 and vapour pressure) -3.5 (from HENRYWIN***v.3.20, predicted from structure using Bond Method).	ECHA, 2017b

*There is some uncertainty in the precise value for water solubility (Chou et al. (1979a). However, all available measurements and predictions agree that the substance is very poorly water soluble.

The following modelling programs are individual models in EPI Suite™:

** Estimation Program Interface Suite for Windows

*** Calculates the Henry's Law constant (air/water partition coefficient) using both the group contribution and the bond contribution methods. Since the group contribution method is not applicable to DP due to fragments missing in the database, only the result of the bond contribution method is shown.

19. DP is produced by the Diels-Alder reaction of two moles of hexachlorocyclopentadiene with one mole of 1,5-cyclooctadiene. DP monoadduct (DPMA) may be formed through partial reactions during the synthesis of DP through the diadduct Diels-Alder process and can therefore be present as an impurity in commercial DP (Sverko et al., 2010). DP compounds with a lower number of chlorine atoms such as mono-dechlorinated DP (DP-1Cl or DP-Cl₁₁) may also occur as impurities in the commercial substance (Li et al. 2013b).

20. Quantitative structure-activity relationship (QSAR) modelling (Sverko et al., 2011; Feo et al., 2012) and testing done by the manufacturer (EHSI, 2004) indicate that DP has characteristics typical of POPs (EHSI, 2004; Sverko et al., 2011; Feo et al., 2012). Physico-chemical properties of DP and analogue substances listed in the Convention are given in Tables 1 and 2 in the UNEP/POPS/POPRC.17/INF/9.

1.2 Conclusion of the POPs Review Committee regarding Annex D and E information

21. At its fifteenth meeting, the POPs Review Committee evaluated the proposal by Norway to list DP (CAS No. 13560-89-9) and its *syn*-isomer (CAS No. 135821-03-3) and *anti*-isomer (CAS No. 135821-74-8) under the Stockholm Convention on Persistent Organic Pollutants and concluded that DP and its *syn*-isomer and *anti*-isomer met the screening criteria specified in Annex D (UNEP/POPS/POPRC.15/7, Decision POPRC-15/4). The Committee decided to review the proposal further and to prepare a draft risk profile in accordance with Annex E to the Convention.

22. The POPs Review Committee considered the draft risk profile at its sixteenth meeting and adopted decision POPRC-16/1, by which it decided to defer its decision on the draft risk profile (UNEP/POPS/POPRC.16/INF/19) to its seventeenth meeting. In its decision, the Committee noted that, while information on persistence, bioaccumulation and the potential for long-range environmental transport was conclusive, the Committee had been unable to reach agreement that the information on adverse effects was sufficient to reach a conclusion on the risk profile. The

Committee also decided to establish an intersessional working group to review and update the draft risk profile; and invited Parties and observers to submit to the Secretariat additional information relating to the adverse effects of Dechlorane Plus before 1 March 2021.

1.3 Data sources

23. The revised draft risk profile is based on the following data sources:

- (a) The proposal submitted by Norway (UNEP/POPS/POPRC.15/3);
- (b) The draft risk profile prepared by the working group (UNEP/POPS/POPRC.16/2) and the outcome of the POPRC-16 (UNEP/POPS/POPRC.16/INF/19);
- (c) Annex E information and comments by Parties and Observers received in response to the invitation for comments on the draft risk profile. Annex E information was provided by: Australia, Belarus, Canada, Colombia, Costa Rica, Egypt, Federal State Autonomous Institute Research Institution 'Environmental Industrial Policy Center (EIPC), Germany, Hungary, Mexico, Monaco, Netherlands, New-Zealand, Peru, Romania, Republic of Korea, Thailand, State of Palestine, Qatar, International Pollutants Elimination Network (IPEN) and Alaska Community Action on Toxics (ACAT). In addition, information was provided by the Society of Indian Automobile Manufacturers (SIAM) and the Aerospace Industries Association of Canada (AIAC);
- (d) Reports and other grey literature as well as information from peer-reviewed scientific journals;
- (e) Documents supporting the identification of DP as a Substance of Very High Concern (SVHC) under the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) regulation in the European Union (EU) (ECHA, 2017b);
- (f) Screening assessment report and risk management approach from Canada (Canada, 2019a,b).

1.4 Status of the chemical under international conventions

24. DP is currently not known to be included in any international Conventions; however as described below and in Table 3 of UNEP/POPS/POPRC.17/INF/9 regulatory processes have been initiated or are in place in several countries including Canada, Egypt, EU and its member states and Norway. In addition, certain limitations and approval, reporting and/or notification requirements for production, import and/or use apply in New Zealand, Thailand,) The Russian Federation and the United States.

25. DP is listed on Canada's Domestic Substances List (DSL) (ECCC, 2019). A final screening assessment of DP was published by Environment and Climate Change Canada (ECCC) and Health Canada in spring 2019 (Canada, 2019a). The assessment concludes that DP meets the criteria for toxicity to the environment, as it is entering or may enter the environment in a quantity or concentration or under conditions that have or may have an immediate or long-term harmful effect on the environment or its biological diversity. The proposed regulatory approach is to amend the *Prohibition of Certain Toxic Substances Regulations, 2012* to prohibit the manufacture, import, use, sale or offer for sale of DP and all products containing the substance (Canada, 2019b).

26. In the EU, based on an Annex XV dossier and a Risk Management Options Analysis prepared by the United Kingdom, DP (including its *syn*- and *anti*-isomers) was identified as a SVHC and added to the REACH Candidate List in January 2018 as it is considered to be very persistent and very bioaccumulative (ECHA, 2017a).

27. In Norway, DP was added to the list of priority substances in January 2019 with a national goal to phase out the use by 2020 (Norwegian Environment Agency, 2019b).

28. Egypt reports that DP is banned by Decree No. 55 year 1996 of the Minister of Trade.

29. According to EIPC, DP is on the "Unified List of Chemical Substances" and its presence in this register provides evidence that DP is used but not produced in the Russian Federation national territory (Annex E, 2021). Information on importers, REACH Only Representatives and tonnages is confidential.

2. Summary information relevant to the risk profile

2.1 Sources

2.1.1 Production, trade, stockpiles

30. DP has been manufactured by Hooker Chemicals and Plastics Corporation (now Occidental Chemical Company (OxyChem)) in Niagara Falls, New York in the US since the 1960s (Shen et al., 2011). DP has been classified as a high production volume chemical (>500 tonnes) in the US (Sverko et al., 2011), and the annual production was estimated to be 450–5000 tonnes since 1986 (Hoh et al., 2006), although data on production was

withheld for some years as confidential business information. Available information suggests that production at the Oxychem facility ceased in mid-2016, but this has not been formally confirmed by the US EPA (ECHA 2021a, Annex E info IPEN, 2021). Manufacturing in China by Anpon Electrochemical Co. in Huai'an has been reported to be 300–1000 tonnes per year since 2003 (Wang et al., 2010a). The resulting global annual production of DP is thus 750–6000 tonnes with an average of 1980 t per year (Hansen et al. 2020). Ren et al. (2008) indicated that the global annual production was approximately 5000 tonnes. Anpon Electrochemical Company Ltd. was recently acquired by ADAMA, which may now be the sole global manufacturer (ADAMA, 2019).

31. DP is available for sale from several vendors globally (PubChem, 2021, see also Annex E, Peru). In the EU, DP has been registered under the REACH Regulation by two registrants (one for each manufacturer) and is manufactured in and / or imported to the European Economic Area, at ≥ 10 to < 100 tonnes per annum (ECHA, 2021a). The major registrant since 2013, Occidental Chemical Belgium BVBA, representing the manufacturer OxyChem US, submitted a notification in December 2017 of "cease of manufacture or import" under REACH and no longer produces or imports DP to the EU (ECHA, 2021a). The Dutch company ADAMA Agriculture BV has, since 2016, been the registrant for the sole importer/distributor Velsicol, who represents Anpon Electrochemical Co. The maximum volume imported in the EU was 300-400 tonnes/year. Sweden registered use of 11 tonnes of DP in 2005 and 5 tonnes in 2006 (Kaj et al., 2010). Hungary reports that DP is used by two Hungarian companies at volumes below 2 tonnes/ year (Annex E, 2021).

32. In Canada, based on surveys conducted under section 71 of CEPA for the years 2011 and 2016, between 1000 and 10 000 kg of DP, including DP in some products, was imported by a few companies (Canada, 2019a,b). Based on information gathered from the Statistical Survey of Chemicals under the Chemicals Control Act, no DP was manufactured in the Republic of Korea since 2010. No DP was imported in 2010, but use was around 40 tonnes. In 2016, 11 tonnes were imported, and 17 kg was used. DP has not been exported since 2010, when 0.9 tonnes of DP was exported. In 2018, 0.63 tonnes of DP were imported, and 0.39 tonnes of DP were used (Annex E information, Republic of Korea).

33. Available information also suggests ongoing use in the Russian Federation (Annex E, 2021).

2.1.2 Uses

34. DP is currently marketed as an alternative/replacement for commercial decabromodiphenyl ether (c-decaBDE) and as a substitute for mirex used as a FR (Hoh et al., 2006). Known uses include electrical and electronic equipment and devices, optical products, food packaging, building and construction materials, tapes and adhesives, fabrics, textiles, apparel, cosmetics, explosives/ fireworks, plastic products, paints and greases (ECHA, 2021a, see Table 4-6 in UNEP/POPS/POPRC.17/INF/9). EPIC also reports that DP is used as a pesticide (Annex E, 2021). Known FR uses include electronic wiring and cables, automobiles, plastic roofing materials, and hard plastic connectors in televisions and computer monitors, wire coatings, and furniture (Zhang et al., 2015, Annex E, 2019, 2020). It is also used extensively in the aerospace industry as a FR and in other applications (AIAC, 2020). DP is often used as an additive to various formulations and can be found in many aircraft components such as cabin interior panels, ducting, engines and other components. In a recent study from China, DP was detected in various building materials (Hou et al., 2018). Samples of wallpaper (non-woven, PVC, paper), latex paint, boards (e.g. laminated floor, fiber board, solid wood), glue, sealant, PVC line pipes and foam (such as sound absorbing foam and expanded polystyrene panels) contained DP in the range non-detect to 5.1 ng/g (Hou et al., 2018). The Society of Indian Automobile Manufacturers (SIAM), on behalf of their members, state that dechlorane plus is used as an alternative to c-decaBDE in automobile manufacturing (SIAM, 2020). In Canada, DP is reported to be used as a FR in automobile manufacturing and DP is known or expected to be used as a FR in wiring and cable jacketing, electronics, roofing materials, and hard plastic connectors (Canada, 2019b). Annex E information from the Netherlands confirms use in automobiles in the powertrain, cooling, chassis and bodywork parts (Annex E, 2019). According to data from the EU DP the most important sectors using DP are the automotive and the aviation sectors which accounts for 57-75 % and 10 % of all use in the EU, while other uses, including in computers, electronics and imported articles etc. account for 33-15% of all use in the EU (ECHA, 2021a, see Table 7-8 in UNEP/POPS/POPRC.17/INF/9).

35. DP is used in many polymeric systems as a FR. These systems are typically either thermoplastics or thermosets. Examples of thermoplastics that may contain DP include nylon (Weil and Levchik 2009; KEMI, 2019), polyester (KEMI, 2019), acrylonitrile butadiene styrene (ABS), natural rubber, polybutylene terephthalate (PBT), polypropylene, and styrene butadiene rubber (SBR) block co-polymer (OxyChem, 2007). DP may be used in thermosets such as epoxy and polyester resins, polyurethane foam, polyethylene, ethylene propylene diene monomer rubber, polyurethane rubber, silicon rubber, and neoprene (OxyChem, 2007). The amount of DP in these materials ranges from 8% in PBT to up to 40% in silicon rubber (OxyChem, 2007).

36. According to one manufacturer (OxyChem, 2007), DP is manufactured for use solely by industrial customers. However, EU data indicate that DP is commonly used in the manufacturing of both professional and consumer products. According to REACH registration data, uses of DP in the EU/ EEA region include uses in formulations or re-packing, at industrial sites and by professional workers. DP is also contained in some articles used by consumers

(ECHA, 2021a). Under REACH, DP is registered for industrial use in the following product categories; polymer preparations and compounds, semiconductors, adhesives and sealants. Registered end uses from formulation and packaging include: 1) Adhesives, sealants, 2) Polymer preparations and compounds and 3) Semiconductors. Registered end uses at industrial sites and by professional workers include: 1) Mixture of preparations and/or re-packaging (excluding alloys), 2) Manufacture of plastics products, including compounding and conversion and 3) Manufacture of computer, electronic and optical products, electrical equipment. In the service life of articles at consumer use stage use in the following article categories have been indicated in the registration dossier: 1) vehicles, 2) machinery, mechanical appliances, 3) electrical/electronic article, 4) fabrics, textiles and apparel and 5) plastic articles. In addition, use in glued articles in aircraft parts are indicated. A minor use of DP is in the development of fireworks, but this is not a registered use in the EU/EEA.

37. In 2011, a 220-tonne sample of representatively composed e-waste was processed in a Swiss recycling facility with the different output streams being analysed. The highest DP mass flows were in fine-grained plastics and in plastics from consumer products such as monitors and laptop casings (Taverna et al., 2017).

38. DP was detected in plastic casings from two TVs and one computer, but not in circuit boards from similar equipment collected from a recycling facility in China (Li et al., 2019c). In addition, DP was detected in recycled ABS from the Guangdong Province in China (Cao et al., 2020).

39. PyroVex® SG is an additive FR that, according to the supplier, contains > 99% DP (Velsicol Chemicals LLC, 2020). PyroVex® SG is typically mixed into plastics and other materials and can be used alone or be formulated with complementary chemistries to meet performance requirements. It is marketed for use in flame retarded wire and cable, and as a FR for electrical connectors used in computers and other electronic devices. PyroVex® SG grade 515, 25 and 35 is the same chemical mixture only with different average particle size.

40. During a public EU consultation on the ninth draft recommendation for inclusion of DP in Annex XIV of REACH, held from September to 5 December 2018, a number of companies identified applications in which they use DP as an additive FR. Identified applications were: in the aviation and automotive industries, in polymeric compounds for cable insulation and minor use in fireworks (ECHA, 2021a).

2.1.3 Releases and emissions to the environment

41. DP and its isomers are not known to be unintentionally produced. The occurrence of DP and its isomers in the environment is a result of human activities.

42. A number of activities such as production, use, recycling and waste handling, as well as landfill leachate and run-off and wastewater treatment, can lead to releases of DP to the environment (Wang et al., 2016; Wang and Kelly, 2017). DP from industrial sources is expected to be released primarily to wastewater (a pathway to surface water and the soil environment). DP may also to some extent be released from commercial- and consumer products to the atmosphere. DP is likely highly removed by adsorption to biosolids in wastewater treatment systems (CECBP 2008 as cited in Canada, 2019a).

43. DP has been detected in sludge (e.g. de la Torre et al., 2011; Barón et al., 2014a; Norwegian Environment Agency, 2018b; 2019a; Ji et al., 2018) and biosolids (Davis et al., 2012) from wastewater treatment plants (WWTPs). In addition, sludge from WWTPs contaminated with DP can be used as soil enrichment (Wu et al., 2017) or fertilizer (Dai, 2011 reviewed in Ji et al., 2018). Furthermore, *syn*- and *anti*-DP was detected in the particulate fraction of storm water in Oslo, Norway (Norwegian Environment Agency, 2018b; 2019a).

44. In samples from Canadian WWTPs investigated for chlorinated FRs, the detection frequency of DP was greater than 90% in both wastewater and biosolids samples, while median removal efficiencies were between 51% and 66% for total DP (Shanmuganathan et al., 2018). In a conventional WWTP in Shanghai DP levels were detected in wastewater with a mean concentration of 0.46 ng/L (range: 0.05 to 1.40 ng/L). Sludge contained DP in the range of 1.1 to 2.0 ng/g. The fraction of *anti*-DP (f_{anti}) was consistently higher than that of *syn*-DP (f_{syn}). Average f_{syn} values ranged from 0.16 to 0.33 and was in the range of two commercial DP mixtures. The annual release of DP via sewage sludge from all WWTPs in Shanghai were estimated to be 164.8 g (Xiang et al., 2014).

45. DP is used as a FR in electrical and electronic equipment and recycling of e-waste is reported to be a source of DP release to the environment (Wang et al., 2016). Very high levels of DP in soil (3327 ng/g) were detected at a recycling site in China suggesting that e-waste recycling activities serve as an important source of DP emissions in China (Yu et al., 2010). In a Swiss study, the overall DP concentration in e-waste was 33 ± 11 mg/kg. This translates into an annual DP mass flow of 2.3 ± 0.9 tonnes/year in all Swiss e-waste generated in 2011 (Taverna et al., 2017).

46. Indoor dust is the sink and carrier of many pollutants including DP released from consumer products. Indoor dust is also considered to be the main source of pollutants in WWTPs receiving only domestic wastewater (Katsoyiannis and Samara, 2004; Xiang et al., 2013). DP has been detected in indoor dust in the range 0.08 to 124 000 ng/g (Newton et al., 2015; Wong et al., 2017, Cao et al., 2014, Lee et al., 2020; Corsolini et al., 2021). For additional

information on releases and emissions from manufacture, WWTPs, e-waste recycling, landfills etc., and indoor dust and air see Table 10 and 16, respectively, in UNEP/POPS/POPRC.17/INF/9 and section 2.3.2 on human exposure.

47. Based on production, usage and disposal data Hansen et al. (2020) prepared two global atmospheric emission scenarios with a detailed geographical distribution. The total DP emission to air was estimated to be 0.02 t/year and 3.2 t/year in a low and high emission scenario, respectively, reflecting the uncertainties in production volumes and emission factors. Furthermore, 0.41-21.8 tonnes/year were predicted to be released to water, and 0-0.004 tonnes/year direct to soil. The validity of the emission estimates was tested by implementing them in the Danish Eulerian Hemispheric Model, an advanced chemistry-transport model, and by evaluating the predicted atmospheric concentrations with all available Arctic measurements from peer-reviewed studies. The high emission estimate was found to be the most probable as the predicted concentration range for the high emission scenario was found to be in line with the measured range, whereas the predicted concentrations for the low emission estimate was more than a factor of 100 lower than the measurements. The estimates reported in this study were based on a global annual production of 75-6,000 tonnes/year (average assumed to be 1,980 tonnes/year) and with two manufacturing plants (one in the United States and one in China).

2.2. Environmental fate

48. The environmental distribution and fate of DP is discussed in Canada (2019a) and ECHA (2017b). In the assessment by Canada (2019a), level III fugacity modelling using the updated EQC model (v1.0, 2012) was applied to describe the fate of DP entering the environment from industrial sources and from commercial and consumer products. The results show that DP is expected to predominantly reside in soil and/or sediment, depending on the compartment of releases.

49. The Canadian assessment further details that the very low water solubility (2.85×10^{-7} mg/L), low vapour pressure (6.57×10^{-11} Pa at 25°C) and very high partition coefficients (log K_{ow} of 8.78, estimated log K_{oc} of 6.65) suggest that DP released into the environment will be less likely to partition into and/or remain in air and water, moving instead to the sediments and soil. If released to air, a small fraction (less than 1%) of DP is expected to remain in air in gas phase, with most of the substance depositing to soil and water with further partitioning to sediment. The high partition coefficients also indicate that DP released into surface water from wastewater is expected to adsorb to the organic fraction of suspended solids and sediments, with less than 4% remaining in water. However, the small mass of DP remaining in air and in water has the potential for dispersion and some transport (e.g., particle transport). On the basis of its high log K_{oc}, once in the sediment, DP is not expected to be mobile and may remain in this compartment with little degradation. When DP is released to soil (i.e., through biosolids application to agricultural lands), the majority of the mass fraction is expected to become adsorbed to soil (99.9%) on the basis of its high estimated log K_{oc} and hydrophobic nature. On the basis of its low vapour pressure, DP is not expected to evaporate (volatilize) from dry soil surfaces and is therefore likely to remain in soil. In addition, low degradation is expected in soil; therefore, DP is likely to remain in this compartment, with loss processes driven by soil burial or surface runoff. The results of Level III fugacity modelling support the expectation that DP predominantly resides in soil or sediment, depending on the compartment of release.

2.2.1 Persistence

50. DP is chemically stable in various environmental compartments with minimal or no abiotic degradation (reviewed in Wang et al., 2016, ECHA, 2017b, Canada, 2019a). Due to a very low water solubility and high log K_{ow}, DP is expected to bind to organic carbon in soil and sediments (Wang et al., 2016), reducing the bioavailability of DP for microorganisms and hence the potential for biodegradation. There are no measured half-life data for degradation of DP in surface water, sediment or soil (ECHA, 2017b). However, physical-chemical properties of DP were predicted using three different models (EPISuite, SPARC and Absolv) and the estimated half-lives in water, soil and sediment were predicted to be 180, 3650 and 1621 days, respectively (Zhang et al., 2016).

51. DP does not contain any functional groups that are susceptible to hydrolysis and hydrolysis is not expected to be a relevant degradation process (Canada, 2019a; ECHA, 2017b). Data on photodegradation of DP in air (e.g. Sverko et al., 2008; Wang et al., 2011; Li et al., 2013b; Wang et al., 2013b; Tao et al., 2015), water (Chou et al., 1979b) and soil (only modelled data), are reviewed in ECHA (2017b) and Canada (2019a). Most of the experimental data are from controlled lab studies that cannot be directly related to natural conditions due e.g. to the use of solvents and differences in light intensities.

52. The studies on photo-degradation suggest that *anti*-DP might be more photodegradable in air than *syn*-DP. Furthermore, DP is considered to be relatively photo-stable in air under natural conditions although photodegradation and formation of lower chlorinated dechloranes and other degradation products are reported in some of the studies. Under natural conditions sorption of DP to airborne particles is expected, which would lower the photolysis rate and result in a longer half-life in air (Canada, 2019a). Furthermore, the study of Wang et al. (2013b) indicates that DP should be photo-stable in the environment and only undergo limited degradation under natural light conditions at the terrestrial surface. Wang et al. (2013b), who investigated the photodegradation of DP (*anti*-DP,

syn-DP, and commercial DP) in *n*-nonane by irradiation using a xenon lamp, observed rapid photodegradation by dechlorination under 200–750 nm light. During the first 5 minutes >50% was degraded, and after 30 minutes 99% had been degraded. The degradation was much slower in 280–750 nm light, and only 20% loss was observed after 4 hours. The quantum yields of dechlorination by-products at 200–280 nm (UV-C) were about 2–3 orders of magnitude higher than at 280–320 nm, and no yields were detected in the 320–750 nm range (Wang et al., 2013b). Hence, both in visible light (>400 nm) and UV-A light (320–400 nm), DP exhibited negligible degradation in *n*-nonane, while UV-C (200–280 nm) played a significant role in the photodegradation of DP and might provide a potentially effective approach to eliminate DP. As the majority of UV exposure in the environment is UV-A light with a wavelength in the 315–400 nm range, these findings indicate that DP should be photo-stable in the environment and only undergo limited degradation under natural light conditions at the terrestrial surface. A sequential degradation process where one to four chlorines are removed from DP and exchanged with hydrogen was observed and suggested that the main overall reaction was photoreduction, leading to photodechlorination of DP (Wang et al., 2013b). Dechlorination was also observed by Li et al. (2013b). In this study, three photolytic degradation experiments were performed by exposing solutions of *anti*-DP, *syn*-DP, and commercial DP to UV light. In addition to *anti*- and *syn*-DP-Cl₁₁, at least two unknown products were identified in all samples following exposure to UV-light, and in the test substance (Li et al., 2013b).

53. According to ECHA (2017b) photolysis is unlikely to be a significant fate process in natural waters, since light is attenuated with increasing water depth and shading. Radical reactions may also be inhibited by humic substances. They conclude that available information suggests that phototransformation in water is a potential but insignificant removal process for DP. The only controlled laboratory study on photo-degradation in water is by Chou et al. (1979b) who measured the phototransformation of DP in an aqueous solution with 5 % acetonitrile using a mercury lamp emitting wavelengths >290 nm for 168 hours. In Canada (2019a) a half-life for DP of >24 years is reported for this study, while ECHA (2017b) report that the rate constant for phototransformation in water was calculated as $<6 \times 10^{-4}$ /h, equivalent to a half-life of >48 days (Chou et al., 1979b). According to ECHA (2017b) the findings from this study are not directly representative of natural conditions as the test solution included a solvent and the initial concentration of DP was 1 ppm (1 mg/L), which is significantly greater than the reported solubility of DP in pure water. For phototransformation in soil ECHA (2017b) reports that no data were reported by the registrant, and that similar to water, this is unlikely to be a significant removal pathway.

54. Generally, model predictions support experimental findings that aerobic and anaerobic biodegradation of DP is very limited and that DP is expected to be persistent in water, soil, and sediment (Canada, 2019a). Predictions using the BIOWIN v4.10 model (US EPA, 2012), Catalogic (2012) and TOPKAT (2004) all indicate that biodegradation of DP will be very slow (Canada, 2019a; ECHA, 2017b). ECHA (2017b) concludes, based on the estimated data from the BIOWIN v4.10 model and associated uncertainties, that "DP is unlikely to be biodegradable". The BIOWIN results obtained for DP are the same as a number of hexachloro-norbornene-containing analogue chemicals and indicate that DP is as persistent as analogue chemicals such as mirex, chlordane, heptachlor, dieldrin/endrin, aldrin and endosulfan that are already listed as POPs under the Stockholm Convention. Furthermore, the two possible microbial degradation pathways predicted for DP are the same as for the POP analogues. According to ECHA (2017b) it is unlikely that metabolic rates for these pathways will be more rapid for DP, which is significantly less water soluble than the already listed POPs. The validity of the BIOWIN v4.10 model is discussed in detail in ECHA (2017b). Despite some uncertainties BIOWIN v4.10 is generally considered as a valid model for DP.

55. In line with these modelling results, laboratory tests discussed in Canada (2019a) have shown that DP is not likely to biodegrade under aerobic conditions; an activated sludge biodegradation test (modified MITI OECD 301C) reported 0.6% biodegradation in 2 weeks (US EPA, 2011; see also Japan J-CHECK, 2020a) and a 21-day test using wastewater biosolids found 0% biodegradation (US EPA, 2009).

56. Monitoring studies provide additional evidence that DP is persistent and that biodegradation in sediment is limited. DP was found to be persistent in suspended sediment with a half-life of approximately 17 years (Sverko et al., 2008). The study of Qiu et al. (2007) suggests that DP can still be present over thirty years after initial deposition. In this study, DP was measured in a sediment core from central Lake Ontario, Canada, and detected in a layer corresponding to around 1980. A linear trend ($r_2 = 0.739$) of increasing f_{anti} values with time was observed, from an average of 0.76 in surficial (recent) sediments to >0.90 in the deeper layers corresponding to around 1980, suggesting that the *anti*-DP could be more persistent than the *syn*- isomer in sediment. However, the variation of f_{anti} in commercial batches over this time period is not known, and the findings could also reflect the isomerisation of *syn*-DP to *anti*-DP. Similar to the findings of Qiu et al. (2007) other studies on DP and f_{anti} values in freshwater (Wang et al., 2010) and marine sediment (Fang et al., 2014), respectively report possible stereoselective depletion of *syn*-DP in freshwater sediment and an enrichment of *anti*-DP. The enrichment of *anti*-DP is explained as likely being due to preferential biodegradation of the *syn*- isomer in the sediment.

57. Some data on the degradation of DP in soil are also available. In a controlled laboratory study that was not considered in the Canadian or the EU assessment (Canada 2019a; ECHA, 2017b), degradation of DP in top layer agricultural soil with a pH of 7.1 and consisting of 64.2% silt, 25.6% sand, 10.2% clay, and 3.12% organic matter, was investigated by spiking the soil with DP in acetone to obtain initial DP concentrations of 0.1, 1, and 10 mg/kg.

After 260 days of incubation at 25 °C 4.2-8.2% of the initial DP had degraded (Cheng et al., 2019). During the same time the labile fraction of DP decreased from 21.25% to 8.2%, 23.8% to 11.7%, and 25.2% to 16.3% in 0.1, 1, and 10 µg/g DP soil, respectively, indicating that the bioavailability of DP decreased under the influence of aging. Model simulations developed to investigate the transformation indicated that the transformation rates were inconstant and statistically different from one another over time. The half-lives of DP in soils spiked with 0.1, 1, and 10 µg/g DP were extrapolated to 1325, 1909, and 2948 days, respectively. As the authors considered that very little bound-residue DP is formed at the start of the aging process, the loss of DP to bound residues was neglected, and the half-lives was simplified to and calculated as the time required for the combined concentration of labile and stable-adsorbed fractions to reduce to half its initial value (Cheng et al., 2019). It has also been reported that *syn*-DP varied little with depth down to 100 cm in soil sampled close to the Chinese manufacturing facility (Wang et al., 2010a), possibly suggesting that isomer-specific degradation or preferential adsorption does not play a significant role in soil. The DP concentration was highest in surface soil (0-5 cm; 837 µg/kg dry weight (dw)), decreasing to 9.16 µg/kg dw at a depth of 60-70 cm and 3.84 µg/kg dw at a depth of 90-100 cm. The f_{anti} value varied little with soil depth but ranged from 0.75 in the surficial layer to 0.67 in the deepest layer (90-100 cm). The study authors make conflicting statements about their finding. On the one hand, they say their findings imply a stereoselective depletion of the *syn*- isomer in soil in comparison with the measured f_{anti} value of the Chinese commercial product, at 0.60. On the other hand, they also state that isomer specific microbial degradation or preferential adsorption does not play a significant role in soil.

58. DP is expected to be persistent in water, soil, and sediment. Based on its binding to particles in air and available laboratory studies and modelling data DP is also considered to be persistent in air under natural conditions. Modelling studies of degradation potential and microbial metabolic pathways suggests that biodegradation of DP is likely to be very slow, and that the probability that DP will degrade any faster than analogue substances listed under the Stockholm Convention is low. Monitoring data show limited degradation in soil and sediments over time and support the conclusion that DP is persistent. The persistency of DP in soil was also confirmed in a controlled laboratory study (Cheng et al., 2019).

2.2.2 Bioaccumulation and toxicokinetics

59. The log K_{ow} for DP is reported to be 9.3 (OxyChem, 2004b). The high log K_{ow} and the very low water solubility (Table 2) indicates that DP is very hydrophobic and partitions to organic matter. This is further supported by the relatively high log sediment-water partition coefficient (log K_p) of 6.65. These properties make it difficult to perform aqueous laboratory studies as it is difficult to maintain stable exposure levels in water. Therefore, aqueous exposure is expected to be of limited importance in terms of bioaccumulation potential. However, significant uptake has been shown to occur in fish in a number of field studies (Guo et al., 2017; Malak et al., 2018; Kurt-Karakus et al., 2019).

60. Although dietary exposure is expected to be more relevant for a substance with physiochemical properties as DP, one fish bioaccumulation study using aqueous exposure reported bioconcentration factor (BCF) values exceeding the bioconcentration criteria set by the Stockholm Convention (BCF >5000 L/kg) (Wang et al., 2019). In this study, carp were exposed continuously for 32 days to DP (range 0.14 to 0.24 ng/L) through water, followed by 32 days depuration. Dosing was carried out using four packages, each containing 1g commercial DP powder ($\geq 99\%$ purity) within filter paper enclosed in a filter screen and placed in a hollow steel ball in the bottom of the aquaria. The authors estimated that a steady state water concentration was reached within 3 days, and the equilibrium would subsequently be maintained. Maximum concentrations in muscle for *syn*- and *anti*-isomer were observed at day 32 with the concentrations of 440 ± 28 ng/g wet weight (ww) and 830 ± 44 ng/g ww, respectively. Equilibrium was reached and the BCF was calculated based on wet weight concentrations in muscle and ratio of uptake and elimination constants. The reported BCF values were 5700 and 9300 L/kg for *syn*- and *anti*-DP, respectively and the depuration half-life was 6.3 and 7.2 days for *syn*- and *anti*-DP, respectively (Wang et al., 2019). The information provided by J-Check English version, on a bioaccumulation test conducted in 1974 with carp exposed to water concentrations of DP at 2.7 and 0.27 µg/L for 8 weeks shows a BCF of 87-121 L/kg (Japan J-Check, 2020b). The concentrations were significantly in excess of the water solubility of the substance. As a consequence, the recorded values may be an underestimate of the true BCF values. However, as the form of exposure may have been a mix of oral and aqueous, this cannot be confirmed. No information on levels observed in fish was provided.

61. Depuration half-life is an important measure of bioaccumulation. A depuration half-life around 8-10 days is indicative of a lipid-normalised and growth-corrected BCF above 5 000 L/kg according to the UK Environment Agency (EA) (UK EA, 2012). In a dietary bioaccumulation laboratory test using juvenile rainbow trout, the depuration half-life (whole body minus liver) for DP was reported to be 30-40 days for the *anti*-isomer and 50-70 days for the *syn*-isomer, which is highly indicative of a very bioaccumulative substance (Tomy et al., 2008). Neither isomer reached steady state after 49 days of exposure (Tomy et al., 2008). Higher bioaccumulation of the *anti*-DP isomer was also observed in several aquatic food webs and organisms as reviewed in Wang et al., (2016). However, a laboratory study in carp showed tissue-specific variations (Tang et al., 2018). Furthermore, a comparison of modelled studies (Larisch and Goss, 2018) and measured oral uptake in fish (Tomy et al., 2008) showed good agreement, indicating that bioaccumulation of super hydrophobic chemicals like DP in fish via food can be predicted. Uptake of DP is slow

but will eventually result in substantial bioconcentration as the substance is not metabolised in fish (Larisch and Goss, 2018). Depuration half-life observed in Wang et al., (2019) for carp muscle was lower than what was observed for trout whole body minus liver in Tomy et al., (2008) and could be due to different fish species and experimental setups.

62. Bioaccumulation of *anti*-DP was observed in juvenile marine green macroalgae *U. pertusa*, following a 21-day exposure to 10^{-8} mol DP/L (equal to ~ 6.53 $\mu\text{g/L}$) and 21-day depuration (Gong et al., 2018). Accumulated DP levels were 127 and 206 ng/g ww on day 7 and 21, respectively, and the half-life was 1.46 and 14.5 days for *syn*- and *anti*-DP, respectively.
63. In rats exposed to commercial DP 25 by gavage for 90 days at different doses (0, 1, 10, and 100 mg/kg/d), DP preferentially accumulated in liver rather than muscle. At high doses (10 and 100 mg/kg/d) *syn*-DP were the predominant isomer in tissues. The elimination half-life for *syn*-DP was about 179 days in liver, 44 days in muscle and 24 days in serum, and for *anti*-DP, 54 days in muscle and 25 days in serum. Depuration time for *anti*-DP for the liver was not calculated due to a non-significant increase in liver after depuration (Li et al., 2013b). In toxicokinetic studies, after single exposure, oral absorption is thought to be limited in rats which may lead to a maximum absorption of 5% to 20% at 1 mg/kg bw into systemic circulation (Brock et al. 2010, ECHA 2017b).
64. In birds, studies indicate that DP is bioavailable and transferred to eggs (e.g. Li et al., 2021b; Smithe et al., 2020). In paired samples of eggs and plasma from bald eagles from Canada, DP was more abundant in eggs compared to plasma with geometric mean of 0.28 and 0.43 ng/g ww in eggs from inland and Great Lakes, respectively, versus plasma levels of 0.02 ng/g ww (Guo et al., 2018). In a lab study oral absorption of DP in hen was around 65% (Li et al., 2021b).
65. Several field studies report biomagnification factors (BMF) and trophic magnification factors (TMF) for DP in various food webs. Factors, such as, temperature, time of sampling, reproduction status, migration, age and tissue versus whole body calculations may affect the calculation of TMF (Borgå et al., 2012; Franklin, 2016). DP biomagnifies in freshwater, marine and terrestrial food webs, as BMFs (Tomy et al., 2007; Wu et al., 2018; Sun et al., 2015; 2017, Chen et al., 2021) and TMFs (Sun et al., 2015; Kurt-Karakus et al., 2019; Na et al., 2017) are reported to be greater than 1 in several studies, including in one out of two studies from Antarctica (Na et al., 2017; Kim et al., 2021). For prairie animals in the remote Xilingol Prairie in Inner Mongolia in China calculated BMFs for ectotherms and endotherms were in the range 0.146–88.0 and 0.866–17.2, respectively (Chen et al., 2021). Ectotherms were found to selectively enrich *syn*-DP, and stereoselective enrichment increased moving up the food web. The highest DP BMF was for lizards (88.0), which eat insects (Chen et al., 2021). Sample numbers were low in this study and therefore there is some uncertainty in the derived values. However, also other studies indicate bioaccumulation of DP in terrestrial food webs. In aquatic and terrestrial food webs of several insect-dominated food chains, including insects – toads, insects – lizards, and insects- bird in or around a pond at an e-waste recycling site, TMF for DP was <1 in the aquatic food web and >1 in the terrestrial food web (Liu et al., 2020). BMFs for Σ DP in four insectivorous birds range from 9.5 to 23 (Liu et al., 2020). Bioaccumulation and maternal transfer have been observed in amphibians collected from a highly contaminated site in South China (Wu et al., 2018). Amphibians occupy an important trophic position in the food web between aquatic organisms and terrestrial biota. Based on the known predator-prey relationship between frog and insects, BMFs above 1 (1.8 – 2.7) for DP were reported in this study (Wu et al., 2018). In a study of biomagnification of DP in a freshwater reservoir in the vicinity of an electronic recycling facility in South China, the TMFs of the DP isomers were found to be 2-3 times greater than those of PBDE congeners and comparable to or lower than those of the highly recalcitrant PCB congeners in the same food web (Wu et al. 2010). In an aquatic food web study from China, TMFs were reported in the range 1.9 – 3.1. In this study, the biomagnification potential (TMF >1) was also reported for DPMA, *anti*-Cl₁₁-DP and *anti*-Cl₁₀-DP (Wang et al., 2015), and this should also be taken into account when evaluating bioaccumulation for DP. For more information on these substances, see section 1.1 Chemical identity. Three terrestrial studies are available showing BMFs above 1 (Yu et al., 2013; She et al., 2013; Wu et al., 2018). Furthermore, in a study investigating levels of DP and other flame-retardants in livers of redlisted and endangered birds in Korea, Jin et al. (2016) found a significant positive relationship ($r^2= 0.22$ and $p=0.031$) between the concentrations of DP and trophic position ($\delta^{15}\text{N}$) in three residential and carnivorous predatory species, the Eurasian eagle owls (*B. bubo*), common kestrels (*F. tinnunculus*), and collared scops owls (*O. lempiji*), indicating for biomagnification of DP in these birds. For further details on the bioaccumulation studies, see Table 9 in UNEP/POPS/POPRC.17/INF/9.
66. The isomer composition of DP in environmental samples can be different from the technical products because of their biota isomer-selective uptake or elimination, bioaccumulation, and stereospecific photodegradation (Wang et al., 2015; and reviewed in Wang et al., 2016). Trophic levels (Peng et al., 2014; Tomy et al., 2007; Wang et al., 2015), exposure concentrations (Li et al., 2013a,b; Mo et al., 2013; Peng et al., 2015), life cyclic stages (Klosterhaus et al., 2012; Sührling et al., 2014), the type of wildlife and tissue (Peng et al., 2014; Zhang et al., 2011; Zheng et al., 2014a) and stereoselective excretion (Li et al., 2019) are the main factors for the stereoselective bioaccumulation of DP in organisms.

67. Studies in birds indicate that uptake, distribution and elimination kinetics of DP and its isomers are complex. Chickens bred in the vicinity of an e-waste recycling facility exposed to DP through the environment from sand and food showed preferential accumulation of the *anti*-DP isomer (Zheng et al., 2014a). Lipid content of the tissue was the main factor in the tissue distribution, although the degree of blood perfusion and the tissue function were also important factors. Furthermore, the study revealed tissue specific accumulation of *anti*-DP, with higher f_{anti} levels in brain, fat and liver (0.65-0.64) compared to (0.54-0.59) for other tissues. The study also indicate that *anti*-Cl₁₁-DP was absorbed through the diet rather than metabolised from DP *in vivo* based on the similar ratios for *anti*-Cl₁₁-DP to *anti*-DP in soil and chicken (Zheng et al., 2014a). This information is supported by the fact that *syn*-Cl₁₁-DP and *anti*-Cl₁₁-DP were detected in commercial DP-25, hence, it can be inferred that these chemicals originated from commercial products (Li et al., 2013b).

68. There are some studies on biotransformation of DP (Tomy et al., 2007; 2008; Ren et al., 2009; Sverko et al., 2010). DP has been shown not to metabolise easily in biota (Tomy et al., 2008; Xian et al., 2011) and the results from Tomy et al. suggest that enzyme-induced metabolism of DP in fish might be low, if it does occur. No hydroxylated or sulfonated metabolites of DP was observed in fish liver extracts or in human serum (Tomy et al., 2007; Ren et al., 2009). Degradation products of DP such as decachloropentacyclooctadiene (DP-Cl₁₀) or undecachloropentacyclooctadiene (DP-Cl₁₁) have been detected in bird eggs (Guerra et al., 2011; Muñoz-Arnanz et al., 2011, 2012; Zheng et al., 2014a), and rat (Li et al., 2013) but some studies suggest they are formed through biotic or abiotic processes prior to uptake or even through impurities in the commercial product (Sverko et al., 2008, 2010; Tomy et al., 2008; Zheng et al., 2010, 2014b; Li et al., 2013). Two additional dechlorinated analogues were also detected in both quails and the test substance (Li et al., 2013a,b). In a study where DP was injected into the yolk sacs of eggs from Japanese quail, no biotransformation of DP was shown (Briels et al., 2018). This was also observed in chicken eggs at day 18, however, at pipping the mass of *syn*- and *anti*-DP in neonate chicks (including the remaining yolks) declined significantly by 5.9% and 15%, respectively, indicating some metabolic activity during the later developmental stage (Li et al., 2019).

69. DP was first detected in archived fish (walleye) from Lake Erie sampled from 1980-2000, which suggested that DP was taken up by the fish (Hoh et al., 2006). Available monitoring studies show that DP is widely dispersed in the environment (reviewed in Canada, 2019a; ECHA 2017b). Global evidence reveals that uptake of DP can occur in various biota samples within aquatic and terrestrial food webs as well as the human body (reviewed in Wang et al., 2016). Field monitoring data suggest that DP is bioavailable and can exceed levels in biota that are of concern based on critical body burden considerations related to baseline narcosis (ECHA 2017b).

70. DP was detected in human milk samples collected in two Canadian cities (Siddique et al., 2012), in human serum (Brasseur et al., 2014; Yan et al., 2012; Ren et al., 2009; Chen et al., 2015) cord serum (Ben et al., 2014) and adipose tissue (Pan et al., 2020; Yin et al., 2020) as well as in human hair (Chen et al., 2015). Furthermore, very high DP concentrations in blood and hair from workers of a manufacturing facility have been reported (Zhang et al., 2013). See Tables 12 and 16 in UNEP/POPS/POPRC.17/INF/9 for further details on detections in biota and human samples.

71. In summary, a long depuration half-life, which is indicative of a BCF above 5000 L/kg, has been reported for DP. The log K_{ow} for DP is greater than 5. Although aqueous exposure is expected to be of limited importance in terms of bioaccumulation potential of DP a BCF above 5000 L/kg has been reported for DP isomers in a fish study. In addition, BMFs and TMFs > 1 have been reported for several organisms and food webs. Many studies have revealed the presence of DP in wildlife and humans. Field monitoring data suggest that DP is bioavailable and can achieve high body burdens (ECHA, 2017b). These lines of evidence support the conclusion that DP is bioaccumulative. Additionally, DP was recently identified as a very bioaccumulative (vB) substance in the EU (ECHA, 2017a).

2.2.3 Potential for long-range transport

72. Due to its high hydrophobicity and low vapour pressure (see Section 1.1), DP in the atmosphere adsorbs primarily to airborne particles (Sverko et al., 2011; AMAP, 2017, Canada, 2019a). Monitoring studies have reported mean fractions of particle-bound DP in air as high as 99 % (Hoh et al., 2006). Measurements in seawater confirm the propensity of DP to adsorb to particles. In the East-Greenland Sea, particle-bound DP accounted for, on average, 97% of *syn*-DP and 80% of *anti*-DP. In Atlantic seawater, the particle bound fraction was on average 58% for *syn*-DP and 75% for *anti*-DP (Möller et al., 2010).

73. The modelled half-life (QSAR estimate) of DP in air is in one study reported to be 13.68 hours (Sverko et al., 2011), i.e. below the criterion of two days set in Annex D (d) (iii) of the Convention. The Canadian assessment reports estimated half-lives in air of 0.468 and 160.12 days for atmospheric oxidation and ozone reaction, respectively, using the AOPWIN 2010 model (12-hr day) (Canada, 2019a). However, air modelling that is based on gas phase reactions may underestimate the half-life in air. In the environment, chemicals may undergo LRET when bound to articles as shown already for other chemicals with similar physico-chemical properties such as decaBDE (Brevik et al., 2006; UNEP/POPS/POPRC.10/10/Add.2.; POPRC-10/2). As described in more detail in ECHA (2017b), the LRET of substances with low vapour pressure that adsorb strongly to particulates in the air is likely to be governed by the fate of the particulates to which they bind and can undergo LRET to remote regions when atmospheric conditions permit

(e.g. during dry periods). In line with this, the Canadian DP assessment concludes that "On the basis of the measurements obtained from remote regions and modelling results, DP, via sorption to particles, also has potential for long-range transport and deposition in remote areas." (Canada, 2019a). Further, as shown by Okonski et al. (2014) particle size likely affects the LRET of DP via air. Okonski et al. (2014) estimated wet and dry deposition of DP and three other FRs and found that smaller size fractions gave atmospheric deposition fluxes for DP that were around a factor of 1.7 times lower than estimates obtained considering bulk atmospheric particulates only. According to the authors, this is a result of the high proportion of particles <0.49 µm diameter that may be transported longer distances and that are less easily deposited or scavenged than larger particulate fractions.

74. Modelling studies performed with the OECD P_{OV} and LRTP Screening Tool, a software tool for screening chemicals for persistence and LRET potential (Wegmann et al., 2009), and comparing DP to DP analogues and benchmark chemicals including α-HCH, HCB, PCB-101, PCB-180, PCB-28, HBCD, atrazine, *p*-cresol, chlordane plus, Dec 602, Dec 603 Dec 604, suggest that DP has transport and persistence properties within the range for listed POPs (Sverko et al., 2011), and that DP may be deposited to some degree in remote regions (Canada, 2019a). However, the results generated using the OECD P_{OV} and LRTP Screening Tool are associated with uncertainty, largely because most of the input parameters used in the model are estimated (ECHA, 2017b; Sverko et al., 2011).

75. DP has been detected in many different environmental matrices and biota in remote regions; in Arctic air, snow, soil, sediment, water and biota (reviewed in AMAP, 2017; Vorkamp et al., 2019a,b; Canada, 2019a); in Antarctic air, soil, lichens, mosses, limpet, amphipods, cod, Antarctic icefish, kelp gulls, penguins, skuas, giant petrels, killer whales, southern elephant seals and Antarctic fur seals (Möller et al., 2010; Muñoz-Arnanz et al., 2016; Gao et al., 2018; Kim et al., 2015; 2018; 2021; Aznar-Aleman et al., 2019; Roscales et al., 2021); in air, soil and lichen at the Tibetan Plateau, a remote mountain region in Asia (Yang et al., 2016a; Liu et al., 2018); and in air, water, soil, plant and animal muscle and hair in the remote Xilingol Prairie in Inner Mongolia, China (Chen et al., 2021).

76. In Arctic biota, most detections have been made in species in the marine food web (polar bear, ringed seal, harp seal, hooded seal, beluga whale, killer whale, narwhal, glaucous gull, common gull, black guillemot eggs, common eider, European shag, kittiwake, mink, and cod (*anti*-DP only)). Detections also include terrestrial and freshwater species, i.e. reindeer dung, peregrine falcon eggs and landlocked Arctic char (reviewed in AMAP 2017; Canada, 2019a; see also Norwegian Environment Agency, 2018a; Dreyer et al., 2019; Letcher et al., 2018; Green 2019; Verrault et al., 2018; Vorkamp et al., 2015; 2018; 2019a; Houde et al., 2017; Simond et al., 2017; Schlabach et al., 2011; Fernie et al. 2017).

77. As discussed in further detail in section 2.3.1 below, while concentrations in remote regions are generally low, they are not always lower than DP levels in source regions (see Table 11 UNEP/POPS/POPRC.17/INF/9). Reported detection frequencies vary from non-detect to 100% detection. Collectively, the available monitoring data suggest that DP can be transported over large geographical distances to remote areas where it deposits, transfers to the receiving environment and is taken up in biota (reviewed in Sverko et al., 2011; AMAP 2017 and Canada 2019a; see also Möller et al., 2010; 2011; 2012; Na et al., 2015; Yang et al., 2013; Wang et al., 2010b).

78. Möller et al. (2010) were the first to publish environmental monitoring data demonstrating the potential for DP to undergo LRET. In this study, marine boundary layer air and surface seawater samples were collected during a sea expedition from the East-Greenland Sea via the Northern and Southern Atlantics toward Antarctica. DP concentrations ranged from 0.05 to 4.2 pg/m³ in the atmosphere and from non-detect to 1.3 pg/L in seawater. The results indicate that transport may occur both via air and seawater and show preferential partitioning to particles in both air and seawater (Möller et al., 2010). In the East-Greenland Sea, the authors observed decreasing concentrations of DP in air with increasing latitude. The authors suggest that the finding reflect stereoselective depletion of *anti*-DP likely caused by UV degradation during LRET. The trend of declining levels with increasing latitude was not reflected in the seawater samples. In the study, Western Europe was identified as a source region for DP in the marine environment. The analysis of air mass back trajectories showed a mix of oceanic, Arctic and continental air masses. The higher levels of DP in seawater observed at some northern sites were probably due to water masses originating from the Atlantic and Arctic Oceans, in combination with freshwater inputs from melting land ice.

79. On a sea expedition from the East China Sea to the Arctic (Möller et al., 2011), DP concentrations ranged from 0.01 to 1.4 pg/m³ in air and from 0.006 to 0.4 pg/L in seawater. The Asian continent was identified as a source of DP to this marine environment, and the air-seawater exchange indicated deposition of DP from air to water. DP concentrations in air in the Chukchi/Bering Sea region were generally low compared to the source region. Decreasing DP concentrations with increasing latitude were also observed in this study; however, concentrations were near detection limits.

80. In yet another study of DP in marine boundary layer air by the same authors, atmospheric samples were collected during a sampling expedition from the East Indian Archipelago toward the Indian Ocean and further to the Southern Ocean (Möller et al., 2012). *Syn*- and *anti*-DP were detected in all marine atmospheric samples at concentrations from 0.26 to 11 pg/m³. Measured concentrations were in the range of previously reported DP concentrations in the marine atmosphere, with the highest concentrations detected in the East Indian Archipelago (see Table 10 UNEP/POPS/POPRC.17/INF/9).

81. As discussed by Vorkamp et al. (2019a), the highest DP levels in Arctic air to date were detected in Alaska in 2012 (approximately 110 pg/m³), but the findings were not confirmed in a subsequent study (Sverko et al., 2011; Rauert et al., 2018). Reported mean DP concentrations in Greenland air in 2012, 2014 and the period 2014-2016 were 6.7, 0.64 and 4.2 pg/m³, respectively (Vorkamp et al., 2015; 2019a). However, it was discussed that the high values observed might reflect influence by local sources. DP was also positively detected in atmospheric samples from other Arctic sites: Alert in the Canadian Arctic (Xiao et al., 2012), Longyearbyen in the Norwegian Arctic (Salamova et al., 2014), and Little Fox Lake in the Canadian Sub-Arctic (Yu et al. 2015). Mean DP concentrations reported for these sites were ~0.75, ~1.2 and ~0.25 pg/m³, respectively. DP was also detected in air from Pallas in Northern-Finland at a mean concentration of 0.039 pg/m³ (Haglund et al., 2016).

82. LRET of DP via air is also documented by studies from other remote regions than the Arctic. The studies by Möller et al. (2010; 2012) demonstrate ongoing long-range atmospheric transport of DP to the Antarctic and further show that the DP levels in Antarctic air are comparable to those of other FRs such as polybrominated diphenyl ethers (PBDEs) and polybrominated biphenyls (PBBs). These findings are supported by detections of DP in Antarctic biota as further described in section 2.3.1 Environmental levels and trends (see Gao et al., 2018; Kim et al., 2015; 2018; 2021; Aznar-Alemayn et al., 2019; Roscales et al., 2021). DP also undergoes LRET to remote mountain areas. In a study on lichen from the southeast Tibetan Plateau, DP concentrations appeared to decrease with increasing altitude (Yang et al., 2013). According to the authors, the distribution pattern for DP observed in this study, combined with knowledge about the monthly average surface wind vector field, provides evidence that DP transport into the area is mainly driven by the Indian monsoon systems and can be attributed to LRET. Yang et al. (2013) further note that the lower DP levels observed at higher altitudes indicate that DP transported into the area via the atmosphere is cold trapped by the mountains.

83. The Global Atmospheric Passive Sampling (GAPS) network has been deploying passive air samplers quarterly since 2005. A retrospective analysis of GAPS samples from 2005 and 2006 covering 67 different sampling sites indicates that DP is globally distributed in air with total DP levels ranging from non-detect to 30 pg/m³. Elevated levels were reported for source areas; The median value at urban sites (~0.9 pg/m³) was approximately 10 times greater than at background, rural, and agricultural sites (Shuster et al., 2021). A study from 2018, probing legacy and alternative flame retardants in the air of 10 provincial capital cities across China using high-volume active air samplers reported a median DP concentration of 1 pg/m³, with concentrations ranging between 0.1 to 53 pg/m³ (Zhao et al., 2021). DP mainly existed in the PM_{2.5} phase (66% ± 25%) and was dominated by *anti*-DP (~60%). The total *anti*-DP concentration averaged at 0.4 ± 1.0 pg/m³, while that of *syn*-DP was 1.1 ± 2.9 pg/m³. When comparing their data to the results of a 2013/2014 survey of flame retardants in atmospheric particles in Chinese cities by Liu et al. (2016b). Zhao et al. (2021) found that levels of DP in air in Chinese cities have decreased by 50 % from 2013/2014 to 2018. Zhao et al. (2021) further report that their data from 2018 were an order of magnitude lower than those reported in a national air study sampled in 2005 (~15 pg/m³, Ren et al., 2008) and data from in northeast China between 2008 and 2013 (~7 pg/m³, Li et al., 2016) and suggests that current DP levels in Chinese cities are in a low range. The findings by Shuster et al. (2020) and Zhao et al. (2021) are comparable to DP levels reported in settlements in the European Arctic by Salamova et al. (2014) and are indicative of median global atmospheric DP levels of around ~1 pg/m³.

84. As shown in Table 12 in document UNEP/POPS/POPRC.17/INF/9 DP has been detected in muscle, liver, guano, eggs, plasma, preen oil and feathers from different species of birds around the world (Gauthier and Letcher 2009; Guerra et al., 2011; Munoz-Arnanz et al., 2011; 2012; Baron et al., 2014b; 2015; Jin et al. 2016; Vorkamp et al., 2018; Løseth et al., 2019; Briels et al., 2019; Mo et al., 2019; Desjardins et al., 2019; Zhu et al., 2020), including in migratory birds and eggs of migratory birds (e.g. Zhu et al., 2020 Vorkamp et al., 2018;). Vorkamp et al. (2018) detected DP in eggs of Greenland peregrine falcons which migrate from the Caribbean and South America to the Arctic for the summer. The DP detected in these eggs could reflect exposure experienced at the wintering and breeding grounds, as well as during migration. According to Jin et al. (2016) concentration levels of DP and other flame retardants in birds depend on the feeding habits and migration patterns of the species. In their study, residential predatory birds showed significantly higher concentrations of DP, PBDEs and other FRs than migratory predators and non-predatory birds because of differences in diet and habitat. Furthermore, DP levels in birds in remote and rural areas are generally in the lower range, whereas elevated concentrations have been reported in urban environments and for carnivorous predatory birds, see Table 12 in document UNEP/POPS/POPRC.17/INF/9 and references therein. For example, in the study by Jin et al. (2016) DP levels up to 3052 and 4724 ng/g lw was reported in livers of Common Kestrels and Common Buzzards in Korea while Chen et al. (2013) reported concentrations of up to 3820 and 2400 ng/g lw in livers of Common Kestrels and Eurasian sparrowhawks from an urban environment in China.

85. Birds have previously been identified as biovectors for the transport and deposition of POPs to ecosystems in remote regions through deposition of guano, feather loss and decaying carcasses (Evenset et al., 2007; Blais et al., 2005) and may represent an additional transport pathway for DP to remote regions. In Ellasjøen at Svalbard, seabird guano was, based on measurements of PCB, found to account for approximately 14% of the contaminant inventory of the lake catchment area, approximately 80% of the contaminant inventory of the lake itself and was suggested to be

approximately thirty times more efficient as a contaminant transport pathway compared to atmospheric LRET (Evenset et al., 2007).

86. In a study by Desjardins et al. (2019), deposition of guano by urban-adapted ring-billed gulls from a colony on Deslauriers Island near the city of Montreal, Canada were postulated to increase exposure locally (Desjardins et al., 2019). The total amount of halogenated FRs deposited by the entire ring-billed gull colony (64 980 gulls, both sexes combined) through guano was estimated to be 1 g during a 28-day period. DP was the predominant halogenated FR detected and *anti*- and *syn*-DP accounted for 2-3% and 1-2% of the total concentration halogenated FRs, respectively. The authors propose that the migratory ring-billed gulls from this urban-adapted colony could be an underestimated biovector of halogenated FRs to nearby ecosystems.

87. As discussed, Möller et al. (2010) postulated that the fraction of *anti*-DP in environmental samples (i.e. $f_{anti} = \text{anti}/(\text{syn} + \text{anti})$ based on concentrations) decreases with increasing distance from the source as a result of more rapid degradation of the *anti*-DP isomer than the *syn*-DP isomer in UV-light. However, these findings may also reflect isomerisation of the *anti*- to *syn*-DP (Sverko et al., 2011). Available monitoring studies provide some evidence of changes in isomer ratios with increasing distance from the source (ECHA, 2017b), but observations from the Arctic are generally inconclusive in this regard (AMAP, 2017). More specifically, DP isomer ratios in Arctic air have been reported in some studies to be in the same range as found in commercial mixtures (Xiao et al., 2012; Salamova et al., 2014; Vorkamp et al., 2015), but have in other studies (Möller et al., 2010; 2011; Carlsson et al., 2018; Na et al., 2015) been reported to differ from the commercial mixtures. Findings from Antarctica are consistent with a change in DP isomer ratio during LRET. In a study from the Fildes Peninsula in Antarctica, Gao et al. (2018) estimated the mean f_{anti} values in soil and lichen to 0.37 and 0.24, respectively. These f_{anti} values are lower than those of commercial products ($f_{anti} = 0.64-0.80$) and the authors conclude that the findings confirms that long-range atmospheric transport is a main source of DP, and that the DP burdens could be driven by a stereoselective enrichment of *syn*-DP during atmospheric transport as postulated by Möller et al. (2010). However, as the concentration of DP in soil and lichen differed between the sampling sites, the authors also suggest that the DP levels may have been influenced by anthropogenic and animal activities in the area (Gao et al., 2018).

2.3 Exposure

2.3.1 Environmental levels and trends

88. DP is globally distributed and is detected worldwide in many different environmental matrices and biota, and at different types of locations spanning from production sites and recycling facilities to urban, rural and remote areas (de la Torre et al., 2010; Sverko et al., 2011; Wang et al., 2016; Saini et al., 2020; Zafar et al., 2020; Schuster et al., 2021). An overview of available data is provided in Tables 10 and 12 in document UNEP/POPS/POPRC.17/INF/9. Elevated DP levels are reported in source areas (e.g. Saini et al., 2020; Schuster et al., 2021,) and are higher in urban centres compared to rural, agricultural and remote sites (Syed et al., 2020; Schuster et al., 2021) and also differ between cities globally (Saini et al., 2020). As described also in section 2.2.3 above, Schuster et al. (2021) report that the median value at urban sites ($\sim 0.9 \text{ pg/m}^3$) was approximately 10 times greater than at background, rural, and agricultural sites. Similar findings were reported by de la Torre et al. (2009) who found that DP levels from urban locations in Spain were 15 times higher than those found at remote sites, suggesting urban zones as important sources of DP to the environment.

89. DP levels in remote regions and in ocean water and air are generally lower than levels reported in source regions near production sites and urban centres, although high levels in some instances also have been observed at remote sites (reviewed in Sverko et al., 2011; Canada, 2019a). As further detailed below, due to limited data, the temporal and spatial trends for DP are equivocal.

90. Several DP monitoring studies have been carried out in the vicinity of DP manufacturers and e-waste recycling sites, including in the US, China and Pakistan (Sverko et al., 2011; Ji et al., 2018; Iqbal et al., 2017). High DP levels have been reported in various matrices from these areas (Wang et al., 2016), with reported concentrations up to several ng/m^3 close to known production or e-waste recycling sites (Sverko et al., 2011). The soil concentration of DP near the Anpon Electrochemical Co. production facility in China was found to be 0.83-1200 ng/g dw but decreased by an order of magnitude within 7.5 km (Wang et al., 2010a). Other studies have reported average soil concentrations of 1490 ng/g and 35.6 ng/g for this site (Wang et al., 2010c; Zhang et al., 2015). Atmospheric DP concentrations near the DP production plant in, China were found to be as high as 7737 to 26,734 pg/m^3 (Wang et al., 2010a). DP has also been detected in air and precipitation at Sturgeon Point, in Niagara River suspended sediment samples and in the Great Lakes watershed near the OxyChem manufacturing facility in the US (Shen et al., 2011; Olukunle et al., 2018). Total DP levels in air at Sturgeon Point were reported to be 2.70 pg/m^3 by Olukunle et al. (2018) and 3.1 pg/m^3 by (Liu et al., 2016), while Hoh et al. (2006) in one instance measured concentrations up to 490 pg/m^3 . Total DP levels in precipitation from the same site were 323 pg/m^3 and 890 pg/m^3 , as reported by Olukunle et al. (2018) and by Salamova and Hites (2011), respectively. Examination of sediments collected in this area between 1980 and 2006 showed a declining concentration trend for DP (Shen et al., 2011). A declining concentration trend for DP was also reported previously for Niagara River suspended sediments (Sverko et al., 2008)

and are consistent with declines observed in a sediment core collected near the river mouth on the Niagara River bar (Sverko et al., 2010). The suspended sediment trends observed by Shen et al. (2011) were also reflected in open lake sediment cores, which generally showed a decline in DP concentrations since 1980 in the Niagara basin. Similar findings were reported for the Mississauga basin (Shen et al., 2010). It is not clear if the decline in DP in Niagara River suspended sediment samples represent changes in production volumes at the Oxychem manufacturing site or whether it reflects emission controls adopted over the past 20 years at the site. Reported DP levels in surface sediments in Lake Ontario (2.230–586.0 ng/g dw) and Lake Erie (0.061–8.620 ng/g dw) in North-America, down-stream of the Oxychem manufacturing plant, are at the high end of reported concentrations globally, along with levels in surface sediments from some locations in China, including the Huai'an area where there is also a DP manufacturing plant (1.860–8.00 ng/g dw), the Dongjiang River (0.80–19.40 ng/g dw) Qiantang River (<0.009–1.10 ng/g dw) and mangrove sediments of the Pearl River Estuary (0.013–1.504 ng/g dw) (Sverko et al., 2008; Wang et al., 2010; He et al., 2014; Sun et al., 2013; 2016, as cited in Niu et al. 2020). Surface sediments in source areas are likely important reservoirs of DP in the environment (Niu et al. 2020), but monitoring data also provide evidence for uptake in biota and elevated levels in aquatic species in these areas (reviewed in Zafar et al. 2020, see also Table 12 in UNEP/POPS/POPRC.17/INF/9). For example, DP levels in muscle tissues of freshwater fish species from the Liaohe River in China ranged from below detection limit to 470 pg/g lw (Ren et al. 2013).

91. A very high DP concentration (3327 ng/g) was found in soil at an e-waste recycling site in Qingyuan, China. DP levels fell dramatically with increasing distance away from the recycling site (Yu et al., 2010). High levels (average 3398 ng/g) of DP were also reported in soil samples from areas with informal e-waste recycling activities in Karachi, Pakistan (Iqbal et al. (2017). The high DP levels in soil reported at recycling sites by Iqbal et al. (2017) and Yu et al. (2010) are comparable with levels reported from other e-waste recycling sites in Asia (South/ Southeast China) (Li et al., 2018; Ge et al., 2020) and much higher than the DP levels in soil collected in close vicinity to the DP manufacturing facility in Huai'an City, China (Wang et al., 2010a,c; Zhang et al. 2015). In the recycling town Guiyu, in southeastern China total DP levels in soil were 3.8–2100 ng/g, and DP was also found in sediment (1100–7200 ng/g dw), road dust (14–1100 ng/g dw) and PM_{2.5} particles (180 pg/m³ *syn*-DP and 170 pg/m³ *anti*-DP) (Li et al., 2018). Elevated DP levels in air (15–85 ng/m³) were also reported in the e-waste area of Karachi, and are higher than reported levels found in air close to the DP manufacturing site (7.7–26.7 ng/m³) in China (Iqbal et al., 2017; Wang et al., 2010a; see also review by Zafar et al., 2020). Elevated levels have also been reported in biota near e-waste sites. DP concentrations from 19 to 9630 ng/g lw have been reported in aquatic species including water snake and mud carp from an e-waste recycling workshop of South China (Wu et al., 2010). In another study from e-waste recycling site at South Longtang Town China, two species, northern snakehead and Crucian carp were studied for maternal transfer and sex-related accumulation of DP. The concentrations of hepatic DP for northern snakehead and Crucian carp were 260–1920 ng/g lw and 340 to 1670 ng/g lw, respectively (Wu et al. 2013).

92. According to Iqbal et al. (2017) elevated levels of DP at informal recycling sites may originate from emissions associated open-air burning-, acid stripping-, manual dismantling-, and shredding of cables, televisions, plastic roofing and connectors. Iqbal et al. (2017) furthermore suggest that sites with informal e-waste recycling may have higher environmental impact than manufacturing sites and urban environments, stressing the importance of chemical management strategies across the entire life cycle of electronics (Iqbal et al., 2017). However, Ge et al. (2020) found elevated concentration levels of DP (range of total DP was 21.8–18 000 ng/g and the median levels of *syn*- and *anti*-isomers was 211 ng/g and 504 ng/g, respectively) in soil samples from within a newly (2015) established recycling facility that uses modern technology. The study compared soil samples from the recycling facility with samples from the surrounding area (range of total DP was 1.76–4 050 ng/g and the median levels of *syn*- and *anti*-isomers was 2.81 ng/g and 8.52 ng/g respectively) which had lower levels of DP in soil. DP has also been detected wildlife living at or in close vicinity of e-waste sites (e.g. Li et al., 2014; Wu et al., 2013; 2018, Peng et al. 2019). Peng et al. (2019) found that birds (kingfishers) feeding close to e-waste recycling sites in South China had 10- to 1 000- times higher levels of total DP concentrations than kingfishers from non-e-waste sites.

93. DP levels in urban areas and urban industrialized areas are reported in several studies and from several global regions. In Pakistan atmospheric concentrations of legacy and emerging flame retardants were monitored in eight major cities across the country (Karachi, Lahore, Faisalabad, Rawalpindi, Multan, Quetta, Peshawar and Gilgit) using a high volume-active air sampler (gas + particle phase). *Anti*- and *syn*-DP were detected in all cities with concentrations ranging from 0.60–12.0 pg/m³ (3.08 pg/m³) and 0.80–42.3 pg/m³ (10.8 pg/m³), respectively. Sum DP concentrations ranged from 1.4–54.2 pg/m³ with an average of 13.8 pg/m³ (Syed et al., 2020). Earlier studies from Pakistan have reported sum DP concentrations in urban air of 88.0, 0.41 and 2.48 pg/m³ in the vicinity to River Ravi, River Chenab and Indus Basin, respectively (Syed et al., 2013; Ali et al., 2015; Mahmood et al., 2015). DP has also been detected in sediments from 14 estuaries from South China that are influenced by a wide range of anthropogenic activities. Reported sum DP concentrations ranged from 0.025 to 1.63 ng/g dw. Estuarine sediments from the Pearl River had the highest DP levels (mean of 0.850 ± 0.089 ng/g dw). *Anti*-Cl₁₁-DP, the dechlorination product of *anti*-DP, was also detected in 52.9% of the 14 estuarine sediments with concentrations ranging from non-detect to 0.047 ng/g dw. The Pearl River Estuary, located in the Pearl River Delta region of South China, has become one of the most developed and urbanized regions in China and is a reservoir for man made contaminants discharged from industrial waste, municipal sewage and upstream runoff and e-waste recycling activities (Xie et al., 2021 and

references therein). In another study from the Pearl River Delta in China, the concentration of DP in surface sediments was higher than for other dechloranes including Mirex, Dec-604, Dec-603, Dec-602 and Chlordene Plus (CP) with a reported mean level of $3090 \pm 5,490$ and minimum and maximum concentrations of 0.00824 and $33 \mu\text{g/g dw}$ (Li et al., 2021a). Spatially, the DP concentrations in sediment were found to be significantly and positively correlated with the municipal gross domestic product, population and sewage discharge of the cities in the Pearl River Delta, but insignificantly related to their industrial outputs, indicating that the DP in the Pearl River Delta mainly originated from urban activities and not industrial ones. Also, other studies highlight urban centers as an important source for emissions and releases of DP. Ren et al. (2008), measured DP in air across 97 Chinese urban and rural sites. DP was detected in 51 of these sites, with a mean air concentration in urban centers of $15.6 \pm 15.1 \text{ pg/m}^3$. The levels in urban centers were approximately 5 times greater than those measured in rural areas $3.5 \pm 5.6 \text{ pg/m}^3$ and are comparable to the levels reported in urban centres in Pakistan by Syed et al. (2020). A more recent study from Chinese cities reported average air levels of 2.04 pg/m^3 and 1.5 pg/m^3 for *syn*- and *anti*-DP, respectively (Liu et al., 2016b). A study from the Great Lakes, US, showed concentrations of DP ranging from 0.31 – 4.80 pg/m^3 at an urban site in Chicago (Hoh et al., 2006). In a study from Canada, urban-feeding ring-billed gulls breeding on Deslauriers Island in an industrialized section of the St. Lawrence River, downstream from Montreal were found to contain high levels of DP. The DP hepatic mean level was 230 ng/g lw . *Anti*- and *syn*-DP were detected in 100% and 93% of the investigated livers, respectively (Gentes et al., 2012). In a study of redlisted and endangered migratory and residential birds in Korea, Jin et al. (2016) found that more than half of the bird samples (65%) contained detectable concentrations of DP. Sum DP concentrations ranged from below the limit of quantification to 4734 ng/g lw with a mean concentration of 172 ng/g lw . Overall, sum DP levels were, according to the authors, similar to sum PBDE concentrations. There was also evidence for potential biomagnification in birds (see details in Section 2.2.2).

94. In a study of halogenated FRs in eels from five Latvian lakes, nine dechlorane-related compounds (i.e. mirex, *syn*-DP, *anti*-DP, Dec 602, Dec 603, Dec 604, hexachlorocyclopentadienyldibromocyclooctane, Cl_{10}DP and Cl_{11}DP) were analyzed. The mean total concentration of dechlorane-related compounds was 0.62 ng/g lw and the geographical distribution was nearly uniform among the five lakes (Zacs et al. 2018). The concentrations of *syn*- and *anti*-DP ranged from non-detect to 0.45 ng/g lw , and from non-detect to 0.89 ng/g lw , respectively, with average concentrations of 0.14 ng/g lw , and 0.24 ng/g lw , respectively. The contribution of *anti*-DP to the total concentration of dechlorane-related compounds was approximately 33%, while *syn*-DP accounted for approximately 10% on average. The composition of mixtures containing *syn*- and *anti*- DP showed an f_{anti} value that was close to the composition of the OxyChem® DP commercial product, indicating the release of DP from disposal of consumer products as the probable source. In another study from the Baltic region, DP was widely detected in Baltic biota, with high levels in some birds (particularly eider) (de Wit et al., 2020). In this study a suite of chemicals of emerging concern, including DP, were analysed in blue mussel, viviparous eelpout, Atlantic herring, grey seal, harbor seal, harbor porpoise, common eider, common guillemot and white-tailed eagle from the Baltic Proper, sampled between 2006 and 2016. Both DP isomers were found in all species. Mean reported *syn*- and *anti*-DP levels were highest in common eider liver (16 and 52 ng/g lw , respectively). Mean reported *syn*- and *anti*-DP levels were also high in grey seal blubber (6.0 and 26 ng/g lw , respectively) and white-tailed eagle (3.7 and 7.8 ng/g lw , respectively). Lower means were found in herring muscle (0.035 and 0.070 ng/g lw for *syn*- and *anti*-DP), harbor seal blubber (0.46 and 0.092 ng/g lw for *syn*- and *anti*-DP), harbor porpoise (0.04 and 0.074 ng/g lw for *syn*- and *anti*-DP), eider eggs (0.10 and 0.21 ng/g lw for *syn*- and *anti*-DP) and guillemot eggs (0.13 and 0.40 ng/g lw for *syn*- and *anti*-DP). The lowest levels were found in blue mussel and viviparous eelpout with reported *syn*- and *anti*-DP means below the reported limits of detection.

95. DP has also been detected in stranded plastic foams from an island located in South China Sea along with other FRs and plasticizers, including PBDEs, organophosphate esters (OPEs), emerging brominated FRs, and DP (Cao et al. 2020). In an assessment of plastic additives in drowned herring gulls ($n=10$) that had been washed ashore in Skulfsfjord in Troms, Northern Norway, DP and other dechloranes were among the additives found with high detection rates (Herzke et al., 2019). The cause of death for these birds was most probably linked to professional fishing activities in the area. Plastic was found in 20 % of the investigated stomach contents. In general, the results did not indicate a relationship between gastric contents (plastic occurrence in the stomach) and the concentration plastic additives in the liver of these herring gulls. However, only limited incidences of ingested plastic were found.

96. As described above and in section 2.2.3, DP is detected in remote areas globally. See also Tables 10 and 11 in UNEP/POPS/POPRC.17/INF/9. A review of emerging Arctic contaminants conducted by AMAP concluded that atmospheric concentrations of DP in the Arctic were comparable to those of PBDEs (AMAP, 2017). While DP concentrations in remote regions are generally low, they are not always lower than in source regions. For example, in the study investigating halogenated FRs in air and soil from Mt. Gongga on the eastern Tibetan Plateau, DP levels ranging from below detection limit to 11.5 pg/m^3 and 48.3 pg/g in air and soil, respectively, were reported (Liu et al., 2018). The Tibetan Plateau is the world's highest elevation plateau and is considered as one of the most remote and isolated areas on earth, and ideal for the study of LRET of POPs (Yang et al., 2016a). According to the authors (Liu et al., 2018), the concentrations of DP in air and soil on the Tibetan Plateau were higher than in the Great Lakes region (0.14 – 4.0 pg/m^3) (Sverko et al., 2011), but within the range of air samples from China (not detected to 66 pg/m^3) (Ren et al., 2008), and much lower than those observed near a production facility in China (7737 – $26,734 \text{ pg/m}^3$)

(Wang et al., 2010a). In another study of lichen from the southeast Tibetan Plateau, DP was detected in 89% of the samples. Concentrations ranged from 27 to 843 pg/g, with an average concentration of 167 pg/g (Yang et al., 2016a). The concentrations reported were comparable, but slightly higher than in tree bark from Shenzhen, Hangzhou and Tianjin in China (Qiu and Hites, 2008) and much lower than in tree barks from South Korea and New York in the US (Qiu and Hites, 2008) but significantly higher than in moss from Ny-Ålesund at Svalbard (Na et al., 2015). In a study investigating concentrations in air and tree bark in the Great Lakes basin, DP was found in all of the atmospheric and tree bark samples, including the ones from the more remote sites (Salamova and Hites, 2010). In a study of brominated and chlorinated flame retardants in about 40 samples of tree bark from 12 locations around the globe, DP concentrations varied from 0.89 to 48 ng/g lipid weight. Both the *syn*- and *anti*-DP isomers were detected in all the tree bark samples analyzed (Salamova and Hites, 2013). The findings by Salamova and Hites in 2013 shows DP to be present at multiple locations globally and lend support to other environmental data indicating that DP is a worldwide pollutant.

97. In Antarctic soil, lichen and moss samples, DP was detected in all samples (100%) (Gao et al., 2018; Kim et al., 2018). The DP concentrations in Antarctic soil were between one and several orders of magnitude lower than the levels reported in surface soil samples collected from an industrial region (0.0336–4.65 ng/g) and an e-waste recycling area (nd–47.4 ng/g) in South China (Yu et al., 2010). On the other hand, DP levels in both lichen and moss from Antarctica were higher than previously measured in mosses sampled in Ny-Ålesund in Arctic Norway in 2012 (Na et al., 2015). The average concentrations of DP in lichen from Antarctica were similar to those in reeds in northeastern China (0.63 ng/g wet weight (ww)) (Wang et al., 2012) and in lichen in the southeast Tibetan Plateau (167 pg/g) (Yang et al., 2016a) but lower than in tree bark from areas in the northeastern US (0.03–115 ng/g) and South Korea (1.4 ng/g) (Qiu and Hites, 2008).

98. In a Norwegian screening assessment of emerging Arctic contaminants, DP was detected in all species monitored. The detection frequency within the species tested was 100%, except in common eiders where it was 80% (Norwegian Environment Agency, 2018a). *Syn*- and *anti*-DP concentrations in organisms from Svalbard were in the range from non-detect to 6.9 ng/g ww with reported average values of 0.04 and 0.07 ng/g ww for common eider (eggs), 0.01 and 0.03 ng/g ww for kittiwakes (eggs), 0.24 and 1.8 ng/g ww for glaucous gull (eggs), 0.63 and 4.5 ng/g ww for polar bear (blood). Moreover, DP was also detected in European shag (eggs) from Røst, in mink (liver) from Sommarøy and in common gulls from Tromsø. Reported *syn*- and *anti*-DP levels in these species were 0.22 and 1.4 ng/g ww in European shag, 0.21 and 1.2 ng/g ww in mink, and 0.1 and 0.63 ng/g ww in common gulls, respectively. In another study conducted for the Norwegian Environment Agency, DP levels were below the limit of quantification in cod livers from Arctic sites at Svalbard and Kjøfjord in the Outer Varangerfjord and the Tromsø harbour area (Green et al., 2019). In contrast, reported *syn*- and *anti*-DP in cod livers from the Bergen harbour area, a non-Arctic site on the west coast of Norway, were 0.178 and 0.203 µg/kg ww. *Syn*- and *anti*-DP levels in cod livers from the Inner-Oslofjord in Southern Norway were 0.135 and 0.231 µg/kg ww, respectively.

99. In the previously mentioned report by Vorkamp et al. (2019a) on DP in air and biota from Greenland, *anti*-DP isomer was detected in 92% of all air samples from 2014 to 2016 and in 46% of the air samples collected in 2014. As already indicated above in section 2.2.3, the mean reported concentrations in air in this study were 4.2 pg/m³ and 0.64 pg/m³ for the period 2014 to 2016 and for 2014, respectively. Except for glaucous gull, DP levels were close to or below limits of quantification in biota samples. Neither of the two DP isomers were found in ringed seal samples from Thule. In ringed seals from East Greenland, Arctic char and narwhals, only the *anti*-DP isomer was present. *Syn*- and *anti*-DP were both detected in 100% of all glaucous gull samples from Thule at mean concentrations of 0.22 and 0.89 ng/g lw (0.012 and 0.049 ng/g ww), respectively. Detection frequencies for *syn*- and *anti*-DP in glaucous gull samples from East Greenland were 87.5 and 100%, with mean concentrations of 0.24 and 0.93 ng/g lw for *syn*- and *anti*-DP. A geographical comparison based on samples of glaucous gull and ringed seal showed relatively uniform DP levels. The DP levels measured in this study were generally comparable to concentrations reported for some of the same species (e.g. ringed seals, glaucous gull and Arctic char) from other locations in the Arctic. However, DP concentrations in glaucous gull liver samples from Greenland reported in this study were a factor of 5-10 lower than in corresponding samples from the Canadian Arctic (Verreault et al., 2018). In the study from the Canadian Arctic, *syn*- and *anti*-DP were detected in 65% and 59% of male liver samples from 2012, respectively (Verreault et al., 2018). In females, the detection frequency of both isomers was 21%. Mean concentrations were only calculated for male birds and were 0.18 and 0.16 ng/g ww for *syn*- and *anti*-DP, respectively.

100. DP was also previously detected in air and biota samples from Greenland along with other FRs (Vorkamp et al., 2015). Mean *syn*- and *anti*-DP concentrations in air were 2.3 and 5.2 pg/m³. While the detection frequency of both isomers in air was 46%, *anti*- and *syn*-DP was detected in 95% and 100% of biota samples, respectively. In biota, ringed seals from East Greenland had the highest DP concentrations with reported mean blubber concentrations of 0.096 and 0.42 ng/g ww for *syn*- and *anti*-DP, respectively, but were not statistically different from West Greenland ringed seal samples. Reported mean concentrations for *syn*- and *anti*-DP in other species were: 0.14 and 0.67 ng/g ww in black guillemot eggs, 0.023 and 0.11 ng/g ww in glaucous gull liver, 0.019 and 0.071 ng/g ww in ringed seal blubber from West Greenland and 0.021 and 0.055 ng/g ww in polar bear adipose tissue.

101. In another study, covering multiple locations in the Canadian Arctic, DP was only detected sporadically in ringed seals (Houde et al., 2017). Detectable levels of *syn*- and *anti*-DP ranged between 0.04 and 0.41 ng/g lw and 0.04–6.3 ng/g lw, respectively.

102. Minke whales and beluga whales from the St. Lawrence Estuary as well as beluga from the Canadian Arctic have also been studied (Simond et al., 2017). In 2013, the mean DP concentration for these three populations was 0.31 (minke whales, St. Lawrence Estuary), 0.44 (beluga, St. Lawrence Estuary) and 1.28 ng/g lw (beluga, Arctic). The highest concentration of 1.28 ng/g lw in the Arctic beluga population was different from the generally observed contaminant distribution, which usually showed higher concentrations at more southern locations, and could not be explained. Concentrations in beluga from St. Lawrence estuary and the Canadian Arctic were roughly the same order of magnitude as observed in two killer whales from Greenland with quantifiable levels of *anti*-DP (Vorkamp et al., 2019).

103. Research on four avian species on King George Island showed that DP was present in Gentoo penguin, Adelle penguin, South Polar skua and Brown skua (Kim et al., 2015). Reported concentrations of DP were 0.250–0.329 ng/g lw in the penguin tissues and 2.12–11.1 ng/g-lw in the skua tissues. Dec 603 and 604 were not detected, whereas Dec 602 was detected in the highest concentrations in all samples, followed by *anti*-DP and *syn*-DP. In another Antarctic study, Gao et al. (2018), reported mean DP concentrations of 0.233 ± 0.089 , 0.201 ± 0.086 and 0.15 ± 0.147 ng/g dw in soil from three different locations (i.e. a coastal area, an inland area and Ardely Island) on the Fildes Peninsula. Reported levels in lichen from the same locations were 0.449 ± 0.213 , 0.337 ± 0.172 and 1.513 ± 1.376 pg/g dw, respectively. DP was detected in all samples. Kim et al. (2018) reported the presence of DP in lichen and mosses sampled from 16 different sampling sites spread across the South-Shetland Islands, King George Island and Anvers Island in Antarctica. Similar to Gao et al. (2018) and based on geographical differences in DP levels and f_{anti} values, the authors propose that LRET, human activities, melting glacier water and biological activities, e.g. from penguins, are possible sources of DP in the area (Kim et al., 2018). In yet another study by Kim et al. (2021), DP was detected in limpet (0.398–11.1 ng/g lw), antarctic cods (0.0–0.548 ng/g lw), amphipods (0.227–8.71 ng/g lw), antarctic icefish (2.50 ng/g lw), gentoo penguins (0.009–11.8 ng/g lw), chinstrap penguins (0.0–3.00 ng/g lw), south polar skua (0.731 ng/g lipid), and kelp gull (0.028 ng/g lw) collected from the Barton Peninsula and Maxwell Bay, King George Island. Compared with the results from King George Island in 2008/09 (Kim et al., 2015), the DP levels in the penguin samples had slightly increased, although the sample size of the previous study was small. Aznar-Alemay et al. (2019) investigated DP and other dechloranes in dead seals from the South Shetland Islands on the Antarctic Peninsula. Only Dec 602 and *anti*-DP were found. *Anti*-DP was detected in adipose tissue from southern elephant seals and in Antarctic fur seals at average concentrations of 0.60 ng/g lw and 0.16 ± 0.16 ng/g lw, respectively, but were below the limit of quantification in muscle, brain and fur. Dec 602 appeared in higher concentrations than *anti*-DP, especially in adipose tissue (Aznar-Alemay et al., 2019). *Syn*- and *anti*-DP have also been detected in integument biopsies (epidermis, dermis and blubber) of killer whales from the Ross Sea, Antarctica (0.0068–0.0741 ng/g lw; with mean and median concentrations of 0.0192 and 0.0119 ng/g lw) (Muñoz-Arnanz et al., 2016).

104. Several studies have detected 1,3- or 1,5-DPMA in environmental samples such as sediment and fish (e.g. Sverko et al., 2010; Tomy et al., 2013; Sühling et al., 2014; Wang et al., 2015; Wolschke et al., 2015; Rjabova et al., 2016). It is possible that failure to use a non-destructive clean-up procedure during sample preparation could lead to under-reporting of this substance (Rjabova et al., 2016). In some cases, the concentrations of DPMA-isomers were greater than the total DP concentration in the same samples, suggesting the possibilities of greater bioaccumulation of 1,3-DPMA due to its smaller molecular size (See Table 13 in UNEP/POPS/POPRC.17/INF/9). 1,3-DPMA (but not 1,5-DPMA) was detected in lake trout at an average concentration of 34 ± 43 µg/kg lw, which was around ten-fold higher than concentrations of DP in the same samples reported by Tomy et al. (2007). In peregrine falcon eggs harvested in Spain and Canada, DPMA was detected in 24 out of 25 eggs, with concentrations ranging from 1.7 to 469 ng/g lw and 1.2 to 1660 ng/g lw in peregrine falcon eggs from Spain and Canada, respectively (Guerra et al., 2011). The corresponding DP levels were in the range 0.3 to 3.6 and 7.5 to 209 ng/g lw, respectively. DPMA has also been detected in tissues (except blood) of brown skua from King George Island, Antarctic indicating potential LRET. The detection frequencies were 21%, 11%, and 7.1% for DPMA, Dec-604 and *anti*-DP with the highest levels of 136, 165, and 8.9 pg/g dw, respectively (Wolschke et al., 2015).

105. Spatial trends of DP in remote regions were reported in five separate studies from the Arctic (Möller et al., 2010; 2011; Vorkamp et al., 2015; 2019a,b). As discussed in section 2.4.4 above, Möller et al. (2010;2011) observed decreasing concentrations of DP in air, but not in water, with increasing latitude. Biota data from Greenland show either fairly uniform levels or no spatial trends; Vorkamp et al. (2015) found that DP levels were not statistically different in ringed seal samples from East and West Greenland, while Vorkamp et al. (2019a) found a fairly uniform occurrence of DP in glaucous gull and ringed seal samples from different locations. In a study from Antarctica, Roscales et al. (2016) investigated differential exposure of giant petrels to POPs across a wide latitudinal gradient. DP was analyzed in plasma of Southern giant petrels breeding on Livingston, Marion, and Gough islands and in Northern giant petrels from Marion Island. Breeding locality was a major factor explaining petrel exposure to POPs compared with species and sex. DP was found at very low levels compared with legacy POPs. Mean DP concentrations were very close to the limit of quantification at all three breeding localities, and in many samples *syn*- and *anti*-DP were below the limit of quantification. Overall, pollutant levels in giant petrels decreased significantly with latitude, but the

relative abundance (%) of the more volatile POPs increased towards Antarctica. Principal component analysis showed a slight increasing tendency in DP presence from southern to northern breeding localities. No significant relationships were found between *anti*-DP concentrations and stable isotope signatures of C and N.

106. Temporal trend data for DP are also equivocal. Air samples (gas and particle phase) from Harbin, China indicated significant and increasing trends for both *syn*- and *anti*-DP in the period 2008 to 2013 (Li et al., 2016). Reported doubling times were 2.3 and 1.8 years, respectively. The time it took for *syn*- and *anti*-DP concentrations to double was significantly different, but the doubling times of both DP isomers were shorter than reported for 2-ethylhexyl 2,3,4,5-tetrabromobenzoate and bis(2-ethylhexyl) tetrabromophthalate, suggesting a more rapid increase for DP than the other FRs in the study. Air parcel backward trajectories and source contribution suggested a strong input from local sources (Li et al., 2016).

107. In a study from North America, DP concentrations in air (gas and particle phase) increased from 2005 to 2013, with doubling times of 4–6 years at three sites (Chicago, Sleeping Bear Dunes and Eagle Harbor) in the US Great Lakes area. At the two sites closest to a known production site for DP, Cleveland and Sturgeon Point, levels were stable and unchanged in the same time period (Liu et al., 2016a). A study from Canadian sites in the Great Lake Basin (Burnt Island, Egbert and Point Petre) reported trends in levels of *syn*- and *anti*-DP in air from 2005 to 2014 at two of the sites, Burnt Island and Point Petre (Shunthirasingham et al., 2018). The concentration of *anti*-DP in air was observed to decrease over time, with reported half-lives of 3.4 years at Burnt Island and 13 years at Point Petre. The declines leveled off after 2010. *Syn*-DP levels increased over time at Burnt Island with a doubling time of 7.6 years but decreased at Point Petre with a half-life of 6.6 years. In yet another study from the Great Lakes area, Olukule et al. (2018) report that *syn*-, *anti*-, or total DP concentrations in the air (vapor + particle phase) and in precipitation have not changed over the period 2005–2015 at four sites near the Great Lakes; Sturgeon Point, Point Petre, Cleveland and Chicago. Salamova and Hites (2011) assessed the temporal trends of DP at US Great Lakes sites over the time period of 2005–2009 and observed an overall (vapor, particle, and precipitation phases combined) doubling time of 9.5 years for *anti*-DP and no significant trends for *syn*-DP and total DP. The increasing levels of DP observed in Chicago, Sleeping Bear Dunes and Eagle Harbor by Liu et al. (2016a) and for *syn*-DP at Burnt Island by Shunthirasingham et al. (2018) are according to the authors consistent with the results presented by Salamova and Hites (2011). The results reported by Liu et al. (2016a), suggest a more rapid doubling time in DP levels in the atmosphere around the Great Lakes from 2005–2013 than reported previously by Salamova and Hites (2011) for the period 2005–2009. However, although the monitoring sites in the Great Lakes area are in the same region, they are not identical.

108. When compared to the global data, levels in air in Chicago and at Sturgeon Point are at the higher end of the range and are similar to the average DP level (3.5 pg/m³) from Taihu, China (Qiu et al., 2010). The increasing levels of DP observed in the Great Lakes region which are reported in some studies (Li et al., 2016; Liu et al., 2016a; Olukunle et al., 2018; Salamova and Hites, 2011; Shunthirasingham et al., 2018), and in a typical urban city in China (Li et al., 2016) may indicate that DP is being increasingly used. However, a more recent study by Zhao et al. (2021) shows that DP levels in air in Chinese cities sampled in 2018 were an order of magnitude lower than reported concentrations from a national air study (~15 pg/m³) sampled in 2005 (Ren et al., 2008), and in northeast China (~7 pg/m³) between 2008 and 2013 (Li et al., 2016). Using a multiple linear regression model of DP concentrations to isolate the variabilities due to sampling date and population near the sampling site, Olukunle et al. (2018) showed that DP concentrations in precipitation, gas and particle phase air samples from the Great Lakes were changing as a function of sampling date. On the other hand, and as described earlier, decreases of DP were observed in suspended sediments, a sediment core, and lake trout samples of Lake Ontario, following peaks in the 1980s (Shen et al., 2011). The reasons for the different time trends in air, precipitation, sediment and fish in the Great Lakes area are not clear, but as indicated by Vorkamp et al. (2018) it is possible that the different matrices studied (and also the study sites) reflect different emission sources.

109. There is also some information on time trends in biota. A recent study by Vorkamp et al. (2018), the only study available on temporal trends of DP in the Arctic, reported DP concentrations in the range 0.984 to 37.9 ng/g lw during the period 1986 to 2014 in peregrine falcon eggs from South Greenland. A non-significant increasing trend (annual change 3.3%) was observed (Vorkamp et al., 2018). In belugas from the St. Lawrence Estuary, DP increased from 1997 to about 2000 and decreased subsequently, possibly with a second peak around 2010 (Simond et al., 2017). No significant change in DP concentrations was found in eggs of white storks or black kites collected in 1999, 2003, 2011 and 2013 from the Doñana Natural Space in Spain (Báron et al., 2015). In a study investigating FRs in Indo-Pacific humpback dolphins and finless porpoises from the South China Sea, Zhu et al. (2014) found a positive temporal trend in the ratio of DP to BDE-209 between 2003 and 2011 in dolphin samples and between 2003 and 2012 in porpoise samples. The finding was by the authors attributed to an increasing use of PBDE alternatives in the South China region following the restriction of the production and use of PBDE commercial mixtures. A study reporting levels and trends of different FRs in the German and polar environments from 2005 to 2015/2016 found declining trends for DP in some of the investigated biological matrices from Germany (i.e. tree leaves, tree shoots, herring gull eggs) and increasing trends in bream (filet) and blue mussels (soft body) from Germany (Dreyer et al., 2019). In the same period, a small declining trend was also observed in riverine suspended matter. However, the standard deviation in this sample set was quite large. As noticed by the authors, herring gulls feed in the marine environment, e.g. on blue

mussels, but may also have a terrestrial diet. Thus, one possibility is that the seemingly contrasting time trends reported by Dreyer et al. (2019) reflect differences in DP emission and deposition between terrestrial and aquatic environments.

110. Rauert et al. (2018) report air monitoring data from 48 global sites across all five UN regions that were collected using polyurethane foam passive air samplers. The study generally showed low detection for all FRs at background sites. PBDEs showed higher mean concentrations and detection frequencies than the novel brominated and chlorinated FRs which also included DP (Rauert et al., 2018). *Syn*-DP was below the detection limit at all 48 sites and *anti*-DP at all sites except Paris, France, where high levels of DP were reported in 2014 (19 pg/m³ in the first and fourth of the year, 7.5 pg/m³ in the second quarter two and 116 pg/m³ in third quarter). When compared to monitoring data from 2005 and 2008 to 2009, respectively, Paris had elevated concentrations of novel FRs and reduced PBDE levels. *Anti*-DP and the FRs β -tetrabromoethylcyclohexane and bis(2-ethyl-1-hexyl) tetrabromophthalate were the dominating novel FRs. Rauert et al. (2018) propose that "the increasing concentrations of the replacement FRs and the reduced levels of the PBDEs may be indicating the shift away from the use of PBDEs to alternative FRs in this urban area".

2.3.2 Human exposure

111. Humans may be exposed to DP from indoor dust, food, indoor and ambient air, water, soil, sediment, *in utero*, and breast milk. DP has been detected in indoor air and domestic dust samples from several countries indicating exposure from products and manufactured items containing DP (Shoeib et al., 2012; Johnson et al., 2013; Wong et al., 2017; See Table 14 in UNEP/POPS/POPRC.17/INF/9). Higher levels of DP in air from urban compared to rural environments have been observed (Ren et al., 2008; Chen et al., 2011). A regular distribution pattern of DP in dust with particle size was observed with stronger enrichment on fine particles than coarse particles, which indicates potential human exposure (Cao et al., 2014). As reported in section 2.1.3 above, in indoor dust in Beijing, China, highest DP levels were found in pooled samples from classrooms (231 ng/g), which was higher than in house dust from Guanzhou, China (mean 18.9 ng/g) (Cao et al., 2014). In indoor dust from different microenvironments (i.e. home, workplace, car) in Greater Cairo, Egypt. The highest level and detection frequency (100%) were found in car dust, with *syn*- and *anti*- DP reaching 5 and 6 ng/g, respectively, followed by workplace and homes (Hassan and Shoeib, 2015). In the Republic of Korea, DP was detected in almost all samples of indoor dust from homes (0.30 – 530 ng/g; median 9.1 ng/g), offices (3.7 – 100 ng/g; median 35 ng/g), and daycare centers (6.2 – 56 ng/g; median 12/ng/g) (Lee et al., 2020). The estimated daily intake (EDI) via dust ingestion for toddlers and adults in the worst scenario (95th percentile) were 0.20 and 0.02 ng/kg bw/day for Σ DP (Lee et al., 2020). Air sampling in 15 homes in India found DP levels ranging from 0.2 to 5.43 pg/m³ (median 2.81 pg/m³) and 0.52–62.7 pg/m³ (median 1.62 pg/m³) for urban and suburban sites, respectively (Yadav et al., 2020). Cequier et al. (2015) found no correlation between DP levels in indoor air, dust and serum levels for 46 Norwegian women. However, both levels in air, dust and serum were low in these samples. Mean DP levels in serum were 2.6 ng/g lw and dust samples from the corresponding homes had mean *syn*- and *anti*-DP levels of 9 and 19 ng/g, respectively while the air concentrations were even lower (Cequier et al., 2014). High levels of DP were detected in ventilation dust from aircraft cabins produced between 1986 and 2008 with a range of 31 to 9600 ng/g for *anti*-DP (Allen et al., 2013). *Syn*- and *anti*-isomers of DP were detected in most indoor dust samples collected from three Antarctic research stations, with both isomers showing the highest concentrations at the Gabriel de Castilla, Spain (GCS) Station at 163 \pm 93.6 and 48.5 \pm 61.1 ng/g, respectively (Corsolini et al., 2021).

112. Ghelli et al. (2021) present a new review of DP and related compounds in food, including fish and seafood, dairy products, egg and egg products, meat and meat products, animal fat and vegetable oil, and other foods. From the analysis of the various articles, DP was quantified with values that would indicate slightly higher contamination levels in the fish and seafood category compared to other foods, see Table 15 in UNEP/POPS/POPRC.17/INF/9 for levels reported in food. In food products from Belgium, DP was found in 5% of the 1289 samples (firm cheese, quail eggs and pork, with mean DP levels of 339, 637 and 331 pg/g ww, respectively) (Poma et al., 2018). In 175 samples of 35 different food products from the retail market in the Republic of Korea, the mean DP was in the range from ND to 170 pg/g ww, and *syn*- and *anti*-DP were detected in 83.4 and 79.4% of the food products, respectively (Kim et al., 2014). Daily intake of DP was estimated to be 11.2 \times 10³ pg/day, which was 3 orders of magnitude higher than for other dechloranes (Dec 602, 603, 604 and Mirex). The highest level of DP was found in bovine liver (DP 170 pg/g ww) followed by spanish mackerel, oysters and dried anchovy with 135, 81 and 78 pg/g ww, respectively. However, in this study, DP was detected in all food products except onion. This result shows the wide-spread exposure of DP in the environment of Korea (Kim et al., 2014). In a study from Lebanon of 58 food samples representing fatty food groups, the estimated daily dietary intake for the adult population had lower and upper boundaries of 1344 and 1718 pg/day for DP (Malak et al., 2019). In a market basket study of 123 food samples from Japan, DP was detected in Groups III (sugar and confectionary), V (legumes and their products e.g. soy bean, tofu, fried bean curd, natto, chickpea), X (fish, shellfish and their products), and XI (meat and eggs) at concentrations of 3.3, 2.8, 1.9, and 1.5 pg/g ww, respectively (Kakimoto et al., 2014). The atmospheric concentration of DP in Osaka city was 11 pg/m³. The estimated daily intake of DP for an average Asian adult via inhalation and diet was (750 pg/day) approximately one percent of that estimated for Σ PBDE (62 ng/day) (Kakimoto et al., 2014). DP was frequently detected in a study

of US and Chinese baby food samples, including infant formula, cereal, and puree. The highest levels were found at 4,000 pg/g and 430 pg/g, in a baby formula sample from China and a cereal sample from the U.S., respectively. Median concentrations of total DP in U.S. baby food samples were 16 pg/g fresh weight in formula (n=12); 7.8 pg/g fresh weight in cereal (n=15); and 7.3 pg/g fresh weight in puree (n=8). Median concentrations of total DP in Chinese baby food samples were 20 pg/g fresh weight in formula (n=9); 5.9 pg/g fresh weight in cereal (n=12); and 5.8 pg/g fresh weight in puree (n=6) (Liu et al., 2014.)

113. In a study comparing human exposure at a production facility and an e-waste recycling facility in Southern China, high levels of DP were found in food products produced in proximity to the e-waste recycling site and production facility, where vegetables contained from 305 ng/g ww (paikchoi cabbage) to 2720 ng/g ww (green onion) DP (Wang et al., 2013a). Grains (common wheat, maize and soybeans) had levels of 1370, 760 and 498 ng/g ww, respectively, and common carp and other local fish species had levels from 56.8 to 1110 ng/g ww (Wang et al., 2013a). Under the assumptions that consumed food was produced locally estimated dietary intake dose accounted for more than 99% and 93% of the daily intake to people from the production site and e-waste recycling site, respectively (Wang et al., 2013a). For the production site, the estimated dietary exposure dose was 1.1×10^{-2} mg/kg/d for workers, residents and children and the estimated dietary exposure dose was lower for the e-waste recycling site at 3.0×10^{-5} mg/kg/d (Wang et al., 2013a). Total estimated exposure dose (dietary, dermal, inhalation) in people from the e-waste recycling site was 0.03×10^{-3} mg/kg/d, which is 300 times lower than at the manufacturing site (Wang et al., 2013a). In another study from an e-waste recycling area in China, free-range chicken eggs harvested in 2010, 2013 and 2016 contained high levels of FRs, including DP. No temporal trend was observed for DP levels. The highest levels of DP were observed in eggs from 2016, mean and SD was 5413 ± 1978 ng/g lw (Huang et al., 2018).

114. DP has been detected in human serum from several countries. In a study from France mean Σ PBDEs (BDE-47, -99, -100, -153 and -154) levels (4.32 ± 2.99 ng/g lw) were in the range typical of Western Europe levels, but lower than the mean Σ 5 dechlorane compounds (Dechlorane 603>DP>Mirex>Dechlorane 602>Chlordene Plus) levels (6.24 ± 4.16 ng/g lw) (Brasseur et al., 2014). Levels of DP observed were lower in serum from Canada, France, Korea and Norway compared to occupationally exposed people in China with median of 2.39, 1.20, 0.73, 1.3 and 42.6 ng/g lw, respectively (Zhou et al., 2014; Brasseur et al., 2014; Kim et al., 2016; Cequier et al., 2015; Ren et al., 2009). In a study from China analyzing pooled human serum samples from different age groups, including 305 subjects from the south coast of Laizhou Bay, Shandong Province, China, there was no relationship between DP levels and age. However, the younger volunteers, age 20 to 29 years, had the highest serum concentrations with mean DP around 7 ng/g lw compared to mean DP for all groups 3.6 ng/g lw (Wang et al., 2014). A time trend of Dechlorane compounds was investigated in seven pooled serum samples from Korean adults to cover the period from 2006 to 2013. No distinct trend was observed for Dechlorane compounds (Kim et al., 2016). In plasma samples of German young adults from the environmental specimen bank collected from 1995 to 2017, no significant trend was observed for DP (Fromme et al., 2020).

115. High levels of DP have been observed from occupational exposure at a DP manufacturing plant compared to nearby areas in eastern China. The DP concentrations ranged from 89.8 to 2958 ng/g lw in whole blood and 4.08 to 2159 ng/g dw in hair. For the workers directly engaged in the DP manufacturing process, their DP levels were significantly higher than in most individuals of the two control groups (Zhang et al., 2013). In another study, DP serum levels in 70 occupationally exposed workers from an e-waste recycling workshops and control group of 3 residents of an urban area in South China were measured. The DP levels were significantly higher in the workers (22–2200 ng/g with median of 150 ng/g lw) than in the urban residents (2.7–91 ng/g with median of 4.6 ng/g lw). The DP concentrations in females were found to be associated with their age but this relationship was not found for males (Yan et al., 2012). Hair samples from female workers at e-waste dismantling facilities in South China contained median and mean concentration of 23.5 ng/g and 61.3 ng/g DP respectively (Qiao et al., 2019). A study of 15 occupationally exposed workers at e-waste recycling sites in Bangladesh found a median DP exposure of 2.3 ng/h/silicone wristband, approximately 130 times higher than non-occupationally exposed US residents (Wang et al., 2020b).

116. In humans, DP has been detected in cord serum (Ben et al., 2014) and in human milk (Siddique et al., 2012) showing transfer to offspring at different developmental stages. Maternal transfer of bioaccumulative substances *in utero* represents a potential risk to embryonic development and may represent the largest source of FRs input to offspring during the first few years of life. In 450 human milk samples from Norway, the Netherlands and Slovakia, DP had detection frequencies in the range of 3 to 9% and 20 to 26% for *syn*- and *anti*-DP, respectively. The concentrations of *syn*-DP ranged between 0.05 and 2.85 ng/g lw and of *anti*-DP between 0.004 and 1.60 ng/g lw (Čechová et al., 2017). Higher detection frequency was observed for 87 human milk samples from Canada (76 and 91% for *syn*- and *anti*-DP, respectively), with mean *syn*- and *anti*-DP concentrations of 0.27 and 0.7 ng/g lw (Siddique et al., 2012). In another Canadian study of 105 milk and 102 maternal serum samples, *syn*- and *anti*-DP detection frequencies were 40 and 50% for milk samples and 77 and 78% for maternal serum samples, respectively (Zhou et al., 2014). However, concentrations of DP in milk samples were lower than in the previous study from Canada (i.e. Siddique et al., 2012). Total DP in maternal serum was 2.37 ng/g lw, which was lower than previously found in two Chinese studies (Ben et al., 2013; Ren et al., 2009), where the median value of total DP in maternal serum

samples of mothers who were not exposed to e-waste recycling activities was 13.7 ng/g lw in one study (Ren et al., 2009), and 4.0 ng/g lw in the other study (Ben et al., 2013). The total DP levels in milk samples were also higher in the Chinese study (Ben et al., 2013) with a median of 2.19 ng/g lw. *Syn*-DP and *anti*-DP were detected in all milk (n = 44) and serum samples (n = 45) (Ben et al., 2013). The Cl₁₁-DP detection frequency in breast milk and serum was 45% and 84%, respectively, and DP levels in blood and milk from residents who had been living in Taizhou for >20 years (R(20) group) were significantly higher than those who had lived in that city for <3 years (R(3) group) (Ben et al., 2013). For overview see Table 16 in UNEP/POPS/POPRC.17/INF/9.

117. In another publication from the same study of 72 residents of the e-waste recycling area of Wenling, China, DP was detected in placenta and cord serum indicating prenatal exposure of DP across the placenta (Ben et al., 2014). In contrast to human milk samples which have been reported to have a f_{anti} ratio close to that of commercial DP mixtures (0.6–0.8) (Zhou et al., 2014; Siddique et al., 2012; Ben et al., 2013), the concentration ratio between the cord serum and maternal serum was estimated to be 0.45 for *syn*- and 0.35 for *anti*-DP, suggesting the placenta partially limited DP transfer with a greater extent for *anti*-DP (Ben et al., 2014).

2.4 Hazard assessment for endpoint of concern

118. Available studies show that DP has the potential to elicit physiological changes or toxic effects in different organisms. Oxidative stress has been observed after exposure to DP in all organisms tested; e.g. freshwater protozoan (Dou et al., 2015), macroalgae (Gong et al., 2013;2018), fish (Chen et al., 2017; Hang et al., 2013; Kang et al., 2016), marine bivalves (Barón et al., 2016, Gagne et al., 2017), earthworm (Zhang et al., 2014; Yang et al., 2016b), birds (Li et al., 2013a) and mice (Wu et al., 2012). The organism may be able to handle oxidative stress/damage by activating/ upregulating antioxidant defence and repair mechanisms. However, reactive oxygen species are important signalling molecules that could trigger unwanted effects by activating signalling cascades regulating cell death and cell survival and by causing damage to DNA, protein, telomeres, cells, tissues etc. Oxidative stress is linked to aging and disease. Li et al. (2020a) reported a positive association between DNA-damage and morphological deformities in carp embryos exposed to DP. As further described below, reported effects include oxidative stress to most tested organisms, neurodevelopmental toxicity in zebrafish and histochemical alterations in juvenile carp brain (Chen et al., 2017; Li et al., 2019b). In addition, studies have indicated effects to the sex and thyroid related genes in zebrafish (Kang et al., 2016) and for thyroid hormone pathway mammalian cells *in vitro* (Zhu et al., 2022). Some epidemiological investigations found associations for DP and thyroid pathways (Ben et al., 2014; Guo et al. 2018b; 2019). Alteration in gene expression indicating immune modulating effects in carp has been observed (Li et al., 2019b). In mammalian and human cells, DP has been shown to activate peroxisome proliferator-activated receptor gamma and inhibit insulin signalling *in vitro* (Peshdary 2019; 2020). DP has been shown to impair liver function by causing liver damages related to inflammatory infiltration and hepatocellular swelling in mice at high dose (Wu et al., 2012; 2013). A few studies in rodents and mammalian cells, indicate that DP at low and environmentally relevant doses promotes adipose tissue dysfunction, and alters insulin secretion, gut microbiota and biomarkers associated with metabolic disorder and development of type 2 diabetes (Peshdary et al., 2020; Zhang et al., 2020; Wu et al., 2021). However, it should be mentioned that the database on toxicity is limited, and long-term studies are largely lacking.

2.4.1 Toxicity to aquatic organisms

119. DP has very low water solubility (< 44 ng/L, see Table 2) and will partition to particles, sediment and biota in the aquatic environment. Zhen et al., (2018) showed that total DP levels in natural water could be higher than just the dissolved concentration due to DP adsorbed to particulate matter, e.g. in Laizhou Bay water, median DP was 11.7 ng/L for sum filtered and particulate, the range of DP in filtered and particulate water samples ranged from not detected (nd) to 10 ng/L and 0.890 to 346 ng/L, respectively (Zhen et al., 2018). For DP, results from most available empirical aquatic toxicity studies are associated with some uncertainty concerning dosing, mainly because test concentrations exceeded the DP water solubility limit. However, all studies use solvent as a vehicle, which increases the solubility of DP. Chen et al., (2019) observed a measured water concentration of 55 µg/L with a nominal concentration of 60 µg/L using DMSO as solvent. Detailed information of conditions in the aquatic toxicity studies and measured concentrations in water are provided in Table 17 and 10, respectively, in UNEP/POPS/POPRC.17/INF/9.

120. DPs had a significant inhibitory effect on sludge anaerobic fermentation to generate the intermediate valuable product short-chain fatty acids (SCFA). The effects were observed at two environmental background doses of DPs 30.3±1.2, 303.4±13.1 mg/kg TSS, and the two doses that covered potential high levels expected from industrial workshop wastewater (606.8±29.5, and 3034.1±101.7 mg/kg TSS). The underlying mechanism investigation indicated DPs promoted the disintegration of sludge but inhibited the process of hydrolysis and acidification. The analysis of microbial community characteristics indicated that DPs reduced the level of *Firmicutes* and *Actinobacteriathae*, which were the key acid producing bacteria. At the genus level, DPs reduced the relative abundance of *Proteiniclasticum* and *Mycobacterium* (Zhang et al., 2021).

121. Chronic exposure studies for benthic and sediment dwelling organisms are lacking. Recognizing the lack of effects data for DP in important compartments, such as soil and sediment, the use of analogue data can be considered as an approach to addressing key gaps in the hazard profile. In selecting analogues, factors such as structure, reactivity, metabolism and bioavailability can be considered. The Canadian screening assessment used toxicity data on the DP analogue chlordane, due to its structural and functional similarities to DP (OECD QSAR Toolbox, 2016), to fill the data-gap as a worst-case-scenario for sediment dwelling organisms and estimated a predicted no effect concentration (PNEC) of 0.0129 mg/kg dw for DP (Canada, 2019a).

122. Two studies on aquatic algae have investigated effects on photosynthetic and antioxidant endpoints at low doses of DP (Gong et al., 2013; 2018). The freshwater algae *P. subcapitata* was exposed to nominal DP concentrations of 13.51, 135.1 and 1 351 ng/L for up to 72 h (Gong et al., 2013). No significant effects were observed on esterase activity or reactive oxygen species (ROS) levels at 72 h, although transient increases were observed at earlier timepoints (Gong et al., 2013). At the highest dose (135.1 and 1 351 ng/L), average chlorophyll a content was significantly increased at 48 and 72 h (Gong et al., 2013). No measurement of growth was made, so the significance of the chlorophyll a content increase is not known. The authors concluded that DP showed low toxicity and had a marginal effect on *P. subcapitata*. In juvenile marine green macroalgae *U. pertusa*, the uptake of DP affected physiological responses of photosynthesis and caused oxidative stress (Gong et al., 2018). Following a 21-day exposure to 10^{-8} mol DP/L and 21-day depuration, *anti*-DP was prone to bioaccumulate. Physiological effects were observed after 14 days exposure to DP doses of 10^{-8} to 10^{-6} mol/L (equal to ~6.53 to 653 µg/L). Antioxidant enzyme activities (superoxide dismutase (SOD) and catalase), oxidative damage to lipids (malondialdehyde, MDA) and chlorophyll fluorescence parameters were affected in a dose and time dependent manner after 1, 7 and 14 days of DP exposure. The effect on photosynthesis and SOD were significantly different from controls at all concentrations tested, while effects on catalase and MDA were significant for the two highest doses. The authors concluded that DP affects photosynthesis in the marine algae, as observed by a low rate of light energy utilization, and that this effect may be ascribed to oxidative damage caused by the uptake of DP (Gong et al., 2018). Growth was not measured in the test, so the significance of the observed effect is not known. However, it is notable that marine micro and macroalgae constitute an important part of the marine food web as being primary producers and food source for a number of species. In addition, macroalgal forests provide habitat for many species.

123. Mediterranean mussels were exposed to DP concentrations of 0, 5.6, 56, and 100 µg/L for 6 days by spiked agarose. Concentrations measured in water immediately after dosing were 0, 0.4 ± 0.3 , 0.3 ± 0.2 and 0.7 ± 0.5 µg/L, and measured levels in mussel after 23 h exposure were 4.7 ± 3.1 , 8.8 ± 2.1 and 21 ± 9.1 µg/mussel, respectively (Barón et al., 2016). DNA strand-breaks were observed in hemocytes at all doses. No clear dose-response was observed, although all doses were significantly different from the negative control; 13, 23 and 18%, respectively. Increased micronuclei formation was observed only for highest dose (100 µg/L) tested (Barón et al., 2016). In a more long-term study, fasting blue mussels were exposed to nominal concentrations of 0, 0.001, 0.01, 0.1 and 1.0 µg DP/L for 29 days, which resulted in ΣDP tissue concentrations of 0, 0.12, 0.98, 7.26 and 57.8 ng/g ww, respectively. Following DP exposure, no histopathological lesions were found in gonads, and no change in hemocyte DNA strand breakage, phagocytosis rate, and viability was observed. Lipid peroxidation in gills was found to increase by 82% and 67% compared to control at the 0.01 and 1.0 µg DP/L dose, respectively (only significant for 0.01 µg/L), while cyclooxygenase activity (COX) was significantly decreased by 44% compared to control at the 1 µg/L dose (Gagné et al., 2017). The relationship between endpoints and survival are not known. The effect on COX could have an impact on the control of spawning process and thus reproduction in blue mussels (Matsutani and Nomura, 1987). These two studies indicate that DP exposure cause oxidative stress at low µg/L concentrations.

124. Acute toxicity fish studies reported (HPV summaries and referred to by ECHA (2017b)) were not considered reliable. However, no mortality was observed in a bioaccumulation study of the freshwater fish species *Lepomis macrochirus* exposed to 0.1 mg/L DP (using acetone as solvent at 0.001%) for 30 days (ECHA 2017b), implying a NOEC > 0.102 mg/L, 30 d.

125. Four studies with embryo/larval zebrafish with waterborne DP exposure have been identified (Hang et al., 2013; Noyes et al., 2015; Chen et al., 2017; Kang et al., 2016). No significant adverse effects on hatchability, survival or malformation were seen in the Noyes et al. (2015), Chen et al. (2017) and Kang et al. (2016) studies following exposure to DP at a nominal concentration of up to 4.18 mg/L, when zebrafish embryos were exposed in the blastula or early gastrula phase (4-6 hpf). Kang et al. (2016) exposed zebrafish embryos to nominal DP concentrations of 0.4, 0.8 or 1.6 mg/L from 4 to 144 hours post fertilization (hpf). Measured water concentrations of DP were 140, 248, 267 µg/L DP at 0 h.

126. In a study conducted by Li et al. (2020a), 300 carp embryos exposed to DP with nominal concentrations of 30, 60, and 120 µg/L (using 0.1% DMSO as solvent) from 3 to 120 hpf displayed significant time- and dose-dependent increased mortality rate, delayed hatching time, reduced hatching rate, and decreased body length. At 120 hpf, significant dose-dependent increased DNA damage was observed for all doses. In the 120 µg/L group, the mortality rate increased to more than 50% between 48 and 72 hpf and continued to increase to between 70 and 80% at 120 hpf. At 96 hpf, mortality rate was approximately 30% for the 60 µg/L group while the rate was below 10% for the control.

Morphological deformities were significantly increased compared to the control at 120 hpf at all doses. DNA damage measured as % tail DNA at 24 hpf increased significantly from below 10 to approximately 30, 45, and 55 % for the solvent control, 30, 60, and 120 µg/L group, respectively. A positive association was observed between DNA-damage and morphological deformities. Observations were supported by alteration in gene-expression for key morphology genes and antioxidant markers. Further, in a less well documented study, Hang et al. (2013, conference paper), reported significant time- and dose-dependent increases in malformations (spine side curve, cardiac edema, tail deformation) in zebrafish embryos exposed to DP at nominal concentrations of 0.037 to 0.37 mg/L from 8 hpf (late gastrula phase) up to 7 days (168 hpf). Due to limited documentation, the results from Hang et al. (2013) should be treated with caution.

127. Short-term exposure studies with embryo/larval zebrafish suggest that DP can induce oxidative stress and neurobehavioral changes (Hang et al., 2013; Noyes et al., 2015; Chen et al., 2017). Noyes et al. (2015) observed a significant hyperactive response to dark-stimuli-activation compared to controls at 120 hpf after exposure to nominal concentration of 6.4 µM DP (4.18 mg/L). The level of response was much lower upon exposure to DP than phosphate ester and chlorinated phosphate ester flame retardants in the same test system. Chen et al., (2017) observed significant increase in spontaneous movements at 24 hpf for all nominal doses (15, 30 and 60 µg/L), dose- and time-reduced distances of swimming after touch-induced stimuli, significant decreased free-swimming activity and reduced swimming speed during each dark and light period. These neurobehavioral changes may be linked to axonal and muscular lesions. DP significantly inhibited primary motor neuron axonal growth and induced cell apoptosis and lesions in muscle fibres of the tail region of larvae in a dose dependent manner at 96 hpf. Axonal growth-related gene-expression (α 1-tubulin and gap43) was significantly increased in the highest dose (60 µg/L), oxidative stress markers such as ROS and MDA as well as messenger ribonucleic acid (mRNA) levels of apoptosis related genes were increased at the two highest doses (30 µg/L and 60 µg/L) (Chen et al., 2017).

128. Co-exposure of zebrafish embryos to nominal concentrations of DP (60 µg/L) and/or 3-methyl phenanthren (3-MP) (5 or 20 µg/L) at 6 to 120 hpf resulted in elevated bioaccumulation of both compounds and synergistic effects on axonal growth reduction, apoptotic markers in muscle and brain Ca^{2+} homeostasis as well as some of the neurobehavioral tests (Chen et al., 2019). ROS levels in whole fish were significantly elevated compared to control from all exposures at 96 hpf, but not significantly altered by co-exposure compared to single exposure. As in the previous study (Chen et al., 2017), a significant increase in spontaneous movement occurred at 24 hpf with 60 µg/L DP, but co-exposure with 5 or 20 µg/L 3-MP gave a significant reduction compared to control. DP and 3-MP co-exposure showed a synergistic or additive effect in reduction of swimming distances after touch-induced stimuli and free-swimming activity. Furthermore, synergistic or additive effect were observed for reduction in ventral axonal length growth, increased number of apoptotic cells in the tail region and induction of axon related gene expression. Brain Ca^{2+} homeostasis was investigated at 96 hpf, and a significant synergistic increase was observed in brain intracellular Ca^{2+} levels, and a synergistic reduction in Ca^{2+} ATPase activity was observed after co-exposure. Furthermore, a synergistic effect was observed in reduced expression of genes related to Ca^{2+} homeostasis. Relative to DP exposure alone, co-exposure with low and high-dose 3-MP increased the accumulation of DP by 14 and 82%, respectively. The larval body burden of DP (60 µg/L) was 583.2 ± 33.5 ng/g ww while co-exposure with 5 and 20 µg/L 3-MP increased DP levels to 665 ± 33.5 and 1061 ± 85.7 ng/g ww, respectively. Relative to 3-MP alone, co-exposure with DP (60 µg/L) increased 3-MP levels by 45 and 47% for the 5 and 20 µg/L 3-MP groups, respectively (Chen et al., 2019). Elevated bioaccumulation of DP resulting from mixture exposure may represent a significant contribution of the synergistic effects observed in combined chemical exposure. Total organic carbon (TOC) content was not reported, which could affect DP concentration levels.

129. Studies also show that DP can cross the blood-brain barrier in fish (Zhang et al., 2011), and *anti*-DP had a high persistence in the brain compared to the liver (Zhang et al., 2011). Stress responses and histopathology changes of the brain and liver were observed in juvenile common carp (n= 30 per group) exposed to nominal concentrations of 0, 30, 60 or 120 µg/L DP for 1, 15 or 30 days (Li et al., 2019b). Tissue morphology (cell abnormality rate) was significantly changed in the groups exposed for 60 and 120 µg/L for 15 and 30 days. In liver, unclear contours, vacuolisation and nuclear lysis were observed, and in the brain, abnormalities were observed in the granular layers, nuclear cell structures, disordered arrangement, microthrombic red blood cells, increased numbers of glia cells and nodulations. Liver SOD, glutathione (GSH) and MDA activity levels decreased with increasing dose and exposure time while the opposite pattern was observed in the brain (Li et al., 2019b). DP also altered expression of immune regulating genes in the brain and liver. CYP1B1 gene expression was significantly upregulated at all timepoints and concentrations in both the liver and brain. In addition, CYP2B and CYP3A1 and the apoptosis related factors bax and bcl-2 were altered in liver. Both the pro-inflammatory cytokines IL-6 and IL-1 β and the anti-inflammatory cytokine IL-10 transcription levels were significantly increased compared to control at all doses at 30 days in liver, while in the brain, IL-6 was significantly upregulated with the two highest concentrations, IL-1 β with highest concentration only, and IL-10 with 30 µg/L (Li et al., 2019b). This is the first study indicating potential immune modulating effects of DP. These results indicate that DP exposure perturbs metabolism in the liver and brain, inhibits antioxidant enzyme activity, increases lipidperoxidation, promotes inflammation and induces cell apoptosis in juvenile carp (Li et al., 2019b).

130. In another study in juvenile common carp exposed to nominal concentrations of 0, 30, 60 or 120 µg/L DP for 4 weeks, DP was shown to negatively affect intestinal barrier, increased the relative abundance of pathogenic bacteria and changed the balance of intestinal flora (Li et al., 2020b). mRNA levels of genes involved in tight binding protein genes in the gut of common carp were significantly reduced by all exposure concentrations and histomorphology, indicated that DP evidently shortened the intestinal folds and damaged the intestinal epithelium layer (Li et al., 2020b). Effects on growth or mortality as a result of DP exposure were not reported for these two studies (Li et al., 2019b, 2020b), and doses were chosen based on results from Chen et al., (2017) and Kang et al., (2016), where no developmental abnormalities were observed for measured doses up to 267 µg/L in those studies.

131. In juvenile Chinese sturgeon treated with DP at doses of 1, 10, and 100 mg/kg ww for 14 days via a single intraperitoneal dose, liver proteomics indicate that DP had effects on the generalized stress response, small G-protein signal cascades, Ca²⁺ signalling pathway and metabolic process, and may induce apoptosis in the liver (Liang et al., 2014). Intraperitoneal exposure route is considered not physiologically relevant and therefore the results have the limitations.

132. In zebrafish receiving doses of 0, 0.3, 1, or 3 µg/g ww of DP by gavage on day 0 and 2, measured tissue concentrations were 5, 30, 44 and 420 ng/g ww, respectively, on day 6. An increase of catalase activity was observed in the liver, indicating oxidative stress response to the liver on day 6 (Kang et al., 2016). Transcriptional responses of both thyroid and sex hormone related genes in the brain were altered, suggesting possible thyroid and sex hormone disrupting potentials of DP (Kang et al., 2016). On day 6, significant induction of CYP19b (brain type of aromatase) was observed at all doses at body residual levels that were environmentally relevant, other TH related genes were not altered. In addition, the brain estrogen receptor, alpha mRNA, was significant elevated at the lowest dose (0.3 µg/g) only. Estrogen and androgen receptor genes and steroidogenic genes were evaluated in testis, but no significant alteration in gene transcription was observed. Plasma thyroxine (T4) concentrations increased in a dose-dependent manner but were not statistically significant different from the control. However, observation was supported by the up-regulation of corticotropin releasing hormone and TSH-β genes in brain at the highest dose (3 µg/g ww), which are both involved in the fine-tuning of the thyroid hormone-pathways in non-mammalian vertebrates (De Groef et al., 2006).

133. Maternal uptake and transfer of DP has been seen in several species of fish (e.g. Wu et al., 2013; Sühring et al., 2015; Peng et al., 2012; Zhang et al., 2011; Zeng et al., 2014) and in frogs (Wu et al., 2018). DP and other FRs were found in developing embryos of female sharks, demonstrating maternal transfer *in utero* (Marler et al., 2018).

2.4.2 Toxicity in terrestrial organisms

134. Oxidative stress responses and neurotoxicity have been observed in earthworms (Zhang et al., 2014; Yang et al., 2016b). No acute toxicity or significant change in body weight were observed in earthworms exposed to DP at nominal doses of 0.1, 0.5, 6.25 and 12.5 mg/kg for 28 days (Yang et al., 2016) or up to 50 mg/kg for 14 days (Zhang et al., 2014). Oxidative stress was indicated by alteration in markers such as SOD, MDA, glutathione, glutathione-peroxidase, catalase and 8-hydroxy-2-deoxyguanosine (8OHdG) in tissue as well as tail DNA in comet assays of isolated coelomocytes. Acetylcholinesterase (AChE) and cellulase activity of earthworms was significantly reduced even by the low dose indicating neurotoxic effects in earthworms (Yang et al., 2016). The overall 28-d no-effect concentration (NOEC) for these biomarker responses is <0.1 mg/kg (ECHA, 2017b). The linkage between these biomarkers and adverse effects is not established. No information regarding long term adverse effects in earthworms is available.

135. Crump et al. (2011) studied concentration-dependent effects of DP using *in vivo* and *in ovo* toxicity approaches in domestic chicken embryonic hepatocytes and chicken embryos. DP was injected into eggs prior to incubation and no overt toxic effects were observed up to the maximum dose of 3 µM in hepatocytes, and up to the highest nominal dose of 500 ng/g/egg for pipping success of the young.

136. Li et al. (2013a) studied the effects of DP on male common quails continuously exposed to commercial DP by gavage for 90 days at dose concentrations ranging from 1 to 100 mg/kg bw/d. Mortality, body and liver weight was not altered in any of the exposure groups. Liver enzyme activity and oxidative stress were measured. The authors reported DP effects on some measures of enzyme activity (e.g., significant decrease of 7-pentoxoresorufin-O-demethylase (PROD) in all exposed groups relative to the control, significant increase in erythromycin N-demethylase (ERND) and the antioxidant enzyme catalase in the high-dose exposed groups relative to control). Furthermore, the study found DP was more prone to accumulate in liver (vs. serum, muscle), and *syn*-DP accumulated (vs. *anti*-DP) in the two high-exposure groups.

137. In a field study of northern fulmars from the Faroe Islands, liver DP concentration was positively correlated with plasma TT4:FT4 ratios, and UDP-glucuronyltransferase-1 mRNA levels. The study indicates an association between liver levels of DP and thyroid homeostasis in northern fulmar but the low levels of DP in liver and the low sample size suggest that these associations should be interpreted with caution (n=9) (Mortensen et al., 2022).

138. The Canadian screening assessment used toxicity data from Mirex, also based on its structural and functional similarities to DP, as a worst-case scenario to fill the data-gap for terrestrial (plant) organisms, with estimated DP predicted no effect concentrations (PNEC) of 0.075 mg/kg dw (Canada, 2019a).

2.4.3 Mammalian toxicology

139. Available assessments and laboratory studies with mammals suggests that, DP is not carcinogenic, mutagenic or toxic to reproduction (as reviewed in ECHA, 2017b; Canada 2019a), although the data are limited. In their alternative assessments for decaBDE, in reference to DP, US EPA 2014 stated "There is potential for carcinogenicity based on analogy to chlordane and decaBDE, the latter for expression of adverse effects in longer term studies." (US EPA, 2014). Other biomarkers of exposure and/or potential adverse effects in mammals have been reported such as oxidative stress, liver impairment and endocrine effects as described below.

140. In terms of *in vitro*, no mutagenic and no toxic effect of DP was observed in a standard Ames test (Patil 2019) on five different strains of *Salmonella typhimurium* nor in a micronucleus test on human blood lymphocytes (Tekale 2019), as observed in experiments described in OxyChem 2004b. No mutagenic and no toxic effect of DP was observed in a standard Ames plate incorporation test in *Salmonella typhimurium* TA 98, TA 100, TA 1535, TA 1537, and TA 1538 with and without metabolic activation by rat liver S9-mix at concentrations up to 10,000 µg/plate. An additional test (Patil 2019) was performed according to OECD 471 with the same results also for strain TA 102 and the former assessed strains TA1537, TA1535, TA98, TA100. The *in vitro* test on mammalian cells also did not show any potential to induce micronuclei or clastogenic or aneuploidic potential in isolated cultured human peripheral blood lymphocytes. Details of these experiments are available at ECHA 2021b. One *in vivo* genotoxicity study was identified. Mice were orally administered 0, 500, 2000 or 5000 mg/kg-bw/day of DP via gavage for 10 days (Wu et al. 2012). Liver samples were collected for a comet assay and the genotoxicity result was negative.

141. Acute toxicity studies in experimental animals suggest low concern for acute toxicity via the oral, inhalation and dermal routes of exposure. No adverse health effects were observed in any of the identified repeated-dose oral toxicity studies, testing dose levels up to 5000 mg/kg-bw/day for 10 or 28 days (as summarized in ECHA, 2017b; Canada 2019a; OxyChem 2004b and further described below). Toxicokinetics studies showed limited or low absorption from the intestine (Brock et al., 2020; ECHA, 2017b). However, there are some data gaps, for example, there are no long-term studies exceeding 90 days, which might be important given the apparently slow uptake and long elimination half-life of the substance. Therefore, in an alternative assessment of decaBDE, the potential for systemic effects of DP for repeated dose studies was rated as moderate based on the high potential for bioaccumulation and potential for expression of adverse effects in longer term studies by analogy to decaBDE (US EPA, 2014). The dosing vehicles might also limit exposure (e.g. due to the presence of undissolved micro-crystals), such that the high doses might not truly reflect the degree of exposure of the organisms (ECHA, 2017b). The 5000 mg/kg bw dose was described as particularly viscous (Brock et al., 2010), and is an extremely high dose (5 times the limit dose of the OECD test guideline).

142. Some of the earlier studies provided in Oxychem (2004b) indicate some liver and ovary effects. In a 13-week oral feeding study in rats, absolute and relative liver weight was significantly increased in males only, at 500 mg/kg, but no associated histopathology was observed (Oscarson, 1975, as described in Oxychem (2004b)). In a repeated-dose inhalation toxicity study, rats were exposed to 0, 640 or 1524 mg DP as dust/m³, 6 h per day, 5 days per week for 28 days (Bishop 1975). Significant increase in absolute and relative liver weight was observed at both doses. Corresponding hepatocytomegaly (swelling of liver cells with signs of cytotoxicity and necrosis) of centrolobular hepatocytes were observed for male rats at both doses and in some females at the higher dose. Effects on lung tissues were observed as increased number of macrophages in alveoli, and significant increase in absolute lung weight was observed for both doses in females and highest dose for males. A lowest-observed-adverse-effect concentration (LOAEC) of 640 mg/m³ was identified (Bishop 1975, as described in Oxychem 2004b). In a repeat-dermal study, male and female rabbits were dermally exposed to DP in 3% aqueous methylcellulose at 0, 500 or 2000 mg/kg bw/d, for 5 days per week, for 4 weeks on 20% of the total body surface (Trzyna 1975). None to minimal erythema occurred at the application site. In females, significant dose-related decrease in absolute and relative (to body and brain weight) ovary weight was observed at both doses, and a decrease in absolute and relative liver weight was observed at the highest dose. However, no other histopathology effects were observed in any of the organs (Trzyna 1975, as described in OxyChem 2004b). However, reliability of these studies is uncertain as discussed in ECHA (2017b) and indicated in the HPV summaries. The description of the studies is also limited.

143. In a combined repeated-dose (RDT)/ developmental/reproductive toxicity (DART) screening test, conducted according to OECD guideline 422, Crl:CD (SD) rats were exposed to DP in corn oil at 0, 750, 1500 or 5000 mg/kg-bw/day by oral gavage (Brock et al. 2010). For the RDT phase, animals (10/sex/dose) were treated for 28 days, for the DART phase animals (20/sex/dose) were treated for 63 and 64 days. No treatment-related effects were observed on clinical signs of toxicity, body weights, food consumption, neurobehavioral and functional observational battery evaluations. No effects were observed on haematology, urinalysis, coagulation or clinical chemistry parameters. No dose-response related changes in organ weights (heart, liver, testes, ovaries, and thyroid/parathyroid glands) were observed. Mortalities and lesions, including histopathology, were observed across all dose groups including controls,

which were linked to gavage administration errors as indicated by gavage injury and administration of test substance in the thoracic/pericardial cavity. As a consequence, it is not clear whether dosing was correct. The 5000 mg/kg vehicle in corn oil was described as particularly viscous. The authors identified a no-observable-effect level (NOEL) of 5000 mg/kg-bw/day for both the RDT and DART screening test. However, some effects were observed but considered not treatment relevant by the authors due to lack of dose-response relationship, which could be due to gavage-related errors. A significant increase in absolute and relative ovary weight was observed in the 1500 mg/kg bw/d group only (38.9 and 38.4%, respectively), and a significant increase in absolute and relative thyroid/parathyroid weight was observed in females in the 750 mg/kg bw/d group only (21.8 and 30.7%, respectively). A non-significant increase in the number of mean early resorption sites (1.8) and postimplantation loss index (13.90%) in the 1500 mg DP/kg bw/d group compared to control (0.8 and 5.39%, respectively) were observed. Furthermore, pup body weight was significantly increased at doses ≥ 1500 mg/kg bw/d. The mean pup weight was increased by 3, 15 and 13% compared to control for the 750, 1500 and 5000 mg/kg/d groups, however these results were considered unrelated to treatment by the study authors. This increased pup weight may be a result of DP's ability to activate the peroxisome proliferator-activated receptor PPAR- γ as described in the paragraph below (Peshdary et al., 2019) as activation of this signal pathway has been linked to pregnancy/fetal outcomes by strengthening of uterus function (Hewitt et al., 2006; Díaz et al., 2012;), but also with effects on the thyroid axis.

144. Wu et al. (2012) report liver impairments in male mice at sublethal doses following a 10-day oral gavage exposure at 500, 2000 or 5000 mg/kg-bw per day. Oxidative stress and altered metabolism was observed at all doses tested. Relative liver weight was significantly increased (15%) in the 2000 mg/kg group only. Oxidative stress to the liver was shown by significant increase in SOD activity and the oxidative DNA-damage marker 8OHdG at all doses, as well as increased catalase (CAT) activity at 2000 mg/kg. However, no increase in DNA strand-breaks was observed in a comet assay. Metabolite analysis indicates that DP alters hepatic carbohydrate-, lipid and fat-, nucleotide- and amino acid metabolism. 18 metabolites were significantly altered with fold change $\geq \pm 1.5$ fold compared to control. Microarray profile of gene expression alterations supported the observations from metabolite analysis and further showed significant alterations in genes involved in signal transduction, endocrine pathways (insulin-, adipocytokine- and PPAR signalling), signal molecules, Gap junction, p53 signaling- and immune system pathways (Wu et al., 2012). Interestingly, Dechlorane 602, which has structural similarities with DP, (see Table 2 in UNEP/POPS/POPRC.17/INF/9), has been shown to alter immune response by dysregulating T-helper-1 and -2 balance in mice and altering immune and transmitter-related metabolism after exposure at low dose (10 $\mu\text{g}/\text{kg}$ bw for 7 days) (Feng et al., 2016; Tao et al., 2018). Furthermore, a medical study by Zhou et al. (2021) found that Dechlorane 602 exacerbates house-dust mite-induced airway inflammation in two mouse models and highlighted some of the biochemical pathways involved. However, while there is some evidence for potential liver impairment, the significance of these findings are unclear (ECHA, 2017b).

145. In a 90-day oral study with rats exposed to DP at 0, 1, 10 or 100 mg/kg/d, DP preferentially accumulated in liver with *syn*-DP as the dominating stereoisomer (Li et al., 2013b). Clinical serum parameters such as alanine aminotransferase, alkaline phosphatase and total bile acids were reduced significantly at highest dose, and serum glucose significantly increased. In the liver, gene-expression of several key enzymes were altered such as n-acetyltransferase, sulfotransferases and CYP2B1 (Li et al., 2013b). However, no significant changes in absolute body or liver weight or liver histopathology were observed. A non-significant increase in thyroid stimulating hormone (TSH) was also observed but other serum thyroid hormones were measured and unaltered. It should be noted that background contaminations gave 100% detection of DP in the control tissues with increasing levels during the experiment (at day 90, levels in serum was 45 ng/ml, muscle and liver levels were 200 and 3100 ng/g lw, respectively), which may have impacted the dose-responses on endpoints such as clinical serum parameters, serum thyroid hormones and liver mRNA expressions.

146. Recently two studies in rodents indicate that low and environmentally relevant doses of DP may alter adipocytes and gut microbiota in ways that are associated with increased risk of metabolic disorder and development of type 2 diabetes (Peshdary et al., 2020; Zhang et al., 2020). In mice, exposure to DP (10, 100 or 1000 $\mu\text{g}/\text{kg}/\text{d}$) for 28 days by gavage increased serum insulin and amylase levels (at 10 and 100 $\mu\text{g}/\text{kg}/\text{d}$) and altered lipid tissue. At 1000 $\mu\text{g}/\text{kg}$, serum lipase was significantly decreased after 28 days. Insulin levels were increased significantly in the 10 and 100 $\mu\text{g}/\text{kg}/\text{d}$ group compared to vehicle control after 28 days, although glucose levels were not affected. DP promoted white adipocyte tissue hypertrophy and dysfunction, which was characterized by increased adipocyte size and elevated serum levels of leptin and plasminogen activator inhibitor-1 as well as potential local adipose tissue inflammation. In epididymal and inguinal adipose tissue, mRNA expression levels of adipogenic genes were significantly decreased. Furthermore, DP exposure induced "whitening" of brown adipose tissue in the control diet group observed as a dose dependent decrease in mRNA expression of uncoupling protein 1 (Peshdary et al., 2020). In only the high-fat diet group, DP promoted glucose intolerance in mice independent of weight gain in a U-shaped dose-response evident from day 7 (Peshdary et al., 2020). In another study, rats were fed 5 mg DP/kg/d through pregnancy and/or lactation. In both offspring and dams, species diversity and richness of gut microbiota was reduced and their production of metabolites of short-chain fatty acids (SCFA) were significantly increased in all DP exposure groups (Zhang et al., 2020; Wu et al., 2021). Low doses of DP appeared to promote SCFA metabolism by the gut microbiota *in vivo* and *in vitro* in a human intestinal microbial ecosystem (SHIME). DP exposure has a greater impact

on the relative abundance of *Firmicutes* in the intestinal flora, such as *Ruminiclostridium*, *Ruminococcus* and *Sporolactobacillus* (Wu et al., 2021).

147. In mechanistic *in vitro* studies, DP was shown to inhibit insulin signalling in differentiated murine and human preadipocytes after 48 h exposure to 10, 100 or 1000 nM DP in a dose dependent manner in human adipose tissue, significant at 1000 nM (Peshdary et al., 2020). Peshdary et al., (2019) showed that DP activated the peroxisome proliferator-activated receptor (PPAR- γ) and induced adipogenesis (observed as lipid accumulation and upregulation of adipogenesis mRNA and protein markers) of both murine and human preadipocytes. These effects were only observed at the highest dose tested (10 μ M) but not at the lower doses tested in that study and the study of Peshdary et al., (2020). Some DP-mediated adipogenic endpoints were independent of PPAR- γ activation, suggesting that other potential modes of actions of DP may be involved in the observed adipogenesis *in vitro* (Peshdary et al., 2019).

148. In a recent mechanistic *in vitro* study, binding affinities of four dechloranes to the two thyroid hormone receptors (TR α and TR β) were determined by fluorescence competitive binding assay. Results demonstrated that dechloranes could interfere with thyroid function by binding with TRs and acting as TR-antagonists (Zhu et al., 2022). DP, DP 602 and DP 603 bind to the two TRs (TR α and TR β) with the relative potency (RP) value ranging from nd to 0.0667. Dechloranes could also inhibit T3 dependent transcriptional activity and T3 dependent proliferation of G3 cells in T screen. The LOEC for DP on T3-promoted GH3 cell proliferation was 50 nmol/L (32.7 ng/mL).

2.4.4 Human epidemiology

149. Epidemiology studies indicate possible associations between serum DP levels and effects on thyroid- and sex-hormones. Presented epidemiological investigations following the case-control design, have a lower level of evidence compared to cohort studies. The study design and covariates also differ between the studies, and in some studies levels of DP are highly correlated with other serum flame retardants and thus weakening the results observed for DP. Information on DP levels, group size and other relevant information is provided in Table 18, UNEP/POPS/POPRC.17/INF/9. DP concentration in maternal sera from the 20-year residents' group was 2 to 3-fold higher than the 3-year residents' group (Ben et al., 2014). *Syn*- and *anti*-DP were detected in placenta and umbilical cord serum samples, indicating that they could translocate from maternal to foetal tissues. Levels of TSH were significantly lower in maternal serum in the group that had lived in the area for 20 years (n= 44), than for those whose mothers had been residents to the area for 3 years or less (n=22), (p=0.046), but correlation analysis showed insignificant association between THs and DP levels in the 20-year residents' group. When concentrations of PBDEs were used as a control variable, the DP concentrations were positively associated with total triiodothyronine concentrations (TT3) in sera from mothers who had lived in the area for over 20 years (r = 0.37 and p = 0.020 for *syn*-DP, r = 0.360 and p = 0.024 for *anti*-DP), but no association was observed with the other serum THs. (Ben et al., 2014).

150. Another study compared serum levels of adults living in an e-waste recycling region (n=54) and a control region (n=58). Lower levels of TSH, thyroid binding globulin and mRNA expression of thyroid receptor (TR α) and higher level of mRNA gene expression iodothyronine deiodinase 1 (ID1) were observed in highly exposed individuals (Guo et al., 2019). For serum DP levels, a nonmonotonic association was observed for serum TSH and a negatively association observed for serum thyroxine binding globulin (TBG) and gene expression of ID1 and TR α was positively and negatively associated with serum DP levels, respectively (Guo et al., 2019). Association between serum DP levels and gene expression of ID1 was also observed in children (n= 112) from the same area (Guo et al., 2020). No association was observed between serum TH hormones and DP levels, but a nonmonotonic association was observed between level of *anti*-DP and TBG. Positive correlation was observed between serum DP levels and gene expression of ID1 (Guo et al., 2020). Another study of pregnant women (n=64) in Wenling, China observed no association between serum thyroid hormones and DP levels. However, a marginal positive relationship between DP serum level and apolipoprotein A was observed (p= 0.095) for total DP and for *syn*-DP (p=0.045) (Yin et al., 2020). In a study from the US, a positive association (b= 0.089, p= 0.023) between *anti*-DP levels in the subjects' wrist band and blood levels of TSH was found for women (n= 62) (Wang et al., 2020a). Overall, some association between DP exposure and thyroid hormone pathway are indicated in available epidemiology studies which include both children and adult in more exposed areas and adults with lower exposure.

151. In another epidemiology study comparing groups from e-waste dismantling (n= 54) and control area (n=58), Guo et al (2018b), showed significant negative association between DP serum levels and follicle stimulating hormone (FSH) in both females and males. In males, a positive association between anti-DP and testosterone and a non-monotonic association between testosterone and *syn*-DP serum levels were observed (Guo et al., 2018b). In a generalized additive model, a significant nonmonotonic association was observed for estrogen and DP in female. However, the study did not report menstrual cycle as one of the listed confounders so the female data may have some limitations.

2.4.5 Other concerns

152. The DP by-products 1,3- and 1,5-Dechlorane Plus monoadduct (DPMA) have in some cases been detected in greater levels than DP in environmental samples (Sverko et al., 2011; Tomy et al., 2013; Guerra et al., 2011), and also in Antarctic biota (Wolschke et al., 2015) (see paragraph 89 for details). DPMA may also be under-reported because of destructive sample preparation methods (Rjabova et al., 2016). Based on predictive QSAR modelling studies, DPMA is potentially both bioaccumulative and persistent. No information is available on its toxicity to humans and the environment, but based on its structural similarity to aldrin and heptachlor it might be epoxidated in the environment to form substances that could be neurotoxic and/or cause hepatotoxicity (ECHA 2017b). For additional information see Figure 1 and Table 19 in UNEP/POPS/POPRC.17/INF/9.

3. Synthesis of information

153. DP is a commercially available additive chlorinated FR that is marketed and used as a replacement for already regulated FRs in plastic polymers. Known applications include the aviation and automotive industries, electrical wire and cable coatings, plastic roofing materials, connectors in TV and computer monitors.

154. DP is detected in the environmental samples, humans and wildlife at different locations around the globe. It is a global contaminant and has been found in remote regions far from sites of production and use, including in the Arctic, the Antarctic and high-mountain regions in Tibet. DP has also been detected in dust, sludge and wastewater from WWTPs at sites far from production and e-waste recycling sites, indicating emission and exposure from consumer products.

155. Available monitoring data provides evidence for ongoing LRET. Bioaccumulation and trophic magnification are reported in several studies, including in biota in remote regions. DP is environmentally persistent, and although existing environmental levels in remote regions are generally low, there is reason for concern. DP levels in remote regions are in some studies reported to be in the same range as in source regions. DP is persistent in the environment and bioaccumulates in biota. The levels are likely to increase over time as a result of continued use and could possibly lead to adverse effects in biota in the future if no action is taken.

156. Although DP has been produced and used for almost half a century, environmental detection of DP was first reported in 2006 in the Great Lakes Basin in North America (Hoh et al., 2006). Retrospective studies have shown its presence in historic samples before that. Further research has identified DP as a global pollutant (Wang et al., 2016). Due to limited data, temporal and spatial trends for DP are equivocal; while increasing trends are observed in some studies, other studies report unchanged levels (see section 2.3.1).

157. The dataset on toxicity and ecotoxicity is limited, particularly regarding exhibited apical effects and long-term studies are largely lacking, however, available short-term toxicity data indicates concern for potential adverse effects to the environment and humans. One relevant mode of action could be the induction of oxidative stress that influences several biological processes. Effects on the developing nervous system and brain have been indicated in fish. A dose dependent increase in DNA strand breaks (Comet assay), mortality rate, delayed hatching time, reduced hatching rate, decreased body length, and increased morphological deformities were observed in a carp embryo study. Furthermore, one co-exposure study on zebrafish indicated that other pollutants may increase bioaccumulation, resulting in greater toxicity. DP has been shown to potentially impair liver function in mice. At low dose insulin secretion was altered, while in combination with high fat diet DP glucose intolerance was also induced. In addition, adipocyte dysfunction as well as alterations in biomarkers associated with insulin resistance were observed in mice. DP has been shown to activate peroxisome proliferator-activated receptor gamma and inhibit insulin signalling *in vitro* in mammalian adipose cells, and to bind to TH-receptors and inhibit T3-mediated cell growth *in vitro*. A few case-control epidemiology studies found some associations with DP exposure with sex- and thyroid hormone pathway in humans. From the available information it is not currently possible to determine with certainty whether DP causes adverse effects on human health.

158. Adding to the concern are the DP by-products, such as 1,3- or 1,5-DPMA which have structural similarities to substances such as aldrin and heptachlor that are already listed by the Stockholm Convention (Table 19, UNEP/POPS/POPRC.17/INF/9). There are no toxicity data for these monoadducts, but in some cases these are detected in greater levels than DP in environmental samples (Sverko et al., 2011; Tomy et al., 2013; Guerra et al., 2011).

159. DP is transferred to developing offspring during pregnancy via blood, and after delivery via breast feeding. Maternal transfer to eggs has been described for fish, bird and amphibians leading to exposure during sensitive life stages. High concentrations of DP have been detected in environmental samples and humans living near DP production plants and e-waste recycling sites.

160. Due to its POP properties and its widespread production and use, global action is warranted to prevent further releases of DP.

4. Concluding statement

161. DP is emitted into the environment from human activities, e.g. from manufacturing, use and disposal and management of waste. DP is persistent, bioaccumulative and undergoes LRET, making emissions and releases of this substance a transboundary pollution problem. Globally, DP is detected in humans, wildlife and environmental samples in all global regions, including in the Arctic and Antarctic.

162. In humans, DP has been detected in hair, umbilical cord blood, serum and breast milk. High concentrations of DP have been detected in environmental samples and humans living near e-waste recycling sites and production plants. DP is also detected in baby food, dust, sludge, wastewater from WWTPs and marine plastics debris indicating emission and exposure from consumer products throughout their lifecycle.

163. Available scientific literature suggests that there is a risk of adverse effects in developing aquatic species and organisms due to oxidative stress that may affect important biological processes. In aquatic organisms, effects on the developing nervous system and brain have been indicated as well. Furthermore, observed effects on photosynthetic activity in marine macroalgae suggest possible impacts on aquatic ecosystems mediated by negative effects on primary production. Evidence of potential impairment of liver function and low dose effects on adipose tissue, and the endocrine modulating effects involving the insulin and peroxisome proliferator-activated receptor gamma pathways gives cause for concern for adverse effects related to metabolic disorders in humans.

164. Mixture toxicity effects such as increased toxicity and bioaccumulation have been indicated for DP in one fish study and suggest an additional cause for concern both for environmental effects and effects on human health. Obesity and metabolic syndromes, such as, type 2 diabetes are increasing worldwide. Several chemicals present in humans have been shown to alter adipose tissue and biomarkers related to this syndrome, and exposure to DP may be a risk factor for these conditions. Climate warming may exacerbate the mobilization and release of DP in polar environments as DP deposited in sea ice, glaciers, and permafrost are released into freshwater and marine environments. The combined effect of climate change and other environmental stressors adds to the risk by affecting exposure levels, the vulnerability and adaptability of organisms, particularly in the polar regions.

165. The greatest uncertainty in the review of DP relates to the significance of its adverse effects. Although data are limited and chronic toxicity studies are lacking, there are studies showing that DP has adverse effects on the environment and that it can be potentially toxic to mammals and humans. There are some traditional toxicological studies for mammals where no adverse effects have been observed even at unrealistically high concentrations. Studies from the peer review literature on endpoints outside of those typically submitted to regulatory authorities indicate concerns which have not been fully addressed in standardized traditional tests. In addition, specific studies have reported hazards such as effects on developing neurosystem and brain in fish, photosynthesis and endocrine pathways related to obesity and metabolism, which may be particularly relevant in the assessment of both human health and environmental effects.

166. Based on evidence for persistence, bioaccumulation and adverse effects of DP observed in some organisms and its widespread occurrence in the global environment including at remote regions and mindful of Article 1 referring to the precautionary approach as set forth in Principle 15 of the Rio Declaration on Environment and Development and Article 8 para 7 (a) stating that a lack of full scientific certainty should not prevent the proposal from proceeding, it is concluded that DP is likely, as a result of its long-range environmental transport, to lead to significant adverse human health and/or environmental effects such that global action is warranted.

References

- AIAC, Aerospace Industries Association of Canada (2020). Letter to the Stockholm Convention secretariat on applications of Dechlorane Plus (DP) and its syn- and anti-isomers in the aerospace industry submitted as a comment to the second draft of the Dechlorane Plus risk profile. 30 March 2020.
- ADAMA (2019). ADAMA Reports on Progress Towards Anpon Acquisition. Available at: <https://www.adama.com/en/449/adama-reports-on-progress-towards-anpon-acquisition>.
- Ali, U., et al. (2015). Assessing the combined influence of TOC and black carbon in soil–air partitioning of PBDEs and DPs from the Indus River Basin, Pakistan. *Environ. Pollut.* 201, 131–140.
- Annex E (2019). Submissions of Annex E (risk profile) information on Dechlorane Plus. 2 December 2019. Available at: <http://chm.pops.int/tabid/8225/Default.aspx>.
- Annex E (2020). Comments on the draft risk profiles on Dechlorane Plus. 30 March 2020. Available at: <http://chm.pops.int/tabid/8366/Default.aspx>.
- Allen JG, Stapleton HM, Vallarino J, McNeely E, McClean MD, Harrad SJ, Rauert CB, Spengler JD (2013). Exposure to flame retardant chemicals on commercial airplanes. *Environ Health* 12:17.
- AMAP (2017). AMAP Assessment 2016: Chemicals of Emerging Arctic Concern. Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway. xvi+353pp.
- Aznar-Alemany Ò, Yang X, Alonso MB, Costa ES, Torres JPM, Malm O, Barceló D, Eljarrat E (2019). Preliminary study of long-range transport of halogenated flame retardants using Antarctic marine mammals *Sci Total Environ.* 650(Pt2):1889-1897. doi: 10.1016/j.scitotenv.2018.09.287.
- Barón E, Dissanayake A, Vilà-Cano J, Crowther C, Readman JW, Jha AN, Eljarrat E, Barceló D (2016). Evaluation of the Genotoxic and Physiological Effects of Decabromodiphenyl Ether (BDE-209) and Dechlorane Plus (DP) Flame Retardants in Marine Mussels (*Mytilus galloprovincialis*). *Environ Sci Technol.* 50(5):2700-8.
- Barón E, Bosch C, Mániz M, Andreu A, Sergio F, Hiraldo F, Eljarrat E, Barceló D (2015). Temporal trends in classical and alternative flame retardants in bird eggs from Doñana Natural Space and surrounding areas (south-western Spain) between 1999 and 2013. *Chemosphere.* 138:316-23.
- Barón E, Santín G, Eljarrat E, Barceló D (2014a). Occurrence of classic and emerging halogenated flame retardants in sediment and sludge from Ebro and Llobregat river basins (Spain). *J Hazard Mater.* 30; 265:288-95.
- Barón E, Mániz M, Andreu AC, Sergio F, Hiraldo F, Eljarrat E, Barceló D (2014b). Bioaccumulation and biomagnification of emerging and classical flame retardants in bird eggs of 14 species from Doñana Natural Space and surrounding areas (South-western Spain). *Environ Int.* 68:118-26. doi: 10.1016/j.envint.2014.03.013.
- Ben Y-J, Li X-H, Yang Y-L, Li L, Di J-P, Wang W-Y, Zhou R-F, Xiao K, Zheng M-Y, Tian Y, Xu X-B (2013). Dechlorane Plus and its dechlorinated analogs from an e-waste recycling center in maternal serum and breast milk of women in Wenling, China. *Environmental Pollution.* 173, 176-181.
- Ben Y-J, Li X-H, Yang Y-L, Li L, Zheng M-Y, Wang W-Y, Xu X-B (2014). Placental Transfer of Dechlorane Plus in Mother–Infant Pairs in an E-Waste Recycling Area (Wenling, China.) *Environ. Sci. Technol.* 48, 5187–5193.
- Blais JM, Kimpe LE, McMahon D, et al. (2005). Arctic seabirds transport marine-derived contaminants. *Science.* 2005;309(5733):445. doi:10.1126/science.1112658.
- Borgå K, Kidd KA, Muir DC, Berglund O, Conder JM, Gobas FA, Kucklick J, Malm O, Powell DE (2012). Trophic magnification factors: considerations of ecology, ecosystems, and study design. *Integr Environ Assess Manag.* 8(1):64-84.
- Brasseur C, Pirard C, Scholl G, De Pauw E, Viel JF, Shen L, Reiner EJ, Focant JF (2014). Levels of dechloranes and polybrominated diphenyl ethers (PBDEs) in human serum from France. *Environ. Int.* 65, 33–40.
- Breivik K, Wania F, Muir DCG, Alae M, Backus S, Pacepavicius G (2006). Empirical and modeling evidence of the long-range atmospheric transport of decabromodiphenyl ether. *Environmental Science and Technology.* 40, 4612–4618.
- Briels N, Torgersen LN, Castaño-Ortiz JM, Løseth ME, Herzke D, Nygård T, Bustnes JO, Ciesielski TM, Poma G, Malarvannan G, Covaci A, Jaspers VLB (2019). Integrated exposure assessment of northern goshawk (*Accipiter gentilis*) nestlings to legacy and emerging organic pollutants using non-destructive samples. *Environ Res.* 178:108678. doi: 10.1016/j.envres.2019.108678.
- Briels N, Løseth ME, Ciesielski TM, Malarvannan G, Poma G, Kjærøvik SA, Léon A, Cariou R, Covaci A, Jaspers VLB (2018). In ovo transformation of two emerging flame retardants in Japanese quail (*Coturnix japonica*). *Ecotoxicol Environ Saf.* 149:51-57.

- Brock WJ, Schroeder RE, McKnight CA, Van Steenhouse JL and Nyberg JM (2010). Oral repeatdose and reproductive toxicity of the chlorinated flame retardant Dechlorane Plus. *Int. J. Toxicol.*, 29, 582–593.
- Cao X, Wang L, Zhang Y, Li Y, Zhu C, Zheng X, Sun Y, Li D, Mo L, Mai B (2020). Occurrence of organic pollutants in plastics on beach: Stranded foams can be sources of pollutants in islands. *Sci Total Environ.* 14; 707:136119.
- Cao Z, Xu F, Covaci A, Wu M, Wang H, Yu G, Wang B, Deng S, Huang J, Wang X (2014). Distribution patterns of brominated, chlorinated, and phosphorus flame retardants with particle size in indoor and outdoor dust and implications for human exposure. *Environmental Science and Technology*, 48, 8839–8846.
- Canada (2019a). Dept. of the Environment. Dept. of Health. Screening Assessment Certain Organic Flame Retardants Grouping 1,4:7,10-Dimethanodibenzo[a,e]cyclooctene,1,2,3,4,7,8,9,10,13,13,14,14-dodecachloro-1,4,4a,5,6,6a,7,10,10a,11,12,12 adodecahydro-Dechlorane Plus (DP), Chemical Abstracts Service Registry Number 13560-89- 9. Available at: <https://www.canada.ca/en/environment-climate-change/services/evaluating-existing-substances/screening-assessment-certain-organic-flame-retardants-substance-grouping-dimethanodibenzo-cyclooctene-dodecachloro-dodecahydro-dechlorane-plus-dp-.html>.
- Canada (2019b). Dept. of the Environment. Dept. of Health. Certain Organic Flame Retardants Grouping Risk Management Approach For 1,4:7,10-Dimethanodibenzo[a,e]cyclooctene, 1,2,3,4,7,8,9,10,13,13,14,14-dodecachloro1,4,4a,5,6,6a,7,10,10a,11,12,12a-dodecahydro- Dechlorane Plus (DP), Chemical Abstracts Service Registry Number 13560-89- 9. Available at: <https://www.canada.ca/en/environment-climate-change/services/evaluating-existing-substances/certain-organic-flame-retardants-grouping-risk-management-approach-dimethanodibenzo-cyclooctene-dodecachloro-dodecahydro-dechlorane-plus-dp.html>.
- Carlsson P, Vrana B, Sobotka J, Borgå K, Bohlin Nizzetto P, Varpe Ø (2018). New brominated flame retardants and dechlorane plus in the Arctic: Local sources and bioaccumulation potential in marine benthos. *Chemosphere.* 211: 1193-1202.doi: 10.1016/j.chemosphere.2018.07.158.
- CATALOGIC (Computer Model) (2012). Version 5.11.6. Bourgas (BG): Bourgas Prof. Assen Zlatarov University, Laboratory of Mathematical Chemistry.
- CECBP - California Environmental Contaminant Biomonitoring Program (2008). Brominated and chlorinated organic chemical compounds used as flame retardants. Materials for the December 4-5, 2008 Meeting of the California Environmental Contaminant Biomonitoring Program Scientific Guidance Panel (SGP). 32 pp.
- Cequier E, Marce RM, Becher G, Thomsen C (2015). Comparing human exposure to emerging and legacy flame retardants from the indoor environment and diet with concentrations measured in serum. *Environment Intern.* 74, 54-59.
- Cequier E, Ionas AC, Covaci A, Marce RM, Becher G, Thomsen C (2014). Occurrence of a broad range of legacy and emerging flame retardants in indoor environments in Norway. *Environ. Sci. Technol.* 48, 6827–6835.
- Čechova E, Vojta S, Kukučka P, Kočan A, Trnovec T, Murinov LP, de Cock M, van de Borc M, Askevold J, Eggesbod M, Scheringer M (2017). Legacy and alternative halogenated flame retardants in human milk in Europe: Implications for children's health. *Environment International* 108, 137–145.
- Chen W, Bao J, Bu T, Jin H, Liu Y, Li T, Wang H, Zhao P, Wang Y, Hu J, Jina J (2021). Dechlorane Plus Biomagnification and Transmission through Prairie Food Webs in Inner Mongolia, China. *Environmental Toxicology and Chemistry* 40:413-421. DOI: 10.1002/etc.4923.
- Chen X, Dong Q, Chen Y, Zhang Z, Huang C, Dong Q, Roper C, Tanguay RL, Zhu Y, Zhang Y (2019). Neurodevelopmental toxicity assesment of alkyl phenanthrene and Dechlorane Plus co-exposure in zebrafish. *Ecotoxicology and Environmental Safety.* 180, 762-769.
- Chen X, Dong Q, Chen Y, Zhang Z, Huang C, Zhu Y, Zhang Y (2017). Effects of Dechlorane Plus exposure on axonal growth, musculature and motor behavior in embryo-larval zebrafish. *Environ Pollut.* 224:7-15. doi: 10.1016/j.envpol.2017.03.011.
- Chen K, Zheng J, Yan X, Yu L, Luo X, Peng X, Yu Y, Yang Z, Mai B (2015). Dechlorane Plus in paired hair and serum samples from e-waste workers: correlation and differences. *Chemosphere* 123, 43–47.
- Chen SJ, Tian M, Wang J, Shi T, Luo Y, Luo XJ, Mai BX (2011). Dechloran Plus (DP) in air and plants at an electronic waste (e-waste) site in South China. *Environ. Pollut.* 159 (5), 1290-1296.
- Cheng Y, Ding J, Xie X, Ji X, Zhang Y (2019). Validation and Application of a 3-Step Sequential Extraction Method to Investigate the Fraction Transformation of Organic Pollutants in Aging Soils: A Case Study of Dechlorane Plus. *Environ Sci Technol.* 5;53(3):1325-1333.
- Chou TW, Liu DHW, Mabey WR, Mitoma C, and Smith JH (1979a). As cited in: HPV Challenge- Robust Summaries & Test Plans: Dechlorane Plus. Final revision as identified by the submitter, June 8, 2009. Vapour Pressure and Water

Solubility. Metabolism and Environmental Screening Studies on Dechlorane Plus. SRI International. SRI Project No. LSC-8060.

Corsolini S, Metzdorff A, Baroni D, Roscales JL, Jiménez B, Cerro-Gálvez E, Dachs J, Galbán-Malagón C, Audy O, Kohoutek J, Přibylková P, Poblete-Morales M, Avendaño-Herrera R, Bergami E, Pozo K (2021). Legacy and novel flame retardants from indoor dust in Antarctica: Sources and human exposure. *Environ Res.* 196:110344. doi: 10.1016/j.envres.2020.110344.

Chou TW, Liu DHW, Mabey WR, Mitoma C and Smith JH (1979b). Metabolism and environmental screening studies on Dechlorane Plus. SRI International, SRI Project LSC-8060, 25 April 1979. Testing laboratory: SRI International, Menlo Park, California, USA. Report no.: LSC-8060. Owner company: Hooker Chemical Corporation, Hooker Research Center. Report date: 1979-04-25.

Crump D, Chiu S, Gauthier LT, Hickey NJ, Letcher RJ, Kennedy SW (2011). The effects of Dechlorane Plus on toxicity and mRNA expression in chicken embryos: A comparison of in vitro and in ovo approaches. *Comp Biochem Physiol C* 154:129-134.

Davis EF, Klosterhaus SL, Stapleton HM (2012). Measurement of flame retardants and triclosan in municipal sewage sludge and biosolids. *Environ Int.* 40:1-7.

De Groef B, Van der Geyten S, Darras VM, Kühn ER (2006). Role of corticotropin-releasing hormone as a thyrotropin-releasing factor in non-mammalian vertebrates. *Gen. Comp. Endocrinol.* 146, 62-68.

de la Torre A, Pacepavicius G, Shen L, Reiner E, Jiménez B, Alae M, Martínez MA (2010). Dechlorane Plus and related compounds in Spanish Air. *Organohalogen Compounds*. Vol. 72, 929-932.

Desjardins CF, Mazerolle MJ, Verreault J (2019). Is the urban-adapted ring-billed gull a biovector for flame retardants? *Environ Pollut.* 244:109-117. doi: 10.1016/j.envpol.2018.10.003.

de Wit CA, Bossi R, Dietz R, Dreyer A, Faxneld S, Garbus SE, Hellström P, Koschorreck J, Lohmann N, Roos A, Sellström U, et al. (2020). Organohalogen compounds of emerging concern in Baltic Sea biota: Levels, biomagnification potential and comparisons with legacy contaminants. *Environment International*, 144: 106037. <https://doi.org/10.1016/j.envint.2020.106037>.

de Wit C A, Kierkegaard A, Ricklund N, Sellstrom U (2011). Emerging brominated flame retardants in the environment. In E. Eljarrat & D. Barcelo (Eds.), *Brominated flame retardants* (pp. 241–286). Berlin: Springer.

Díaz M, Bassols J, López-Bermejo A, Gómez-Roig MD, de Zegher F, Ibáñez L (2012). Placental Expression of Peroxisome Proliferator- Activated Receptor γ (PPAR γ): Relation to Placental and Fetal Growth. *J Clin Endocrinol Metab*, 97(8): E1468–E1472.

Dou J, Jin Y, Wu B, Li M (2015). Potential genotoxicity and risk assessment of a chlorinated flame retardant, Dechlorane Plus. *Chemosphere* 135: 462–466.

Dreyer A, Neugebauer F, Lohmann N, Rüdell H, Teubner D, Grotti M, Rauert C, Koschorreck J (2019). Recent findings of halogenated flame retardants (HFR) in the German and Polar environment. *Environ Pollut.* 253:850-863. doi:10.1016/j.envpol.2019.07.070.

ECCC - Environment and Climate Change Canada (2019). Domestic Substances List. <https://www.canada.ca/en/environment-climate-change/services/canadian-environmental-protection-act-registry/substances-list/domestic.html>.

ECHA (2021a). Annex XV Restriction Report. Proposal for a restriction. Substance name: 1,6,7,8,9,14,15,16,17,17,18,18- Dodecachloropentacyclo[12.2.1.16,9.02,13.05,10]octadeca-7,15-diene (“Dechlorane Plus”TM) [covering any of its individual anti- and syn-isomers or any combination thereof]. Available at: <https://echa.europa.eu/restrictions-under-consideration/-/substance-rev/28201/term>.

ECHA (2021b). Registration dossier 1,6,7,8,9,14,15,16,17,17,18,18-dodecachloropentacyclo[12.2.1.16,9.02,13.05,10]octadeca-7,15-diene. EC / List no.: 236-948-9, CAS no.: 13560-89-9. <https://echa.europa.eu/registration-dossier/-/registered-dossier/11906/7/7/1> (Accessed 29.6.2021).

ECHA (2017a). Agreement of the Member State Committee in accordance with Article 59(8): 1,6,7,8,9,14,15,16,17,17,18,18-Dodecachloropentacyclo- [12.2.1.16,9.02,13.05,10]octadeca-7,15-diene (“Dechlorane Plus”TM) [covering any of its individual anti- and syn-isomers or any combination thereof] is identified as a substance of very high concern because it meets the criteria of Article 57 (e) of Regulation (EC) 1907/2006 (REACH) as a substance which is very persistent and very bioaccumulative (vPvB), in accordance with the criteria and provisions set out in Annex XIII of REACH Regulation. Available at: <https://echa.europa.eu/documents/10162/15b88a69-2162-9385-4089-7c247cba4da6>.

ECHA (2017b). Member State Committee support document for identification of 1,6,7,8,9,14,15,16,17,17,18,18-Dodecachloropentacyclo- [12.2.1.16,9.02,13.05,10]octadeca-7,15-diene (“Dechlorane Plus”TM) [covering any of its

- individual anti- and syn-isomers or any combination thereof] as a substance of very high concern because it meets the criteria of Article 57 (e). Adopted on 30 November 2017. Available at: <https://echa.europa.eu/documents/10162/acd1ca10-4d98-5537-3d5c-54cf8f3570c4> .
- Evenset A, Carroll J, Christensen GN, Kallenborn R, Gregor D, Gabrielsen GW (2007). Seabird guano is an efficient conveyer of persistent organic pollutants (POPs) to Arctic lake ecosystems. *Environ Sci Technol.* 41(4): 1173-9.
- Fang M, Kim JC, Chang YS (2014). Investigating Dechlorane Plus (DP) distribution and isomer specific adsorption behavior in size fractionated marine sediments. *Science of the Total Environment*, 481, 114–120.
- Feng Y, Tian J, Xie HQ, She J, Xu SL, Xu T, Tian W, Fu H, Li W, Tao W, Wang L, Chen Y, Zhang S, Zhang W, Guo TL, Zhao B (2016). Effects of acute low-dose exposure to the chlorinated flame retardant Dechlorane 602 and the Th1 and Th2 Immune responses in adult male mice. *Environ Health Perspect.* 124 (9): 1406-1413.
- Feo ML, Barón E, Eljarrat E, Barceló D (2012). Dechlorane Plus and related compounds in aquatic and terrestrial biota: a review. *Anal Bioanal Chem.* 404(9): 2625-37. doi: 10.1007/s00216-012-6161-x. Epub 2012 Jun 14. Review.
- Fernie KJ, Chabot D, Champoux L, Brimble S, Alae M, Martenson S, Chen D, Palace V, Bird DM, Letcher RJ. 2017. Spatiotemporal patterns and relationships among the diet, biochemistry, and exposure to flame retardants in an apex avian predator, the peregrine falcon. *Environmental Research.* 158:43-53. <https://doi.org/10.1016/j.envres.2017.05.035>
- Franklin J (2016). How reliable are field-derived biomagnification factors and trophic magnification factors as indicators of bioaccumulation potential? Conclusions from a case study on per- and polyfluoroalkyl substances. *Integr Environ Assess Manag.* 12(1):6-20.
- Fromme H, Thomsen C, Aschenbrenner B, Haug LS, Weber T, Kolossa-Gehring M, Völkel W, Schober W (2020). Time trend of exposure to dechloranes: Plasma samples of German young adults from the environmental specimen bank collected from 1995 to 2017. *International Journal of Hygiene and Environmental Health.* 229, 113593. <https://doi.org/10.1016/j.ijheh.2020.113593>.
- Gagné PL, Fortier M, Fraser M, Parent L, Vaillancourt C, Verreault J (2017). Dechlorane Plus induces oxidative stress and decreases cyclooxygenase activity in the blue mussel. *Aquat Toxicol.* 188:26-32.
- Gao H, Na G, Yao Y, Li R, Gao Y, Zhang Z, Yao Z (2018). Distribution Characteristics and Source of Dechloranes in Soil and Lichen of the Fildes Peninsula (Antarctica). *Int J Environ Res Public Health.* 15(10).
- Gauthier LT, Letcher RJ (2009). Isomers of Dechlorane Plus flame retardant in the eggs of herring gulls (*Larus argentatus*) from the Laurentian Great Lakes of North America: temporal changes and spatial distribution. *Chemosphere*;75(1):115-20. doi: 10.1016/j.chemosphere.2008.11.030.
- Ge X, Ma S, Zhang X, Yang Y, Li G, Yu Y (2020). Halogenated and organophosphorous flame retardants in surface soils from an e-waste dismantling park and its surrounding area: Distributions, sources, and human health risks. *Environ Int.* 139:105741.
- Gentes ML, Letcher RJ, Caron-Beaudoin E, Verreault J (2012). Novel flame retardants in urban-feeding ring-billed gulls from the St. Lawrence River, Canada. *Environ Sci Technol.* 2012;46(17):9735–9744. doi:10.1021/es302099f.
- Ghelli E, Cariou R, Dervilly G, Pagliuca G, Gazzotti T (2021). Dechlorane Plus and Related Compounds in Food -A Review. *Int. J. Environ. Res. Public Health*, 18, 690. <https://doi.org/10.3390/ijerph18020690>.
- Gong N, Shao K, Han X, Zhang Y, Sun Y (2018). Enrichment and physiological responses of dechlorane plus on juvenile marine macroalgae (*Ulva pertusa*). *Chemosphere* 205:594-600.
- Gong N, Shao K, Hang X, Zhao L, Zhong R, Zhao E, Li G and Sun Y (2013). Effect of Dechlorane Plus on *Pseudokirchneriella subcapitata* by flow cytometry. *Organohalogen Compounds*, 75, 200-204. <http://www.dioxin20xx.org/pdfs/2013/86.pdf>.
- Green NW, Schøyen M, Hjermmann DØ, Øxnevad S, Ruus A, Beylich B, Lund E, Tveiten L, Jenssen MTS, Håvardstun J, Ribeiro AL, Doyer I, Rundberget JT and Bæk K (2019). Contaminants in coastal waters of Norway 2018. NIVA report 7412-2019. pp. 235.
- Guerra P, Fernie K, Jiménez B, Pacepavicius G, Shen L, Reiner E, Eljarrat E, Barceló D, Alae M (2011). Dechlorane plus and related compounds in peregrine falcon (*Falco peregrinus*) eggs from Canada and Spain. *Environ Sci Technol.* 45(4):1284-90.
- Guo L-C, Liu T, Yang Y, Yu S, Gao Y, Huang W, Xiao J, Ma W, Rutherford S, Zhang Y (2020). Changes in thyroid hormone related proteins and gene expression induced by polychlorinated biphenyls and halogen flame retardants exposure of children in a Chinese e-waste recycling area. *Science of the Total Environment* 742, 140597. <https://doi.org/10.1016/j.scitotenv.2020.140597>.

- Guo LC, Yu S, Wu D, Huang J, Liu T, Xiao J, Huang W, Gao Y, Li X, Zeng W, Rutherford S, Ma W, Zhang Y, Lin L (2019). Disruption of thyroid hormone regulated proteins and gene expression by polychlorinated biphenyls, polybrominated diphenylethers and new flame retardants in residents to an e-waste region. *Environ. Poll.* 254: 112925.
- Guo J, Simon K, Romanak K, Bowerman W, Vernier M (2018a). Accumulation of flame retardants in paired eggs and plasma of bald eagles. *Environmental Pollution* 237: 499-507.
- Guo LC, Pan S, Yu S, Liu T, Xiao J, Zhu B, Qu Y, Huang W, Li M, Li X, Zeng W, Rutherford S, Lin L, Zhang Y, Ma W (2018b). Human Sex Hormone Disrupting Effects of New Flame Retardants and Their Interactions with Polychlorinated Biphenyls, Polybrominated Diphenyl Ethers, a Case Study in South China. *Environ. Sci. Technol.* 2018, 52, 13935–13941.
- Guo J, Venier M, Salamova A, Hites RA (2017). Bioaccumulation of Dechloranes, organophosphate esters, and other flame retardants in Great Lakes fish. *Sci Total Environ.* 583:1-9.
- Haglund P, Kaj L, Brorström-Lundén E (2016). Analysis of new brominated flame retardants in human samples and background air. For Swedish EPA (Naturvårdsverket), contract no. 2219-14-006.
- Hang XM, Jiang Y, Liu Y, Jia HL and Sun YQ (2013). An integrated study on the toxicity of Dechlorane Plus in zebrafish. *Organohalogen Compounds*, 75, 1085-1089. <http://dioxin20xx.org/organohalogen-compounds-database-search/?swpquery=Dechlorane> (<http://www.dioxin20xx.org/pdfs/2013/5154.pdf>).
- Hansen KM, Fauser P, Vorkamp K, Christensen JH (2020). Global emissions of Dechlorane Plus. *Sci Total Environ.* 2020 Nov 10;742:140677. doi: 10.1016/j.scitotenv.2020.140677. Epub 2020 Jul 2. PMID: 32721756.
- Hassan Y, Shoeib T (2015). Levels of polybrominated diphenyl ethers and novel flame retardants in microenvironment dust from Egypt: An assessment of human exposure. *Science of the Total Environment.* 505, 47–55.
- Herzke D, Rostkowski P, Harju M, Borgen AR, Christensen-Dalsgaard Signe (2019). Assessment of additives used in plastic in seabirds. NILU report 1/2019. Available at: [NILU-report-1-2019.pdf](http://nilu.no/reports/1-2019.pdf) (unit.no).
- Hewitt DP, Mark PJ, Waddell BJ (2006). Placental Expression of Peroxisome Proliferator-Activated Receptors in Rat Pregnancy and the Effect of Increased Glucocorticoid Exposure. *Biology of Reproduction* 74, 23–28.
- Hoh E, Zhu LY, and Hites RA (2006). Dechlorane plus, a chlorinated flame retardant, in the Great Lakes. *Environ Sci Technol* 40 (4), 1184–1199.
- Houde M, Wang X, Ferguson SH, Gagnon P, Brown TM, Tanabe S, Kunito T, Kwan M, Muir DCG (2017). Spatial and temporal trends of alternative flame retardants and polybrominated diphenyl ethers in ringed seals (*Phoca hispida*) across the Canadian Arctic. *Environ Pollut.* 223:266-276. doi: 10.1016/j.envpol.2017.01.023. Epub 2017 Jan 26.
- Hou M, Wang Y, Zhao H, Zhang Q, Xie Q, Zhang X, Chen R, Chen J (2018). Halogenated flame retardants in building and decoration materials in China: Implications for human exposure via inhalation and dust ingestion. *Chemosphere.* 203:291-299.
- HPV summaries. USEPA High Production Volume Information System concerning Dechlorane Plus, last updated 2009. Provided as Annex E submission by IPEN-ATCA. Available at U.S. EPA's ChemView: <https://chemview.epa.gov/chemview>.
- Huang C-C, Zeng Y-H, Luo X-J, Tang B, Liu Y-E, Rena Z-H, Mai B-X (2018). Level changes and human dietary exposure assessment of halogenated flame retardant levels in free-range chicken eggs: A case study of a former e-waste recycling site, South China / *Science of the Total Environment* 634, 509–515.
- Iqbal M, Syed JH, Breivik K, Chaudhry MJI, Li J, Zhang G, Malik RN (2017). E-waste driven pollution in Pakistan: the first evidence of environmental and human exposure to flame retardants (FRs) in Karachi City. *Environ Sci Technol* 51(23):13895–13905.
- J-CHECK, Japan Chemicals Collaborative database (2020a): https://www.nite.go.jp/chem/jcheck/template.action?ano=7755&mno=4-0296&cno=13560-89-9&request_locale=en, (Accessed 26.05.2020).
- J-CHECK, Japan Chemicals Collaborative database (2020b): https://www.nite.go.jp/chem/jcheck/template.action?ano=28771&mno=4-0296&cno=13560-89-9&request_locale=en (Accessed 26.05.2020).
- Ji X, Xie X, Ding J, Cheng Y, He H, Huang Y, Qin L, Zhu H, Zhao C, Li A, and Han C (2018). Chlorinated flame retardant Dechlorane Plus: environmental pollution in China. *Environ. Rev.* 26: 273–285.

- Jia H, Sun Y, Liu X, Yang M, Wang D, Qi H, Shen L, Sverko E, Reiner EJ, Li YF (2011). Concentration and bioaccumulation of dechlorane compounds in coastal environment of northern China. *Environ Sci Technol.* 1;45(7):2613-8.
- Jin X, Lee S, Jeong Y, Yu JP, Baek WK, Shin KH, Kannan K, Moon HB (2016). Species-specific accumulation of polybrominated diphenyl ethers (PBDEs) and other emerging flame retardants in several species of birds from Korea. *Environ Pollut.* 219:191-200. doi: 10.1016/j.envpol.2016.10.040.
- Johnson PI, Stapleton HM, Mukherjee B, Hauser R, Meeker JD (2013). Associations between brominated flame retardants in house dust and hormone levels in men. *Sci Total Environ* 445-446:177-184.
- Kaj L, Norstrom K, Egelrud L, Remberger M, Lilja K, Brorstrom-Lunden E (2010). Results from the Swedish National Screening Programme 2009, Subreport 2: Dechlorane Plus. Report B1950 Swedish Environmental Research Institute IVL, on assignment from the Swedish Environmental Protection Agency.
- Kakimoto K, Nagayoshi H, Takagi S, Akutsu K, Konishi Y, Kajimura K, Hayakawa K, Toriba A (2014). Inhalation and dietary exposure to Dechlorane Plus and polybrominated diphenyl ethers in Osaka, Japan. *Ecotoxicology and Environmental Safety.* 99, 69–73.
- Kang H, Moon HB, Choi K (2016). Toxicological responses following short-term exposure through gavage feeding or water-borne exposure to Dechlorane Plus in zebrafish (*Danio rerio*). *Chemosphere.* 146:226-32. doi: 10.1016/j.chemosphere.2015.12.024.
- Katsoyiannis A, Samara C (2004). Persistent organic pollutants (POPs) in the sewage treatment plant of Thessaloniki, northern Greece: occurrence and removal. *Water Res.* 38:2685–98.
- KEMI (2019). Swedish Chemicals Agency Commodity Guide (database on Internet). CAS no 13560-89-9 Dechlorane Plus. (cited June 2013). Available from: <https://webapps.kemi.se/varuguiden/AmneMaterial.aspx>.
- Kim JT, Choi YJ, Barghi M, Kim JH, Jung JW, Kim K, Kang JH, Lammel G, Chang YS (2021). Occurrence, distribution, and bioaccumulation of new and legacy persistent organic pollutants in an ecosystem on King George Island, maritime Antarctica. *J Hazard Mater.* Vol. 405: 2021, 124141, doi: 10.1016/j.jhazmat.2020.124141.
- Kim JT, Choi YJ, Barghi M, Yoon YJ, Kim JH, Kim JH, Chang YS (2018). Occurrence and distribution of old and new halogenated flame retardants in mosses and lichens from the South Shetland Islands, Antarctica. *Environ Pollut.* 235:302-311.
- Kim JT, Son MH, Kang JH, Kim JH, Jung JW, Chang YS (2015). Occurrence of Legacy and New Persistent Organic Pollutants in Avian Tissues from King George Island, Antarctica. *Environ Sci Technol.* 17;49(22):13628-38. doi: 10.1021/acs.est.5b03181.
- Kim J, Son M, Shin E, Choi S-D, Chang Y-S (2016). Occurrence of Dechlorane compounds and polybrominated diphenyl ethers (PBDEs) in the Korean general population. *Environmental Pollution* 212; 330-336.
- Kim J, Son M, Kim J, Suh J, Kang Y (2014). Assessment of Dechlorane compounds in foodstuffs obtained from retail markets and estimation of dietary intake in Korean population. *J. Hazard. Materials.* 275, 19-25.
- Klasmeier J, Matthies M, MacLeod M, Fenner K, Scheringer M, Stroebe M, Le Gall AC, McKone TE, van de Meent D, Wania F (2006). Application of multimedia models for screening assessment of long-range transport potential and overall persistence. *Environ Sci Technol* 40:53-60.
- Klosterhaus SL, Stapleton HM, La Guardia MJ, Greig DJ (2012). Brominated and chlorinated flame retardants in San Francisco Bay sediments and wildlife. *Environ Int.* 15; 47:56-65.
- Kurt-Karakus PB, Muir DC, de Jourdan B, Teixeira C, Martindale JE, Embers H, Wang X, Keir M, Backus S (2019). Bioaccumulation of selected halogenated organic flame retardants in Lake Ontario. *Environmental Toxicology and Chemistry—Volume 38, Number 6—pp. 1198–1210.*
- Larisch W, Goss KU (2018). Modelling oral up-take of hydrophobic and super-hydrophobic chemicals in fish. *Environ Sci Process Impacts.* 20(1):98-104.
- Lee HK, Kang H, Lee S, Kim S, Choi K, Moon HB (2020) Human exposure to legacy and emerging flame retardants in indoor dust: A multiple-exposure assessment of PBDEs, *Sci Total Environ* 719: 137386.
- Letcher RJ, Morris AD, Dyck M, Sverko E, Reiner EJ, Blair DAD, Chu SG, Shen L (2018). Legacy and new halogenated persistent organic pollutants in polar bears from a contamination hotspot in the Arctic, Hudson Bay Canada. *Sci Total Environ.* 2018 Jan 1;610-611:121-136. doi: 10.1016/j.scitotenv.2017.08.035.
- Li H, Song A, Liu H, Li Y, Liu M, Sheng G, Peng P, Ying G (2021a). Occurrence of Dechlorane series flame retardants in sediments from the Pearl River Delta, South China. *Environ Pollut.* 279:116902. doi: 10.1016/j.envpol.2021.116902.

- Li Z-R, Luo X-J, Lin L, Zheng Y-H, Mai B-X (2021b). Effect of laying sequence and selection of maternal tissues in assessment of maternal transfer of organohalogenated contaminants during chicken egg formation: A pilot study. *Environmental Pollution* 270, 116157. <https://doi.org/10.1016/j.envpol.2020.116157>.
- Li B, Chen J, Du Q, Wang B, Qu Y, Chang Z (2020a). Toxic effects of dechlorane plus on the common carp (*Cyprinus carpio*) embryonic development. *Chemosphere* 249, 126418. <https://doi.org/10.1016/j.chemosphere.2020.126481>.
- Li B, Chen J, Wang S, Qi P, Chang X, Chang Z (2020b). Effects of dechlorane plus on intestinal barrier function and intestinal microbiota of *Cyprinus carpio* L. *Ecotoxicology and Environmental Safety*. 204: 111124.
- Li ZR, Luo XJ, Luo YL, Zeng YH, Mai BX (2019a). Comparative study of dechlorane plus (DP) in adult chickens and developing embryos: Stereo-selective bioaccumulation of DP in chickens. *Environ Pollut*. 247:550-555.
- Li B, Qi P, Qu Y, Wang B, Chen J, Chang Z (2019b). Effects of dechlorane plus on oxidative stress, inflammatory response, and cell apoptosis in *Cyprinus carpio*. *Drug and Chemical toxicol*. DOI: 10.1080/01480545.2019.1701001.
- Li TY, Ge JL, Pei J, Bao LJ, Wu CC, Zeng EY (2019c). Emissions and Occupational Exposure Risk of Halogenated Flame Retardants from Primitive Recycling of E-Waste. *Environ Sci Technol*. 53(21):12495-12505.
- Li N, Chen XW, Deng WJ, Giesy JP, Zheng HL (2018) PBDEs and Dechlorane Plus in the environment of Guiyu, Southeast China: a historical location for E-waste recycling (2004, 2014). *Chemosphere* 199:603–611.
- Li WL, Liu LY, Song WW, Zhang ZF, Qiao LN, Ma WL, Li YF (2016). Five-year trends of selected halogenated flame retardants in the atmosphere of Northeast China. *Sci Total Environ*. 1;539 :286-293.
- Li L, Wang W, Lv Q, Ben Y, Li X (2014). Bioavailability and tissue distribution of Dechloranes in wild frogs (*Rana limnocharis*) from an e-waste recycling area in Southeast China. *J of Environ. Sci*. 26: 636–642.
- Li Y, Yu L, Zhu Z, Dai J, Mai B, Wu J, Wang J (2013a). Accumulation and effects of 90-day oral exposure to Dechlorane Plus in quail (*Coturnix coturnix*). *Environ Toxicol Chem*. 32(7):1649-54.
- Li Y, Yu L, Wang J, Wu J, Mai B, Dai J (2013b). Accumulation pattern of Dechlorane Plus and associated biological effects on rats after 90 d of exposure. *Chemosphere*. 90(7):2149-56.
- Liang X, Li W, Martyniuk CJ, Zha J, Wang Z, Cheng G, Giesy JP (2014). Effects of dechlorane plus on the hepatic proteome of juvenile Chinese sturgeon (*Acipenser sinensis*). *Aquat Toxicol*. 148:83-91.
- Liu Y, Luo X, Zeng Y, Tu W, Deng M, Wu Y, Mai B (2020). Species-specific biomagnification and habitat-dependent trophic transfer of halogenated organic pollutants in insect-dominated food webs from an e-waste recycling site. *Environment International* 138, 105674. <https://doi.org/10.1016/j.envint.2020.105674>.
- Liu X, Bing H, Chen Y, Li J, Wu Y, Zhang G (2018). Brominated flame retardants and dechlorane plus on a remote high mountain of the eastern Tibetan Plateau: implications for regional sources and environmental behaviors. *Environ Geochem Health*. 40(5):1887-1897.
- Liu LY, Salamova A, Venier M, Hites RA (2016a). Trends in the levels of halogenated flame retardants in the Great Lakes atmosphere over the period 2005-2013. *Environ Int*. 92-93:442-9.
- Liu, D., et al. (2016b). Occurrence and concentrations of halogenated flame retardants in the atmospheric fine particles in Chinese cities. *Environ. Sci. Technol*. 50 (18), 9846–9854.
- Liu L-Y, Salamova A, Hites RA (2014). Halogenated flame retardants s in Baby Food from the United States and China and the estimated dietary intakes by infants. *Environ Sci Technol* 48:9812-9818.
- Løseth ME, Briels N, Flo J, Malarvannan G, Poma G, Covaci A, Herzke D, Nygård T, Bustnes JO, Jenssen BM, Jaspers VLB (2019). White-tailed eagle (*Haliaeetus albicilla*) feathers from Norway are suitable for monitoring of legacy, but not emerging contaminants. *Sci Total Environ*. 647:525-533. doi:10.1016/j.scitotenv.2018.07.333.
- Mahmood, A., et al. (2015). Dietary exposure and screening-level risk assessment of polybrominated diphenyl ethers (PBDEs) and dechloran plus (DP) in wheat, rice, soil and air along two tributaries of the River Chenab, Pakistan. *Chemosphere* 118, 57–64.
- Malak IA, Cariou R, Guiffard I, Vénisseau A, Dervilly-Pinel G, Jaber F, Le Bizec B (2019). Assessment of Dechlorane Plus and related compounds in foodstuffs and estimates of daily intake from Lebanese population. *Chemosphere*. 235, 492-497.
- Malak IA, Cariou R, Vénisseau A, Dervilly-Pinel G, Jaber F, Babut M, Le Bizec B (2018). Occurrence of Dechlorane Plus and related compounds in catfish (*Silurus spp.*) from rivers in France. *Chemosphere*. 207:413-420.
- Marler H, Adams DH, Wu Y, Nielsen CK, Shen L, Reiner EJ, Chen D (2018). Maternal Transfer of Flame Retardants in Sharks from the Western North Atlantic Ocean. *Environ Sci Technol*. 20;52(22):12978-12986.

- Matsutani T, Nomura T (1987). In vitro effects of serotonin and prostaglandins on release of eggs from the ovary of the scallop, *Patinopecten yessoensis*. *Gen. Comp. Endocrinol.* 67, 111–118.
- Mo L, Zheng X, Zhu C, Sun Y, Yu L, Luo X, Mai B (2019). Persistent organic pollutants (POPs) in oriental magpie-robins from e-waste, urban, and rural sites: Site-specific biomagnification of POPs. *Ecotoxicol Environ Saf.* 186:109758. doi: 10.1016/j.ecoenv.2019.109758.
- Mo L, Wu JP, Luo XJ, Sun YX, Zheng XB, Zhang Q, Zou FS, Mai BX (2013). Dechlorane Plus flame retardant in kingfishers (*Alcedo atthis*) from an electronic waste recycling site and a reference site, South China: influence of residue levels on the isomeric composition. *Environ Pollut.* 174:57-62.
- Mortensen Å-K, Verreault J, François A, Houde M, Giraudo M, Damd M, Jenssen BJ (2022). Flame retardants and their associations with thyroid hormone-related variables in northern fulmars from the Faroe Islands. *Sci.Tot. Environ.* 806 (2022) 150506. <https://doi.org/10.1016/j.scitotenv.2021.150506>
- Muñoz-Arnanz J, Panti C, Bainsi M, Marsili L, Lauriano G, Panigada S, Fossi, MC, Jiménez B (2016). Assessment of contamination by POPs in killer whales (*Orcinus orca*) from the Ross Sea, Antarctica. *Dioxin* 2016.
- Muñoz-Arnanz J, Roscales JL, Vicente A, Aguirre JI, Jiménez B (2012). Dechlorane Plus in eggs of two gull species (*Larus michahellis* and *Larus audouinii*) from the southwestern Mediterranean Sea. *Anal Bioanal Chem.* 404(9):2765-73.
- Muñoz-Arnanz J, Sáez M, Hiraldo F, Baos R, Pacepavicius G, Alae M, Jiménez B (2011). Dechlorane plus and possible degradation products in white stork eggs from Spain. *Environ Int.* 37(7):1164-8.
- Möller A, Xie Z, Cai M, Sturm R, Ebinghaus R (2012). Brominated flame retardants and dechlorane plus in the marine atmosphere from Southeast Asia toward Antarctica. *Environ Sci Technol.* 46(6):3141-8.
- Möller A, Xie Z, Cai M, Zhong G, Huang P, Cai M, Sturm R, He J, Ebinghaus R (2011). Polybrominated diphenyl ethers vs alternate brominated flame retardants and dechloranes from East Asia to the Arctic. *Environmental Science and Technology*, 45:6793-6799.
- Möller A, Xie Z, Sturm R, Ebinghaus R (2010). Large-scale distribution of dechlorane plus in air and seawater from the Arctic to Antarctica. *Environ Sci Technol.* 44(23):8977-82.
- Na G, Yao Y, Gao H, Li R, Ge L, Titaley IA, Santiago-Delgado L, Massey Simonich SL (2017). Trophic magnification of Dechlorane Plus in the marine food webs of Fildes Peninsula in Antarctica. *Mar Pollut Bull.* 117(1-2):456-461.
- Na G, Wei W, Zhou S, Gao H, Ma X, Qiu L, Ge L, Bao C, Yao Z (2015). Distribution characteristics and indicator significance of Dechloranes in multi-matrices at Ny-Ålesund in the Arctic. *J Environ Sci (China)*. 1;28 :8-13.
- Newton S, Sellström U, de Wit CA (2015). Emerging flame retardants, PBDEs, and HBCDDs in indoor and outdoor media in Stockholm, Sweden. *Environ Sci Technol.* 49(5):2912-20.
- Niu N, Feng L, Lin Y, Li X, Zhang, D and Yao S (2020). The sources of Dechlorane Plus (DP) in surface sediment from Bohai Sea and the northern part of the Yellow Sea, China: Evidence from the fractional abundance of anti-DP (f anti) combined with lignin biomarker. *Regional Studies in Marine Science.* 39. 101437. 10.1016/j.rsma.2020.101437.
- Norwegian Environment Agency (2019a). Environmental Contaminants in an Urban fjord. Report L. Nr. 7410-2019/M-1441.
- Norwegian Environment Agency (2019b). Prioritetslisten. Published 18.01.2019. Available at: <http://www.miljostatus.no/prioritetslisten> (Accessed 27.03.2019).
- Norwegian Environment Agency (2018a). Screening program 2017. AMAP Assessment compounds. Report M-1080.
- Norwegian Environment Agency (2018b). Environmental Contaminants in an Urban fjord. Report M-1131.
- Noyes PD, Haggard DE, Gonnerman GD and Tanguay RL (2015). Advanced morphological behavioral test platform reveals neurodevelopmental defects in embryonic zebrafish exposed to comprehensive suite of halogenated and organophosphate flame retardants. *Toxicol. Sci.*, 145, 177-195.
- Olukunle OI, Lehman DC, Salamova A, Venier M, Hites RA (2018). Temporal trends of Dechlorane Plus in air and precipitation around the North American Great Lakes. *Sci Total Environ.* 15; 642:537-542.
- OxyChem® Dechlorane Plus® (2013). Available at: <https://www.oxy.com/OurBusinesses/Chemicals/ResponsibleCare/Documents/Dechlorane%20Plus.pdf>.
- OxyChem® Dechlorane Plus® Manual. (2007). Ver. 07-27-2007. Available from: https://www.oxy.com/OurBusinesses/Chemicals/Products/Documents/dechloranepius/dechlorane_plus.pdf.

- OxyChem® Dechlorane Plus® (2004). Available at: <https://www.oxy.com/pages/search.aspx?k=dechlorane%20plus>
- OxyChem test-plan for Dechlorane plus (2004b). Available at: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.578.5473&rep=rep1&type=pdf>.
- Pan H-Y, Li J-F-T, Li X-H, Yang Y-L, Qin Z-F, Li J-B, Li Y-Y (2020). Transfer of dechlorane plus between human breast milk and adipose tissue and comparison with legacy lipophilic compounds. *Environmental Pollution*, 265: 115096. <https://doi.org/10.1016/j.envpol.2020.115096>.
- Peng Y, Wu J, Luo X, Zhang X, Giesy JP, Mai B (2019). Spatial distribution and hazard of halogenated flame retardants and polychlorinated biphenyls to common kingfisher (*Alcedo atthis*) from a region of South China affected by electronic waste recycling. *Environ Int.* 130:104952.
- Peng Y, Wu JP, Tao L, Mo L, Zheng XB, Tang B, Luo XJ, Mai BX (2015). Accumulation of Dechlorane Plus flame retardant in terrestrial passerines from a nature reserve in South China: the influences of biological and chemical variables. *Sci Total Environ.* 1, 514:77-82.
- Peng H, Wan Y, Zhang K, Sun J, Hu J (2014). Trophic transfer of dechloranes in the marine food web of Liaodong Bay, north China. *Environ Sci Technol.* 20;48(10):5458-66.
- Peng H, Zhang K, Wan Y, Hu J (2012). Tissue distribution, maternal transfer, and age-related accumulation of dechloranes in Chinese sturgeon. *Environ Sci Technol.* 18;46(18):9907-13.
- Peshdary V, Styles G, Rigden M, Caldwell D, Kawata A, Sorisky A, Atlas E (2020). Exposure to low doses of Dechlorane Plus promotes adipose tissue dysfunction and glucose intolerance in male mice. *Endocrinol.* 161(7):1-15.
- Peshdary V, Calzadilla G, Landry A, Sorisky A, Atlas E (2019). Dechlorane Plus increases adipogenesis in 3T3-L1 and human primary preadipocytes independent of peroxisome proliferator-activated receptor γ transcriptional activity. *Int J Obes (Lond).* 43(3):545-555.
- Poma G, Malysheva SV, Gosciny S, Malarvannan G, Voorspoels S, Covaci A, Van Loco J (2018). Occurrence of selected halogenated flame retardants in Belgian foodstuff. *Chemosphere.* 194:256-265.
- PubChem, 2021. Dechlorane Plus. <https://pubchem.ncbi.nlm.nih.gov/compound/Dechlorane-plus>
- Qiao L, Zheng X-B, Zheng J, Chen S-J, Zhong C-Q, Chen J-H, Yang Z-Y, Mai B-X (2019). Legacy and Currently Used Organic Contaminants in Human Hair and Hand Wipes of Female E-Waste Dismantling Workers and Workplace Dust in South China. *Environ. Sci. Technol.* 53, 5, 2820–2829 <https://doi.org/10.1021/acs.est.8b05503>
- Qiu X, Hites RA (2008). Dechlorane plus and other flame retardants in tree bark from the northeastern United States. *Environ Sci Technol.* 42(1):31-6.
- Qiu XH, Marvin CH and Hites RA (2007). Dechlorane plus and other flame retardants in a sediment core from Lake Ontario. *Environ. Sci. Technol.* 41 (17), 6014–6019.
- Qiu X, Zhu T, Hu J (2010). Polybrominated diphenyl ethers (PBDEs) and other flame retardants in the atmosphere and water from Taihu Lake, East China. *Chemosphere.* 80(10):1207-12.
- Rauert C, Schuster JK, Eng A, Harner T (2018). Global Atmospheric Concentrations of Brominated and Chlorinated Flame Retardants and Organophosphate Esters. *Environ Sci Technol.*;52(5):2777-2789. doi: 10.1021/acs.est.7b06239.
- Ren G, Wang Z, Yu Z, Wang Y, Ma S, Wu M et al. (2013). Primary investigation on contamination pattern of legacy and emerging halogenated organic pollutions in freshwater fish from Liaohe River, Northeast China. *Environ Pollut* 172:94–99.
- Ren GF, Yu ZQ, Ma ST, Li HR, Peng PG, Sheng GY, Fu JM (2009). Determination of Dechlorane Plus in serum from electronics dismantling workers in South China. *Environ Sci Technol* 43 (24), 9453–9457.
- Ren NQ, Sverko E, Li YF, Zhang Z, Harner T, Wang DG, Wan XN and Mccarry BE (2008). Levels and isomer profiles of Dechlorane Plus in Chinese air. *Environ. Sci. Technol.* 42 (17), 6476–6480.
- Rjabova J, Bartkevics V, Zacs D (2016). The occurrence of Dechlorane Plus and related norbornene-based flame retardants in Baltic wild salmon (*Salmo salar*). *Chemosphere* 147, 210-217.
- Roscales JL, González-Solís J, Zango L, Ryan PG, Jiménez B (2016). Latitudinal exposure to DDTs, HCB, PCBs, PBDEs and DP in giant petrels (*Macronectes* spp.) across the Southern Ocean. *Environ Res.* 148:285-294. doi: 10.1016/j.envres.2016.04.005.
- Saini A, Harner T, Chinnadhurai S, Schuster JK, Yates A, Sweetman A, Aristizabal-Zuluaga BH, Jiménez B, Manzano CA, Gaga EO, Stevenson G, Falandysz J, Ma J, Miglioranza KSB, Kannan K, Tominaga M, Jariyasopit N, Rojas NY, Amador-Muñoz O, Sinha R, Alani R, Suresh R, Nishino T, Shoeib T (2020). GAPS-megacities: A new global platform for investigating persistent organic pollutants and chemicals of emerging concern in urban air. *Environ Pollut.* 267:115416. doi: 10.1016/j.envpol.2020.115416.

- Salamova A, Hites RA (2011). Dechlorane plus in the atmosphere and precipitation near the Great Lakes. *Environ Sci Technol.* 45(23):9924-30.
- Salamova A, Hermanson MH, Hites RA (2014). Organophosphate and halogenated flame retardants in atmospheric particles from a European Arctic site. *Environ Sci Technol.* 48(11):6133-40.
- Schlabach M, Remberger M, Brorström-Lunden E, Norström K, Kaj L, Andersson H, Herzke D, Borgen A, Harju M (2011). Brominated flame retardants (BFR) in the Nordic Environment. *TemaNord 2011:528*. Nordic Council of Ministers, Copenhagen.
- Schwemmer P, Corman AM, Koch D, Borrmann RM, Koschorreck J, Voigt CC, Garthe S (2019). Charakterisierung der Habitatwahl von Silbermöwen *Larus argentatus* durch GPS-Datenlogger zur Einschätzung der Schadstoffbelastung an der deutschen Nordseeküste. *Vogelwarte* 57 (1), 13-30.
- Schuster JK, Harner T and Sverko E (2021). Dechlorane Plus in the Global Atmosphere. *Environ. Sci. Technol. Lett.* 2021, 8, 1, 39-45. <https://doi.org/10.1021/acs.estlett.0c00758>.
- Shanmuganathan M, Zhang Z, Sverko E, Brymer R, Gill B, Smyth SA, Marvin CH (2018). Analysis of halogenated flame retardants in Canadian wastewater treatment plants using gas chromatography–tandem mass spectrometry (GC-MS/MS). *Water Quality Research Journal* 53.4: XX-XX.
- She YZ, Wu JP, Zhang Y, Peng Y, Mo L, Luo XJ, Mai BX (2013). Bioaccumulation of polybrominated diphenyl ethers and several alternative halogenated flame retardants in a small herbivorous food chain. *Environ Pollut.* 174:164-70.
- Shunthirasingham C, Alexandrou N, Brice KA, Dryfhout-Clark H, Su K, Shin C, Park R, Pajda A, Noronha R, Hung H (2018). Temporal trends of halogenated flame retardants in the atmosphere of the Canadian Great Lakes Basin (2005-2014). *Environ Sci Process Impacts.* 20(3):469-479.
- Shen L, Reiner EJ, Helm PA, Marvin CH, Hill B, Zhang X, Macpherson KA, Kolic TM, Tomy GT, Brindle ID (2011). Historic trends of dechloranes 602, 603, 604, dechlorane plus and other norbornene derivatives and their bioaccumulation potential in Lake Ontario. *Environ Sci Technol.* 45(8):3333-40.
- Shoeib M, Harner T, Webster GM, Sverko E, Cheng, Y (2012). Legacy and current-use flame retardants in house dust from Vancouver, Canada. *Environ Pollut* 169:175-182.
- Siddique S, Xian QM, Abdelouahab N, Takser L, Phillips SP, Feng YL, Wang B, Zhu JP (2012). Levels of dechlorane plus and polybrominated diphenylethers in human milk in two Canadian cities. *Environ. Int.* 39 (1), 50–55.
- Simond AE, Houde M, Lesage V, Verreault J (2017). Temporal trends of PBDEs and emerging flame retardants in belugas from the St. Lawrence Estuary (Canada) and comparisons with minke whales and Canadian Arctic belugas. *Environ Res.* 156:494-504.
- Smythe TA, Mattioli LC, Letcher RJ (2020). Distribution behaviour in body compartments and in ovo transfer of flame retardants in North American Great Lakes herring gulls. *Environmental Pollution.* 262, 114306. <https://doi.org/10.1016/j.envpol.2020.114306>
- SIN LIST (2020). International Chemical Secretariat's (ChemSec) Substitute It Now (SIN) list since 2014. Available at: <https://sinsearch.chemsec.org/chemical/13560-89-9>
- Sühling R, Freese M, Schneider M, Schubert S, Pohlmann JD, Alae M, Wolschke H, Hanel R, Ebinghaus R, Marohn L (2015). Maternal transfer of emerging brominated and chlorinated flame retardants in European eels. *Sci Total Environ.* 530-531:209-218.
- Sühling R, Byer J, Freese M, Pohlmann JD, Wolschke H, Moller A, Hodson PV, Alae M, Hanel R, Ebinghaus R (2014). Brominated flame retardants and Dechloranes in European and American eels from glass to silver life stages *Chemosphere* 116, 104–111.
- Sun YX, Hu YX, Zhang ZW, Xu XR, Li HX, Zuo LZ, Zhong Y, Sun H, Mai BX (2017). Halogenated organic pollutants in marine biota from the Xuande Atoll, South China Sea: Levels, biomagnification and dietary exposure. *Mar Pollut Bull.* 118(1-2):413-419.
- Sun YX, Zhang ZW, Xu XR, Hu YX, Luo XJ, Cai MG, Mai BX (2015). Bioaccumulation and biomagnification of halogenated organic pollutants in mangrove biota from the Pearl River Estuary, South China. *Mar Pollut Bull.* 15:99(1-2):150-6.
- Sverko E, Tomy GT, Reiner EJ, Li YF, McCarry BE, Arnot JA, Law RJ and Hites RA (2011). Dechlorane Plus and related compounds in the environment: a review. *Environ. Sci. Technol.* 45 (12), 5088–5098.
- Sverko E, Reiner EJ, Tomy GT, McCrindle R, Shen L, Aresenault G, Zaruk D, MacPherson KA, Marvin CH, Helm PL, McCarry BE (2010). Compounds structurally related to Dechlorane Plus in sediments and biota from Lake Ontario (Canada). *Environ. Sci. Technol.* 44, 574–579.

- Sverko E, Tomy GT, Marvin CH, Zaruk D, Reiner E, Helm PA, Hill B, McCarry BE (2008). Dechlorane plus levels in sediment of the lower Great Lakes. *Environ Sci Technol.* 42(2):361-6.
- Syed JH, Iqbal M, Breivik K, Chaudhry MJI, Shah Nawaz M, Abbas Z, Nasir J, Rizvi SHH, Taqi MM, Li J, Zhang G. (2020). Legacy and emerging flame retardants (FRs) in the urban atmosphere of Pakistan: Diurnal variations, gas-particle partitioning and human health exposure. *Sci Total Environ.* 2020 Nov 15;743:140874. doi: 10.1016/j.scitotenv.2020.140874.
- Syed, J.H., et al. (2013). Levels, profile and distribution of dechlorane plus (DP) and polybrominated diphenyl ethers (PBDEs) in the environment of Pakistan. *Chemosphere* 93 (8), 1646–1653.
- Tang B, Luo XJ, Huang CC, Sun RX, Wang T, Zeng YH, Mai BX (2018). Stereoselective bioaccumulation of syn- and anti-Dechlorane plus isomers in different tissues of common carp (*Cyprinus carpio*). *Sci Total Environ.* 616-617:1339-1346.
- Tao W, Tian J, Xu T, Xu L, Xie HQ, Zhou Z, Guo Z, Fu H, Yin X, Chen Y, Xu H, Zhang S, Zhang W, Ma C, Ji F, Yang J, Zhao B (2018). Metabolic profiling study on potential toxicity in male mice treated with Dechlorane 602 using UHPLC-ESI-IT-TOF-MS. *Environ Pollut.* 246: 141-147.
- Tao W, Zhou Z, Shen L and Zhao B (2015). Determination of dechlorane flame retardants in soil and fish at Guiyu, an electronic waste recycling site in south China. *Environmental Pollution*, 206, 361-368.
- Taverna R, Gloor R, Maier U, Zennegg M, Figi R, Birchler E (2017). Stoffflüsse im Schweizer Elektronikschrott. Metalle, Nichtmetalle, Flammschutzmittel und polychlorierte Biphenyle in elektrischen und elektronischen Kleingeräten. Bundesamt für Umwelt, Bern. Umwelt-Zustand Nr. 1717 (full report in German: <https://www.bafu.admin.ch/uz-1717-d>; summary in English: <https://www.bafu.admin.ch/uz-1717-e>).
- Tomy GT, Sverko E, Palace V, Rosenberg B, McCrindle R, McAlees A, Smith LA, Byer J, Pacepavicius G, Alae M, McCarry BE (2013). Dechlorane plus monoadducts in a Lake Ontario (Canada) food web and biotransformation by lake trout (*Salvelinus namaycush*) liver microsomes. *Environ Toxicol Chem.* 32(6):1376-81.
- Tomy GT, Thomas CR, Zidane TM, Murison KE, Pleskach K, Hare J, Arsenault G, Marvin CH, Sverko E (2008). Examination of isomer specific bioaccumulation parameters and potential in vivo hepatic metabolites of syn- and anti-Dechlorane Plus isomers in juvenile rainbow trout (*Oncorhynchus mykiss*). *Environ Sci Technol.* 42(15):5562-7.
- Tomy GT, Pleskach K, Ismail N, Whittle DM, Helm PA, Sverko E, Zaruk D, Marvin CH (2007). Isomers of dechlorane plus in Lake Winnipeg and Lake Ontario food webs. *Environ Sci Technol.* 1;41(7):2249-54.
- TOPKAT, Toxicity Prediction by Komputer Assisted Technology, Prediction Module. (2004). Version 6.2. San Diego (CA): Accelrys Software Inc.
- UK EA (2012). Depuration Rate Constant as an Indicator of Bioaccumulation Potential. Environment Agency, Bristol, UK. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/291525/LIT_7371_44228a.pdf.
- US EPA (1978). Reviews of the Environmental Effects of Pollutants: XII. Hexachlorocyclopentadiene.
- US EPA, US Environmental Protection Agency (2009). HPV Challenge; Robust Summaries and Test Plans: Dechlorane Plus, cover letter dated on March 3, 2009 (doc. no. 201-16774), IUCLID report dated November 7, 2008.
- US EPA, US Environmental Protection Agency (2011). Screening-level hazard characterization – Dechlorane plus (CASRN 13560-89-9).
- US EPA, US Environmental Protection Agency (2012). Estimation Programs Interface Suite™ for Microsoft® Windows, v4.11. U.S. Environmental Protection Agency, Washington, DC, USA. Available online at <http://www.epa.gov/opptintr/exposure/pubs/episuitedi.htm> as of November 2012.
- US EPA, US Environmental Protection Agency (2014). An Alternatives Assessment for the Flame Retardant Decabromodiphenyl Ether (DecaBDE), January 2014. Available at: <https://www.epa.gov/saferchoice/partnership-evaluate-flame-retardant-alternatives-decabde-publications>.
- US EPA, US Environmental Protection Agency (2020). The Toxic Substances Control Act (TSCA) Chemical Substance Inventory. Available at: <https://www.epa.gov/tscainventory> (accessed 17.04.2020).
- Verreault J, Letcher RJ, Gentes ML, Braune BM (2018). Unusually high Deca-BDE concentrations and new flame retardants in a Canadian Arctic top predator, the glaucous gull. *Sci Total Environ.*;639:977-987. doi:10.1016/j.scitotenv.2018.05.222.
- Velsicol Chemicals LLC (2020). PyroVex® Flame Retardants. Excellent Fire Retardant Additives from an experienced manufacturer. Available at: <https://www.velsicol.com/images/pdf/Pyrovex/sds/pyrovex-sg-sds.pdf> (accessed 06.02.2020).

- Vorkamp K, Rigét F, Sanderson H, Bossi R, Hansen KM, Skov H (2019a). POP/PBT characterisation of dechlorane plus and novel brominated flame retardants based on data from Greenland. Aarhus University, DCE – Danish Centre for Environment and Energy, 80 pp. Scientific Report from DCE – Danish Centre for Environment and Energy No. 339. Available at: <https://dce2.au.dk/pub/SR339.pdf> (accessed 20.04.2020).
- Vorkamp K, Balmer J, Hung H, Letcher RJ, Rigét FF, de Wit CA (2019b). Current-use halogenated and organophosphorous flame retardants: A review of their presence in Arctic ecosystems, *Emerging Contaminants*, Volume 5, Pages 179-200, ISSN 2405-6650, <https://doi.org/10.1016/j.emcon.2019.05.004>.
- Vorkamp K, Falk K, Møller S, Rigét FF, Sørensen PB (2018). Regulated and Unregulated Halogenated Flame Retardants in Peregrine Falcon Eggs from Greenland. *Environ Sci Technol*. 16;52(2):474-483.
- Vorkamp K, Bossi R, Rigét FF, Skov H, Sonne C, Dietz R. (2015). Novel brominated flame retardants and dechlorane plus in Greenland air and biota. *Environmental Pollution*. 196:284-291.
- Wang S, Romanak KA, Hendryx M, Salamova A, Venier M (2020a). Association between Thyroid Function and Exposures to Brominated and 3 Organophosphate Flame Retardants in rural Central Appalachia. *Environ. Sci. Technol.*, 2020 54(1):325-334. DOI: 10.1021/acs.est.9b04892.
- Wang Y, eris A, Rifat MR, Ahmed SI, Aich N, Nguyen LV, Urik J, Eljarrat E, Vrana B, Jantunen LM, Diamond ML (2020b). Measuring exposure of e-waste dismantlers in Dhaka Bangladesh to organophosphate esters and halogenated flame retardants using silicone wristbands and T-shirts, *Sci Total Environ* 720 :137480.
- Wang D, Jia H, Hong WJ, Xue X, Sun Y, Li YF, Guo W (2019). Uptake, depuration, bioaccumulation, and selective enrichment of dechlorane plus in common carp (*Cyprinus carpio*). *Environ Sci Pollut Res Int*. doi:10.1007/s11356-019-07239-8. (Epub ahead of print) PubMed PMID: 31865567.
- Wang Q, Kelly BC (2017). Occurrence, distribution and bioaccumulation behaviour of hydrophobic organic contaminants in a large-scale constructed wetland in Singapore. *Chemosphere*. 183:257-265.
- Wang P, Zhang Q, Zhang H, Wang T, Sun H, Zheng S, Li Y, Liang Y, Jiang G. (2016). Sources and environmental behaviors of Dechlorane Plus and related compounds - A review. *Environ Int*. 88:206-220.
- Wang DG, Guo MX, Pei W, Byer JD, Wang Z (2015). Trophic magnification of chlorinated flame retardants and their dechlorinated analogs in a fresh water food web. *Chemosphere*. 118:293-300.
- Wang Y, Xu M, Jin, He S, Li M, Sun Y (2014). Concentrations and relationships between classes of persistent halogenated organic compounds in pooled human. *Science of the Total Environment* 482–483, 276–282.
- Wang DG, Alae M, Byer JD, Brimble S, Pacepavicus G (2013a). Human health risk assessment of occupational and residential exposures to dechlorane plus in the manufacturing facility area in China and comparison with e-waste recycling site. *Sci. Total Environ*. 445-446, 329-336.
- Wang S, Huang J, Yang Y, Yu G, Deng S, Wang B (2013b). Photodegradation of Dechlorane Plus in n-nonane under the irradiation of xenon lamp. *J Hazard Mater*. 15; 260: 16-23.
- Wang L, Jia HL, Liu, XJ, Yang M, Hong WJ, Sun YQ, Qi H, Song WW, Lin J, Li YF (2012). Dechloranes in a river in northeastern China: Spatial trends in multi-matrices and bio-accumulation in fish (*Enchelyopuselongatus*). *Ecotoxicol. Environ. Saf*. 84, 262–267.
- Wang DG, Yang M, Qi H, Sverko E, Ma WL, Li YF, Alae M, Reiner EJ and Shen L (2010a). An Asia-specific source of Dechlorane Plus: concentration, isomer profiles, and other related compounds. *Environ. Sci. Technol*. 44 (17): 6608–6613.
- Wang XP, Gong P, Yao TD, Jones KC (2010b). Passive air sampling of organochlorine pesticides, polychlorinated biphenyls, and polybrominated diphenyl ethers across the Tibetan Plateau. *Environmental Science and Technology*. 44: 2988–2993.
- Wang B, Iino F, Huang J, Lu Y, Yu G, Morita M (2010c). Dechlorane Plus pollution and inventory in soil of Huai'an City, China. *Chemosphere*. 80(11):1285-90.
- Wegmann, F.; Cavin, L.; MacLeod, M.; Scheringer, M.; Hungerbühler, K. (2009) The OECD software tool for screening chemicals for persistence and long-range transport potential. *Environmental Modeling and Software* 24(2), 228-237.
- Weil ED, Levchik, SV (2009). *Flame Retardants for Plastics and Textiles: Practical Applications*. Cincinnati, Ohio: Hanser Publishers. 297 pp.
- Wolschke H, Meng X-Z, Xie Z, Ebinghaus R, Cai M (2015). Novel flame retardants (N-FRs), polybrominated diphenyl ethers (PBDEs) and dioxin-like polychlorinated biphenyls (DL-PCBs) in fish, penguin, and skua from King George Island, Antarctica. *Marine Pollution Bulletin* 96, 513–518.

- Wong F, Suzuki G, Michinaka C, Yuan B, Takigami H, de Wit CA (2017). Dioxin-like activities, halogenated flame retardants, organophosphate esters and chlorinated paraffins in dust from Australia, the United Kingdom, Canada, Sweden and China. *Chemosphere* 168: 1248-1256.
- Wu Z, Pan Y, Ma F, Ren Q, Wu J, Jiang Y, He Y, Yu Y, Zhang G (2021). Dechlorane Plus exposure on gut microbiome evaluated by using both *in vivo* and *in vitro* assays. *International Biodeterioration & Biodegradation*. 163, 105255. <https://doi.org/10.1016/j.ibiod.2021.105255>.
- Wu JP, Chen XY, Si-Kang W, Sun Y, Feng WL, Tao L, Luo XJ, Mai BX (2018). Dechlorane Plus flame retardant in a contaminated frog species: Biomagnification and isomer-specific transfer from females to their eggs. *Chemosphere*. 211:218-225.
- Wu Q, Li H, Kuo DTF, Chen S, Mai B, Li H, Liu Z, Deng M, Zhang H, Hu X, Geng X, Chen Y (2017). Occurrence of PBDEs and alternative halogenated flame retardants in sewage sludge from the industrial city of Guangzhou, China. *Environ Pollut*. 220(Pt A):63-71.
- Wu JP, She YZ, Zhang Y, Peng Y, Mo L, Luo XJ, Mai BX (2013). Sex-dependent accumulation and maternal transfer of Dechlorane Plus flame retardant in fish from an electronic waste recycling site in South China. *Environ. Pollut*. 177: 150–155.
- Wu Z, Huang L, Gao Y, Guo Q, Sun Y (2013). The effects of Dechlorane Plus on toxicity. *Appl. Mechanics and Materials*. Vols.380-384, 4163-4166.
- Wu B, Liu S, Guo X, Zhang Y, Zhang X, Li M, Cheng S (2012). Responses of mouse liver to dechlorane plus exposure by integrative transcriptomic and metabolomic studies. *Environ Sci Technol*. 46(19):10758-64.
- Wu JP, Zhang Y, Luo XJ, Wang J, Chen SJ, Guan YT, Mai BX (2010). Isomer-specific bioaccumulation and trophic transfer of Dechlorane Plus in the freshwater food web from a highly contaminated site, South China. *Environ Sci Technol*. 15;44(2):606-11.
- Xian Q, Siddique S, Li T, Feng YL, Takser L, Zhu J (2011). Sources and environmental behavior of dechlorane plus--a review. *Environ Int*. 37(7):1273-84.
- Xiao H, Shen L, Su Y, Barresi E, Dejong M, Hung H, Lei YD, Wania F, Reiner EJ, Sverko E, Kang SC (2012a). Atmospheric concentrations of halogenated flame retardants at two remote locations: The Canadian High Arctic and the Tibetan Plateau. *Environmental Pollution*, 161:154-161.
- Xiang N, Chen L, Meng XZ, Li YL, Liu Z, Wu B, Dai L, Dai X (2014). Polybrominated diphenyl ethers (PBDEs) and dechlorane plus (DP) in a conventional wastewater treatment plant (WWTP) in Shanghai: seasonal variations and potential sources. *Sci Total Environ*. 15; 487:342-9.
- Xiang N, Zhao X, Meng XZ, Chen L (2013). Polybrominated diphenyl ethers (PBDEs) in a conventional wastewater treatment plant (WWTP) from Shanghai, the Yangtze River Delta: implication for input source and mass loading. *Sci Total Environ*. 461-462:391-6.
- Xie J, Sun Y, Cheng Y, Chen Y, Chen L, Xie C, Dai S, Luo X, Zhang L, Mai B (2021). Halogenated flame retardants in surface sediments from fourteen estuaries, South China. *Mar Pollut Bull*. 164:112099. doi: 10.1016/j.marpolbul.2021.112099.
- Yadav IC, Devi NL, Kumar A, Li J, Zhang G (2020) Airborne brominated, chlorinated and organophosphate ester flame retardants inside the buildings of the Indian state of Bihar: Exploration of source and human exposure, *Ecotoxicol Environ Saf* 191:110212.
- Yan X, Zheng J, Chen KH, Yang J, Luo XJ, Yu LH, Chen SJ, Mai BX, Yang ZY (2012). Dechlorane Plus in serum from e-waste recycling workers: influence of gender and potential isomer-specific metabolism. *Environ. Int*. 49, 31–37.
- Yang R, Zhang S, Li X, Luo D, Jing C (2016a). Dechloranes in lichens from the southeast Tibetan Plateau: Evidence of long-range atmospheric transport. *Chemosphere*. 144:446-51.
- Yang Y, Ji F, Cui Y, Li M (2016b). Ecotoxicological effects of earthworm following long-term Dechlorane Plus exposure. *Chemosphere*. 144:2476-81.
- Yang R., Zhang S, L, A, Jiang G, Jing C (2013). Atitudinal and spatial signature of POPs in soil, lichen, conifer needles, and bark of the southeast tibetan plateau: Implications for sources and environmental cycling. *Environmental Science and Technology*, 47, 12736–13743.
- Yin JF, Li JFT, Li XH, Yang YL, Qin ZF (2020). Bioaccumulation and transfer characteristics of dechlorane plus in human adipose tissue and blood stream and the underlying mechanisms. *Sci Total Env*. 700: 134391. <https://doi.org/10.1016/j.scitotenv.2019.134391>.

- Yu Y, Hung H, Alexandrou N, Roach P, Nordin K (2015). Multiyear measurements of flame retardants and organochlorine pesticides in air in Canada's Western sub-Arctic. *Environ. Sci. Technol.* 49, 8623-8630.
- Yu L, Luo X, Zheng X, Zeng Y, Chen D, Wu J, Mai B (2013). Occurrence and biomagnification of organohalogen pollutants in two terrestrial predatory food chains. *Chemosphere.* 93(3):506-11.
- Yu Z, Lu S, Gao S, Wang J, Li H, Zeng X, Sheng G, Fu J (2010). Levels and isomer profiles of Dechlorane Plus in the surface soils from e-waste recycling areas and industrial areas in South China. *Environ Pollut.* 158(9):2920-5.
- Zacs D, Ikkere LE, Bartkevics V (2018). Emerging brominated flame retardants and dechlorane-related compounds in European eels (*Anguilla anguilla*) from Latvian lakes. *Chemosphere.* 197:680-690. doi:10.1016/j.chemosphere.2018.01.105.
- Zafar MI, Kali S, Ali M, Riaz MA, Naz T, Iqbal MM, Masood N, Munawar K, Jan B, Ahmed S, Waseem A, Niazi MBK (2020). Dechlorane Plus as an emerging environmental pollutant in Asia: a review. *Environ Sci Pollut Res Int.* 27(34):42369-42389. doi: 10.1007/s11356-020-10609-2. Epub 2020 Aug 31. PMID: 32864714.
- Zeng Y-H, Luo X-J, Tang B, Zheng X-B and Mai B-X (2014). Gastrointestinal absorption, dynamic tissue-specific accumulation, and isomer composition of dechlorane plus and related analogs in common carp by dietary exposure. *Ecotoxicology and Environmental Safety*, 100, 32-38.
- Zhang J, Zhao J, Sun Y, Xin M, Zhang D, Bian R (2021). Mechanisms of emerging pollutant Dechlorane Plus on the production of short-chain fatty acids from sludge anaerobic fermentation. *Environ Sci Pollut Res.* 28, 34902-34912. <https://doi.org/10.1007/s11356-021-13101-7>.
- Zhang G, Ren Q, Ma S, Wu J, Yang X, Yu Y (2020). Intergenerational transfer of Dechlorane Plus and the associated long-term effects on the structure and function of gut microbiota in offspring. *Environ Int.* 2020; 141: 105770. doi:10.1016/j.envint.2020.105770.
- Zhang X, Sühling R, Serodio D, Bonnell M, Sundin N, Diamond ML (2016). Novel flame retardants: Estimating the physical-chemical properties and environmental fate of 94 halogenated and organophosphate PBDE replacements. *Chemosphere.* 144:2401-7.
- Zhang Q, Zhu C, Zhang H, Wang P, Li Y, Ren D, Jiang G (2015). Concentrations and distributions of Dechlorane Plus in environmental samples around a Dechlorane Plus manufacturing plant in East China. *Sci. Bull.* 60(8):792-797.
- Zhang L, Ji F, Li M, Cui Y, Wu B (2014). Short-term effects of Dechlorane Plus on the earthworm *Eisenia fetida* determined by a systems biology approach. *J Hazard Mater.* 273:239-46. doi: 10.1016/j.jhazmat.2014.03.018.
- Zhang H, Wang P, Li Y, Shang H, Wang Y, Wang T, Zhang Q, Jiang G (2013). Assessment on the Occupational Exposure of Manufacturing Workers to Dechlorane Plus through Blood and Hair Analysis. *Environ. Sci. Technol.* 47, 10567-10573.
- Zhang Y, Wu JP, Luo XJ, Wang J, Chen S-J and Mai BX (2011). Tissue distribution of Dechlorane Plus and its dechlorinated analogs in contaminated fish: High affinity to the brain for anti-DP. *Environmental Pollution*, 159, 3647-3652.
- Zhao S, Tian L, Zou Z, Liu X, Zhong G, Mo Y, Wang Y, Tian Y, Li J, Guo H and Zhang G (2021). Probing Legacy and Alternative Flame Retardants in the Air of Chinese Cities. *Environmental Science & Technology Article ASAP.* DOI: 10.1021/acs.est.0c07367.
- Zhao L, Gong N, Mi D, Luan C, Shao K, Jia H, Sun Y (2014). Kinetics of stereoselective enrichment of Dechlorane Plus in *Ulva Pertusa*. *Chemosphere.* 111:580-6.
- Zhen X, Tang J, Liu L, Wang X, Li Y, Xie Z (2018). From headwaters to estuary: Distribution and fate of halogenated flame retardants (HFRs) in a river basin near the largest HFR manufacturing base in China. *Sci. Total Environ.* 621: 1370-1377. doi:10.1016/j.scitotenv.2017.10.091.
- Zheng XB, Luo XJ, Zeng YH, Wu JP, Mai BX (2014a). Sources, gastrointestinal absorption and stereo-selective and tissue-specific accumulation of Dechlorane Plus (DP) in chicken. *Chemosphere.* 114:241-6.
- Zheng XB, Luo XJ, Zeng YH, Wu JP, Chen SJ, Mai BX (2014b). Halogenated flame retardants during egg formation and chicken embryo development: maternal transfer, possible biotransformation, and tissue distribution. *Environ Toxicol Chem.* 33(8):1712-9.
- Zheng J, Wang J, Luo XJ, Tian M, He LY, Yuan JG, Mai BX, Yang ZY (2010). Dechlorane Plus in human hair from an e-waste recycling area in South China: comparison with dust. *Environ Sci Technol.* 15:44(24):9298-303.
- Zhou P, Zheng T, Li Y, Zhang X, Feng J, Wei Y, Wang H, Yao Y, Gong F, Tian W and Sun L (2021). Chlorinated Flame-Retardant Dechlorane 602 Potentiates Type 2 Innate Lymphoid Cells and Exacerbates Airway Inflammation. *Environ Sci Technol.* 55(2):1099-1109. doi: 10.1021/acs.est.0c03758.
- Zhou SN, Siddique S, Lavoie L, Takser L, Abdelouahab N, Zhu J (2014). Hexachloronorborene-based flame retardants in humans: Levels in maternal serum and milk. *Environ Int* 66:11-17.

Zhu J, Zhao L, Guo L (2022). Dechloranes exhibit binding potency and activity to thyroid hormone receptors, *Journal of Environmental Sciences*. 112, 16-24. <https://doi.org/10.1016/j.jes.2021.04.030>.

Zhu C, Sun Y, Li D, Zheng X, Peng X, Zhu T, Mo L, Luo X, Xu X, Mai B (2020). Evidence for complex sources of persistent halogenated compounds in birds from the south China sea. *Environ Res*;185:109462.

Zhu B, Lai NL, Wai TC, Chan LL, Lam JC, Lam PK (2014). Changes of accumulation profiles from PBDEs to brominated and chlorinated alternatives in marine mammals from the South China Sea. *Environ Int*. 66:65-70.
