## Schedule 4

Schedule	4					
Inspected Items		Package Type	Number of Packages per Lot (N)	Number of Packages Opened for Sampling (n)	Quantity of Collected Specimens (kg)	Number of Specimens
Microorganisms		Not specified	150 151 ~ 1,200 1,201	3 5 8	0.3 0.3 0.3	1 1 1
Irradiation		Not specified	50 51 ~ 500 501 ~ 3,200	2 3 5	0.5 <sup>*1</sup> 0.5 <sup>*1</sup> 0.5 <sup>*1</sup> 0.5 <sup>*1</sup> 0.5 <sup>*1</sup>	1 1 1 1
Radioactive substances		Not specified	3,201 50 51 ~ 150 151 ~ 500 501 ~ 3,200 3,201 ~ 35,000 35,001	3 5 8 13 20 32	1 1 1 1 1 1	1 1 1 1 1 1
Acid value, Peroxide value		Not specified	50 51 ~ 500 501 ~ 3,200 3,201	2 3 5 8	1.5 1.5 1.5 1.5	1 1 1 1
	(i) Distributed homogeneously	Not specified	1	1	0.3	1
Additives	(ii) Distributed heterogeneously	Not specified	50 51 ~ 500 501 ~ 3,200 3,201	2 3 5 8	0.3 0.3 0.3 0.3	1 1 1 1
Agricultural chemicals	(i) Dehydrated vegetables, dried fruits, tea (excluding matcha)	Not specified	5,201 50 51 ~ 150 151 ~ 500 501 ~ 3,200 3,201 ~ 35,000 35,001	3 5 8 13 20 32	0.3 0.3 0.3 0.3 0.3 0.3	1 1 1 1 1 1
	(ii) Cabbage (excluding Brussels sprouts), Chinese cabbage*2	Not specified	Not specified	4	A quarter each is collected from 4 individual cabbage.	1
	(iii) Processed foods (excluding simple processing)	Not specified	150 151 ~ 1,200 1,201	3 5 8	1 1 1	1 1 1
	(iv) Other than (i), (ii) and (iii)	Not specified	50 51 ~ 150 151 ~ 500 501 ~ 3,200 3,201 ~ 35,000 35,001	3 5 8 13 20 32	1 1 1 1 1 1	1 1 1 1 1 1
Residual hazardous substances in livestock and aquatic foods	(i) Paralytic shellfish poison	Not specified	150 151 ~ 1,200 1,201	3 5 8	0.5 0.5 0.5	1 1 1
	(ii) Diarrhetic shellfish poison	Not specified	150 151 ~ 1,200 1,201	3 5 8	0.5 <sup>*3</sup> 0.5 <sup>*3</sup> 0.5 <sup>*3</sup>	1 1 1
	(iii) Pufferfish being mixed	Not specified	150 151 ~ 1,200 1,201	3 5 8	Take two pieces from each carton and one piece shall be regarded as one specimen.	6 10 16
	(iv) Dried seaweeds	Not specified	150 151 ~ 1,200 1,201	3 5 8	0.3 0.3 0.3	1 1 1
	(v) Other than (i), (ii), (iii) and (iv)	Not specified	150 151 ~ 1,200 1,201	3 5 8	0.5 0.5 0.5	1 1 1
Patulin* <sup>4</sup> and DON	(i) Products in bags with its net weight about 20 kg or more	In bags	280 281 ~ 500 501 ~ 1,200 1,201 ~ 3,200 3,201	32 50 80 130 (65×2) 210 (70×3)	1 1 1 2 (1×2) 3 (1×3)	1 1 1 2 3
	(ii) Products in cans or cartons with its net weight 4.5 kg or more	In cans or cartons	50 51 ~ 500 501	2 4 (2×2) 6 (2×3)	0.5 1 (0.25×2)×2 1.5 (0.25×2)×3	1 2 3
	(iii) Other than (i) and (ii)	Packaged in small containers	50 51 ~ 500	2 (2×1) 3 (3×1)	The minimum weight of one sample is 150 g. If the weight of the contents of one sample	1 1
			501 ~ 3,200 3,201	6 (3×2) 9 (3×3)	is less than 150 g, the contents of other containers are added to make one sample of 150 g.	2

<sup>\*1:</sup> Seafood (squilla) shall be regarded as 1.

\*2: Excluding those finely chopped, such as julienned or shredded.

\*3: For diarrhetic shellfish poison of shellfish such as freshwater clam, when weight is less than 10 g as shelled, 0.25 is applied.

\*4: For Patulin, use methods (ii) or (iii).

<sup>\*</sup> For collecting specimens of products in bulk cargo such as grains, beans, follow the procedures below:

A. Specimen collection upon loading onto a silo or a barge (hereinafter referred to as silo, etc.)

When loading onto a silo, select a single arbitrary silo, etc. as one lot. Use means such as autosamplers to collect specimens that are representative of the entire lot. Collect a total of 10 kg or more of the specimen in 15 collections over appropriate intervals, and divide them up to obtain 1 specimen (of 1 kg or more).

B. Specimen collection on a barge

Collect a total of 10 kg or more of the specimen from a total of 15 positions in the upper, middle and lower parts of an arbitrary barge.

Then mix all specimens together and divide them up to obtain 1 specimen (1 kg or more).

C. Specimen collection from a container

Collect a total of 10 kg or more of the specimen from a total of 15 positions in the upper, middle and lower parts of an arbitrary container.

Then mix all specimens together and divide them up to obtain 1 specimen (1 kg or more).