

【資料 6】

参考文献

1 海外文献

No.	著者	タイトル	掲載箇所・年	備考
1	World Health Organization Inter National Agency For Research On Cancer	IARC MONOGRAPHS ON THE EVALUATION OF CARCINOGENIC RISKS TO HUMANS Volume100F(Chemical Agent and Related Occupation)	2012	IARCのオルトートルイジンのモノグラフ (※翻訳中)
2	Kaoru Sekihashi, Ayumu Yamamoto, Yukie Matsumura, Shunji Ueno, Mie Watanabe-Akanuma, Fekadu Kassied, Siegfried Knasmüller, Shuji Tsuda, Yu F. Sasaki	Comparative investigation of multiple organs of mice and rats in the comet assay	Mutation Research 2002 vol517	複数の化学物質について、マウス及びラットの複数の臓器を使用したアッセイでの比較実験
3	Natalie Danford	The genetic toxicology of ortho-toluidine	Mutation Research 1991 vol258	オルトートルイジンに係る代謝物、遺伝毒性について
4	Terry V. Zensera, Vijaya M. Lakshmi, Pong Fu Hsu, Bernard B. Davis	Metabolism of N-acetylbenzidine and initiation of bladder cancer	Mutation Research 2002 vol506-507	N-アセチルベンジンの代謝経路と膀胱がんのイニシエーション
5	Tom Sorahan, Linda Hamilton, John R Jackson	A further cohort study of workers employed at a factory manufacturing chemicals for the rubber industry ,with special reference to the chemicals 2- mercaptobenzothiazole(MBT),aniline,phenyl-B-naphthylamine and o-toluidine	Occupational and Environmental Medicine 2000 vol57	ゴム工業に用いる化学物質を製造する工場作業従事者のコホート研究
6	Elizabeth Ward, Arvind Carpenter, Steven Markowitz, Dennis Roberts, William Halperin	Excess Number of Bladder Cancers in Workers Exposed to Ortho-Toluidine and Aniline	Journal of the National Cancer Institute 1991 vol83	オルト-トルイジン及びアニリンにばく露した労働者の膀胱がん発症例
7	Giovanni F. Rubino, Giovanni Scansetti, Giorgio Piolatto, and Enrico Pira	The Carcinogenic Effect of Aromatic Amine:An Epidemiological Study on the Role of o-Toluidine and 4,4'-Methylene bis(2-Methylaniline)in Inducing Bladder Cancer in Man	Environmental Research 1982 vol27	男性に膀胱がんを誘発する際のo-トルイジン及び4,4'-メチレンビス(2-メチルアニリン)の役割に関する疫学研究
8	Elizabeth M.Ward,Gabriele Sabbioni D.Gayle DeBord,Alexander W.Teass,Kenneth K Brown,Glenn G.Talaska,D ennis R.Robert,Avima M.Ruder,Robert P.Streicher	Monitoring of Aromatic Amine Exposures in Workers at a Chemical Plant With a Known Bladder Cancer Excess	Journal of the National Cancer Institute 1996 vol88	米国ニューヨーク州のゴム化学工場で働く労働者のオルトートルイジンばく露濃度と膀胱がん発症事例 (※翻訳中)
9	G Korinth,T Weiss,S Penkert,K H Schaller,J Angerer,H Drexler	Percutaneous absorption of aromatic amines in rubber industry workers:impact of impaired skin and skin barrier creams	Occup Environ Med 2007 Vol64	ドイツ自動車工場で働く労働者のオルトートルイジンばく露濃度と膀胱がん発症事例 (※翻訳中)

2 国内文献

No.	著者	タイトル	掲載箇所・年	備考
1	独立行政法人製品評価技術基盤機構（委託先：財団法人化学物質評価研究機構）	有害性評価書 Ver.1.0 No.202 オルトートルイジン	2004	オルトートルイジンの環境中の生物及びヒト健康に対する影響
2	独立行政法人製品評価技術基盤機構、財団法人化学物質評価研究機構（委託元：独立行政法人新エネルギー・産業技術総合開発機構）	化学物質の初期リスク評価書 Ver.1.0 No.202 オルトートルイジン	2008	オルトートルイジンの環境中の生物及びヒト健康に対するリスク評価
3	環境省環境保健部環境リスク評価室	化学物質の環境リスク評価第1巻 [23]オルトートルイジン	2002	オルトートルイジンのばく露評価、健康リスク評価

Pub Med 検索条件

“bladder cancer o-toluidine” OR “urinary cancer o-toluidine”

Pub Med 検索結果

1. Biomonitoring of human exposure to arylamines.
Richter E.
Front Biosci (Elite Ed). 2015 Jan 1;7:193-207. Review.
2. Bladder cancer incidence among workers exposed to o-toluidine, aniline and nitrobenzene at a rubber chemical manufacturing plant.
Carreón T, Hein MJ, Hanley KW, Viet SM, Ruder AM.
Occup Environ Med. 2014 Mar;71(3):175-82. doi: 10.1136/oemed-2013-101873.
Epub 2013 Dec 24.
3. Establishing a total allowable concentration of o-toluidine in drinking water incorporating early lifestage exposure and susceptibility.
English JC, Bhat VS, Ball GL, McLellan CJ.
Regul Toxicol Pharmacol. 2012 Nov;64(2):269-84. doi: 10.1016/j.yrtph.2012.08.011.
Epub 2012 Aug 21.
4. Exposure to o-toluidine, aniline, and nitrobenzene in a rubber chemical manufacturing plant: a retrospective exposure assessment update.
Hanley KW, Viet SM, Hein MJ, Carreón T, Ruder AM.
J Occup Environ Hyg. 2012;9(8):478-90. doi: 10.1080/15459624.2012.693836.
5. DNA adducts of ortho-toluidine in human bladder.
Böhm F, Schmid D, Denzinger S, Wieland WF, Richter E.
Biomarkers. 2011 Mar;16(2):120-8. doi: 10.3109/1354750X.2010.534556.
Epub 2010 Nov 30. Erratum in: Biomarkers. 2011 May;16(3):288.
6. Determination of 2,5-toluylenediamine (2,5-TDA) and aromatic amines in urine after personal application of hair dyes: kinetics and doses.
Schettgen T, Heinrich K, Kraus T, Gube M.
Arch Toxicol. 2011 Feb;85(2):127-33. doi: 10.1007/s00204-010-0563-3.
Epub 2010 Jun 15.
7. Increased bladder cancer risk among workers exposed to o-toluidine and aniline: a reanalysis.
Carreón T, Hein MJ, Viet SM, Hanley KW, Ruder AM, Ward EM.
Occup Environ Med. 2010 May;67(5):348-50. doi: 10.1136/oem.2009.051136.
Epub 2009 Nov 2.

8. Monoarylamines in the general population--a cross-sectional population-based study including 1004 Bavarian subjects.
Kütting B, Göen T, Schwegler U, Fromme H, Uter W, Angerer J, Drexler H.
Int J Hyg Environ Health. 2009 May;212(3):298-309.
doi: 10.1016/j.ijheh.2008.07.004. Epub 2008 Sep 12.
9. Bladder cancer risks in workers manufacturing chemicals for the rubber industry.
Sorahan T.
Occup Med (Lond). 2008 Oct;58(7):496-501. doi: 10.1093/occmed/kqn104.
Epub 2008 Aug 25.
10. Percutaneous absorption of aromatic amines in rubber industry workers: impact of impaired skin and skin barrier creams.
Korinth G, Weiss T, Penkert S, Schaller KH, Angerer J, Drexler H.
Occup Environ Med. 2007 Jun;64(6):366-72. Epub 2006 Dec 20.
11. Hemoglobin adducts of the human bladder carcinogen o-toluidine after treatment with the local anesthetic prilocaine.
Gaber K, Harréus UA, Matthias C, Kleinsasser NH, Richter E.
Toxicology. 2007 Jan 5;229(1-2):157-64. Epub 2006 Oct 29.
12. NMR-based model reveals the structural determinants of mammalian arylamine N-acetyltransferase substrate specificity.
Zhang N, Liu L, Liu F, Wagner CR, Hanna PE, Walters KJ.
J Mol Biol. 2006 Oct 13;363(1):188-200. Epub 2006 Aug 15.
13. Determination of three carcinogenic aromatic amines in urine of smokers and nonsmokers.
Riedel K, Scherer G, Engl J, Hagedorn HW, Tricker AR.
J Anal Toxicol. 2006 Apr;30(3):187-95.
14. Corrections to: Markowitz SB, Levin K. Continued epidemic of bladder cancer in workers exposed to ortho-toluidine in a chemical factory. *J Occup Environ Med.* 2004;46:154-160.
Markowitz SB.
J Occup Environ Med. 2005 Sep;47(9):875-7.
15. Continued epidemic of bladder cancer in workers exposed to ortho-toluidine in a chemical factory.
Markowitz SB, Levin K.
J Occup Environ Med. 2004 Feb;46(2):154-60.

16. Correlation between induction of DNA fragmentation in urinary bladder cells from rats and humans and tissue-specific carcinogenic activity.
Robbiano L, Carrozzino R, Bacigalupo M, Corbu C, Brambilla G.
Toxicology. 2002 Sep 30;179(1-2):115-28.
17. A further cohort study of workers employed at a factory manufacturing chemicals for the rubber industry, with special reference to the chemicals 2-mercaptobenzothiazole (MBT), aniline, phenyl-beta-naphthylamine and o-toluidine.
Sorahan T, Hamilton L, Jackson JR.
Occup Environ Med. 2000 Feb;57(2):106-15.
18. T25: a simplified carcinogenic potency index: description of the system and study of correlations between carcinogenic potency and species/site specificity and mutagenicity.
Dybing E, Sanner T, Roelfzema H, Kroese D, Tennant RW.
Pharmacol Toxicol. 1997 Jun;80(6):272-9.
19. Monitoring of aromatic amine exposures in workers at a chemical plant with a known bladder cancer excess.
Stephens ER.
J Natl Cancer Inst. 1997 May 21;89(10):734-5; author reply 735-6. .
20. Monitoring of aromatic amine exposures in workers at a chemical plant with a known bladder cancer excess.
Ward EM, Sabbioni G, DeBord DG, Teass AW, Brown KK, Talaska GG, Roberts DR, Ruder AM, Streicher RP.
J Natl Cancer Inst. 1996 Aug 7;88(15):1046-52.
21. Workers' response to risk notification.
Boal WL, Friedland J, Schulte PA.
Am J Ind Med. 1995 Apr;27(4):471-83.
22. A reexamination of recent publications suggesting o-toluidine may be a human bladder carcinogen.
Freudenthal RI, Anderson DP.
Regul Toxicol Pharmacol. 1995 Feb;21(1):199-202; discussion 203-5.
23. A re-examination of the cause of excess bladder cancers in chemical plant workers.
Freudenthal RI, Anderson DP.
J Natl Cancer Inst. 1994 Jan 5;86(1):59-62.

24. Biological monitoring for occupational exposures to o-toluidine and aniline.
Teass AW, DeBord DG, Brown KK, Cheever KL, Stettler LE, Savage RE, Weigel WW, Dankovic D, Ward E.
Int Arch Occup Environ Health. 1993;65(1 Suppl):S115-8.
25. Reevaluating the carcinogenicity of ortho-toluidine: a new conclusion and its implications.
Sellers C, Markowitz S.
Regul Toxicol Pharmacol. 1992 Dec;16(3):301-17.
26. Renal cell cancer among paperboard printing workers.
Sinks T, Lushniak B, Haussler BJ, Snieszek J, Deng JF, Roper P, Dill P, Coates R.
Epidemiology. 1992 Nov;3(6):483-9.
27. Incidence of bladder cancer in a cohort of workers exposed to 4-chloro-o-toluidine while synthesising chlordimeform.
Popp W, Schmieding W, Speck M, Vahrenholz C, Norpoth K.
Br J Ind Med. 1992 Aug;49(8):529-31.
28. An alternative hypothesis for bladder cancer among workers exposed to ortho-toluidine and aniline.
Acquavella JF, Wilson JD, Conner P, Bannister R.
J Natl Cancer Inst. 1991 Nov 20;83(22):1686-7.
29. Bladder cancer in workers exposed to aniline.
Tannenbaum SR.
J Natl Cancer Inst. 1991 Oct 16;83(20):1507-8.
30. [The carcinogenic effect of chlordimeform and 4-chloro-o-toluidine].
Popp W, Norpoth K.
Dtsch Med Wochenschr. 1991 Sep 20;116(38):1454. German.
31. [4-Chloro-o-toluidine-induced bladder cancer].
Stasik MJ.
Dtsch Med Wochenschr. 1991 Sep 20;116(38):1444-7. Review. German.
32. NIOSH alert: request for assistance in preventing bladder cancer from exposure to o-toluidine and aniline.
Centers for Disease Control (CDC).
MMWR Morb Mortal Wkly Rep. 1991 May 31;40(21):353-4. No abstract available.

33. Preventing bladder cancer from exposure to o-toluidine and aniline.
[No authors listed]
Am Ind Hyg Assoc J. 1991 May;52(5):A260, A262.

34. Excess number of bladder cancers in workers exposed to ortho-toluidine and aniline.
Ward E, Carpenter A, Markowitz S, Roberts D, Halperin W.
J Natl Cancer Inst. 1991 Apr 3;83(7):501-6.

35. Carcinomas of the urinary bladder in a 4-chloro-o-toluidine cohort.
Stasik MJ.
Int Arch Occup Environ Health. 1988;60(1):21-4.

36. Haemoglobin adducts of aromatic amines in people exposed to cigarette smoke.
Bryant MS, Vineis P, Skipper PL, Tannenbaum SR.
IARC Sci Publ. 1988;(89):133-6.

37. Identification and quantitative determination of aniline and toluidines in urine.
el-Bayoumy K, Donahue JM, Hecht SS, Hoffmann D.
Cancer Res. 1986 Dec;46(12 Pt 1):6064-7.

38. A mortality survey of men engaged in the manufacture of organic dyes.
Ott MG, Langner RR.
J Occup Med. 1983 Oct;25(10):763-8.

39. Bioassay for carcinogenicity of 3,2'-dimethyl-4-nitrosobiphenyl, O-nitrosotoluene, nitrosobenzene and the corresponding amines in Syrian golden hamsters.
Hecht SS, El-Bayoumy K, Rivenson A, Fiala ES.
Cancer Lett. 1983 Oct;20(3):349-54.

40. Comparative carcinogenicity of o-toluidine hydrochloride and o-nitrosotoluene in F-344 rats.
Hecht SS, El-Bayoumy K, Rivenson A, Fiala E.
Cancer Lett. 1982 May-Jun;16(1):103-8.

41. The carcinogenic effect of aromatic amines: an epidemiological study on the role of o-toluidine and 4,4'-methylene bis (2-methylaniline) in inducing bladder cancer in man.
Rubino GF, Scansetti G, Piolatto G, Pira E.
Environ Res. 1982 Apr;27(2):241-54.

42. Metabolism of ortho-, meta-, and para-toluidine in the adult male rat.
Cheever KL, Richards DE, Plotnick HB.
Toxicol Appl Pharmacol. 1980 Dec;56(3):361-9.

43. Bioassay of o-toluidine hydrochloride for possible carcinogenicity.
National Toxicology Program.
Natl Cancer Inst Carcinog Tech Rep Ser. 1979;153:1-147.

44. Bioassay of 5-Chloro-o-toluidine for Possible Carcinogenicity (CAS No. 95-79-4).
National Toxicology Program.
Natl Toxicol Program Tech Rep Ser. 1979;187:1-107.

45. [Industrial hygiene and status of health of workers engaged in the production of O-toluidine].
Khlebnikova MI, Gladkova EV, Kurenko LT, Pshenitsyn AV, Shalin BM.
Gig Tr Prof Zabol. 1970 Aug;14(8):7-10. Russian.

* 黄色でマークした文献については、資料6により配付済みの文献。