



Table 1 - continued

Clinical signs  
Male, Female

Sex	Group and dose	Clinical sign	Days																							
			264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285		
Male	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10		
		No abnormality	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8		
		Dyscoria	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		Corn	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	
		Mass	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	4 mg/kg	Number of examined	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	
		No abnormality	7	7	7	7	7	7	7	7	7	7	6	6	6	6	6	6	6	6	6	6	6	6	6	
		Dead	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		Imminently sacrifice	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		Bradypnea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		Hypoactivity	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		Soiled lower abdomen	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		Shuffling gait	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		Reddish urine	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		Hypothermia	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		Pale skin	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
		Corn	2	2	2	2	2	2	2	2	2	2	3	3	3	3	3	3	3	3	3	3	3	3	3	
		20 mg/kg	Number of examined	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8
			No abnormality	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
			Dead	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Corn	3		3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3		
Swelling of limbs	0		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
100 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10		
	No abnormality	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	8	8	8	8	8	8	8		
	Dead	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	Bradypnea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	Hypoactivity	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	Soiled lower abdomen	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	Soiled perinaris	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	Corn	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	2	2	2	2	2		







Table 1 - continued  
Clinical signs  
Male, Female

Sex	Group and dose	Clinical sign	Days												
			352	353	354	355	356	357	358	359	360	361	362	363	364
Male	Control	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10	10
		No abnormality	6	5	5	5	5	4	4	4	5	5	5	5	5
		Dyscoria	0	0	0	0	0	0	0	0	0	0	0	0	0
		Corn	3	4	4	4	4	5	5	5	4	4	4	4	4
		Mass	1	1	1	1	1	1	1	1	1	1	1	1	1
	4 mg/kg	Number of examined	8	8	8	8	8	8	8	8	8	8	8	8	8
		No abnormality	5	4	4	4	4	4	4	4	4	6	6	6	6
		Dead	0	0	0	0	0	0	0	0	0	0	0	0	0
		Imminently sacrifice	0	0	0	0	0	0	0	0	0	0	0	0	0
		Bradypnea	0	0	0	0	0	0	0	0	0	0	0	0	0
		Hypoactivity	0	0	0	0	0	0	0	0	0	0	0	0	0
		Soiled lower abdomen	0	0	0	0	0	0	0	0	0	0	0	0	0
		Shuffling gait	0	0	0	0	0	0	0	0	0	0	0	0	0
Reddish urine		0	0	0	0	0	0	0	0	0	0	0	0	0	
Hypothermia	0	0	0	0	0	0	0	0	0	0	0	0	0		
	Pale skin	0	0	0	0	0	0	0	0	0	0	0	0		
Corn	3	4	4	4	4	4	4	4	4	2	2	2	2		
20 mg/kg	Number of examined	8	8	8	8	8	8	8	8	8	8	8	8		
	No abnormality	5	5	5	5	5	5	5	5	5	5	5	5		
	Dead	0	0	0	0	0	0	0	0	0	0	0	0		
	Corn	3	3	3	3	3	2	2	2	2	2	2	2		
	Swelling of limbs	1	1	1	1	1	1	1	1	1	1	1	1		
100 mg/kg	Number of examined	10	10	10	10	10	10	10	10	10	10	10	10		
	No abnormality	7	7	7	7	7	5	5	5	6	6	6	6		
	Dead	0	0	0	0	0	0	0	0	0	0	0	0		
	Bradypnea	0	0	0	0	0	0	0	0	0	0	0	0		
	Hypoactivity	0	0	0	0	0	0	0	0	0	0	0	0		
	Soiled lower abdomen	0	0	0	0	0	0	0	0	0	0	0	0		
	Soiled perianis	0	0	0	0	0	0	0	0	0	0	0	0		
	Corn	3	3	3	3	3	5	5	5	4	4	4	4		



































