

The efficient and speedy production of this book was carried out by Amanda May, Suzanne Fuzzey and Diana Beaven of the RCP publications department, and the text was expertly copyedited by Dr Mary Firth. To all of them, our thanks.

Contents

1	Tobacco smoking in Britain: an overview	1
1.1	History of tobacco use in Britain	1
	<i>Earliest records</i>	1
	<i>16th and 17th centuries: the age of the pipe</i>	1
	<i>The 18th century: the age of snuff</i>	2
	<i>The 19th century: the age of the cigar</i>	3
	<i>The 20th century: the age of the cigarette</i>	4
1.2	Trends in smoking prevalence in Britain	4
1.3	Risk factors and determinants of cigarette smoking	8
	<i>Gender</i>	8
	<i>Age</i>	8
	<i>Socio-economic status</i>	9
	<i>Region of residence</i>	10
	<i>Risk factors for smoking in children</i>	11
1.4	Smoking in pregnancy	12
	<i>Trends in smoking in pregnancy</i>	12
	<i>Determinants of smoking in pregnancy</i>	13
	<i>Smoking cessation in pregnancy</i>	14
1.5	Morbidity and mortality caused by smoking	15
	<i>Deaths from smoking</i>	15
	<i>Years of life lost</i>	16
	<i>Deaths from smoking by disease</i>	16
	<i>Mortality and the individual smoker</i>	18
	<i>Admissions to hospital</i>	19
	<i>General practitioner consultations</i>	21
	<i>Conclusions</i>	21
1.6	The costs of smoking in Britain	21
	<i>The health burden of smoking</i>	22
	<i>The costs of smoking to society</i>	22
	References	25
2	Physical and pharmacological effects of nicotine	27
2.1	Nicotine receptors and subtypes	27
	<i>Introductory perspective: general features of nicotinic acetylcholine receptors</i>	27
	<i>Heterogeneity of neuronal nicotinic acetylcholine receptors</i>	28
	<i>Nomenclature of nicotinic acetylcholine receptor subunits</i>	29
	<i>Prevalence and distribution of nicotinic acetylcholine receptor subunits</i>	
	<i>Subunit composition of native nicotinic acetylcholine receptors</i>	30
	<i>Pharmacological distinctions</i>	31
	<i>Conclusions</i>	35
2.2	Pharmacology and pharmacokinetics of nicotine	35
	<i>Chemistry of nicotine in tobacco smoke</i>	35
	<i>Absorption of nicotine from tobacco products</i>	36
	<i>Nicotine metabolism</i>	38
	<i>Cardiovascular, endocrine and metabolic effects of nicotine</i>	39
2.3	Pathophysiological effects and toxicity of nicotine	40
	<i>Acute systemic effects</i>	40
	<i>Local toxic effects</i>	41
	<i>Chronic systemic toxicity</i>	41
2.4	Effects of nicotine on mother and fetus in pregnancy	45

<i>Pregnancy</i>	45
<i>The fetus</i>	46
2.5 Animal self-administration and nicotine addiction	46
<i>Nicotine self-administration in rats</i>	47
<i>Summary</i>	48
2.6 Nicotine neurochemistry: nicotine receptor and brain reward systems	49
<i>The neurobiology underlying the positive reinforcing properties of nicotine</i>	50
<i>Other neural responses to nicotine which may contribute to its positive effects on smoking</i>	52
<i>Evidence for desensitisation of neuronal nicotinic receptors</i>	52
<i>Effects on brain 5-hydroxytryptamine pathways</i>	53
References	55
3 Psychological effects of nicotine and smoking in man	67
3.1 The effects of nicotine and smoking on mood and cognition	67
<i>Smokers' perceptions of the psychological effects of smoking</i>	67
<i>Laboratory studies of the effect of nicotine and smoking on mood</i>	68
<i>Laboratory studies of nicotine and smoking effects on cognitive performance</i>	70
<i>Other evidence on the psychological effects of nicotine and its withdrawal</i>	71
<i>Conclusions</i>	72
3.2 The nicotine withdrawal syndrome	72
3.3 Psychological dependence on nicotine and smoking	74
<i>Consumption of nicotine by smokers</i>	74
<i>Markers and measures of dependence in smokers</i>	75
<i>Conclusions</i>	78
References	78
4 Is nicotine a drug of addiction?	83
4.1 The definition of addiction and dependence	83
4.2 Does nicotine use through smoking meet standard diagnostic criteria for addiction?	84
<i>Discussion</i>	87
4.3 The history of social, cultural and political responses to nicotine addiction in Britain	88
<i>Drugs, alcohol and addiction</i>	88
<i>Tobacco and nicotine</i>	89
4.4 How does nicotine addiction compare with addiction to other drugs?	92
<i>Nicotine dosage delivery forms</i>	93
<i>Nicotine dosage delivery by cigarette</i>	96
<i>Incidence, prevalence and risk of progression</i>	97
<i>Remission and relapse</i>	97
<i>Reports of addictiveness by drug abusers</i>	97
<i>Psychoactivity and euphoria</i>	98
<i>Reinforcing effects</i>	98
<i>Physical dependence</i>	99
<i>Tolerance</i>	99
<i>Conclusions</i>	100
4.5 Relevance to society of recognition of nicotine as an addictive drug	100
References	102
5 The natural history of smoking: the smoker's career	107
5.1 Nicotine intake in novice smokers, and the development of dependence	107
<i>Measuring nicotine dependence in adolescents</i>	107
<i>Inhalation and nicotine intake</i>	108
<i>Quit attempts and withdrawal symptoms</i>	108
<i>Time to the first cigarette of the day</i>	110
<i>Subjective effects</i>	110
<i>Gender differences in dependence</i>	111
<i>Initial smoking experiences</i>	111
<i>Progression from experimental to regular smoking</i>	111
<i>Conclusions</i>	111
5.2 Persistent and compulsive smoking in the face of smoking related ill-health	112
<i>Cardiovascular disease</i>	113
<i>Cancer</i>	114

<i>Respiratory disease</i>	114
<i>Smoking cessation in hospital patients</i>	115
<i>Pregnancy</i>	116
<i>Conclusions</i>	116
5.3 Non-dependent smokers	116
<i>Definition and prevalence of non-dependent smoking</i>	117
<i>Conclusions</i>	118
5.4 Smoking cessation rates in Britain	118
<i>Estimation of annual cessation rates</i>	120
<i>Cessation before the age of 60</i>	121
<i>Conclusions</i>	122
References	122
6 Regulation of nicotine intake for smokers, and implications for health	127
6.1 Titration and compensation	127
<i>Cross-sectional studies of self-selected own brand smoking</i>	129
<i>Forced-switching studies</i>	129
<i>Spontaneous brand switching</i>	130
6.2 Epidemiological evidence on the effects of changes in tar and nicotine yields of cigarettes on disease risk	131
<i>The relation between cigarette yield and lung cancer risk</i>	131
<i>Cigarette yields and other cancers</i>	132
6.3 Does the epidemiological evidence support a direct relation between reduction in cigarette yields and disease risk?	135
References	
7 The management of nicotine addiction	139
7.1 General and non-pharmacological approaches	139
<i>Wide-reach approaches</i>	140
<i>Intensive interventions</i>	142
7.2 Nicotine replacement therapy	143
<i>Mode of action</i>	143
<i>Overall efficacy</i>	143
<i>Dose response</i>	144
<i>Dependence potential</i>	144
<i>Additional support</i>	145
<i>Currently available nicotine replacement therapy products</i>	145
<i>Differences between products in efficacy</i>	146
<i>Nicotine replacement therapy combinations</i>	146
7.3 Non-nicotine medications for treating nicotine addiction	147
<i>Antidepressants</i>	147
<i>Clonidine</i>	148
<i>Mecamylamine</i>	149
<i>Bupirone</i>	149
<i>Lobeline</i>	149
<i>Naltrexone</i>	150
<i>Silver acetate</i>	150
<i>Sensory replacement</i>	150
<i>Summary</i>	150
7.4 Evidence-based treatment of nicotine addiction	151
7.5 Nicotine replacement treatment in pregnancy	153
7.6 Cost-effectiveness of treating nicotine addiction	155
<i>Conclusions</i>	158
References	159
8 Regulatory approaches to tobacco products in Britain	165
8.1 The evolution of British laws and voluntary agreements	
8.2 The regulation of nicotine, tar and additives in cigarettes	166
<i>Filters, low tar cigarettes and the Federal Trade Commission test</i>	166
<i>Tobacco substitutes</i>	166
<i>New voluntary agreements on tar reductions</i>	167

<i>The introduction of additives</i>	168
<i>Health warnings</i>	169
<i>Other nicotine containing products</i>	170
<i>Conclusions</i>	170
8.3 Anomalies in current nicotine regulation	171
8.4 Novel nicotine delivery devices from tobacco companies	171
8.5 Tobacco regulation in the USA	173
<i>Industry manipulation of tobacco</i>	174
<i>The Food and Drug Administration tobacco rule</i>	174
<i>The Federal Trade Commission test</i>	175
8.6 Future nicotine regulation in the UK	176
<i>Regulation at a national level</i>	176
<i>Regulation at a European level</i>	176
<i>The scope of future nicotine regulation</i>	177
<i>Conclusions</i>	178
8.7 Economic effects of nicotine regulation	178
<i>Employment</i>	178
<i>Tax revenue</i>	179
<i>Other effects of smoking reduction</i>	180
<i>Conclusions</i>	180
References	180
9 Summary and recommendations	183
9.1 Tobacco and nicotine addiction	183
9.2 Consequences of nicotine addiction	184
9.3 Treatment of nicotine addiction	186
9.4 Regulation of tobacco products	186
9.5 Recommendations	187
Reference	189

Key points

Tobacco smoking in Britain

- Smoking prevalence in Britain has declined during the past 50 years; this trend now appears to be stabilising
- In 1997 in Britain approximately one in four adults were cigarette smokers
- By age 15, one in four British children are regular smokers
- Smoking causes one in every five deaths in Britain, and the loss of more than 550,000 years of life before age 75
- The greatest impact of smoking on mortality is on deaths from lung cancer, ischaemic heart disease and chronic obstructive airways disease
- Passive smoking damages children before and after birth
- Thirty percent of pregnant women in Britain smoke
- Smoking is strongly related to poverty and deprivation
- Smoking costs the NHS an estimated £1.5 billion per year
- No other single avoidable factor accounts for such a high proportion of deaths, hospital admissions or general practitioner consultations
- Smoking is the single most important public health problem in Britain

Physical and pharmacological effects of nicotine

- Nicotine receptors are present in the brain and many other organs vary markedly in their binding, activation and desensitisation characteristics
- Cigarettes deliver rapid doses of nicotine to receptors in the brain
- Animal studies provide strong and consistent evidence that nicotine is addictive
- The addictive effect of nicotine is mediated at least in part by stimulation of dopamine release in the nucleus accumbens
- Pure nicotine has potential adverse effects on the human body but unlike cigarettes does not appear to cause cancer or significant cardiovascular disease
- Pure nicotine may be harmful to the fetus in pregnancy but is likely to be far less hazardous than the effects of smoking.

Psychological effects of nicotine and smoking

- Smoking is widely believed to have positive effects on mood
- Objective evidence suggests that the only improvements in mood resulting from smoking are those arising from the relief of withdrawal symptoms
- Smoking withdrawal symptoms are relieved by nicotine
- Nicotine intake in smokers is stable and consistent over time
- There is strong evidence of psychological dependence on cigarettes
- The major psychological motivation to smoke is the avoidance of negative mood states caused by withdrawal of nicotine

Is nicotine a drug of addiction?

- Nicotine obtained from cigarettes meets all the standard criteria used to define a drug of dependence or addiction
- Historically, and in contrast to addiction to opiates or alcohol, addiction to nicotine has not been recognised as a medical or social problem in Britain
- Nicotine is highly addictive, to a degree similar or in some respects exceeding addiction to 'hard' drugs such as heroin or cocaine
- Most smokers do not smoke out of choice, but because they are addicted to nicotine

The natural history of smoking: the smoker's career

- Addiction to nicotine is established in most smokers during teenage years, in many cases before reaching the age at which it is legal to buy cigarettes
- Teenagers who smoke one or more cigarettes per day demonstrate evidence of addiction similar to that seen in addicted adults, but addiction can be evident at lower levels of smoking
- Addiction to nicotine is usually established in young smokers within about a year of first experimenting with cigarettes
- A small proportion of smokers, approximately 5%, do not appear to be addicted to nicotine
- Once addicted, most smokers are unable to give up smoking even when they develop disease caused by smoking and made worse by continued smoking
- Only about 2% of smokers succeed in giving up in any year
- About 50% of young adult smokers will still be smoking when they are 60

Regulation of nicotine intake for smokers, and implications for health

- Smokers tend to regulate or titrate their nicotine intake to maintain body levels within a preferred range
- Smokers who switch to cigarettes which on machine smoking deliver less nicotine and tar tend to compensate for this by smoking the cigarette more deeply or more intensively
- Smokers of low yield cigarettes actually achieve little, if any, reduction in intake of nicotine and tar, and the health benefit accrued from switching to such cigarettes is, if anything, small
- The availability of low yield cigarettes may actually be counter productive in public health terms if they encourage health conscious smokers to switch to low yield brands instead of giving up completely

Management of nicotine addiction

- Effective interventions to reduce nicotine addiction are available at both population and individual levels
- *The fact that smoking is so common in Britain means that even interventions that have small effects on smoking prevalence can, if widely applied, yield substantial returns in terms of the numbers of people who give up smoking*
- Nicotine replacement therapy approximately doubles the effectiveness of most other currently available smoking cessation interventions
- Smoking cessation interventions, including nicotine replacement therapy, are extremely cost effective, costing society between £212 and £873 per year of life saved in 1996 prices
- The cost-effectiveness of smoking cessation interventions using nicotine replacement therapy compares very favourably with most other medical intervention.
- Effective smoking cessation services should therefore be universally available to smokers through the NHS
- Smoking cessation services must be able to adapt to accommodate new effective therapies and interventions in the future
- Further research into the use and safety of nicotine replacement therapy relative to continued smoking during pregnancy is needed
- Regulatory approaches to tobacco products in Britain
- Cigarettes are extremely damaging to consumers and yet have enjoyed unparalleled freedom from consumer protection regulation
- Much of the regulation applying to tobacco in Britain has been in the form of 'voluntary agreements' with the tobacco industry
- The use of additives in cigarettes has not been subject to appropriate assessments of public health impact
- The policy of progressively reducing tar yields from cigarettes, and of printing tar yields on cigarette packs, is based on flawed measurement methodology and may be ineffective in terms of achieving public health benefits
- Pharmaceutical nicotine delivery products (eg nicotine replacement therapy) are subject to regulation by the Medicines Control Agency and are required to meet the same safety standards as any other drug; however, cigarettes are exempt from these controls
- Cigarettes are tobacco-based nicotine delivery products and should be subject to the same safety standards as any other drug.
- A co-ordinated nicotine regulation framework needs to be established in Britain to resolve anomalies in the sale and promotion of nicotine delivery products, to maximise current and future public health

Main conclusions

- Most smokers do not continue to smoke cigarettes out of choice, but because they are addicted to nicotine
- Nicotine addiction is the underlying cause of the massive burden of premature death and disability caused by smoking in Britain
- Doctors, other health professionals and indeed society as a whole, need to acknowledge nicotine addiction as a major medical and social problem
- Treatment for nicotine addiction should be universally available for all smokers as a routine facility of the National Health Service
- Tobacco products must be made subject to safety regulations that are consistent with the controls that apply to all other drugs available in Britain, and so that they are commensurate with the extent of the damage to individuals and society that smoking causes

This page last updated on October 5, 2000

[Links](#) | [Site map](#) | [Search](#) | [Contact us](#) | [Find us](#)

All material copyright Royal College of Physicians of London