

	大	接触	有効な治療開始 後24時間まで	
	小	標準		
	子宮内膜炎(産褥敗血症)	標準		
	乳幼児の咽頭炎	飛沫	有効な治療開始 後24時間まで	
	乳幼児の肺炎	飛沫	有効な治療開始 後24時間まで	
	乳幼児の猩紅熱	飛沫	有効な治療開始 後24時間まで	
	新生児の溶連菌症(B群溶連菌)	標準		
	溶連菌症(非A非B群)、他の箇所でカバーされ ない	標準		
	多剤耐性(⇒多剤耐性菌)			
わ	Vincent's angina	標準		

標準:標準的な感染予防策、空気:空気感染隔離予防策、飛沫:飛沫感染予防策、接触:接触感染予防策

¹³⁵ Kenyon A, Ridzon R, Luskin-HawkR, et al. A Nosocomial outbreak of Multidrug-Resistant Tuberculosis. Ann Intern Med. 1997;127(1):32-36.

¹³⁶ Pavelchak N, DePersis RP, London M, et al: Identification of factors that disrupt negative air pressurization of respiratory isolation rooms. Infect Control Hosp Epidemiol. 2000;21:191-5.

¹³⁷ 大久保憲, 簣淳夫他. 病院空調設備の設計・管理指針 HEAS-02-2004. 日本医療福祉設備協会.

¹³⁸ Haley CE, McDonald RC, Rossi L, Jones WD, Jr., Haley RW, Luby JP: Tuberculosis epidemic among hospital personnel. Infect Control Hosp Epidemiol. 1989;10:204-10.

¹³⁹ Ryan MG:Developing a respiratory protection program, Understanding the written elements. AAOHN J. 2001;49:293-307.

¹⁴⁰ Drinka PJ, Krause P, Nest L, Goodman BM, Gravenstein S: Risk of acquiring influenza A in a nursing home from a culture-positive roommate. Infect Control Hosp Epidemiol. 2003;24:872-4.

¹⁴¹ Seto WH, Tsang D, Yung RWH, et al: Effectiveness of precautions against droplets and contact in prevention of nosocomial transmission of severe acute respiratory syndrome (SARS). Lancet. 2003;361:1591-1620.

¹⁴² Chang VT, Nelson K. The role of physical proximity in nosocomial diarrhea. Clin Infect Dis. 2000;31:717-22.

¹⁴³ Trick WA, Weinstein RA, et al: Comparison of routine glove use and contact-isolation precautions to prevent transmission of multidrug-resistant bacteria in a long-term care facility. J Am Geriatr Soc. 2004 Dec;52(12):2003-9.

¹⁴⁴ Sally W, et al: Effectiveness of barrier precautions and surveillance cultures to control transmission of multidrug-resistant organisms. A systematic review of the literature. AJIC. 2006;34:484-494.