

-
- alcohol, and chlorhexidine for prevention of infection associated with central venous and arterial catheters. *Lancet* 1991;338:339-343.
- ²⁸⁴ Mimosz O, Pieroni L, Lawrence C et al: Prospective, randomized trial of two antiseptic solutions for prevention of central venous or arterial catheter colonization and infection in intensive care unit patients. *Crit Care Med* 1996;24:1818-1823.
- ²⁸⁵ Humar A, Ostromecki A, Direnfeld J et al: Prospective randomized trial of 10% povidone-iodine versus 0.5% tincture of chlorhexidine as cutaneous antiseptics for prevention of central venous catheter infection. *Clin Infect Dis* 2000; 31: 1001-1007.
- ²⁸⁶ Strand CL, Wajsbort RR, Sturmman K: Effect of iodophor vs iodine tincture skin preparation on blood culture contamination rate. *JAMA* 1993;269:1004-1006.
- ²⁸⁷ Little JR, Murray PR, Traynor PS et al: A randomized trial of povidone-iodine compared with iodine tincture for venipuncture site disinfection: effect on rates of blood culture contamination. *Am J Med* 1999;107:119-125.
- ²⁸⁸ 井上善文, 阪尾 淳, 廣田昌紀, ほか: 中心静脈カテーテル挿入部管理における希ヨードチンキの意義. *外科治療* 2006;94:857-862.
- ²⁸⁹ Tanner J, Woodings D, Moncaster K: Preoperative hair removal to reduce surgical site infection. *Cochrane Database Syst Rev* 2006;2:CD004122.
- ²⁹⁰ Kjonniksen I, Andersen BM, Sondenaa VG, et al: Preoperative hair removal—a systematic literature review. *AORN J* 2002;75:928-938.
- ²⁹¹ Alexander JW, Fischer JE, Boyajian M, et al: The influence of hair-removal methods on wound infections. *Arch Surg* 1983;118:347-352.
- ²⁹² Balthazar ER, Colt JD, Nicholas RL: Preoperative hair removal: a random prospective study of shaving versus clipping. *South Med J* 1982;75:799-801.
- ²⁹³ Maki DG, Band JD: A comparative study of polyantibiotic and iodophor ointments in prevention of vascular catheter-related infection. *Am J Med* 1981;70:739-744.
- ²⁹⁴ Norden CW: Application of antibiotic ointment to the site of venous catheterization: A controlled trial. *J Infect Dis* 1969;120:611-615.
- ²⁹⁵ Thodis E, Bhaskaran S, Pasadakis P, et al: Decrease in *Staphylococcus aureus* exit-site infections and peritonitis in CAPD patients by local application of mupirocin ointment at the catheter exit site. *Perit Dial Int* 1998;18:261-270.
- ²⁹⁶ Sesso R, Barbosa D, Leme IL, et al: *Staphylococcus aureus* prophylaxis in hemodialysis patients using central venous catheter: effect of mupirocin ointment. *Am Soc Nephrol* 1998;9:1085-1092.
- ²⁹⁷ Zakrzewska-Bode A, Muyltjens HL, Liem KD, et al: Mupirocin resistance in coagulase-negative staphylococci, after topical prophylaxis for the reduction of colonization of central venous catheters. *J Hosp Infect* 1995;31:189-193.
- ²⁹⁸ Levin A, Mason AJ, Jindal KK et al: Prevention of hemodialysis subclavian vein catheter infections by topical povidone-iodine. *Kidney Int* 1991;40:934-938.
- ²⁹⁹ 井上善文, 吉川正人, 阪尾淳, ほか: 中心静脈カテーテル挿入部管理におけるポビドンヨードおよびポビドンヨードゲル使用の意義に関する検討—クロルヘキシジンアルコールとの比較. *外科と代謝・栄養* 2004;38:21-27.
- ³⁰⁰ Conly JM, Grieves K, Peters B: A prospective, randomized study comparing transparent and dry gauze dressings for central venous catheters. *J Infect Dis* 1989;159:310-319.
- ³⁰¹ Meylan PR: Increased risk of bacterial colonization of intravenous catheters covered with transparent adhesive polyurethane bandages, compared to classical gauze bandages. *Schweiz Med Wochenschr* 1987;117:2013-2016.
- ³⁰² Vazquez RM, Jarrard MM: Care of the central venous catheterization site: the use of a transparent polyurethane film. *JPEN* 1984;8:181-186.
- ³⁰³ Hoffmann KK, Western SA, Kaiser DL et al: Bacterial colonization and phlebitis-associated risk with transparent polyurethane film for peripheral intravenous site dressings. *Am J Infect Control* 1988;16:101-106.
- ³⁰⁴ Ricard P, Martin R, Marcoux JA: Protection of indwelling vascular catheters: incidence of bacterial contamination and catheter-related sepsis. *Crit Care Med* 1985;13:541-543.