

-
- controlled trial. *Am J Infect Control* 2004;32:291-295.
- ³²⁶ Yebenes JC, Martnez R, Serra-Prat M, et al: Resistance to the migration of microorganisms of a needle-free disinfectable connector. *Am J Infect Control* 2003;31:462-464.
- ³²⁷ Brown JD, Moss HA, Elliott TS: The potential for catheter microbial contamination from a needleless connector. *J Hosp Infect* 1997;36:181-189.
- ³²⁸ Brouza E, Munoz P, Lopez-Rodrigues J, et al: A needleless closed system device (CLAVE) protects from intravascular catheter tip and hub colonization: a prospective randomized study. *J Hosp Infect* 2003;54:279-287.
- ³²⁹ Seymour VM, Dhallu TS, Moss HA, et al: A prospective clinical study to investigate the microbial contamination of a needleless connector. *J Hosp Infect* 2000;45:165-168.
- ³³⁰ Mendelson MH, Short LJ, Schechter CB, et al: Study of a needleless intermittent intravenous-access system for peripheral infusions: analysis of staff, patient, and institutional outcomes. *Infect Control Hosp Epidemiol* 1998;19:401-406.
- ³³¹ Danzig LE, Short LJ, Collins K, et al: Bloodstream infections associated with a needleless intravenous infusion system in patients receiving home infusion therapy. *JAMA* 1995;273:1862-1864.
- ³³² McDonald LC, Banerjee SN, Jarvis WR: Line-associated bloodstream infections in pediatric intensive-care-unit patients associated with a needleless device and intermittent intravenous therapy. *Infect Control Hosp Epidemiol* 1998;19:772-777.
- ³³³ Kellerman S, Shay DK, Howard J, et al: Bloodstream infections in home infusion patients: the influence of race and needleless intravascular access devices. *J Pediatr* 1996;129:711-717.
- ³³⁴ Maragakis LL, Bradley KL, Song X, et al: Increased catheter-related bloodstream infection rates after the introduction of a new mechanical valve intravenous access port. *Infect Control Hosp Epidemiol* 2006;27:67-70.
- ³³⁵ Cookson ST, Ihrig M, O'Mara EM, et al: Increased bloodstream infection rates in surgical patients associated with variation from recommended use and care following implementation of a needleless device. *Infect Control Hosp Epidemiol* 1998;19:23-27.
- ³³⁶ Do AN, Ray BJ, Banerjee SN et al: Bloodstream infection associated with needleless device use and the importance of infection-control practices in the home health care setting. *J Infect Dis* 1999;179:442-448.
- ³³⁷ McDonald LC, Banerjee SN, Jarvis WR: Line-associated bloodstream infections in pediatric intensive-care-unit patients associated with a needleless device and intermittent intravenous therapy. *Infect Control Hosp Epidemiol* 1998;19:772-777.
- ³³⁸ Arduino MJ, Bland LA, Danzig LE et al: Microbiologic evaluation of needleless and needle-access devices. *Am J Infect Control* 1997;26:377-380.
- ³³⁹ Brown JD, Moss HA, Elliott TS: The potential for catheter microbial contamination from a needleless connector. *J Hosp Infect* 1997;36:181-189.
- ³⁴⁰ Mendelson MH, Short LJ, Schechter CB et al: Study of a needleless intermittent intravenous-access system for peripheral infusions: analysis of staff, patient, and institutional outcomes. *Infect Control Hosp Epidemiol* 1998;19:401-406.
- ³⁴¹ Menyhay SZ, Maki DG: Disinfection of needleless catheter connectors and access ports with alcohol may not prevent microbial entry: the promise of a novel antiseptic-barrier cap. *Infect Control Hosp Epidemiol* 2006;27:23-27.
- ³⁴² Luebke MA, Arduino MJ, Duda DL, et al: Comparison of the microbial barrier properties of a needleless and a conventional needle-based intravenous access system. *Am J Infect Control* 1998;26:437-441.
- ³⁴³ Inoue Y, Nezu R, Matsuda H, et al: Experimental study of hub contamination: effect of a new connection device: the I-system. *JPEN* 1992;16:178-180.
- ³⁴⁴ Inoue Y, Nezu R, Matsuda H, et al: Prevention of catheter-related sepsis during parenteral nutrition: effect of a new connection device. *JPEN* 1992;16:581-585.
- ³⁴⁵ Sitges-Serra A: Strategies for prevention of catheter-related bloodstream infections.