

-
- Support Care Cancer 1999;7:391-395.
- ³⁴⁶ Maki DG, Goldman DA, Rhame FS: Infection control in intravenous therapy. Ann Intern Med 1973;79:867-887.
- ³⁴⁷ Ryan JA, Abel RM, Abbott WM et al: Catheter complications in total parenteral nutrition: A prospective study of 200 consecutive patients. 1974;290:757-761.
- ³⁴⁸ Brosnan KM, Parham AM, Rutledge B, et al: Stopcock contamination. Am J Nurs 1988;88:320-324.
- ³⁴⁹ McArthur BJ, Hargiss C, Schoenknecht FD: Stopcock contamination in an ICU. Am J Nurs 1975;75:96-97.
- ³⁵⁰ Walrath JM, Abbott NK, Caplan E et al: Stopcock bacterial contamination in invasive monitoring systems. Heart Lung 1979;8:100-104.
- ³⁵¹ Mueller-Premru M, Gubina M, Kaufmann ME et al: Use of semi-quantitative and quantitative culture methods and typing for studying the epidemiology of central venous catheter-related infections in neonates on parenteral nutrition. J Neo Microbiol 1999;48:451-460.
- ³⁵² Gibilisco PA, Lopez GA, Appleman MD et al: In vitro contamination of "Piggyback/Heparin lock" assemblies: Prevention of contamination with a closed, positive locking device (Click-Lock). JPEN 1986;10:431-434.
- ³⁵³ Ruschman KL, Fulton JS: Effectiveness of disinfectant techniques on intravenous tubing latex injection ports. Intraven Nurs 1993;16:304-308.
- ³⁵⁴ 井上善文, 根津理一郎, 李鐘甲、ほか: カテーテル敗血症予防のための新しい輸液ライン接続システムの開発-実験的・臨床的検討-. 外科と代謝・栄養 1989;23:292-302.
- ³⁵⁵ 内山昭則, 妙中信之, 井上善文, ほか: 薬剤静注時の細菌侵入防止対策; インジェクションプラグの応用. ICU と CCU 1992;16:157-160.
- ³⁵⁶ Salzman MB, Isenberg HD, Rubin LG: Use of disinfectants to reduce microbial contamination of hubs of vascular catheters. J Clin Microbiol 1993;31:475-479.
- ³⁵⁷ Ruschman KL, Fulton JS: Effectiveness of disinfectant techniques on intravenous tubing latex injection ports. J Intraven Nurs 1993;16:304-308.
- ³⁵⁸ Snydman DR, Donnelly-Reidy M, Perry LK et al: Intravenous tubing containing burettes can be safely changed at 72-hour intervals. Infect Control 1987;8:113-116.
- ³⁵⁹ Josephson A, Gombert ME, Sierra MF et al: The relationship between intravenous fluid contamination and the frequency of tubing replacement. Infect Control 1985;6:367-370.
- ³⁶⁰ Maki DG, Botticelli JT, LeRoy ML et al: Prospective study of replacing administration sets for intravenous therapy at 48- vs 72-hour intervals. 72 hours is safe and cost-effective. JAMA 1987;258:1777-1781.
- ³⁶¹ Sitges-Serra A, Linares J, Perez JL et al: A randomized trial on the effect of tubing changes on hub contamination and catheter sepsis during parenteral nutrition. JPEN 1985;9:322-325.
- ³⁶² Raad I, Hanna HA, Awad A, et al: Optimal frequency of changing intravenous administration sets: is it safe to prolong use beyond 72 hours? Infect Control Hosp Epidemiol 2001;22:136-139.
- ³⁶³ Gillies D, O'Riordan L, Wallen M, et al: Timing of intravenous administration set changes: a systematic review. Infect Control Hosp Epidemiol 2004;25:240-250.
- ³⁶⁴ Simon A, Fleischhack G, Wiszniewsky G, et al: Influence of prolonged use of intravenous administration sets in paediatric cancer patients on CVAD-related bloodstream infection rates and hospital resources. Infection 2006;34:258-263.
- ³⁶⁵ Inoue Y, Nezu R, Matsuda H, et al: Prevention of catheter-related sepsis during parenteral nutrition: effect of a new connection device. JPEN 1992;16:581-585.
- ³⁶⁶ Pearson ML, the Hospital Infection Control Practices Advisory Committee. Guidelines for prevention of intravascular device-related infections. Infect Control Hosp Epidemiol 1996;17:435-473.
- ³⁶⁷ Matlow AG, Kitai I, Kirpalani H, et al: A randomized trial of 72- versus 24-hour intravenous tubing set changes in newborns receiving lipid therapy. Infect Control Hosp