

表5-2-6 7セトミ/フェンとクループの併発

		発現数 (%)							①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)	
		薬剤 未使用 ^①	薬剤使用 計		(内訳)		総計	未使用			使用開始	既使用	総計	薬剤未使用	薬剤使用	
			使用開始 ^②	既使用 ^③												
第1病日	朝	10 0.4%	2 2.0%	2 2.0%	—	12 0.4%	0.067	5.64	2,747	99	2,846	0.4%	2.0%			
	昼	4 0.2%	2 0.7%	2 1.1%	0 0.0%	6 0.2%	0.124	4.59	2,554	183	2,834	0.5%	2.7%			
	夜	11 0.5%	3 0.4%	3 0.7%	0 0.0%	14 0.5%	0.218	0.79	2,103	447	2,828	1.0%	3.1%			
第2病日	朝	4 0.2%	2 0.2%	1 1.1%	1 0.1%	6 0.2%	0.385	1.23	2,001	91	2,814	1.2%	3.4%			
	昼	3 0.2%	1 0.1%	0 0.0%	1 0.1%	4 0.1%	0.802	0.66	1,865	132	2,808	1.4%	3.5%			
	夜	1 0.1%	0 0.0%	0 0.0%	0 0.0%	1 0.0%	0.623	0.00	1,748	114	2,804	1.5%	3.5%			
第3病日	朝	0 0.0%	1 0.1%	0 0.0%	1 0.1%	1 0.0%	0.377		1,739	8	2,803	1.5%	3.6%			
	昼	0 0.0%	1 0.1%	0 0.0%	1 0.1%	1 0.0%	0.379		1,729	10	2,802	1.5%	3.6%			
	夜	0 0.0%	0 0.0%	0 0.0%	0 0.0%	0 0.0%			1,716	13	2,801	1.5%	3.6%			
第4病日	朝	0 0.0%	0 0.0%	0 0.0%	0 0.0%	0 0.0%			1,714	2	2,801	1.5%	3.6%			
	昼	0 0.0%	1 0.1%	0 0.0%	1 0.1%	1 0.0%	0.388		1,711	3	2,801	1.5%	3.7%			
	夜	1 0.1%	0 0.0%	0 0.0%	0 0.0%	1 0.0%	0.610	0.00	1,707	4	2,800	1.5%	3.7%			
第5病日	朝	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%			1,706		2,799	1.5%	3.7%			
	昼	0 0.0%	0 0.0%	0 0.0%	0 0.0%	0 0.0%			1,703	3	2,799	1.5%	3.7%			
	夜	0 0.0%	0 0.0%	0 0.0%	0 0.0%	0 0.0%			1,702	1	2,799	1.5%	3.7%			
第6病日	朝	1 0.1%	0 0.0%	—	0 0.0%	1 0.0%	0.319	0.00	1,702		2,799	1.6%	3.7%			
	昼	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%			1,701		2,798	1.6%	3.7%			
	夜	1 0.1%	0 0.0%	—	0 0.0%	1 0.0%	0.318	0.00	1,701		2,798	1.6%	3.7%			
第7病日	朝	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%			1,700		2,797	1.6%	3.7%			
	昼	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%			1,700		2,797	1.6%	3.7%			
	夜	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%			1,700		2,797	1.6%	3.7%			
累積発現率 (参考)		36	13	8	5	49 1.7%						1.6%	3.7%			

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.39 (0.71 , 2.74)	0.336
第1～第2病日	1.36 (0.64 , 2.87)	0.426
第1～第4病日	1.57 (0.79 , 3.11)	0.198

表5-1-7 タミフルと中耳炎の併発

		発現数 (%)						①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)	
		薬剤 未使用 ^①	薬剤使用 (内訳)		既使用 ^③	総計	未使用			使用開始	既使用	総計	薬剤未使用	薬剤使用	
			計	使用開始 ^②											
第1病日	朝	6 0.2%	1 0.3%	1 0.3%	—	7 0.2%	0.829	1.27	2,516	330	—	2,846	0.2%	0.3%	
	昼	3 0.1%	1 0.2%	1 0.3%	0 0.0%	4 0.1%	0.478	1.16	2,204	306	329	2,839	0.4%	0.5%	
	夜	3 0.2%	1 0.1%	1 0.2%	0 0.0%	4 0.1%	0.352	0.55	1,764	437	634	2,835	0.5%	0.6%	
第2病日	朝	6 0.5%	3 0.2%	0 0.0%	3 0.3%	9 0.3%	0.081	0.35	1,166	595	1,070	2,831	1.1%	0.7%	
	昼	1 0.1%	1 0.0%	1 0.2%	0 0.0%	2 0.1%	0.161	0.33	701	459	1,662	2,822	1.2%	0.8%	
	夜	0 0.0%	3 0.1%	1 0.5%	2 0.1%	3 0.1%	0.284	—	490	210	2,120	2,820	1.2%	0.9%	
第3病日	朝	2 0.5%	3 0.1%	0 0.0%	3 0.1%	5 0.2%	0.318	0.24	394	96	2,327	2,817	1.7%	1.0%	
	昼	0 0.0%	1 0.0%	0 0.0%	1 0.0%	1 0.0%	0.861	—	332	60	2,420	2,812	1.7%	1.1%	
	夜	0 0.0%	1 0.0%	0 0.0%	1 0.0%	1 0.0%	0.882	—	303	29	2,479	2,811	1.7%	1.1%	
第4病日	朝	0 0.0%	3 0.1%	0 0.0%	3 0.1%	3 0.1%	0.710	—	295	8	2,507	2,810	1.7%	1.2%	
	昼	1 0.3%	1 0.0%	0 0.0%	1 0.0%	2 0.1%	0.368	0.11	289	6	2,512	2,807	2.0%	1.3%	
	夜	0 0.0%	3 0.1%	0 0.0%	3 0.1%	3 0.1%	0.722	—	286	2	2,517	2,805	2.0%	1.4%	
第5病日	朝	0 0.0%	3 0.1%	0 0.0%	3 0.1%	3 0.1%	0.724	—	284	2	2,516	2,802	2.0%	1.5%	
	昼	0 0.0%	2 0.1%	0 0.0%	2 0.1%	2 0.1%	0.807	—	282	2	2,515	2,799	2.0%	1.6%	
	夜	0 0.0%	0 0.0%	0 0.0%	0 0.0%	0 0.0%	—	—	281	1	2,515	2,797	2.0%	1.6%	
第6病日	朝	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%	—	—	281	—	2,516	2,797	2.0%	1.6%	
	昼	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%	—	—	281	—	2,516	2,797	2.0%	1.6%	
	夜	0 0.0%	2 0.1%	0 0.0%	2 0.1%	2 0.1%	0.809	—	280	1	2,516	2,797	2.0%	1.7%	
第7病日	朝	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%	—	—	280	—	2,515	2,795	2.0%	1.7%	
	昼	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%	—	—	280	—	2,515	2,795	2.0%	1.7%	
	夜	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%	—	—	280	—	2,515	2,795	2.0%	1.7%	
累積発現率 (参考)	22	29	5	24	51	1.8%	—	—	—	—	—	—	2.0%	1.7%	

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	0.68 (0.34 , 1.35)	0.268
第1～第2病日	0.62 (0.26 , 1.48)	0.282
第1～第4病日	0.59 (0.28 , 1.22)	0.154

表5-2-7 アセトアミノフェンと中耳炎

		発現数 (%)						総計	①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)	
		薬剤 未使用 ^①	薬剤使用		(内訳)		既使用 ^③				未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用
			計	使用開始 ^②	既使用 ^③											
第1病日	朝	6 0.2%	1 1.0%	1 1.0%	—	—	7 0.2%	0.236	4.66	2,747	99	2,846	0.2%	1.0%		
	昼	4 0.2%	0 0.0%	0 0.0%	0 0.0%	0 0.0%	4 0.1%	0.659	0.00	2,558	183	2,839	0.4%	1.0%		
	夜	3 0.1%	1 0.1%	1 0.2%	0 0.0%	0 0.0%	4 0.1%	0.615	0.96	2,104	450	2,835	0.5%	1.1%		
第2病日	朝	7 0.3%	2 0.2%	0 0.0%	2 0.3%	9 0.3%	0.708	0.70	2,009	92	730	2,831	0.9%	1.4%		
	昼	2 0.1%	0 0.0%	0 0.0%	0 0.0%	2 0.1%	0.438	0.00	1,868	134	820	2,822	1.0%	1.4%		
	夜	3 0.2%	0 0.0%	0 0.0%	0 0.0%	3 0.1%	0.240	0.00	1,752	114	954	2,820	1.1%	1.4%		
第3病日	朝	1 0.1%	4 0.4%	0 0.0%	4 0.4%	5 0.2%	0.155	6.49	1,741	8	1,068	2,817	1.2%	1.8%		
	昼	1 0.1%	0 0.0%	0 0.0%	0 0.0%	1 0.0%	0.615	0.00	1,730	10	1,072	2,812	1.3%	1.8%		
	夜	0 0.0%	1 0.1%	1 8.3%	0 0.0%	1 0.0%	0.004	0.00	1,717	12	1,082	2,811	1.3%	1.8%		
第4病日	朝	2 0.1%	1 0.1%	0 0.0%	1 0.1%	3 0.1%	0.978	0.78	1,715	2	1,093	2,810	1.4%	1.9%		
	昼	2 0.1%	0 0.0%	0 0.0%	0 0.0%	2 0.1%	0.371	0.00	1,711	2	1,094	2,807	1.5%	1.9%		
	夜	0 0.0%	3 0.3%	1 25.0%	2 0.2%	3 0.1%	0.001	0.00	1,705	4	1,096	2,805	1.5%	2.2%		
第5病日	朝	3 0.2%	0 0.0%	—	0 0.0%	3 0.1%	0.084	0.00	1,705	—	1,097	2,802	1.7%	2.2%		
	昼	1 0.1%	1 0.1%	0 0.0%	1 0.1%	2 0.1%	0.952	1.55	1,699	3	1,097	2,799	1.7%	2.3%		
	夜	0 0.0%	0 0.0%	0 0.0%	0 0.0%	0 0.0%	—	—	1,697	1	1,099	2,797	1.7%	2.3%		
第6病日	朝	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%	—	—	1,697	—	1,100	2,797	1.7%	2.3%		
	昼	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%	—	—	1,697	—	1,100	2,797	1.7%	2.3%		
	夜	1 0.1%	1 0.1%	—	1 0.1%	2 0.1%	0.760	1.54	1,697	—	1,100	2,797	1.8%	2.4%		
第7病日	朝	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%	—	—	1,696	—	1,099	2,795	1.8%	2.4%		
	昼	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%	—	—	1,696	—	1,099	2,795	1.8%	2.4%		
	夜	0 0.0%	0 0.0%	—	0 0.0%	0 0.0%	—	—	1,696	—	1,099	2,795	1.8%	2.4%		
累積発現率 (参考)	36	15	4	11	51	1.8%							1.8%	2.4%		

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.04 (0.55 , 1.95)	0.910
第1～第2病日	0.58 (0.19 , 1.71)	0.321
第1～第4病日	1.14 (0.58 , 2.26)	0.698

表5-1-8 タミフルと筋炎の併発

		発現数 (%)								①②③間の χ ² 検定 p 値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)		
		薬剤 未使用 ^①		薬剤使用		(内訳)		総計				未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用	
				計	使用開始 ^②	既使用 ^③												
第1病日	朝	7	0.3%	2	0.6%	2	0.6%	—	9	0.3%	0.368	2.19	2,516	330	—	2,846	0.3%	0.6%
	昼	4	0.2%	1	0.2%	0	0.0%	1	0.3%	0.515	0.87	2,205	304	328	2,837	0.5%	0.8%	
	夜	10	0.6%	1	0.1%	1	0.2%	0	0.0%	0.039	0.16	1,765	436	631	2,832	1.0%	0.9%	
第2病日	朝	2	0.2%	2	0.1%	1	0.2%	1	0.1%	0.862	0.70	1,158	597	1,066	2,821	1.2%	1.0%	
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	702	454	1,661	2,817	1.2%	1.0%	
	夜	0	0.0%	1	0.0%	0	0.0%	1	0.0%	0.751	0.00	492	210	2,115	2,817	1.2%	1.0%	
第3病日	朝	1	0.3%	0	0.0%	0	0.0%	0	0.0%	0.141	0.00	397	95	2,324	2,816	1.4%	1.0%	
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	335	61	2,419	2,815	1.4%	1.0%	
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	307	28	2,480	2,815	1.4%	1.0%	
第4病日	朝	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0.894	0.00	299	8	2,508	2,815	1.4%	1.0%	
	昼	0	0.0%	1	0.0%	0	0.0%	1	0.0%			293	6	2,516	2,815	1.4%	1.1%	
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%			290	3	2,521	2,814	1.4%	1.1%	
第5病日	朝	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	288	2	2,524	2,814	1.4%	1.1%	
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	286	2	2,526	2,814	1.4%	1.1%	
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	285	1	2,528	2,814	1.4%	1.1%	
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	285	—	2,529	2,814	1.4%	1.1%	
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	285	—	2,529	2,814	1.4%	1.1%	
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	284	1	2,529	2,814	1.4%	1.1%	
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	284	—	2,530	2,814	1.4%	1.1%	
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	284	—	2,530	2,814	1.4%	1.1%	
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	284	—	2,530	2,814	1.4%	1.1%	
累積発現率 (参考)	24		8		4		4		32	1.1%						2,814	1.4%	1.1%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	0.54 (0.22 , 1.33)	0.178
第1～第2病日	0.61 (0.25 , 1.52)	0.291
第1～第4病日	0.54 (0.22 , 1.33)	0.178

表5-2-8 アセトアミノフェンと筋炎の併発

		発現数 (%)								①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)	
		薬剤 未使用 ^①		薬剤使用		(内訳)		総計				未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用
				計	使用開始 ^②	既使用 ^③											
第1病日	朝	8	0.3%	1	1.0%	1	1.0%	—	9	0.3%	0.315	3.49	2,747	99	2,846	0.3%	1.0%
	昼	5	0.2%	0	0.0%	0	0.0%	0	5	0.2%	0.593	0.00	2,556	183	2,837	0.5%	1.0%
	夜	9	0.4%	2	0.3%	2	0.4%	0	11	0.4%	0.316	0.64	2,103	448	2,832	0.9%	1.3%
第2病日	朝	2	0.1%	2	0.2%	1	1.1%	1	4	0.1%	0.272	2.44	2,001	93	2,821	1.0%	1.5%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%	—	—	1,868	131	2,817	1.0%	1.5%
	夜	1	0.1%	0	0.0%	0	0.0%	0	1	0.0%	0.622	0.00	1,751	117	2,817	1.1%	1.5%
第3病日	朝	0	0.0%	1	0.1%	0	0.0%	1	1	0.0%	0.378	—	1,742	8	2,816	1.1%	1.6%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%	—	—	1,732	10	2,815	1.1%	1.6%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%	—	—	1,719	13	2,815	1.1%	1.6%
第4病日	朝	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%	—	—	1,717	2	2,815	1.1%	1.6%
	昼	0	0.0%	1	0.1%	0	0.0%	1	1	0.0%	0.390	—	1,714	3	2,815	1.1%	1.7%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%	—	—	1,710	4	2,814	1.1%	1.7%
第5病日	朝	0	0.0%	0	0.0%	—	—	0	0	0.0%	—	—	1,710	—	2,814	1.1%	1.7%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%	—	—	1,707	3	2,814	1.1%	1.7%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0	0.0%	—	—	1,706	1	2,814	1.1%	1.7%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0	0.0%	—	—	1,706	—	2,814	1.1%	1.7%
	昼	0	0.0%	0	0.0%	—	—	0	0	0.0%	—	—	1,706	—	2,814	1.1%	1.7%
	夜	0	0.0%	0	0.0%	—	—	0	0	0.0%	—	—	1,706	—	2,814	1.1%	1.7%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0	0.0%	—	—	1,706	—	2,814	1.1%	1.7%
	昼	0	0.0%	0	0.0%	—	—	0	0	0.0%	—	—	1,706	—	2,814	1.1%	1.7%
	夜	0	0.0%	0	0.0%	—	—	0	0	0.0%	—	—	1,706	—	2,814	1.1%	1.7%
累積発生率 (参考)	25		7		4		3	32	1.1%						1.1%	1.7%	

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.25 (0.52 , 3.05)	0.618
第1～第2病日	0.95 (0.35 , 2.60)	0.928
第1～第4病日	1.25 (0.52 , 3.05)	0.618

II. 患者家族用調査票についての基本的集計

表 6 発熱状況

	第1病日	第2病日	第3病日	第4病日	第5病日	第6病日	第7病日
人数	2,782	2,740	2,622	2,547	2,429	2,257	2,110
平均値	38.8	38.5	37.4	36.9	36.7	36.6	36.5
90%点	39.7	39.8	38.9	37.8	37.4	37.1	37.0
75%点	39.3	39.3	38.0	37.1	36.9	36.8	36.7
50%点	38.8	38.6	37.2	36.7	36.6	36.5	36.5
25%点	38.3	37.8	36.7	36.4	36.3	36.3	36.2
10%点	37.8	37.0	36.3	36.1	36.0	36.0	36.0

表 7-1 症状の発現の有無

	おびえ 恐怖	幻視幻覚	突然大声 うわごと	怒り出す ニヤリと	指を食物 のように	けいれん	意識消失	ひどい咳
発現なし	2,224	2,262	2,110	2,222	2,434	2,372	2,405	1,423
%	87.4	88.9	82.9	87.3	95.6	93.2	94.5	55.9
発現	221	149	326	215	17	80	39	1,045
%	8.7	5.9	12.8	8.5	0.7	3.1	1.5	41.1
不明	100	134	109	108	94	93	101	77
%	3.9	5.3	4.3	4.2	3.7	3.7	4.0	3.0

表 7-2 症状の発現の有無

	嘔吐下痢	筋肉痛	関節痛	ノド痛み	頭痛	耳の痛み	不機嫌
発現なし	1,576	1,841	1,685	1,221	1,189	2,109	84
%	61.9	72.3	66.2	48.0	46.7	82.9	3.3
発現	891	360	505	1,029	1,041	138	460
%	35.0	14.2	19.8	40.4	40.9	5.4	18.1
不明	78	344	355	295	315	298	2,001
%	3.1	13.5	14.0	11.6	12.4	11.7	78.6

表 7-3 薬剤使用の有無

	解熱剤	抗生物質	シメトリル	タミフル	リリザ
使用なし	1,340	1,801	2,418	184	2,498
%	52.7	70.8	95.0	7.2	98.2
使用	1,176	709	83	2,347	6
%	46.2	27.9	3.3	92.2	0.2
不明	29	35	44	14	41
%	1.1	1.4	1.7	0.6	1.6

表 8-1 症状の発現時期

		おびえ 恐怖	幻視幻覚	突然大声 うわごと	怒り出す ニヤリと	指を食物 のように	けいれん	意識消失	ひどい咳
第1病日	朝	19	8	14	20		5	1	398
	昼	20	8	27	22	1	16	6	93
	夜	101	52	154	61	7	28	11	201
第2病日	朝	5	8	14	11		10	5	62
	昼	24	18	21	29	2	12	3	27
	夜	30	32	68	41		6	5	48
第3病日	朝	1	2	1	7	1		1	42
	昼	4	4	2	6	1		1	16
	夜	9	5	13	6	1	1	2	38
第4病日	朝	1	1	1	3	1		1	27
	昼			2					7
	夜	1	2	2	2				26
第5病日	朝	1	1	2					11
	昼				1	1			7
	夜	1	1	1	1				16
第6病日	朝								4
	昼								1
	夜								6
第7病日	朝								2
	昼								2
	夜								1
計		217	142	322	210	15	78	36	1,035
第1病日	朝	8.8%	5.6%	4.3%	9.5%		6.4%	2.8%	38.5%
	昼	9.2%	5.6%	8.4%	10.5%	6.7%	20.5%	16.7%	9.0%
	夜	46.5%	36.6%	47.8%	29.0%	46.7%	35.9%	30.6%	19.4%
第2病日	朝	2.3%	5.6%	4.3%	5.2%		12.8%	13.9%	6.0%
	昼	11.1%	12.7%	6.5%	13.8%	13.3%	15.4%	8.3%	2.6%
	夜	13.8%	22.5%	21.1%	19.5%		7.7%	13.9%	4.6%
第3病日	朝	0.5%	1.4%	0.3%	3.3%	6.7%		2.8%	4.1%
	昼	1.8%	2.8%	0.6%	2.9%	6.7%		2.8%	1.5%
	夜	4.1%	3.5%	4.0%	2.9%	6.7%	1.3%	5.6%	3.7%
第4病日	朝	0.5%	0.7%	0.3%	1.4%	6.7%		2.8%	2.6%
	昼			0.6%					0.7%
	夜	0.5%	1.4%	0.6%	1.0%				2.5%
第5病日	朝	0.5%	0.7%	0.6%					1.1%
	昼				0.5%	6.7%			0.7%
	夜	0.5%	0.7%	0.3%	0.5%				1.5%
第6病日	朝								0.4%
	昼								0.1%
	夜								0.6%
第7病日	朝								0.2%
	昼								0.2%
	夜								0.1%
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%

表 8-2 症状の発現時期

		嘔吐下痢	筋肉痛	関節痛	ノド痛み	頭痛	耳の痛み	不機嫌
第1病日	朝	125	120	173	484	400	18	131
	夜	107	42	60	91	164	11	80
第2病日	朝	188	102	158	242	287	24	150
	夜	101	36	50	86	76	11	14
第3病日	朝	101	18	24	24	35	8	20
	夜	84	18	14	29	27	10	25
第4病日	朝	56	9	4	14	16	9	9
	夜	35	1	3	9	5	4	7
第5病日	朝	25	4	4	12	9	5	10
	夜	19	1	1	12	4	8	2
第6病日	朝	9	1	2	4	1	3	3
	夜	13		1	1	1	8	3
第7病日	朝	4	1	1	1	1	6	1
	夜	3	1	1	3	1	3	1
計	朝	8	1	2	1	1	2	
	夜	4	1	1	2	2	3	1
計		887	357	500	1,017	1,033	136	460
第1病日	朝	14.1%	33.6%	34.6%	47.6%	38.7%	13.2%	28.5%
	夜	12.1%	11.8%	12.0%	8.9%	15.9%	8.1%	17.4%
第2病日	朝	21.2%	28.6%	31.6%	23.8%	27.8%	17.6%	32.6%
	夜	11.4%	10.1%	10.0%	8.5%	7.4%	8.1%	3.0%
第3病日	朝	11.4%	5.0%	4.8%	2.4%	3.4%	5.9%	4.3%
	夜	9.5%	5.0%	2.8%	2.9%	2.6%	7.4%	5.4%
第4病日	朝	6.3%	2.5%	0.8%	1.4%	1.5%	6.6%	2.0%
	夜	3.9%	0.3%	0.6%	0.9%	0.5%	2.9%	1.5%
第5病日	朝	2.8%	1.1%	0.8%	1.2%	0.9%	3.7%	2.2%
	夜	2.1%	0.3%	0.2%	1.2%	0.4%	5.9%	0.4%
第6病日	朝	1.0%	0.3%	0.4%	0.4%	0.1%	2.2%	0.7%
	夜	1.5%		0.2%	0.1%	0.1%	5.9%	0.7%
第7病日	朝	0.5%	0.3%	0.2%	0.1%	0.1%	4.4%	0.2%
	夜	0.3%	0.3%	0.2%	0.3%	0.1%	2.2%	0.2%
計	朝	0.9%	0.3%	0.4%	0.1%	0.1%	1.5%	
	夜			0.2%	0.1%	0.1%		0.2%
第7病日	朝	0.3%	0.3%	0.2%	0.1%	0.2%		0.2%
	夜	0.5%	0.3%		0.2%	0.2%	2.2%	0.2%
計	朝	0.1%					0.7%	0.4%
	夜	0.1%					1.5%	
計		100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%

表 8-3 薬剤の使用開始時期

		解熱剤	抗生物質	シメトレル	タミフル	リンザ*
第1病日	朝	90	121	12	316	1
	昼	210	123	6	306	
	夜	485	151	15	403	2
第2病日	朝	82	54	16	522	
	昼	132	77	19	427	
	夜	117	29	3	182	1
第3病日	朝	6	11	2	82	
	昼	10	23	4	62	
	夜	9	7	1	22	
第4病日	朝	1	6	3	8	
	昼	3	10		6	
	夜	5	9		2	
第5病日	朝		21	1	1	
	昼	3	15			
	夜		11	1		
第6病日	朝		7			
	昼	1	8			
	夜	1	8		1	
第7病日	朝		7			
	昼	1	6			
	夜		2			
計		1,156	706	83	2,340	4
第1病日	朝	7.8%	17.1%	14.5%	13.5%	25.0%
	昼	18.2%	17.4%	7.2%	13.1%	
	夜	42.0%	21.4%	18.1%	17.2%	50.0%
第2病日	朝	7.1%	7.6%	19.3%	22.3%	
	昼	11.4%	10.9%	22.9%	18.2%	
	夜	10.1%	4.1%	3.6%	7.8%	25.0%
第3病日	朝	0.5%	1.6%	2.4%	3.5%	
	昼	0.9%	3.3%	4.8%	2.6%	
	夜	0.8%	1.0%	1.2%	0.9%	
第4病日	朝	0.1%	0.8%	3.6%	0.3%	
	昼	0.3%	1.4%		0.3%	
	夜	0.4%	1.3%		0.1%	
第5病日	朝		3.0%	1.2%	0.0%	
	昼	0.3%	2.1%			
	夜		1.6%	1.2%		
第6病日	朝		1.0%			
	昼	0.1%	1.1%			
	夜	0.1%	1.1%		0.0%	
第7病日	朝		1.0%			
	昼	0.1%	0.8%			
	夜		0.3%			
計		100.0%	100.0%	100.0%	100.0%	100.0%

表9 薬剤使用状況別の症状の経時的な発現数(%)

表9-1-1 タミフルとおびえ・恐怖

	薬剤 未使用 ^①	発現数(%)								総計	①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率(%)		
		薬剤使用		(内訳)		使用開始 ^②		既使用 ^③					未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用	
		計																	
第1病日	朝	14	0.7%	5	1.6%	5	1.6%	—	—	19	0.8%	0.105	2.51	2,124	305	0	2,429	0.7%	1.6%
	昼	7	0.4%	13	2.2%	7	2.5%	6	2.0%	20	0.8%	0.001	5.93	1,827	283	300	2,410	1.0%	3.8%
	夜	57	4.0%	44	4.6%	26	6.8%	18	3.2%	101	4.2%	0.026	1.17	1,437	383	570	2,390	5.0%	8.3%
第2病日	朝	2	0.2%	3	0.2%	1	0.2%	2	0.2%	5	0.2%	0.998	0.97	898	482	909	2,289	5.2%	8.5%
	昼	3	0.6%	21	1.2%	7	1.8%	14	1.0%	24	1.1%	0.249	1.97	499	397	1,388	2,284	5.7%	9.5%
	夜	11	3.3%	19	1.0%	3	1.9%	16	0.9%	30	1.3%	0.007	0.29	334	162	1,764	2,260	8.9%	10.4%
第3病日	朝	0	0.0%	1	0.1%	0	0.0%	1	0.1%	1	0.0%	0.855	—	251	72	1,907	2,230	8.9%	10.5%
	昼	1	0.5%	3	0.1%	0	0.0%	3	0.2%	4	0.2%	0.585	0.29	197	54	1,978	2,229	9.3%	10.6%
	夜	0	0.0%	9	0.4%	1	5.3%	8	0.4%	9	0.4%	0.092	—	177	19	2,029	2,225	9.3%	11.0%
第4病日	朝	0	0.0%	1	0.0%	0	0.0%	1	0.0%	1	0.0%	0.920	—	170	7	2,039	2,216	9.3%	11.1%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	—	—	164	6	2,045	2,215	9.3%	11.1%
	夜	0	0.0%	1	0.0%	0	0.0%	1	0.0%	1	0.0%	0.926	—	162	2	2,051	2,215	9.3%	11.1%
第5病日	朝	0	0.0%	1	0.0%	—	—	1	0.0%	1	0.0%	0.697	—	162	0	2,052	2,214	9.3%	11.1%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	162	0	2,051	2,213	9.3%	11.1%
	夜	0	0.0%	1	0.0%	—	—	1	0.0%	1	0.0%	0.697	—	162	0	2,051	2,213	9.3%	11.2%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	162	0	2,050	2,212	9.3%	11.2%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	162	0	2,050	2,212	9.3%	11.2%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	—	—	161	1	2,050	2,212	9.3%	11.2%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	161	0	2,051	2,212	9.3%	11.2%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	161	0	2,051	2,212	9.3%	11.2%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	161	0	2,051	2,212	9.3%	11.2%
累積発生率(参考)	95		122		50		72		217	8.9%								9.3%	11.2%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.30 (0.96 , 1.77)	0.092
第1～第2病日	1.30 (0.95 , 1.77)	0.102
第1～第4病日	1.30 (0.95 , 1.77)	0.096

表9-2-1 解熱剤とおびえ・恐怖

		発現数 (%)								①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)			
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^②		既使用 ^③				総計	未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用	
第1病日	朝	16	0.7%	3	3.7%	3	3.7%	—	—	19	0.8%	0.027	5.47	2,320	82	0	2,402	0.7%	3.7%
	昼	12	0.6%	8	2.9%	7	3.5%	1	1.3%	20	0.8%	0.003	5.13	2,103	201	79	2,383	1.3%	6.4%
	夜	68	4.2%	31	4.3%	24	5.3%	7	2.6%	99	4.2%	0.196	1.02	1,635	456	272	2,363	5.4%	10.4%
第2病日	朝	2	0.1%	3	0.4%	0	0.0%	3	0.4%	5	0.2%	0.369	2.95	1,499	68	697	2,264	5.5%	10.7%
	昼	14	1.0%	10	1.1%	4	3.3%	6	0.8%	24	1.1%	0.114	1.11	1,376	121	762	2,259	6.5%	11.8%
	夜	17	1.4%	13	1.3%	3	2.8%	10	1.1%	30	1.3%	0.453	0.98	1,255	107	873	2,235	7.7%	12.9%
第3病日	朝	1	0.1%	0	0.0%	0	0.0%	0	0.0%	1	0.0%	0.559	0.00	1,233	5	967	2,205	7.8%	12.9%
	昼	4	0.3%	0	0.0%	0	0.0%	0	0.0%	4	0.2%	0.095	0.00	1,225	7	972	2,204	8.1%	12.9%
	夜	4	0.3%	5	0.5%	0	0.0%	5	0.5%	9	0.4%	0.783	1.54	1,214	7	979	2,200	8.4%	13.4%
第4病日	朝	0	0.0%	1	0.1%	0	0.0%	1	0.1%	1	0.0%	0.448		1,209	1	981	2,191	8.4%	13.5%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,207	2	981	2,190	8.4%	13.5%
	夜	1	0.1%	0	0.0%	0	0.0%	0	0.0%	1	0.0%	0.549	0.00	1,203	4	983	2,190	8.5%	13.5%
第5病日	朝	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.0%	0.207		1,202	0	987	2,189	8.5%	13.5%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,199	3	986	2,188	8.5%	13.5%
	夜	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.0%	0.208		1,199	0	989	2,188	8.5%	13.6%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			1,199	0	988	2,187	8.5%	13.6%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,198	1	988	2,187	8.5%	13.6%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,197	1	989	2,187	8.5%	13.6%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			1,197	0	990	2,187	8.5%	13.6%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,196	1	990	2,187	8.5%	13.6%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			1,196	0	991	2,187	8.5%	13.6%
累積発生率 (参考)		139		76		41		35		215	9.0%							8.5%	13.6%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.26 (0.94 , 1.69)	0.116
第1～第2病日	1.30 (0.96 , 1.76)	0.095
第1～第4病日	1.24 (0.92 , 1.66)	0.157

表9-1-2 タミフルと幻視幻覚

		発現数 (%)						①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)					
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^② 既使用 ^③				総計	未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用			
第1病日	朝	7	0.3%	1	0.3%	1	0.3%	—	8	0.3%	0.992	0.99	2,090	302	0	2,392	0.3%	0.3%	
	昼	1	0.1%	7	1.2%	5	1.8%	2	0.7%	8	0.3%	0.000	22.08	1,805	278	301	2,384	0.4%	1.5%
	夜	24	1.7%	28	2.9%	11	2.9%	17	3.0%	52	2.2%	0.130	1.76	1,423	381	572	2,376	2.1%	4.4%
第2病日	朝	4	0.4%	4	0.3%	0	0.0%	4	0.4%	8	0.3%	0.149	0.64	907	492	925	2,324	2.5%	4.7%
	昼	3	0.6%	15	0.8%	5	1.3%	10	0.7%	18	0.8%	0.515	1.40	505	398	1,413	2,316	3.1%	5.5%
	夜	9	2.7%	23	1.2%	3	1.8%	20	1.1%	32	1.4%	0.106	0.43	335	167	1,796	2,298	5.7%	6.6%
第3病日	朝	1	0.4%	1	0.0%	0	0.0%	1	0.1%	2	0.1%	0.379	0.12	251	75	1,940	2,266	6.1%	6.6%
	昼	0	0.0%	4	0.2%	0	0.0%	4	0.2%	4	0.2%	0.626		197	53	2,014	2,264	6.1%	6.8%
	夜	0	0.0%	5	0.2%	0	0.0%	5	0.2%	5	0.2%	0.634		177	20	2,063	2,260	6.1%	7.0%
第4病日	朝	0	0.0%	1	0.0%	0	0.0%	1	0.0%	1	0.0%	0.922		170	7	2,078	2,255	6.1%	7.1%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			164	6	2,084	2,254	6.1%	7.1%
	夜	0	0.0%	2	0.1%	0	0.0%	2	0.1%	2	0.1%	0.860		162	2	2,090	2,254	6.1%	7.2%
第5病日	朝	0	0.0%	1	0.0%	—	—	1	0.0%	1	0.0%	0.699		162	0	2,090	2,252	6.1%	7.2%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			162	0	2,089	2,251	6.1%	7.2%
	夜	0	0.0%	1	0.0%	—	—	1	0.0%	1	0.0%	0.699		162	0	2,089	2,251	6.1%	7.3%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			162	0	2,088	2,250	6.1%	7.3%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			162	0	2,088	2,250	6.1%	7.3%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			161	1	2,088	2,250	6.1%	7.3%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			161	0	2,089	2,250	6.1%	7.3%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			161	0	2,089	2,250	6.1%	7.3%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			161	0	2,089	2,250	6.1%	7.3%
累積発現率 (参考)		49		93		25		68		142	5.9%						6.1%	7.3%	

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.40 (0.94 , 2.09)	0.098
第1～第2病日	1.40 (0.93 , 2.11)	0.103
第1～第4病日	1.40 (0.93 , 2.08)	0.103

表9-2-2 解熱剤と幻視幻覚

		発現数 (%)								①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)			
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^②		既使用 ^③				総計	未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用	
																			未使用
第1病日	朝	6	0.3%	2	2.5%	2	2.5%	—	8	0.3%	0.026	9.61	2,284	81	0	2,365	0.3%	2.5%	
	昼	6	0.3%	2	0.7%	0	0.0%	2	2.5%	8	0.3%	0.047	2.50	2,079	199	79	2,357	0.6%	3.2%
	夜	29	1.8%	23	3.2%	13	2.9%	10	3.6%	52	2.2%	0.108	1.80	1,623	450	276	2,349	2.3%	6.2%
第2病日	朝	3	0.2%	4	0.5%	1	1.4%	3	0.4%	7	0.3%	0.297	2.62	1,521	73	703	2,297	2.5%	6.7%
	昼	13	0.9%	5	0.6%	2	1.6%	3	0.4%	18	0.8%	0.210	0.60	1,394	124	772	2,290	3.4%	7.2%
	夜	20	1.6%	12	1.2%	3	2.7%	9	1.0%	32	1.4%	0.293	0.76	1,271	110	891	2,272	4.9%	8.4%
第3病日	朝	1	0.1%	1	0.1%	0	0.0%	1	0.1%	2	0.1%	0.982	1.25	1,246	5	989	2,240	5.0%	8.4%
	昼	4	0.3%	0	0.0%	0	0.0%	0	0.0%	4	0.2%	0.093	0.00	1,237	8	993	2,238	5.3%	8.4%
	夜	4	0.3%	1	0.1%	0	0.0%	1	0.1%	5	0.2%	0.494	0.30	1,225	8	1,001	2,234	5.6%	8.5%
第4病日	朝	0	0.0%	1	0.1%	0	0.0%	1	0.1%	1	0.0%	0.452		1,220	1	1,008	2,229	5.6%	8.6%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,217	3	1,008	2,228	5.6%	8.6%
	夜	1	0.1%	1	0.1%	0	0.0%	1	0.1%	2	0.1%	0.988	1.20	1,213	4	1,011	2,228	5.7%	8.7%
第5病日	朝	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.0%	0.210		1,212	0	1,014	2,226	5.7%	8.8%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,209	3	1,013	2,225	5.7%	8.8%
	夜	0	0.0%	1	0.1%	—	—	1	0.1%	1	0.0%	0.211		1,209	0	1,016	2,225	5.7%	8.9%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			1,209	0	1,015	2,224	5.7%	8.9%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,208	1	1,015	2,224	5.7%	8.9%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,207	1	1,016	2,224	5.7%	8.9%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			1,207	0	1,017	2,224	5.7%	8.9%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			1,206	1	1,017	2,224	5.7%	8.9%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			1,206	0	1,018	2,224	5.7%	8.9%
累積発生率 (参考)		87		54		21		33		141	6.0%							5.7%	8.9%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.22 (0.86 , 1.73)	0.269
第1～第2病日	1.31 (0.90 , 1.90)	0.155
第1～第4病日	1.18 (0.83 , 1.68)	0.360

表9-1-3 タミフルと突然大声・うわごと

		発現数 (%)								①②③間の		期間ごとの オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)		
		薬剤 未使用 ^①		薬剤使用 計		(内訳) 使用開始 ^②		既使用 ^③		総計			χ ² 検定 p値	未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用
第1病日	朝	10	0.5%	4	1.3%	4	1.3%	—	—	14	0.6%	0.108	2.83	2,118	302	0	2,420	0.5%	1.3%
	昼	11	0.6%	16	2.7%	9	3.1%	7	2.3%	27	1.1%	0.000	4.64	1,822	286	298	2,406	1.1%	4.0%
	夜	82	5.7%	71	7.5%	34	9.0%	37	6.5%	153	6.4%	0.079	1.34	1,435	376	568	2,379	6.7%	11.2%
第2病日	朝	5	0.6%	9	0.7%	3	0.6%	6	0.7%	14	0.6%	0.949	1.18	883	470	873	2,226	7.3%	11.8%
	昼	4	0.8%	17	1.0%	6	1.5%	11	0.8%	21	0.9%	0.461	1.21	489	389	1,334	2,212	8.0%	12.7%
	夜	10	3.1%	58	3.1%	8	5.1%	50	2.9%	68	3.1%	0.391	1.02	327	158	1,706	2,191	10.8%	15.4%
第3病日	朝	0	0.0%	1	0.1%	0	0.0%	1	0.1%	1	0.0%	0.851		244	73	1,806	2,123	10.8%	15.5%
	昼	0	0.0%	2	0.1%	0	0.0%	2	0.1%	2	0.1%	0.783		191	53	1,878	2,122	10.8%	15.6%
	夜	1	0.6%	12	0.6%	0	0.0%	12	0.6%	13	0.6%	0.903	1.08	175	16	1,929	2,120	11.3%	16.1%
第4病日	朝	0	0.0%	1	0.1%	0	0.0%	1	0.1%	1	0.0%	0.917		167	7	1,933	2,107	11.3%	16.1%
	昼	0	0.0%	2	0.1%	0	0.0%	2	0.1%	2	0.1%	0.848		161	6	1,939	2,106	11.3%	16.2%
	夜	2	1.3%	0	0.0%	0	0.0%	0	0.0%	2	0.1%	0.006	0.00	159	2	1,943	2,104	12.5%	16.2%
第5病日	朝	0	0.0%	2	0.1%	—	—	2	0.1%	2	0.1%	0.577		157	0	1,945	2,102	12.5%	16.3%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			157	0	1,943	2,100	12.5%	16.3%
	夜	1	0.6%	0	0.0%	—	—	0	0.0%	1	0.0%	0.023	0.00	157	0	1,943	2,100	13.0%	16.3%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			156	0	1,943	2,099	13.0%	16.3%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			156	0	1,943	2,099	13.0%	16.3%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%			155	1	1,943	2,099	13.0%	16.3%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			155	0	1,944	2,099	13.0%	16.3%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			155	0	1,944	2,099	13.0%	16.3%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%			155	0	1,944	2,099	13.0%	16.3%
累積発生率 (参考)		126		195		64		131		321	13.3%							13.0%	16.3%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.43 (1.11 , 1.85)	0.005
第1～第2病日	1.50 (1.16 , 1.94)	0.002
第1～第4病日	1.45 (1.13 , 1.87)	0.004

表9-2-3 解熱剤と突然大声・うわごと

		発現数 (%)						①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)	
		薬剤 未使用 ^①	薬剤使用 計		(内訳)		総計			未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用
			使用開始 ^②	既使用 ^③											
第1病日	朝	14	0	0	0	—	14	0.317	0.00	2,310	84	0	2,394	0.6%	0.0%
	昼	16	11	11	0	0	27	<.0001	5.33	2,100	196	84	2,380	1.4%	3.9%
	夜	90	62	39	23	8.6%	152	0.022	1.62	1,634	450	269	2,353	6.8%	12.2%
第2病日	朝	6	8	1	7	1.1%	14	0.171	2.73	1,475	69	657	2,201	7.2%	13.2%
	昼	12	9	3	6	0.8%	21	0.316	1.21	1,350	119	718	2,187	8.0%	14.1%
	夜	26	41	6	35	4.2%	67	0.008	2.13	1,233	105	829	2,166	9.9%	17.9%
第3病日	朝	0	1	0	1	0.1%	1	0.425	0.00	1,202	5	892	2,099	9.9%	18.0%
	昼	2	0	0	0	0.0%	2	0.324	0.00	1,194	8	896	2,098	10.1%	18.0%
	夜	6	7	0	7	0.8%	13	0.710	1.52	1,184	8	904	2,096	10.5%	18.6%
第4病日	朝	1	0	0	0	0.0%	1	0.565	0.00	1,177	1	905	2,083	10.6%	18.6%
	昼	0	2	0	2	0.2%	2	0.189	0.00	1,173	3	906	2,082	10.6%	18.8%
	夜	1	1	1	0	0.0%	2	0.004	1.28	1,169	4	907	2,080	10.7%	18.9%
第5病日	朝	1	1	—	1	0.1%	2	0.860	1.28	1,168	0	910	2,078	10.8%	19.0%
	昼	0	0	0	0	0.0%	0	0.199	—	1,165	2	909	2,076	10.8%	19.0%
	夜	0	1	—	1	0.1%	1	—	—	1,165	0	911	2,076	10.8%	19.1%
第6病日	朝	0	0	—	0	0.0%	0	—	—	1,165	0	910	2,075	10.8%	19.1%
	昼	0	0	0	0	0.0%	0	—	—	1,164	1	910	2,075	10.8%	19.1%
	夜	0	0	0	0	0.0%	0	—	—	1,163	1	911	2,075	10.8%	19.1%
第7病日	朝	0	0	—	0	0.0%	0	—	—	1,163	0	912	2,075	10.8%	19.1%
	昼	0	0	0	0	0.0%	0	—	—	1,162	1	912	2,075	10.8%	19.1%
	夜	0	0	—	0	0.0%	0	—	—	1,162	0	913	2,075	10.8%	19.1%
累積発現率 (参考)	175	144	61	83	13.3%	319								10.8%	19.1%

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.83 (1.46 , 2.31)	<.0001
第1～第2病日	1.86 (1.47 , 2.36)	<.0001
第1～第4病日	1.83 (1.45 , 2.30)	<.0001

表9-1-4 タミフルと怒り出す

		発現数 (%)								①②③間の χ ² 検定 p値	期間ごと の オッズ比	観察例数 (臨床症候の既発現者を除く)				累積発現率 (%)			
		薬剤 未使用 ^①		薬剤使用 計		(内訳)		総計				未使用	使用開始	既使用	総計	薬剤未使用	薬剤使用		
						使用開始 ^②	既使用 ^③												
第1病日	朝	13	0.6%	7	2.3%	7	2.3%	—	20	0.8%	0.009	3.83	2,117	303	0	2,420	0.6%	2.3%	
	昼	10	0.5%	12	2.1%	6	2.1%	6	2.0%	22	0.9%	0.009	3.82	1,820	284	296	2,400	1.2%	4.3%
	夜	36	2.5%	25	2.6%	10	2.7%	15	2.6%	61	2.6%	0.980	1.05	1,433	377	568	2,378	3.6%	6.9%
第2病日	朝	6	0.7%	5	0.4%	2	0.4%	3	0.3%	11	0.5%	0.570	0.54	908	489	920	2,317	4.3%	7.2%
	昼	7	1.4%	22	1.2%	7	1.8%	15	1.1%	29	1.3%	0.542	0.88	506	396	1,404	2,306	5.6%	8.3%
	夜	6	1.8%	35	1.8%	4	2.4%	31	1.7%	41	1.8%	0.830	1.01	335	164	1,778	2,277	7.3%	10.0%
第3病日	朝	2	0.8%	5	0.3%	2	2.6%	3	0.2%	7	0.3%	0.017	0.32	252	77	1,907	2,236	8.0%	10.2%
	昼	1	0.5%	5	0.2%	1	1.8%	4	0.2%	6	0.3%	0.246	0.48	194	56	1,979	2,229	8.5%	10.4%
	夜	0	0.0%	6	0.3%	1	5.3%	5	0.2%	6	0.3%	0.079	—	174	19	2,030	2,223	8.5%	10.7%
第4病日	朝	0	0.0%	3	0.1%	0	0.0%	3	0.1%	3	0.1%	0.782	—	167	7	2,043	2,217	8.5%	10.8%
	昼	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	—	—	161	6	2,047	2,214	8.5%	10.8%
	夜	1	0.6%	1	0.0%	0	0.0%	1	0.0%	2	0.1%	0.266	0.08	159	2	2,053	2,214	9.1%	10.9%
第5病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	158	0	2,054	2,212	9.1%	10.9%
	昼	0	0.0%	1	0.0%	—	—	1	0.0%	1	0.0%	0.700	—	158	0	2,054	2,212	9.1%	10.9%
	夜	0	0.0%	1	0.0%	—	—	1	0.0%	1	0.0%	0.700	—	158	0	2,053	2,211	9.1%	10.9%
第6病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	158	0	2,052	2,210	9.1%	10.9%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	158	0	2,052	2,210	9.1%	10.9%
	夜	0	0.0%	0	0.0%	0	0.0%	0	0.0%	0	0.0%	—	—	157	1	2,052	2,210	9.1%	10.9%
第7病日	朝	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	157	0	2,053	2,210	9.1%	10.9%
	昼	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	157	0	2,053	2,210	9.1%	10.9%
	夜	0	0.0%	0	0.0%	—	—	0	0.0%	0	0.0%	—	—	157	0	2,053	2,210	9.1%	10.9%
累積発現率 (参考)		82		128		40		88		210	8.7%					2,210	9.1%	10.9%	

薬剤使用開始と症候発現が同時期： 使用開始後に出現と仮定	ハザード比 (95%CI)	p値
第1～第7病日	1.25 (0.90 , 1.75)	0.181
第1～第2病日	1.34 (0.95 , 1.88)	0.095
第1～第4病日	1.25 (0.90 , 1.74)	0.188