

9. Navarro E, Casao MA, Solera J. Diagnosis of human brucellosis using PCR. *Expert Rev Mol Diagn* 2004; 4:115–23.
10. Avdikou I, Maipa V, Alamanos Y. Epidemiology of human brucellosis in a defined area of northwestern Greece. *Epidemiol Infect* 2005; 133: 905–10.
11. Alton GG, Jones LM, Pietz DE. Laboratory techniques in brucellosis. 2nd ed. Geneva: World Health Organization, 1975.
12. Yagupsky P. Detection of brucellae in blood cultures. *J Clin Microbiol* 1999; 37:3437–42.
13. Solomon HM, Jackson D. Rapid diagnosis of *Brucella melitensis* in blood: some operational characteristics of the BACT/ALERT. *J Clin Microbiol* 1992; 30:222–4.
14. Shapiro DS, Wong JD. *Brucella*. In: Murray P, Baron E, Pfaller M, Tenover F, Tenover R, eds. *Manual of clinical microbiology*. Washington DC: American Society for Microbiology, 1999:625–31.
15. Baily GG, Krahn JB, Drasar BS, Stoker NG. Detection of *Brucella melitensis* and *Brucella abortus* by DNA amplification. *J Trop Med Hyg* 1992; 95:271–5.
16. Espy MJ, Uhl JR, Sloan LM, et al. Real-time PCR in clinical microbiology: applications for routine laboratory testing. *Clin Microbiol Rev* 2006; 19:165–256.
17. Tenover FC. Rapid detection and identification of bacterial pathogens using novel molecular technologies: infection control and beyond. *Clin Infect Dis* 2007; 44:418–23.
18. Tolomeo M, Di Carlo P, Abbadessa V, et al. Monocyte and lymphocyte apoptosis resistance in acute and chronic brucellosis and its possible implications in clinical management. *Clin Infect Dis* 2003; 36:1533–8.
19. Boura P, Skendros P, Kountouras J, Zacharioudaki E, Tsapas T. Effect of bacterial extracts on the immunologic profile in chronic relapsing brucellosis patients. *Int J Immunopathol Pharmacol* 1999; 12:103–11.
20. Gotuzzo E, Carrillo C, Guerra J, Llosa L. An evaluation of diagnostic methods for brucellosis—the value of bone marrow culture. *J Infect Dis* 1986; 153:122–5.
21. Navarro E, Segura JC, Castano MJ, Solera J. Use of real-time quantitative polymerase chain reaction to monitor the evolution of *Brucella melitensis* DNA load during therapy and post-therapy follow-up in patients with brucellosis. *Clin Infect Dis* 2006; 42:1266–73.
22. Maas KS, Mendez M, Zavaleta M, et al. Evaluation of brucellosis by PCR and persistence after treatment in patients returning to the hospital for follow-up. *Am J Trop Med Hyg* 2007; 76:698–702.
23. Queipo-Ortuño MI, Colmenero JD, Baeza G, Morata P. Comparison between LightCycler Real-Time Polymerase Chain Reaction (PCR) assay with serum and PCR–enzyme-linked immunosorbent assay with whole blood samples for the diagnosis of human brucellosis. *Clin Infect Dis* 2005; 40:260–4.
24. Kattar MM, Zalloua PA, Araj GE, et al. Development and evaluation of real-time polymerase chain reaction assays on whole blood and paraffin-embedded tissues for rapid diagnosis of human brucellosis. *Diagn Microbiol Infect Dis* 2007; 59:23–32.
25. Debeaumont C, Falconnet PA, Maurin M. Real-time PCR for detection of *Brucella* spp DNA in human serum samples. *Eur J Clin Microbiol Infect Dis* 2005; 24:842–5.
26. Dhand A, Röss JJ. Implantable cardioverter-defibrillator infection due to *Brucella melitensis*: case report and review of brucellosis of cardiac devices. *Clin Infect Dis* 2007; 44:e37–39.
27. Pappas G, Akritidis N, Tsianos E. Effective treatments in the management of brucellosis. *Expert Opin Pharmacother* 2005; 6:201–9.

医薬品 研究報告 調査報告書

識別番号・報告回数		報告日		第一報入手日 2008. 7. 3	新医薬品等の区分 該当なし	機構処理欄
一般的名称 (製造販売承認書に記載なし)		研究報告の公表状況		Pérez-Osorio CE, Zavala-Velázquez JE, Arias León JJ, Zavala-Castro JE. Emerg Infect Dis. 2008 Jul;14(7):1019-23.	公表国 メキシコ	
販売名(企業名) 合成血-LR「日赤」(日本赤十字社) 照射合成血-LR「日赤」(日本赤十字社)						
研究報告の概要 227	○ <i>Rickettsia felis</i>—ヒトへの新興世界的脅威 <i>Rickettsia felis</i>は、TRG <i>rickettsiae</i> (transitional group rickettsiae) に属する新興病原体である。<i>R. felis</i>は、イヌ、ネコ、齧歯類など世界中の様々な外部寄生生物から検出され、ネコノミは最も一般的な媒介生物と考えられている。ヒトにおける<i>R. felis</i>感染症の臨床症状は発疹熱やデング熱などに類似しており、ヒトの感染症は実際よりも少なく推定されている可能性が高い。ヒト症例は、1994年に第1例が米国で報告された後、ヨーロッパやアジアでも報告されている。<i>R. felis</i>は存続のためにノミだけを必要とする可能性があるが、ノミが媒介する紅斑熱リゲチアのライフサイクルにおける動物の役割についてはまだわかっていない。また、鑑別診断においては、ノミが媒介する紅斑熱を考慮すべきである。ヒトにおける<i>R. felis</i>感染症の実際の発現率、臨床症状の範囲、重症度を調べ、さらに公衆衛生への影響を評価するために、今後、試験を実施していく必要がある。					使用上の注意記載状況・ その他参考事項等 合成血-LR「日赤」 照射合成血-LR「日赤」 血液を介するウイルス、 細菌、原虫等の感染 vCJD等の伝播のリスク
	報告企業の意見		今後の対応			
ネコノミが媒介する<i>Rickettsia felis</i>感染症のヒト症例は世界中で報告されているが、症状は発疹熱やデング熱などに類似しており、実際よりも少なく推定されている可能性が高く、今後調査が必要であるとの報告である。		日本赤十字社では、輸血感染症対策として問診時に海外渡航歴の有無を確認し、帰国(入国)後4週間は献血不適としている。また、発熱などの体調不良者を献血不適としている。今後も引き続き、新興・再興感染症の発生状況等に関する情報の収集に努める。				

***Rickettsia felis* as Emergent Global Threat for Humans**

Carlos E. Pérez-Osorio,* Jorge E. Zavala-Velázquez,* Juan José Arias León,*
and Jorge E. Zavala-Castro*

Rickettsia felis is an emergent pathogen belonging to transitional group rickettsiae. First described in 1990, *R. felis* infections have been reported to occur worldwide in fleas, mammals, and humans. Because clinical signs of the illness are similar to those of murine typhus and other febrile illnesses such as dengue, the infection in humans is likely underestimated. *R. felis* has been found throughout the world in several types of ectoparasites; cat fleas appear to be the most common vectors. *R. felis* infection should be considered an emergent threat to human health.

Rickettsia felis is a member of the genus *Rickettsia*, which comprises intracellular pathogens that produce infections commonly called rickettsioses. Although the genus has no recognized subspecies, rickettsiae have traditionally been subdivided into 2 groups: the spotted fever group (SFG) and the typhus group. Infections produced by these 2 groups are clinically indistinguishable; however, groups can be differentiated by outer membrane protein OmpA (absent in the typhus group) and by vector. SFG members are transmitted by ticks; typhus group members, by fleas and lice (1,2). More recently, Gillespie et al. (3) added to this classification by designating the transitional group of rickettsiae and describing an ancestral group of rickettsiae.

In 1990, Adams et al. described a rickettsia-like organism, which resembled *R. typhi*, in the cytoplasm of midgut cells of a colony of cat fleas (1). The new rickettsia received the initial name of ELB agent after the company from which the fleas were obtained (El Labs, Soquel, CA;

USA) (4). The first observations, such as reactivity with antibodies to *R. typhi* (1), the type of vector in which it was first discovered (1), and the apparent absence of OmpA (5), suggested that the new organism belonged to the typhus group of rickettsiae (4).

The molecular characterization of the organism described by Adams and reported by Bouyer et al. in 2001 provided sufficient evidence to support the designation of *R. felis* as a member of the SFG (6), and in 2002, La Scola et al. provided further characterization (7). One noticeable characteristic is the temperature-dependent growth of the bacterium, which requires incubation temperatures of 28°–32°C for optimal growth. However, the most striking characteristic of the novel rickettsia was the plasmid DNA in its genome (8).

World Distribution in Potential Host Vectors

Soon after the initial description of the typhus-like rickettsia, Williams et al. (9) reported that cat fleas collected from opossums in an urban setting in California were infected with the novel rickettsia, but no organism was detected in the tissues of the opossums. Since this report, this organism has been described in infected vectors from 20 countries on 5 continents (9). Not until 2002 did interest in *R. felis* increase, when the United States (9), Brazil (10), Mexico (11), and Spain (12) were among the first countries to describe cat fleas (*Ctenocephalides felis*) infected with *R. felis*. During the following 5 years, 28 additional reports appeared from all over the world (Table 1). These reports describe new potential vectors being infected with the emergent rickettsia, including the following: fleas, such as *C. canis* (13–15), *Anomiopsyllus nudata* (16), *Archaeopsylla erinacei* (15,17), *Ctenophthalmus* sp.

*Universidad Autónoma de Yucatán, Mérida, México

SYNOPSIS

Table 1. Potential vectors infected with *Rickettsia felis* reported worldwide, 1992–2007*

Year	Source of DNA sample	Animal†	Country	Reference
1992	<i>Ctenocephalides felis</i>	Opossum	USA	(9)
2002	<i>C. felis</i>	Cats and dogs	Brazil	(10)
2002	<i>C. felis</i>	Dogs	Mexico	(11)
2002	<i>C. felis</i>	Cats and dogs	Spain	(12)
2003	<i>Haemaphysalis flava</i> , <i>H. kitaokai</i> , and <i>Ixodes ovatus</i>	Unknown (flagging)	Japan	(19)
2003	<i>C. felis</i>	Cats	France	(22)
2003	<i>C. felis</i>	Cats and dogs	UK	(23)
2004	<i>C. felis</i>	Dogs	Peru	(24)
2005	<i>Anomopsyllus nudata</i>	Wild rodents	USA	(16)
2005	<i>C. felis</i>	Cats and dogs	New Zealand	(25)
2005	<i>C. felis</i>	Monkey	Gabon	(26)
2006	<i>C. felis</i> and <i>C. canis</i>	Dogs	Brazil	(13)
2006	<i>C. felis</i> and <i>C. canis</i>	Cats and dogs	Uruguay	(14)
2006	<i>Archaeopsylla erinacei</i> and <i>C. canis</i>	Hedgehog and rodents	Algeria	(15)
2006	<i>A. erinacei</i> and <i>Ctenophtalmus</i> sp.	Rodents and hedgehog	Portugal	(17)
2006	<i>Xenopsylla cheopis</i>	Rodents‡	Indonesia	(18)
2006	<i>C. felis</i> , <i>Rhipicephalus sanguineus</i> , and <i>Amblyomma cajennense</i>	Dogs and horse	Brazil	(20)
2006	Unknown flea	Gerbil	Afghanistan	(27)
2006	<i>C. felis</i>	Cats and dogs	Australia	(28)
2006	<i>C. felis</i>	Cats	Israel	(29)
2006	<i>C. felis</i>	Rodents	Cyprus	(30)
2007	Mites	Wild rodents	South Korea	(21)
2007	<i>C. felis</i>	Cats	USA	(31)
2007	<i>C. felis</i>	Cats	Chile	(32)

*PCR was used to detect *R. felis* infection with 1 noted exception.

†Animal host of potential vectors.

‡Quantitative PCR.

(17), and *Xenopsylla cheopis* (18); ticks, *Haemaphysalis flava* (19), *Rhipicephalus sanguineus* (20), and *Ixodes ovatus* (19); and mites from South Korea (21) (Table 1). Despite the large number of potential vectors reported, the only vector currently recognized is *C. felis* because it has been demonstrated that this flea is able to maintain a stable infected progeny through transovarial transmission (4). In addition, production of antibody to *R. felis* has been noted in animals after they have been exposed to infected cat fleas (9). Other evidence to be considered is the fact that 68.8% of the reports state that the cat flea is the most recurrent vector in which *R. felis* has been detected. These data further support the wide distribution of rickettsiae because they correlate with the worldwide distribution of *C. felis*; this distribution represents a threat to the human population because of lack of host specificity of the cat flea.

R. felis infection is diagnosed by PCR amplification of targeted genes. The genes most commonly amplified by researchers are *gltA* and *ompB*; followed by the 17-kDa gene. Also, 25% of published articles report that *R. felis* was detected by amplifying >2 genes, and all report that amplicons were confirmed as *R. felis* by sequencing. The animal hosts from which the infected ectoparasites were recovered represent a diversity of mammals (Table 1), which included 9 different naturally infested animal

species. However, in 16 of 33 articles, ectoparasites were recovered from dogs. Other hosts for ectoparasites were cats (in 13 of 33 reports); rodents (5 of 33 reports); opossums and hedgehogs (2 reports each); and horses, sheep, goats, gerbils, and monkeys (1 report for each animal species).

In summary, the presence of *R. felis* in a diverse range of invertebrate and mammalian hosts represents a high potential risk for public health and the need for further studies to establish the role of ectoparasites other than *C. felis* as potential vectors. To date, whether any vertebrate may serve as the reservoir of this emergent pathogen has not been determined. However, preliminary data from our laboratory suggest that opossums are the most likely candidates.

World Distribution of Human Cases

In 1994, the first human case of infection with the new cat flea rickettsia was reported in the United States (2). This became the first evidence of *R. felis*' potential as a human pathogen. *R. felis* infection had a similar clinical manifestation as murine typhus (including high fever [39°–40°C], myalgia, and rash). Although the initial idea was that the murine typhus-like rickettsia had a transmission cycle involving cat fleas and opossums (2,5,9), no viable *R. felis* has yet been isolated from a vertebrate host.

Three more cases of *R. felis* infection were reported from southeastern Mexico in 2000. The patients had had contact with fleas or animals known to carry fleas. The clinical manifestations were those of a typical rickettsiosis: all patients had fever and myalgia; but the skin lesions, instead of a rash, were similar to those described for rickettsialpox. In addition, for 3 patients, central nervous system involvement developed, manifested as photophobia, hearing loss, and signs of meningitis (33).

As occurred with the fast-growing reports of the worldwide detection of *R. felis* in arthropod hosts, the reports of human cases of *R. felis* infection increased rapidly in the following years (Table 2). But, in contrast, only 11 articles reported human infection by *R. felis* compared with 32 that reported ectoparasite infection with the new rickettsia. Nevertheless, these findings indicate that an effective surveillance system is urgently needed to distinguish *R. felis* rickettsiosis from other rickettsial infections such as murine typhus and Rocky Mountain spotted fever, and from other febrile illnesses such as dengue. Although PCR is still a method of choice for many laboratories, its high cost prevents many from using the technique, particularly in developing countries. Important advances have been achieved in diagnostics, such as the recent establishment of a stable culture of *R. felis* in cell lines that allows its use as antigen in serologic assays differentiating the cat flea rickettsia from others. Use of this culture in the immunofluorescent assay has enabled detection of additional human cases (38).

The first autochthonous human case in Europe was reported in 2002, which demonstrated that *R. felis* has a potential widespread distribution and is not confined to the Americas. It also confirmed the risk for human disease anywhere in the world. After the first report in Europe of a human infection of *R. felis*, other human cases have appeared in other countries around the world, including Thailand (36), Tunisia (38), Laos (39), and Spain (40); additional cases have been reported in Mexico and Brazil (34). All the data support the conclusion that the incidence of *R. felis* rickettsiosis and the simultaneous worldwide distribution of the flea vector plausibly explain its endemicity.

At present, the involvement of domestic animals (e.g., dogs and cats) or wild animals coexisting in urban areas (e.g., opossums) maintains *R. felis* infection in nature. *C. felis* fleas serve as the main reservoir and likely have a central role in transmission of human illness.

Conclusions

R. felis is an emergent rickettsial pathogen with a worldwide distribution in mammals, humans, and ectoparasites. The clinical manifestations of *R. felis* infections resemble those of murine typhus and dengue, which makes them difficult to diagnose without an appropriate laboratory test. For this reason, infections due to this emergent pathogen are likely underestimated and misdiagnosed. Although *R. felis* may require only fleas for its maintenance in nature, we still do not know the role of animals in the life cycle of flea-borne spotted fever rickettsia. In addition, flea-borne spotted fever should be considered in the differential diagnosis of infectious diseases. Further research should be conducted to determine the actual incidence of *R. felis* infection in humans, the spectrum of clinical signs and symptoms, and the severity of this infection and also to assess the impact on public health.

Acknowledgment

We thank Patricia Croquet-Valdes for her comments and helpful advice.

This research was supported by grants from the CONACYT (44064-M) to J. E. Z.-V.

Mr. Perez-Orsorio is currently professor of microbiology at the Autonomous University of Yucatan. His research interests focus on rickettsial diseases.

References

- Adams JR, Schmidtman ET, Azad AF. Infection of colonized cat fleas, *Ctenocephalides felis* (Bouché), with a rickettsia-like microorganism. *Am J Trop Med Hyg*. 1990;43:400-9.
- Schriefer ME, Sacci JB Jr, Dumlér JS, Bullen MG, Azad AF. Identification of a novel rickettsial infection in a patient diagnosed with murine typhus. *J Clin Microbiol*. 1994;32:949-54.

Table 2. Human cases of *Rickettsia felis* infection reported worldwide, 1994–2006*

Year	No. cases	Method	Country	Reference
1994	1	PCR	USA	(2)
2000, 2006	5	PCR	Mexico	(33)
2001, 2006	3	PCR	Brazil	(34)
2002	2	PCR/serology	Germany	(35)
2003	1	Serology (seroconversion)	Thailand	(36)
2005	3	Serology (Western blot)	South Korea	(37)
2006	8	Serology (IFAT/Western blot)	Tunisia	(38)
2006	1	Serology (seroconversion)	Laos	(39)
2006	33	Serology (IFAT)	Spain	(40)
Total	68			

*IFAT, indirect fluorescent antibody test.

SYNOPSIS

3. Gillespie JJ, Beier MS, Rahman MS, Ammerman NC, Shallow JM, Purkayastha A, et al. Plasmids and rickettsial evolution: insight from *Rickettsia felis*. PLoS One. 2007;2:e266. DOI: 10.1371/journal.pone.0000266
4. Azad AF, Sacci JB Jr, Nelson WM, Dasch GA, Schmidtman ET, Carl M. Genetic characterization and transovarial transmission of a typhus-like rickettsia found in cat fleas. Proc Natl Acad Sci U S A. 1992;89:43–6. DOI: 10.1073/pnas.89.1.43
5. Higgins JA, Radulovic S, Schrieffer ME, Azad AF. *Rickettsia felis*: a new species of pathogenic rickettsia isolated from cat fleas. J Clin Microbiol. 1996;34:671–4.
6. Bouyer DH, Stenos J, Crocquet-Valdes P, Moron CG, Popov VL, Zavala-Velazquez JE, et al. *Rickettsia felis*: molecular characterization of a new member of the spotted fever group. Int J Syst Evol Microbiol. 2001;51:339–47.
7. La Scola B, Meconi S, Fenollar F, Rolain JM, Roux V, Raoult D. Emended description of *Rickettsia felis* (Bouyer et al. 2001), a temperature-dependent cultured bacterium. Int J Syst Evol Microbiol. 2002;52:2035–41. DOI: 10.1099/ijs.0.02070-0
8. Ogata H, Renesto P, Audic S, Robert C, Blanc G, Fournier PE, et al. The genome sequence of *Rickettsia felis* identifies the first putative conjugative plasmid in an obligate intracellular parasite. PLoS Biol. 2005;3:e248. DOI: 10.1371/journal.pbio.0030248
9. Williams SG, Sacci JB Jr, Schrieffer ME, Andersen EM, Fujioka KK, Sorvillo FJ, et al. Typhus and typhuslike rickettsiae associated with opossums and their fleas in Los Angeles County, California. J Clin Microbiol. 1992;30:1758–62.
10. Oliveira RP, Galvão MA, Mafra CL, Chamone CB, Calic SB, Silva SU, et al. *Rickettsia felis* in *Ctenocephalides* spp. fleas, Brazil. Emerg Infect Dis. 2002;8:317–9.
11. Zavala-Velazquez JE, Zavala-Castro JE, Vado-Solis I, Ruiz-Sosa JA, Moron CG, Bouyer DH, et al. Identification of *Ctenocephalides felis* fleas as a host of *Rickettsia felis*, the agent of a spotted fever rickettsiosis in Yucatan, Mexico. Vector Borne Zoonotic Dis. 2002;2:69–75. DOI: 10.1089/153036602321131869
12. Marquiez FJ, Muniain MA, Perez JM, Pachon J. Presence of *Rickettsia felis* in the cat flea from southwestern Europe. Emerg Infect Dis. 2002;8:89–91.
13. Horta MC, Chiebao DP, de Souza DB, Ferreira F, Pinheiro SR, Labruna MB, et al. Prevalence of *Rickettsia felis* in the fleas *Ctenocephalides felis* and *Ctenocephalides canis* from two Indian villages in São Paulo Municipality, Brazil. Ann N Y Acad Sci. 2006;1078:361–3. DOI: 10.1196/annals.1374.071
14. Venzal JM, Perez-Martinez L, Felix ML, Portillo A, Blanco JR, Oteo JA. Prevalence of *Rickettsia felis* in *Ctenocephalides felis* and *Ctenocephalides canis* from Uruguay. Ann N Y Acad Sci. 2006;1078:305–8. DOI: 10.1196/annals.1374.056
15. Bitam I, Parola P, De La Cruz KD, Matsumoto K, Baziz B, Rolain JM, et al. First molecular detection of *Rickettsia felis* in fleas from Algeria. Am J Trop Med Hyg. 2006;74:532–5.
16. Stevenson HL, Labruna MB, Monteneri JA, Kosoy MY, Gage KL, Walker DH. Detection of *Rickettsia felis* in a New World flea species, *Anomiopsyllus nudata* (Siphonaptera: Ctenophthalmidae). J Med Entomol. 2005;42:163–7. DOI: 10.1603/0022-2585(2005)042[0163:DORFIA]2.0.CO;2
17. De Sousa R, Edouard-Fournier P, Santos-Silva M, Amaro F, Bacellar F, Raoult D. Molecular detection of *Rickettsia felis*, *Rickettsia typhi* and two genotypes closely related to *Bartonella elizabethae*. Am J Trop Med Hyg. 2006;75:727–31.
18. Jiang J, Soeatmadji DW, Henry KM, Ratiwayanto S, Bangs MJ, Richards AL. *Rickettsia felis* in *Xenopsylla cheopis*, Java, Indonesia. Emerg Infect Dis. 2006;12:1281–3.
19. Ishikura M, Ando S, Shinagawa Y, Matsuura K, Hasegawa S, Nakayama T, et al. Phylogenetic analysis of spotted fever group rickettsiae based on *gltA*, 17-kDa, and *rOmpA* genes amplified by nested PCR from ticks in Japan. Microbiol Immunol. 2003;47:823–32.
20. Cardoso LD, Freitas RN, Mafra CL, Neves CV, Figueira FC, Labruna MB, et al. Characterization of *Rickettsia* spp. circulating in a silent peri-urban focus for Brazilian spotted fever in Caratinga, Minas Gerais, Brazil [in Portuguese]. Cad Saude Publica. 2006;22:495–501.
21. Choi YJ, Lee EM, Park JM, Lee KM, Han SH, Kim JK, et al. Molecular detection of various rickettsiae in mites (Acari: Trombiculidae) in southern Jeolla Province, Korea. Microbiol Immunol. 2007;51:307–12.
22. Rolain JM, Franc M, Davoust B, Raoult D. Molecular detection of *Bartonella quintana*, *B. koehlerae*, *B. henselae*, *B. clarridgeiae*, *Rickettsia felis*, and *Wolbachia pipiensis* in cat fleas, France. Emerg Infect Dis. 2003;9:338–42.
23. Kenny MJ, Birtles RJ, Day MJ, Shaw SE. *Rickettsia felis* in the United Kingdom. Emerg Infect Dis. 2003;9:1023–4.
24. Blair PJ, Jiang J, Schoeler GB, Moron C, Anaya E, Cespedes M, et al. Characterization of spotted fever group rickettsiae in flea and tick specimens from northern Peru. J Clin Microbiol. 2004;42:4961–7. DOI: 10.1128/JCM.42.11.4961-4967.2004
25. Kelly P, Rolain JM, Raoult D. Prevalence of human pathogens in cat and dog fleas in New Zealand. N Z Med J. 2005;118:U1754.
26. Rolain JM, Bourry O, Davoust B, Raoult D. *Bartonella quintana* and *Rickettsia felis* in Gabon. Emerg Infect Dis. 2005;11:1742–4.
27. Marie JL, Fournier PE, Rolain JM, Briolant S, Davoust B, Raoult D. Molecular detection of *Bartonella quintana*, *B. elizabethae*, *B. koehlerae*, *B. doshiae*, *B. taylorii*, and *Rickettsia felis* in rodent fleas collected in Kabul, Afghanistan. Am J Trop Med Hyg. 2006;74:436–9.
28. Schloderer D, Owen H, Clark P, Stenos J, Fenwick SG. *Rickettsia felis* in fleas, Western Australia. Emerg Infect Dis. 2006;12:841–3.
29. Bauer O, Baneth G, Eshkol T, Shaw SE, Harris S. Polygenic detection of *Rickettsia felis* in cat fleas (*Ctenocephalides felis*) from Israel. Am J Trop Med Hyg. 2006;74:444–8.
30. Psaroulaki A, Antoniou M, Papaeustathiou A, Tournazos P, Loukaides F, Tselentis Y. First detection of *Rickettsia felis* in *Ctenocephalides felis* fleas parasitizing rats in Cyprus. Am J Trop Med Hyg. 2006;74:120–2.
31. Hawley JR, Shaw SE, Lappin MR. Prevalence of *Rickettsia felis* DNA in the blood of cats and their fleas in the United States. J Feline Med Surg. 2007;9:258–62. DOI: 10.1016/j.jfms.2006.12.005
32. Labruna MB, Ogrowska M, Moraes-Filho J, Lep P, Gallegos JL, Lopez J. *Rickettsia felis* in Chile. Emerg Infect Dis. 2007;13:1794–5.
33. Zavala-Velazquez JE, Ruiz-Sosa JA, Sanchez-Elias RA, Becerra-Carmona G, Walker DH. *Rickettsia felis* rickettsiosis in Yucatan. Lancet. 2000;356:1079–80. DOI: 10.1016/S0140-6736(00)02735-5
34. Galvão MA, Zavala-Velazquez JE, Zavala-Castro JE, Mafra CL, Calic SB, Walker DH. *Rickettsia felis* in the Americas. Ann N Y Acad Sci. 2006;1078:156–8. DOI: 10.1196/annals.1374.027
35. Richter J, Fournier PE, Petridou J, Haussinger D, Raoult D. *Rickettsia felis* infection acquired in Europe and documented by polymerase chain reaction. Emerg Infect Dis. 2002;8:207–8.
36. Parola P, Miller RS, McDaniel P, Telford SR III, Rolain JM, Wongsrichanalai C, et al. Emerging rickettsioses of the Thai-Myanmar border. Emerg Infect Dis. 2003;9:592–5.
37. Choi YJ, Jang WJ, Ryu JS, Lee SH, Park KH, Paik HS, et al. Spotted fever group and typhus group rickettsioses in humans, South Korea. Emerg Infect Dis. 2005;11:237–44.
38. Znazen A, Rolain JM, Hammami A, Jemaa MB, Raoult D. *Rickettsia felis* infection, Tunisia. Emerg Infect Dis. 2006;12:138–40.
39. Phongmany S, Rolain JM, Phetsouvanh R, Blacksell SD, Soukhaemum V, Rasachack B, et al. Rickettsial infections and fever, Vientiane, Laos. Emerg Infect Dis. 2006;12:256–62.