

4 その他の患者状態像(医療区分採用項目以外)

■疾患(複数回答) [図表8]

| | 平成20年度調査 | | | | | | | |
|----------------|------------------|-------|------------------|-------|-----------------|-------|----------------|-------|
| | 医療区分1 (n=361) | | 医療区分2 (n=385) | | 医療区分3 (n=51) | | 全 体 (n=797) | |
| 糖尿病 | 39 | 10.8% | 92 | 23.9% | 10 | 19.6% | 141 | 17.7% |
| 不整脈 | 23 | 6.4% | 29 | 7.5% | 5 | 9.8% | 57 | 7.2% |
| 急性心不全 | 2 | 0.6% | 2 | 0.5% | 1 | 2.0% | 5 | 0.6% |
| 慢性心不全 | 33 | 9.1% | 26 | 6.8% | 10 | 19.6% | 69 | 8.7% |
| 高血圧症 | 143 | 39.6% | 144 | 37.4% | 14 | 27.5% | 301 | 37.8% |
| 虚血性心疾患 | 39 | 10.8% | 39 | 10.1% | 7 | 13.7% | 85 | 10.7% |
| 脊髄損傷 | 0 | 0.0% | 6 | 1.6% | 0 | 0.0% | 6 | 0.8% |
| 大腿骨頸部骨折 | 24 | 6.6% | 22 | 5.7% | 5 | 9.8% | 51 | 6.4% |
| 脊椎圧迫骨折 | 33 | 9.1% | 32 | 8.3% | 1 | 2.0% | 66 | 8.3% |
| その他の骨折 | 19 | 5.3% | 26 | 6.8% | 1 | 2.0% | 46 | 5.8% |
| 関節リウマチ | 15 | 4.2% | 9 | 2.3% | 1 | 2.0% | 25 | 3.1% |
| アルツハイマー病 | 26 | 7.2% | 43 | 11.2% | 3 | 5.9% | 72 | 9.0% |
| アルツハイマー病以外の認知症 | 33 | 9.1% | 66 | 17.1% | 7 | 13.7% | 106 | 13.3% |
| 失語症 | 13 | 3.6% | 17 | 4.4% | 5 | 9.8% | 35 | 4.4% |
| 脳性麻痺 | 1 | 0.3% | 0 | 0.0% | 0 | 0.0% | 1 | 0.1% |
| 脳梗塞 | 70 | 19.4% | 91 | 23.6% | 17 | 33.3% | 178 | 22.3% |
| 脳出血 | 10 | 2.8% | 26 | 6.8% | 3 | 5.9% | 39 | 4.9% |
| 片側不全麻痺 | 26 | 7.2% | 35 | 9.1% | 3 | 5.9% | 64 | 8.0% |
| 四肢麻痺 | 8 | 2.2% | 19 | 4.9% | 7 | 13.7% | 34 | 4.3% |
| 仮性球麻痺 | 2 | 0.6% | 4 | 1.0% | 0 | 0.0% | 6 | 0.8% |
| 気管支喘息 | 13 | 3.6% | 9 | 2.3% | 2 | 3.9% | 24 | 3.0% |
| 腸閉塞(イレウス) | 3 | 0.8% | 3 | 0.8% | 0 | 0.0% | 6 | 0.8% |
| 肝不全 | 4 | 1.1% | 2 | 0.5% | 0 | 0.0% | 6 | 0.8% |
| がん(悪性腫瘍) | 16 | 4.4% | 20 | 5.2% | 8 | 15.7% | 44 | 5.5% |
| 腎不全 | 14 | 3.9% | 86 | 22.3% | 7 | 13.7% | 107 | 13.4% |