



図20.昨シーズンとの比較

2006/2007





2007/2008




31

図21.タミフル（リン酸オセルタミビル）服用の有無の性別




32

図22.タミフル（リン酸オセルタミビル）服用の有無の年齢別



33


図23.シンメトレル（塩酸アマンタジン）服用の有無n=77




34

図24.昨シーズンとの比較

2006／2007




2007／2008



35


図25.リレンザ（ザナミビル）使用の有無n=77




36

図26.昨シーズンとの比較

2006/2007




2007/2008



37


図27.アセトアミノフェン服用の有無n=77




38

図28.昨シーズンとの比較

2006／2007




2007／2008



39


図29.異常行動と睡眠の関係n=77




40

図30.昨シーズンとの比較

2006/2007




2007/2008



41

図31.異常行動の分類（複数回答）





42

図32.昨シーズンとの比較

2006/2007


2007/2008




43

図33.タミフル有無と異常行動と睡眠の関係

タミフル服用有り群



タミフル服用無し群




44

突然走り出す・飛び降りのみ の 分析

45

図34.患者の年齢n=41




平均値9.21
中央値7

46


図35.昨シーズンとの比較

2006/2007



平均値9.93
中央値10


2007/2008



平均値9.21
中央値7

47

図36.患者の性別n=41





48

図37.昨シーズンとの比較


2006/2007

2007/2008



49

図38.最高体温n=41



平均値39.34 50
中央値39.2