

グリシドール脂肪酸エステル毒性関連文献

- GE-1 Hendry JA, Homer RF, Rose FL, Walpole AL, 1951, Cytotoxic agents: II. Bis-epoxides and related compounds. *Br J Pharmacol Chemother*, 6:235-255
- GE-2 Arthur L Walpole, 1958, Carcinogenic action of alkylating agents. *Annals New York Academy of Sciences*.
- GE-3 Weil OS, Condra N, Halun C, Striegel JA, 1963, Experimental carcinogenicity and acute toxicity of representative epoxides. *Am Ind Hyg Assoc J* 24 :305-25
- GE-4 Swern D, Wieder R, McDonough M, Meranze DR, Shimkin MB, 1970, Investigation of fatty acids and derivatives for carcinogenic activity. *Cancer Res* 30 :1037-1046
- GE-5 Swern D, Wieder R, McDonough M, Meranze DR, Shimkin MB, 1972, Replication of low-level carcinogenic activity bioassays. *Cancer Res* 32 :880-881
- GE-6 Canter DA, Zeiger E, Haworth S, Lawlor T, Mortelmans K, Speck W, 1986, Comparative mutagenicity of aliphatic epoxides in *Salmonella* *Mutation Res.* 172:105-138
- GE-7 IARC, 1987, Glycidyl oleate. *IARC Monogr Eval Carcinog Risk Chem Man* 11 :183-186
- GE-8 IARC, 1987, Glycidyl stearate. *IARC Monogr Eval Carcinog Risk Chem Man* 11 :187-190
- GE-9 Boogaard PJ, van Elburg PA, de Kloe KP, Watson WP, van Sittert NJ, 1999, Metabolic inactivation of 2-oxiranylmethyl 2-ethyl-2,5-dimethylhexanoate (C10GE) in skin, lung and liver of human, rat and mouse. *Xenobiotica* 29:987-1006

グリシドール毒性関連文献

- G-1 NTP, 1990, NTP Toxicology and carcinogenesis studies of glycidol (CAS No. 556-52-5) in F344/N rats and B6C3F1 mice (gavage studies). Natl Toxicol Program Tech Rep Ser 374:1-229
- G-2 IARC, 2000, Glycidol. IARC Monogr Eval Carcinog Risks Hum 77:469-486
- G-3 Nomeir AA, Silveira DM, Ferrala NF, Markham PM, McComish MF, Ghanayem BI, Chadwick M, 1995, Comparative disposition of 2,3-epoxy-1-propanol (glycidol) in rats following oral and intravenous administration. J Toxicol Environ Health 44:203-217
- G-4 Irwin RD, Eustis SL, Stefanski S, Haseman JK, 1996, Carcinogenicity of glycidol in F344 rats and B6C3F1 mice. J Applied Toxicol 16:201-209
- G-5 Landin HH, Tareke E, Rydberg P, Olsson U, Tornqvist M, 2000, Heating of food and hemoglobin adducts from carcinogens: possible precursor role of glycidol. Food Chem Toxicol 38:963-969
- G-6 Chen Y, Megosh LC, Gilmour SK, Sawicki JA, O'Brien TG, 2000, K6/ODC transgenic mice as a sensitive model for carcinogen identification. Tox. Letters 116: 27-35
- G-7 Guo TL, McCay JA, Brown RD, Musgrove DL, Butterworth L, Munson AE, Germolec DR, White KLJ, 2000, Glycidol modulation of the immune responses in female B6C3F1 mice. Drug Chem Toxicol. 23(3):433-57
- G-8 日本バイオアッセイセンター, 2004, グリシドールの吸入によるがん原性試験結果の概要。日本バイオアッセイセンター, ホームページ
- G-9 Kim JH, Kim KY, Kwon KJ, Go SY, Min KN, Lee, WS, Park SN, Sheen YY, 2006, Genetic toxicity test of glycidol by Ames, micronucleus, comet assays and microarray analysis. J. Applied Pharmacol. 14: 240-245
- G-10 Ramy RE, Ould Elhkim M, Lezmi S, Poul JM, 2007, Evaluation of the genotoxic potential of 3-monochloropropane-1,2-diol (3-MCPD) and its metabolites, glycidol and beta-chlorolactic acid, using the single cell gel/comet assay. Food Chem Toxicol 45:41-48
- G-11 NTP, 2007, Toxicology and carcinogenesis study of glycidol (CAS No. 556-52-5) in genetically modified haploinsufficient p16(Ink4a)/p19(Arf) mice (gavage study). Natl Toxicol Program Genet Modif Model Rep. 2007 Nov;(13):1-81.



