

<本評価書中で使用した略号>

BMI	ボディマス指数
CC16S	クララ細胞蛋白質
Cd	カドミウム
Cd-B	血液中カドミウム量
Cd-F	糞中カドミウム量
Cd-I	摂取カドミウム量
Cd-MT	カドミウム-メタロチオネイン
Cd-U	尿中カドミウム量
Cd/Zn	カドミウム/亜鉛
CI	信頼区間
DMT1	2価金属イオン輸送体1
EC	欧州委員会
EFSA	欧州食品安全機関
FEV1	一秒量
FVC	努力性呼気肺活量
IARC	国際がん研究機構
JECFA	FAO/WHO合同食品添加物専門家会議
ML	最大レベル
MT	メタロチオネイン
MTP1	金属輸送蛋白質1
NAG	N-アセチル-β-D-グルコサミニダーゼ
NHANES	国民健康栄養調査
Ni-Cd	ニッケル-カドミウム
NOAEL	無毒性量
PTWI	暫定耐容週間摂取量
RBP	レチノール結合蛋白質
RfD	参照用量
SIR	標準化罹患比
SMR	標準化死亡比
TDS	トータルダイエットスタディ
TWI	耐容週間摂取量
U.S EPA	米国環境保護庁
WHO	世界保健機関
α1-MG	α1-ミクログロブリン
β2-MG	β2-ミクログロブリン
%FEV1	一秒率
%TRP	尿細管リン再吸収率

<引用文献>

1. 物理、化学的特性
 - 1 - 1 大木道則, 大沢利昭, 田中元治, 千原秀昭偏, 化学大辞典 第1版. 第6刷, 株式会社 東京化学同人, 2001, pp453-454.
2. 採鉱、精練及び用途
 - 2 - 1 Wilson B., Investigation of trace metals in the aqueous environment: Final report(January 1986-December 1987), Houston, Texas Southern University, 1988^a, p.28(Report No.DOE/CH/10255-T1, prepared for the US Department of Energy, Washington).
3. 分布、変化
 - 3 - 1 GESAMP, IMO/FAO/UNESCO/WMO/IAEA/UN/UNEP Joint Group of Experts on the Scientific Aspects of Marine Population: Report of the Fourteenth Session, Vienna,26-30 March, 1984, Vienna, International Atomic Energy Agency(Reports and Studies No.21).
 - 3 - 2 GESAMP, IMO/FAO/UNESCO/WMO/IAEA/UN/UNEP Joint Group of Experts on the Scientific Aspects of Marine Population: Report of the Seventeenth Session, Rome, Geneva, World Health Organization, 1987 (Reports and Studies No.31).
 - 3 - 3 Nriagu J.O., Global inventory of natural and anthropogenic emissions of trace metals to the atmosphere.Nature(Lond.), 1979; 279: 409-411.
 - 3 - 4 Boyle E.A., Sclater F., Edmond J.M., On the marine geochemistry of cadmium. Nature(Lond.), 1976; 263: 42-44.
 - 3 - 5 Martin J.H., Broenkow W.W., Cadmium in plankton:elevated concentrations off Baja California. Science, 1975; 190: 884-885.
 - 3 - 6 Simpson W.R., A critical review of cadmium in the marine environment. Prog. Oceanog., 1981; 10: 1-70.
 - 3 - 7 Förstner U., Cadmium in the environment, Part I In: Nriagu,J.O., ed. Cadmium in polluted sediments, New York, Chichester, John Wiley & Sons, 1980 ; 305-363.
 - 3 - 8 Sangster B., De Groot G., Loeber J.G., Derks H.J.G.M., Krajnc E.I., Savelkoul T.J.F. Urinary excretion of cadmium, protein, beta-2-microglobulin and glucose in individuals living in a cadmium-polluted area. Hum. Toxicol., 1984; 3: 7-21.
 - 3 - 9 Yamagata N., Shigematsu I., Cadmium pollution in perspective. Bull. Inst. Public Health (Tokyo) , 1970; 19: 18-24.
 - 3 - 10 Alloway B.J., Thornton I., Smart G.A., Sherlock J.C., Quinn M.J., Metal availability. Sci. Total Environ, 1988; 75: 41-69.
 - 3 - 11 Lund L.J., Betty E.E., Page A.L., Elliott R.A. Occurrence of naturally high cadmium levels in soils and its accumulation by vegetation. J.environ.Qual., 1981; 10: 551-556.
 - 3 - 12 Davis R.D., Coker E.G. Cadmium in agriculture, with special reference to the utilization of sewage sludge on land, Medmenham, United Kingdom, Water Reseach Centre (Technical Report TR/139) , 1980.
 - 3 - 13 Bryan G.W., Langston W.J., Hummerstone L.G., The use of biological indicators of heavy-metal contamination in estuaries with special reference to an assessment of the biological availability of metals in estuarine sediments from south-west Britain, Citadal Hill, Devon, Marine

- Biological Association of the United Kingdom, 1980; pp73 (Occasional Publication No.1) .
- 3 - 14 Nielsen S.A., Cadmium in New Zealand dredge oysters: geographic distribution. *Int. J. environ. Anal. Chem.*, 1975; 4: 1-7.
- 3 - 15 Buchet J.P., Lauwerys R., Vandevorde A., Pycke J.M., Oral daily intake of cadmium, lead, manganese, copper, chromium, mercury, calcium, zinc and arsenic in Bergium. *Food chem. Toxicol.*, 1983; 21: 19-24.
- 3 - 16 Martin J.H., Elliott P.D., Anderlini V.C., Girvin D., Jacobs S.A., Risebrough R.W., Delong R.L., Gilmartin W.G., Mercury - selenium-bromine imbalance in premature parturient California sea lions. *Mar. Biol.*, 1976; 35: 91-104.
- 3 - 17 Stoneburner D.L., Heavy metals in tissues of stranded short-finned pilot whales. *Sci. Total Environ.*, 1978; 9: 293-297.
- 3 - 18 Nicolson J.K., Osborn D., Kidney lesions in pelagic seabirds with high tissue levels of cadmium and mercury. *J.Zool.Lond.*, 1983; 200: 88-118.
- 3 - 19 MARC, Biological monitoring of environmental contaminants (plants) , London, Monitoring and Assessment Research Centre, Chelsea College, University of London, 1986; pp247 (MARC Report Number 32) .
4. ヒトへの曝露経路と曝露量
- 4 - 1 Peplow D., Edmonds R., Health risks associated with contamination of groundwater by abandoned mines near Twisp in Okanogan County, Washington, USA. *Environ. Geochem. Health.* 2004; 26: 69-79.
- 4 - 2 Lee J.S., Chon H.T., Kim K.W., Human risk assessment of As, Cd, Cu and Zn in the abandoned metal mine site. *Environ Geochem Health.* 2005; 27: 185-191.
- 4 - 3 農林水産省 (2002) , 農作物等に含まれるカドミウムの実態調査について
- 4 - 4 水産庁 (2003) , 水産物に含まれるカドミウムの実態調査について
- 4 - 5 農林水産省 (2007) , 食品中のカドミウムに関する情報, 2.日本のコメに含まれるカドミウム. <http://www.maff.go.jp/cd/html/A12.htm>.
- 4 - 6 Wolink K.A., Fricke F.L., Caper S.G., Braude G.L., Meyer M.W., Satzger R.D., Bonnin E., Elements in major raw agricultural crops in the United States. 1. Cadmium and lead in lettuce, peanuts, potatoes, soybeans, sweet corn, and wheat. *J. Agric. Food Chem.* 1983; 31: 1240-1244.
- 4 - 7 Wolink K.A., Fricke F.L., Caper S.G., Meyer M.W., Satzger R.D., Bonnin E., Gaston C.M., Elements in major raw agricultural crops in the United States. 3. Cadmium, lead, and eleven other elements in carrots, field corn, onion, rice, spinach, and tomatoes. *J. Agric. Food Chem.* 1985; 33: 807-811.
- 4 - 8 Bucke D., Norton M.G., Rolfe, M.S., Field assessment of effects of dumping wastes at sea: II. Epidermal lesions and abnormalities of fish in the outer Thames estuary, London, Ministry of Agriculture, Fisheries and Food, 1983; pp29 (Technical Report No.72) .
- 4 - 9 Varo P., Nuurtamo M., Saari E., Koivistoinen P., Mineral element composition of Finnish Foods. *Acta Agric. Scand.* 1980; 22: Suppl: 127-139.
- 4 - 10 Jorhem L., Mattson P., Slorach S., Lead, cadmium, zinc and certain other metals in foods on the Swedish market. *Vår Föda*, 1984; 36: Suppl. 3.
- 4 - 11 Andersen A., [Lead, cadmium, copper and zinc in the Danish diet], Copenhagen, Statens Levnedsmedelinstytut, 1979; pp89 (Report No.40) (in Danish) .

- 4 - 12 RIVM (1988) In: Ros J.P.M., Sloof W., ed. Integrated criteria document cadmium, Bilthoven, The Netherlands, National Institute of Public Health and Environmental Protection (RIVM-Report No.758476004) .
- 4 - 13 WHO(1992), Environmental Health Criteria 134 Cadmium. Geneva.
- 4 - 14 Friberg L., Piscator M., Nordberg G., Kjellstrom T., Cadmium in the environment, 2nd ed., Cleveland(OH): CRC Press,1974.
- 4 - 15 Elinder C.G., Kjellström T., Friberg L., Lind B., Linnman L., Cadmium in kidney cortex, liver, and pancreas from swedish autopsies. Arch. Environ. Health. 1976; 31: 292-302.
- 4 - 16 Elinder C.G., Kjellström T., Lind B., Linnman L., Piscator M., Sundstedt K., Cadmium exposure from smoking cigaretters: variations with time and country where purchased. Environ Res 1983; 32: 220-227.
- 4 - 17 Friberg L., Vanter M., Assessment of exposure to lead and cadmium through biological monitoring: results of a UNEP/WHO global study. Environ Res 1983; 30: 95-128.
- 4 - 18 Bensryd I., Rylander L., Högstedt B., Aprea P., Bratt I., Fahraeus C., et al., Effect of acid precipitation on retention and excretion of elements in man. Sci .Total. Environ. 1994; 145: 81-102.
- 4 - 19 Nilsson U., Schutz A, Skerfving S, Mattsson S., Cadmium in kidneys in Swedes measured in vivo using X-ray fluorescence analysis. Int. Arch. Occup. Environ. Health. 1995; 67: 405-11.
- 4 - 20 国立医薬品食品衛生研究所食品部 (2000) , 日本におけるトータルダイエツト調査 (食品汚染物の 1 日摂取量) 1977~1999 年度.
- 4 - 21 松田りえ子, 食品中の有害物質等の摂取量の調査及び評価に関する研究, 平成 19 年度総括・分担研究報告書.
- 4 - 22 新田裕史, 日本人のカドミウム曝露量推計に関する研究. 厚生労働科学研究費補助金 (特別研究事業) 総括研究報告書 平成 16 年 3 月.
- 4 - 23 櫻井治彦, 池田正之, 香山不二雄, 大前和幸, 食品中に残留するカドミウムの健康影響評価について. 平成 15 年度 総括・分担研究報告書 厚生労働省. 2004 ; 66-112.

5. ヒトにおける動態及び代謝

- 5 - 1 Horiguchi H., Oguma E., Sasaki S., Miyamoto K., Ikeda Y., Machida M., Kayama F., Comprehensive study of the effects of age, iron deficiency, diabetes mellitus, and cadmium burden on dietary cadmium absorption in cadmium-exposed female Japanese farmers. Toxicol. Appl. Pharmacol, 2004; 196: 114-23.
- 5 - 2 Bunker V. W., Lawson M. S., Delves H. T., Delves H.T., Clayton B. E., The intake and excretion of lead and cadmium by the elderly¹⁻³. Am. J. Clin. Nutr. 1984 ; 39: 803-808.
- 5 - 3 Vanderpool R. A., Reeves P. G., Cadmium absorption in women fed processed edible sunflower kernels labeled with a stable isotope of cadmium, ¹¹³Cd. Environ. Research Section A 2001 ; 87: 69-80.
- 5 - 4 Suzuki S., Lu C. C., A balance study of cadmium - An estimation of daily input, output and retained amount in two subjects. Industrial Health. 1976 ; 14 : 53-65.
- 5 - 5 Flanagan P. R., McLellan J. S., Haist J., Cherian G., Chamberlain M. J., Valberg L. S., Increased dietary cadmium absorption in mice and human subjects with iron deficiency. Gastroenterology 1978 ; 74 : 841-846.
- 5 - 6 McLellan J. S., Flanagan P. R., Chamberlain M. J., Velberg L. S.,

- Measurement of dietary cadmium absorption in humans. *J. Toxicol. Environ. Health*, 1978 ; 4 : 131-138.
- 5 - 7 Newton D., Johnson P., Lally A. E., Pentreath R. J., Swift D. J., The uptake by man of cadmium ingested in crab meat. *Human Toxicol.* 1984 ; 3 : 23-28.
- 5 - 8 Berglund M., Akesson A., Nermell B., Vahter M., Intestinal absorption of dietary cadmium in women depends on body iron stores and fiber intake. *Environ. Health Perspect.* 1994 ; 102 : 1058-1066.
- 5 - 9 Vahter M., Berglund M., Nermell B., Akesson A., Bioavailability of cadmium from shellfish and mixed diet in women. *Toxicol. Appl. Pharmacol.* 1996 : 136 : 332-341.
- 5 - 10 Crews H. M., Owen L. M., Langfoed N., Fairweather-Tait S. J., Fox T. E., Hubbard L., Phillips D., Use of the stable isotope ¹⁰⁶Cd for studying dietary cadmium absorption in humans. *Toxicol. Lett* 112-113. 2000: 201-207.
- 5 - 11 Kikuchi Y., Nomiyama T., Kumagai N., Dekio F., Uemura T., Takebayashi T., Nishiwaki Y., Matsumoto Y., Sano Y., Hosoda K., Watanabe S., Sakurai H., Omae K., Uptake of cadmium in meals from the digestive tract of young non-smoking Japanese female volunteers. *Journal of Occupation Health* 2003 ; 45 : 43-52.
- 5 - 12 Zalups R. K., Ahmad S., Molecular handling of cadmium in transporting epithelia. *Toxicol. Appl. Pharmacol.* 2003; 186: 163-88. Review.
- 5 - 13 小泉直子, カドミウムの生体内動態に関する基礎的研究. *日本衛生学会誌*, 昭和 50 年; 第 30 卷 第 2 号 別冊; 300-324.
- 5 - 14 Elinder C.G., Normal values for cadmium in human tissues, blood, and urine in different countries. In: Friberg L., Elinder C.G., Kjellström T., Nordberg G.F. eds., *Cadmium and health: A toxicological and epidemiological appraisal*. vol I. CRC Press, Inc. Boca Raton, Florida. 1985; 81-102.
- 5 - 15 Yoshida M., Ohta H., Yamauchi Y., Seki Y., Sagi M., Yamazaki K., Sumi Y., Age-dependent changes in metallothionein levels in liver and kidney of the Japanese. *Biological Trace Element Research* 1998 ; 63 : 167-175.
- 5 - 16 Orłowski C., Piotrowski J.K. Biological levels of cadmium and zinc in the small intestine of non-occupationally exposed human subjects. *Human & Experimental Toxicology* 2003 ; 22: 57-63.
- 5 - 17 Satarug S., Baker J.R., Reilly P. B., Moore M. R., Williams D. J., Cadmium levels in the lung, liver, kidney cortex, and urine samples from australians without occupational exposure to metals. *Archives of Environmental Health* 2002 ; 57: 69-77.
- 5 - 18 Garcia F., Ortega A., Domingo J. L., Corbella J., Accumulation of metals in autopsy tissues of subjects living in Tarragona County, Spain. *J. Environ. Sci. Health* 2001 ; A36(9) : 1767-1786.
- 5 - 19 Torra M., To-Figueras J., Rodamilans M., Brunet M., Corbella J., Cadmium and zinc relationships in the liver and kidney of humans exposed to environmental cadmium. *Sci. Total Environ.* 1995 ; 170 : 53-57.
- 5 - 20 Tiran B., Karpf E., Tiran A., Age dependency of selenium and cadmium content in human liver, kidney, and thyroid. *Arch. Environ. Health* 1995; 50: 242-246.
- 5 - 21 Takacs S., Tatar., Trace Elements in the Environment and in Human Organs: Analysis according to domicile and sex. *Z.gesamte Hyg.* 1991; 37: 53-55.
- 5 - 22 Nogawa K., Honda R., Yamada Y., Kido T., Tsuritani I., Ishizaki M.,

- Yamaya H., Critical concentration of cadmium in kidney cortex of humans exposed to environmental cadmium. *Environmental Research*. 1986; 40: 251-260.
- 5 - 23 Sumino K., Hayakawa K., Shibata T., Kitamura S., Heavy metals in normal Japanese tissues. *Arch Environ Health*. 1975; 30: 487-494.
- 5 - 24 Tsuchiya K., Cadmium in human urine, feces, blood, hair, organs, and tissues. In: Tsuchiya K. ed., *Cadmium studies in Japan: A review*. Kodansha Ltd. 1978; pp37-43.
- 5 - 25 Tati M., Katagiri Y., Kawai M., Urinary and fecal excretion of cadmium in normal Japanese: An approach to non-toxic levels of cadmium."In: *Effects and Dose-Response Relationships of Toxic Metals*.(G.F. Nordberg, ed.)Amsterdam: Elsevier Scientific Publishing Co. 1976; pp331-342.
- 5 - 26 Tsuchiya K., Proteinuria of cadmium workers. *J. Occup. Med*. 1976; 18: 463-466.
- 5 - 27 Sugita M., Tsuchiya K., Estimation of variation among individuals of biological half-time of cadmium calculated from accumulation data. *Environmental Research* 1995; 68: 31-37.
- 5 - 28 Kjellström T., Nordberg G. F., A kinetic model of cadmium metabolism in the human being. *Environ. Res*. 1978; 16: 248-269.
- 5 - 29 Nomiyama T., Kikuchi Y., Kumagai N., Dekio F., Uemura T., Hosoda K., Sakurai H., Omae K., Short-term in cadmium in feces, blood and urine after dietary cadmium intake in young Japanese female. *J.Occup. Health* 2002; 44: 429-432.

6. ヒトにおける有害性評価

6.1 急性影響

- 6.1.1 - 1 日本産業衛生学会, 許容濃度等の勧告(2002 年度). 産衛誌 2002; 44: 140-164.

6.2 慢性影響

6.2.1 腎臓への影響

- 6.2.1 - 1 Friberg L., Health hazards in the manufacture of alkaline accumulators with special reference to chronic cadmium poisoning. *Acta Med. Scand*. 1950; Suppl. 240: 1-124.
- 6.2.1 - 2 Adams R.G., Harrison J.F., Scott P., The development of cadmium-induced proteinuria, impaired renal function, and osteomalacia in alkaline battery workers. *Q. J. Med*. 1969; 38: 425-443.
- 6.2.1 - 3 Kazantzis G., Renal tubular dysfunction and abnormalities of calcium metabolism in cadmium workers. *Environ. Health Perspect*. 1979; 28: 155-159.
- 6.2.1 - 4 村田 勇, イタイイタイ病の研究. 日本医師会雑誌, 1971; 65: 15-42.
- 6.2.1 - 5 武内重五郎, 中本 安, イタイイタイ病. 現代内科学大系 1969 年刊追補, 中山書店, 1969; pp366-394.
- 6.2.1 - 6 Aoshima K., Environmental cadmium pollution and its health effects on inhabitants in Japan. Jinzu River basin: Clinical findings in Itai-itai disease. *In Advances in the Prevention of Environmental Cadmium Pollution and Countermeasures*, 13-19, Nogawa K., Kurachi M., Kasuya M. (Eds.), Eiko Laboratory, Kanazawa, 1999.
- 6.2.1 - 7 Järup L., Berglund M., Elinder C.G., Nordberg G., Vahter M., Health

- effects of cadmium exposure—a review of the literature and a risk estimate. *Scand J Work Environ Health*. 1998; 24: Suppl 1:1-51. (訂正稿: *Scand J Work Environ Health* 1998:240)
- 6.2.1 - 8 Tsukahara T., Ezaki T., Moriguchi J., Furuki K., Fukui Y., Ukai H., Okamoto S., Sakurai H., Ikeda M., No significant effect of iron deficiency on cadmium body burden or kidney dysfunction among women in the general population in Japan. *Int Arch Occup Environ Health*, 2003; 76: 275-281.
- 6.2.1 - 9 de Burbure C., Buchet J.P., Leroyer A., Nisse C., Haguenoer J.M., Mutti A., Smerhovsky Z., Cikrt M., Trzcinka-Ochocka M., Razniewska G., Jakubowski M., Bernard A., Renal and neurologic effects of cadmium, lead, mercury, and arsenic in children: evidence of early effects and multiple interactions at environmental exposure levels. *Environ Health Perspect*. 2006; 114: 584-590.
- 6.2.2 カドミウム土壤汚染地域住民における影響
- 6.2.2 - 1 斎藤 寛, 中野篤浩, カドミウム環境汚染と人間の健康. —生態中毒学的アプローチ—“エコトキシコロジー” (大井 玄, 鈴木継美, 井村伸正編), 篠原出版, 1983, pp119-126.
- 6.2.2 - 2 「カドミウムによる土壤汚染地域住民健康調査」検討委員会, カドミウムによる環境汚染地域住民健康調査. 環境保健レポート, 1989; 56: 69-345.
- 6.2.2 - 3 カドミウム汚染地域住民健康影響調査検討会報告書. 平成 14 年 3 月.
- 6.2.2 - 4 神通川流域住民健康調査検討会報告書. 平成 15 年 7 月, 富山県厚生部健康課.
- 6.2.2 - 5 Tohyama C., Shaikh Z.A., Nogawa K., Kobayashi E., Honda R., Urinary metallothionein as a new index of renal dysfunction in "Itai-itai" disease patients and other Japanese women environmentally exposed to cadmium. *Arch. Toxicol*, 1982; 50: 159-166.
- 6.2.2 - 6 樊 建軍, 青島恵子, 加藤輝隆, 寺西秀豊, 加須屋 実, 富山県神通川流域カドミウム環境汚染地域住民の尿細管障害に関する追跡研究 第 1 報 土壤汚染改良事業開始後のカドミウム曝露の変化と尿細管障害の予後. *日衛誌* 1998; 53: 545-557.
- 6.2.2 - 7 Cai Y., Aoshima K., Katoh T., Teranishi H., Kasuya M., Renal tubular dysfunction in male inhabitants of a cadmium-polluted area in Toyama, Japan — an eleven-year follow-up study. *J. Epidemiol*. 2001; 11: 180-189.
- 6.2.2 - 8 Kido T., Honda R., Tsuritani I., Yamaya H., Ishizaki M., Yamada Y., Nogawa K., Progress of renal dysfunction in inhabitants environmentally exposed to cadmium. *Arch. Environ. Health*. 1988; 43: 213-217.
- 6.2.2 - 9 Iwata K., Saito H., Moriyama M., Nakano A., Renal tubular function after reduction of environmental cadmium exposure: a ten-year follow-up. *Arch. Environ. Health*. 1993; 48: 157-163.
- 6.2.2 - 10 生野鉦山周辺地域カドミウム汚染総合調査班報告書; 昭和 47 年 4 月, 1972.
- 6.2.2 - 11 喜田村正次, 小泉直子, 幡山文一, 地域住民の尿中 β_2 -microglobulin 濃度に関する疫学的研究, 食品に含まれるカドミウムの安全性に関する研究, 昭和 52 年度食品衛生調査研究報告書, 1977.
- 6.2.2 - 12 Ellis KJ., Yasumura S., Vartsky D., Cohn SH., Evaluation of biological indicators of body burden of cadmium in humans. *Fundamental & Applied Toxicol*. 1983; 3: 169-174.

- 6.2.2 - 13 Kjellström T., Exposure and accumulation of cadmium in populations from Japan, the United States, and Sweden. *Environ. Health Perspect.* 1979; 28: 169-197.
- 6.2.2 - 14 小林悦子, 環境中カドミウムの人体影響に関する疫学的研究 (第2報) Cd 汚染地居住期間別尿所見. *日本公衛誌*, 1982; 29: 201-207.
- 6.2.2 - 15 Silensen, Kasiske, Laboratory assessment of kidney disease, *The Kidney*, Ed. B. M. Brenner., 2004; pp1127-1128.
- 6.2.2 - 16 加須屋 実, イタイイタイ病を頂点とするカドミウムの人体影響に関する研究の将来展望“カドミウム環境汚染の予防と対策における進歩と成果” (能川浩二、倉知三夫、加須屋 実編). 栄光ラボラトリ, 1999; pp115-119.
- 6.2.2 - 17 斎藤 寛, 蔀 幸三, 永井謙一, 有川 卓, カドミウム環境汚染による慢性カドミウム中毒の研究—カドミウムによる健康影響の早期診断、ならびにカドミウム負荷量と健康影響発現の間の量・効果関係の検討—. “中毒学と栄養学—その方法論的接点—” (鈴木継美、井村伸正、鈴木庄亮編), 篠原出版, 1978; pp85-99.
- 6.2.3 カルシウム代謝及び骨への影響
- 6.2.3 - 1 青島恵子, 岩田孝吉, 加須屋 実, カドミウム環境汚染による健康影響に関する研究 第2報. 富山県神通川流域カドミウム汚染地住民の血清カルシウム、リン、アルカリホスファターゼ値ならびに骨萎縮度について. *日衛誌*. 1988; 43: 864-871.
- 6.2.3 - 2 青島恵子, 加藤輝隆, 寺西秀豊, 堀口兵剛, 加須屋 実, カドミウム腎症におけるカルシウム・リン・ビタミンD代謝異常—富山県神通川流域カドミウム汚染地域に見いだされた近位尿細管障害 34 例の検討. *日衛誌*. 1993; 47: 1009-1020.
- 6.2.3 - 3 Takebayashi S., Jimi S., Segawa M., Kiyoshi Y., Cadmium induces osteomalacia mediated by proximal tubular atrophy and disturbances of phosphate reabsorption. A study of 11 autopsies. *Pathol. Res. Pract.* 2000; 196: 653-663.
- 6.2.3 - 4 原田孝司, 原 耕平, 緒方時雄, 西村敬一, 重野 哲, 西村 昇, 七條利幸, 鉄塚 眞, 佐藤佐由利, 長崎県対馬カドミウム環境汚染地域における経過観察者の主要検査所見の推移. *環境保健レポート*. 1991; 58: 205-211.
- 6.2.3 - 5 Kido T., Nogawa K., Yamada Y., Honda R., Tsuritani I., Ishizaki M., Yamaya H., Osteopenia in inhabitants with renal dysfunction induced by exposure to environmental cadmium. *Int. Arch. Occup. Environ. Health* 1989; 61: 271-276.
- 6.2.3 - 6 Kido T., Honda R., Tsuritani I., Ishizaki M., Yamada Y., Nakagawa H., Nogawa K., Dohi Y., Serum levels of bone Gla-protein in inhabitants exposed to environmental cadmium. *Arch. Environ. Health*. 1991; 46: 43-49.
- 6.2.3 - 7 城戸照彦, 大道正義, 能川浩二, 本多隆文, 釣谷伊希子, 石崎昌夫, 山田裕一, 北川正信, 石川県梯川流域 Cd 汚染地住民 (健康管理対象者) の一症例—臨床及び病理検査成績—. *環境保健レポート*. 1991; 58: 161-165.
- 6.2.3 - 8 中川秀昭, 西条旨子, 森河裕子, 田畑正司, 千間正美, 三浦克之, 由田克士, 奥村義治, 河野俊一, 城戸照彦, 北川正信, 骨病理検索を行った梯川流域カドミウム汚染地域要管理者の1例. *環境保健レポート*. 1993; 60: 130-135.
- 6.2.3 - 9 Honda R., Tsuritani I., Noborisaka Y., Suzuki H., Ishizaki M., Yamada Y., Urinary cadmium excretion is correlated with calcaneal bone mass.

- in Japanese women living in an urban area. *Environ Res.* 2003; 91: 63-70.
- 6.2.3 - 10 骨軟化症研究班. 骨軟化症の診断に関する研究. 環境保健レポート. 1993; 60: 267-273.
- 6.2.3 - 11 吉川靖三, 骨とリン代謝. 日本骨代謝学会誌. 1983; 1: 26-32.
- 6.2.3 - 12 斎藤 寛, 薮 幸三, 古川洋太郎, 塩路隆治, 古山 隆, 吉永 馨, カドミウム腎障害 - 慢性カドミウム中毒およびいわゆるイタイイタイ病の腎病変と骨軟化症. 日本臨床. 1978; 73: 838-848. (日本臨床-1978に記載なし、年数、巻号、合致しない)
- 6.2.3 - 13 Bhattacharyya M.H., Whelton, B.D., Stern P.H., Peterson D.P., Cadmium accelerates bone loss in ovariectomized mice and fetal rat limb bones in culture. *Proc. Natl. Acad. Sci. USA.* 1988; 85: 8761-8765.
- 6.2.3 - 14 Miyahara T., Takata M., Mori-uchi S., Miyata M., Nagai M., Sugure A., Matsushita M., Kozuka H., Kuze S., Stimulative effects of cadmium on bone resorption in neonatal parietal bone resorption. *Toxicology* 1992; 73: 93-99.
- 6.2.3 - 15 Ogoshi K., Nanzai Y., Moriyama T., Decrease in bone strength of cadmium-treated young and old rats. *Arch. Toxicol.* 1992; 66: 315-320.
- 6.2.3 - 16 Alfvén T., Elinder C.G., Carlsson M.D., Grubb A., Hellström L., Persson B., Pettersson C., Spang G., Schütz A., Järup L., Low-level cadmium exposure and osteoporosis. *J. Bone Miner. Res.* 2000; 15: 1579-1586.
- 6.2.3 - 17 Staessen J.A., Roels H.A., Emelianov D., Kuznetsova T., Thijs L., Vangronsveld J., Fagard R., Environmental exposure to cadmium, forearm bone density, and risk of fractures: prospective population study. *Lancet* 1999; 353: 1140-1144.
- 6.2.3 - 18 Horiguchi H., Oguma E., Sasaki S., Miyamoto K., Ikeda Y., Machida M., Kayama F., Environmental exposure to cadmium at a level insufficient to induce renal tubular dysfunction does not affect bone density among female Japanese farmers. *Environ Res.* 2005; 97: 83-92.
- 6.2.4 呼吸器への影響
- 6.2.4 - 1 Sakurai H., Omae K., Toyama T., Higashi T., Nakadate T., Cross-sectional study of pulmonary function in cadmium alloy workers. *Scand. J. Work Environ. Health.* 1982; 8: 122-130.
- 6.2.4 - 2 Mannino D.M., Holguin F., Greves H.M., Savage-Brown A., Stock A.L., Jones R.L., Urinary cadmium levels predict lower lung function in current and former smokers: data from the third national health and nutrition examination survey. *Thorax.* 2004; 59: 194-198.
- 6.2.5 高血圧及び心血管系への影響
- 6.2.5 - 1 鍛冶利幸, 小山 洋, 佐藤雅彦, 遠山千春, 低用量カドミウム曝露と健康影響 (2) 生活習慣病と生殖毒性. 日本衛生学雑誌. 2002; 57: 556-563.
- 6.2.5 - 2 Schroeder H.A., Cadmium as a factor in hypertension. *J Chron Dis.* 1965; 18: 647-656.
- 6.2.5 - 3 Glauser S.C., Bello C.T., Glauser E.M., Blood-cadmium levels in normotensives and untreated hypertensive humans. *THE LANCET*, APRIL 3, 1976; 1: 717-718.
- 6.2.5 - 4 Beevers D.G., Campbell B.C., Goldberg A., Moore M.R., Hawthorne