

Despite low frequency of neurological sequelae, they may last up to 5 years. A team of neurologists, psychiatrists and epidemiologists should be a core of follow up systems. Collaboration with administration and health care personnel is important.

Neurological and medical health checks are of limited value in a mobile community. However, questionnaire surveys should be developed.

3. The role of neurologists in terrorist attacks which use nerve agents is as part of a rescue team.

A rapid diagnosis of the causative agent and serum ChE measurement is essential. Although miosis is a clue with high sensitivity and specificity, it is not an indication of severity and cannot be used for triage of victims. In long-term follow-up, neurologists should work in close collaboration with psychiatrists and epidemiologists. Education of affected and non-affected victims should be done.

#### Acknowledgement

We thank Dr. Leon Prockop (University of South Florida) for editing the manuscript.

#### References

- [1] Bright JE, Inns RH, Tuckwell NJ, Griffiths GD, Marrs TC. A histochemical study of changes observed in the mouse diaphragm after organophosphate poisoning. *Hum Exp Toxicol* 1991;10:9–14.
- [2] Burchfiel JL, Duffy FH. Organophosphate neurotoxicity: chronic effects of Sarin on the electroencephalogram of monkey and man. *Neurobehav Toxicol Teratol* 1982;4:768–78.
- [3] Crowell JA, Parker RM, Bucci TJ, Dacre JC. Neuropathy target esterase in hens after Sarin and soman. *J Biochem Toxicol* 1989;4:15–20.
- [4] Duffy FH, Burchfiel JL, Bartels PH, Gaon M, van Sim M. Long-term effects of an organophosphate upon the human electroencephalogram. *Toxicol Appl Pharmacol* 1979;47:161–76.
- [5] Goldstein BD, Fincher DR, Searle JR. Electrophysiological changes in the primary sensory neuron following subchronic soman and Sarin: iterations in sensory receptor function. *Toxicol Appl Pharmacol* 1987;91:55–64.
- [6] Gupta RC, Deubarn WD. Potential of memantine, D-tubocurarine, and atropine in preventing acute myopathy induced by organophosphate nerve agents: soman, Sarin, tabun and VX. *Neurotoxicology* 1992;13:649–61.
- [7] Husain K, Vijayaraghavan R, Pant SC, Raza SK, Pandey KS. Delayed neurotoxic effect of Sarin in mice after repeated inhalation exposure. *J Appl Toxicol* 1993;13:143–5.
- [8] Inns RH, Tuckwell NJ, Bright JE, Marrs TC. Histochemical demonstration of calcium accumulation in muscle fibers after experimental organophosphate poisoning. *Hum Exp Toxicol* 1990;9:245–50.
- [9] Kadar T, Shapira S, Cohen G, Sahar R, Alkalay D, Raven L. Sarin-induced neuropathology in rats. *Hum Exp Toxicol* 1995;14:252–9.
- [10] Kadokura M, Ushijima S, Ogawa Y, Shimizu H, Agata T, Yamamura T. Post traumatic stress disorder in victims of an attack with Sarin nerve gas on the Tokyo subway system. *Rinsho Seishin Igaku* 2000;29:677–83.
- [11] Kato T, Hamanaka T. Ocular signs and symptoms caused by exposure to Sarin gas. *Am J Ophthalmol* 1996;121:209–10.
- [12] Kawana N, Ishimatsu S, Kanda K. Victims of the terrorist Sarin attack on the Tokyo subway system. *Rinsho Seishin Igaku, Special Issue* 2002;139–45.
- [13] Masuda N, Okuda T, Miyazaki S, Takatsu M, Ito K, Kanetaka T, et al. Clinical and laboratory data in 71 patients with Sarin poisoning. *Teishin Igaku* 1995;47:387–90.
- [14] Matsumoto Regional Comprehensive Medical Council. Health emergency control system in Matsumoto city. *Health Hyg Matsumoto* 2000;22:1–107 [supplement].
- [15] Morita H, Yanagisawa N, Nakajima T, Shimizu M, Hirabayashi H, Okudera H, et al. Sarin poisoning of citizens in Matsumoto. *Lancet* 1995;346:290–3.
- [16] Murata K, Araki S, Yokoyama K, Okumura T, Ishimatsu S, Takasu N, et al. Asymptomatic sequelae to acute Sarin poisoning in the central and autonomic nervous system 6 months after the Tokyo subway attack. *J Neurol* 1997;244:601–6.
- [17] Nakajima T, Ohta S, Fukushima T, Yanagisawa N. Sequelae of Sarin toxicity at one and three years after exposure in Matsumoto, Japan. *J Epidemiol* 1999;9:337–43.
- [18] Nakajima T, Ohta S, Morita H, Midorikawa Y, Mimura S, Yanagisawa N. Epidemiological study of Sarin poisoning occurred in Matsumoto City, Japan. *J Epidemiol* 1998;8:33–41.
- [19] Nakajima T, Sato S, Morita H, Yanagisawa N. Sarin poisoning of a rescue team in the Matsumoto Sarin incident in Japan. *Occup Environ Med* 1997;54:697–701.
- [20] National Police Academy, Police Science Institute. A report on casualties of the subway Sarin incident. *Keisatsu Koron (Police Public Opinion)* 1999;54:37–47.
- [21] Nishiwaki T, Maekawa K, Ogawa Y, Asukai N, Minami M, Omae K, Sarin Health Effects Study Group. Effects of Sarin on the nervous system in rescue team staff members and police officers 3 years after the Tokyo subway Sarin attack. *Environ Health Perspect* 2001;109:1169–1173.
- [22] Nohara M. Ophthalmological health checks after Sarin exposure. In: Matsumoto Regional Comprehensive Medical Council, editor. *Health Hyg Matsumoto*, vol. 22; 2000. p. 42–51. Suppl.
- [23] Nohara M, Segawa K. Ocular symptoms due to organophosphorus gas (Sarin) poisoning in Matsumoto. *Br J Ophthalmol* 1996;80:1023.
- [24] Nozaki H, Hori S, Shinozawa T, Fujishima S, Takuma K, Kimura H, et al. Relationship between pupil size and acetylcholinesterase activity in patients exposed to Sarin vapour. *Intensive Care Med* 1997;23:1005–1007.
- [25] Ohbu S, Yamashita A, Takasu N, Yamaguchi T, Murai T, Nakana K, et al. Sarin poisoning on Tokyo subway. *South Med J* 1997;90:587–93.
- [26] Ohtani T, Iwanami A, Kasai K, Yamasue H, Kato T, Sakaki T, et al. Post traumatic stress disorder symptoms in victims of Tokyo subway attack: a 5 years follow-up study. *Psychiatry Clin Neurosci* 2004;58:624–9.
- [27] Okudera H, Morita H, Iwashita T, Shibata T, Otagiri T, Kobayashi S, et al. Unexpected nerve gas exposure in the city of Matsumoto: report of rescue activity in the first Sarin gas terrorism. *Am J Emerg Med* 1997;15:527–8.
- [28] Okumura T, Suzuki K, Fukuda A, Koyama, Takasu N, Ishimatsu S, et al. The Tokyo subway Sarin attack: disaster management: Part 1. Community emergency response: Part 2. Hospital response: Part 3. National and international responses. *Acad Emerg Med* 1998;5:613–628.
- [29] Okumura T, Takasu N, Ishimatsu S, Miyanoki S, Mitsuhashi A, Kumada K, et al. Report on 640 victims of the Tokyo subway Sarin attack. *Ann Emerg Med* 1996;28:129–35.
- [30] Sekijima Y, Morita H, Yanagisawa N. Two-year follow-up study of victims of Sarin poisoning in Matsumoto, Japan. *Ann Intern Med* 1997;127:1042.
- [31] Singer AW, Jaax NK, Graham JS, McLeod Jr CG. Cardiomyopathy in soman and Sarin intoxicated rats. *Toxicol Lett* 1987;36:243–9.

- [32] Suzuki J, Kohno T, Tsukagoshi M, Furihata T, Yamazaki K. Eighteen cases exposed to Sarin in Matsumoto, Japan. *Int Med* 1997;36:466–70.
- [33] Yanagisawa N, editor. Report for Toxic Gas Attack in Matsumoto. Matsumoto Regional Comprehensive Medical Council; 1995.
- [34] Yokoyama K, Araki S, Murata K, Nishitani M, Okumura T, Ishimatsu S, et al. A preliminary study on delayed vestibulo-cerebellar effects of Tokyo subway Sarin poisoning in relation to gender difference: frequency analysis of postural sway. *J Occup Environ Med* 1998;40:17–21.
- [35] Yokoyama K, Ogura T, Kishimoto M, Hinoshita F, Hara S, Yamada A, et al. Blood purification for severe Sarin poisoning after the Tokyo subway attack. *JAMA* 1995;274:379.