

○糖尿病に関連する疾患、危険因子、生活習慣等の状況

第52表-1 糖尿病合併症の状況（神経障害、性・年齢階級別）

	神経障害 がある	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男女計	ある	0	0	0	0	1	1.8	12	8.3	28	12.5	27	13.5	68	10.5
	ない	6	85.7	13	100.0	55	96.5	130	89.7	189	84.4	168	84.0	561	86.8
	不明	1	14.3	0	0	1	1.8	3	2.1	7	3.1	5	2.5	17	2.6
	一総数一	7	100.0	13	100.0	57	100.0	145	100.0	224	100.0	200	100.0	646	100.0
男 性	ある	0	0	0	0	0	0	8	7.9	17	12.2	13	10.9	38	9.4
	ない	2	66.7	9	100.0	32	97.0	90	89.1	117	84.2	102	85.7	352	87.1
	不明	1	33.3	0	0	1	3.0	3	3.0	5	3.6	4	3.4	14	3.5
	一総数一	3	100.0	9	100.0	33	100.0	101	100.0	139	100.0	119	100.0	404	100.0
女 性	ある	0	0	0	0	1	4.2	4	9.1	11	12.9	14	17.3	30	12.4
	ない	4	100.0	4	100.0	23	95.8	40	90.9	72	84.7	66	81.5	209	86.4
	不明	0	0	0	0	0	0	0	0	2	2.4	1	1.2	3	1.2
	一総数一	4	100.0	4	100.0	24	100.0	44	100.0	85	100.0	81	100.0	242	100.0

第52表-2 糖尿病合併症の状況（網膜症、性・年齢階級別）

	網膜症がある	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男女計	ある	1	14.3	0	0	1	1.8	4	2.8	27	2.8	23	11.5	56	8.7
	ない	5	71.4	13	100.0	55	96.5	137	94.5	191	94.5	172	86.0	573	88.7
	不明	1	14.3	0	0	1	1.8	4	2.8	6	2.8	5	2.5	17	2.6
	一総数一	7	100.0	13	100.0	57	100.0	145	100.0	224	100.0	200	100.0	646	100.0
男 性	ある	0	0	0	0	0	0	1	1.0	18	1.0	12	10.1	31	7.7
	ない	2	66.7	9	100.0	32	97.0	96	95.0	117	95.0	103	86.6	359	88.9
	不明	1	33.3	0	0	1	3.0	4	4.0	4	4.0	4	3.4	14	3.5
	一総数一	3	100.0	9	100.0	33	100.0	101	100.0	139	100.0	119	100.0	404	100.0
女 性	ある	1	25.0	0	0	1	4.2	3	6.8	9	6.8	11	13.6	25	10.3
	ない	3	75.0	4	100.0	23	95.8	41	93.2	74	93.2	69	85.2	214	88.4
	不明	0	0	0	0	0	0	0	0	2	0	1	1.2	3	1.2
	一総数一	4	100.0	4	100.0	24	100.0	44	100.0	85	100.0	81	100.0	242	100.0

第52表-3 糖尿病合併症の状況（腎症、性・年齢階級別）

	腎症がある	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男女計	ある	0	0	0	0	5	8.8	15	10.3	23	10.3	20	10.0	63	9.8
	ない	6	85.7	13	100.0	51	89.5	125	86.2	192	85.7	176	88.0	563	87.2
	不明	1	14.3	0	0	1	1.8	5	3.4	9	4.0	4	2.0	20	3.1
	—総数—	7	100.0	13	100.0	57	100.0	145	100.0	224	100.0	200	100.0	646	100.0
男 性	ある	0	0	0	0	4	12.1	9	8.9	14	10.1	14	11.8	41	10.1
	ない	2	66.7	9	100.0	28	84.8	87	86.1	119	85.6	101	84.9	346	85.6
	不明	1	33.3	0	0	1	3.0	5	5.0	6	4.3	4	3.4	17	4.2
	—総数—	3	100.0	9	100.0	33	100.0	101	100.0	139	100.0	119	100.0	404	100.0
女 性	ある	0	0	0	0	1	4.2	6	13.6	9	10.6	6	7.4	22	9.1
	ない	4	100.0	4	100.0	23	95.8	38	86.4	73	85.9	75	92.6	217	89.7
	不明	0	0	0	0	0	0	0	0	3	3.5	0	0	3	1.2
	—総数—	4	100.0	4	100.0	24	100.0	44	100.0	85	100.0	81	100.0	242	100.0

第52表-4 糖尿病合併症の状況（足壊疽、性・年齢階級別）

	足壊疽がある	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男女計	ある	0	0	0	0	0	0	1	0.7	2	0.9	3	1.5	6	0.9
	ない	6	85.7	13	100.0	56	98.2	141	97.2	213	95.1	192	96.0	621	96.1
	不明	1	14.3	0	0	1	1.8	3	2.1	9	4.0	5	2.5	19	2.9
	—総数—	7	100.0	13	100.0	57	100.0	145	100.0	224	100.0	200	100.0	646	100.0
男 性	ある	0	0	0	0	0	0	0	0	1	0.7	0	0	1	0.2
	ない	2	66.7	9	100.0	32	97.0	98	97.0	132	95.0	115	96.6	388	96.0
	不明	1	33.3	0	0	1	3.0	3	3.0	6	4.3	4	3.4	15	3.7
	—総数—	3	100.0	9	100.0	33	100.0	101	100.0	139	100.0	119	100.0	404	100.0
女 性	ある	0	0	0	0	0	0	1	2.3	1	1.2	3	3.7	5	2.1
	ない	4	100.0	4	100.0	24	100.0	43	97.7	81	95.3	77	95.1	233	96.3
	不明	0	0	0	0	0	0	0	0	3	3.5	1	1.2	4	1.7
	—総数—	4	100.0	4	100.0	24	100.0	44	100.0	85	100.0	81	100.0	242	100.0

第53表-1 心臓病および脳卒中の状況（心臓病の既往、性・年齢階級別）

	心臓病の既往 がある	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		—計—	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男女計	ある	3	0.6	8	1.0	13	1.6	56	4.6	123	9.6	215	17.6	418	7.2
	ない	462	99.4	754	99.0	821	98.4	1,151	95.4	1,153	90.4	1,008	82.4	5,349	92.8
	—総数—	465	100.0	762	100.0	834	100.0	1,207	100.0	1,276	100.0	1,223	100.0	5,767	100.0
男 性	ある	1	0.5	3	1.0	7	2.1	28	5.9	62	10.8	92	18.0	193	8.2
	ない	182	99.5	288	99.0	320	97.9	446	94.1	514	89.2	418	82.0	2,168	91.8
	—総数—	183	100.0	291	100.0	327	100.0	474	100.0	576	100.0	510	100.0	2,361	100.0
女 性	ある	2	0.7	5	1.1	6	1.2	28	3.8	61	8.7	123	17.3	225	6.6
	ない	280	99.3	466	98.9	501	98.8	705	96.2	639	91.3	590	82.7	3,181	93.4
	—総数—	282	100.0	471	100.0	507	100.0	733	100.0	700	100.0	713	100.0	3,406	100.0

第53表-2 心臓病および脳卒中の状況（脳卒中の既往、性・年齢階級別）

	脳卒中の既往 がある	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		—計—	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男女計	ある	1	0.2	4	0.5	6	0.7	22	1.8	67	5.3	129	10.6	229	4.0
	ない	464	99.8	758	99.5	827	99.3	1,183	98.2	1,207	94.7	1,092	89.4	5,531	96.0
	—総数—	465	100.0	762	100.0	833	100.0	1,205	100.0	1,274	100.0	1,221	100.0	5,760	100.0
男 性	ある	1	0.5	3	1.0	4	1.2	8	1.7	43	7.5	70	13.8	129	5.5
	ない	182	99.5	288	99.0	323	98.8	466	98.3	534	92.5	436	86.2	2,229	94.5
	—総数—	183	100.0	291	100.0	327	100.0	474	100.0	577	100.0	506	100.0	2,358	100.0
女 性	ある	0	0	1	0.2	2	0.4	14	1.9	24	3.4	59	8.3	100	2.9
	ない	282	100.0	470	99.8	504	99.6	717	98.1	673	96.6	656	91.7	3,302	97.1
	—総数—	282	100.0	471	100.0	506	100.0	731	100.0	697	100.0	715	100.0	3,402	100.0

第54表-1 糖尿病の状況別、糖尿病合併症の状況（神経障害、性・年齢階級別）

	糖尿病の状況	神経障害の状況	20-29歳		30-39歳		40-49歳		50-59歳	
			人数	%	人数	%	人数	%	人数	%
男女計	HbA1c 5.6%未満	神経障害がある								
		はい	0	0	0	0	0	0	5	0.5
		いいえ	391	99.7	679	99.9	686	99.7	904	99.2
		不明	1	0.3	1	0.1	2	0.3	2	0.2
	—総数—	392	100.0	680	100.0	688	100.0	911	100.0	
	HbA1c 5.6%以上 6.1%未満	神経障害がある								
		はい	0	0	0	0	0	0	1	0.8
		いいえ	4	100.0	27	100.0	48	98.0	118	98.3
		不明	0	0	0	0	1	2.0	1	0.8
	—総数—	4	100.0	27	100.0	49	100.0	120	100.0	
	HbA1c 6.1%以上 または 現在治療を受けている	神経障害がある								
		はい	0	0	0	0	1	3.3	4	4.3
いいえ		2	100.0	6	100.0	29	96.7	87	94.6	
不明		0	0	0	0	0	0	1	1.1	
—総数—	2	100.0	6	100.0	30	100.0	92	100.0		
男 性	HbA1c 5.6%未満	神経障害がある								
		はい	0	0	0	0	0	0	4	1.2
		いいえ	141	99.3	248	100.0	272	100.0	318	98.5
		不明	1	0.7	0	0	0	0	1	0.3
	—総数—	142	100.0	248	100.0	272	100.0	323	100.0	
	HbA1c 5.6%以上 6.1%未満	神経障害がある								
		はい	0	0	0	0	0	0	0	0
		いいえ	3	100.0	7	100.0	9	90.0	45	97.8
		不明	0	0	0	0	1	10.0	1	2.2
	—総数—	3	100.0	7	100.0	10	100.0	46	100.0	
	HbA1c 6.1%以上 または 現在治療を受けている	神経障害がある								
		はい	0	0	0	0	0	0	3	5.0
いいえ		0	0	2	100.0	13	100.0	56	93.3	
不明		0	0	0	0	0	0	1	1.7	
—総数—	0	0	2	100.0	13	100.0	60	100.0		
女 性	HbA1c 5.6%未満	神経障害がある								
		はい	0	0	0	0	0	0	1	0.2
		いいえ	250	100.0	431	99.8	414	99.5	586	99.7
		不明	0	0	1	0.2	2	0.5	1	0.2
	—総数—	250	100.0	432	100.0	416	100.0	588	100.0	
	HbA1c 5.6%以上 6.1%未満	神経障害がある								
		はい	0	0	0	0	0	0	1	1.4
		いいえ	1	100.0	20	100.0	39	100.0	73	98.6
		不明	0	0	0	0	0	0	0	0
	—総数—	1	100.0	20	100.0	39	100.0	74	100.0	
	HbA1c 6.1%以上 または 現在治療を受けている	神経障害がある								
		はい	0	0	0	0	1	5.9	1	3.1
いいえ		2	100.0	4	100.0	16	94.1	31	96.9	
不明		0	0	0	0	0	0	0	0	
—総数—	2	100.0	4	100.0	17	100.0	32	100.0		

60-69歳		70歳以上		— 計 —		20-39歳(再掲)		40-59歳(再掲)		60歳以上(再掲)	
人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
3	0.4	1	0.1	9	0.2	0	0	5	0.3	4	0.2
850	99.3	768	99.6	4,278	99.5	1,070	99.8	1,590	99.4	1,618	99.4
3	0.4	2	0.3	11	0.3	2	0.2	4	0.3	5	0.3
856	100.0	771	100.0	4,298	100.0	1,072	100.0	1,599	100.0	1,627	100.0
1	0.6	3	1.6	5	0.9	0	0	1	0.6	4	1.1
176	98.3	182	97.3	555	98.1	31	100.0	166	98.2	358	97.8
2	1.1	2	1.1	6	1.1	0	0	2	1.2	4	1.1
179	100.0	187	100.0	566	100.0	31	100.0	169	100.0	366	100.0
18	10.3	21	11.8	44	9.1	0	0	5	4.1	39	11.1
154	88.5	151	84.8	429	89.0	8	100.0	116	95.1	305	86.6
2	1.1	6	3.4	9	1.9	0	0	1	0.8	8	2.3
174	100.0	178	100.0	482	100.0	8	100.0	122	100.0	352	100.0
2	0.5	1	0.3	7	0.4	0	0	4	0.7	3	0.4
371	98.9	296	99.0	1,646	99.2	389	99.7	590	99.2	667	99.0
2	0.5	2	0.7	6	0.4	1	0.3	1	0.2	4	0.6
375	100.0	299	100.0	1,659	100.0	390	100.0	595	100.0	674	100.0
0	0	2	2.6	2	0.9	0	0	0	0	2	1.3
73	100.0	74	96.1	211	97.7	10	100.0	54	96.4	147	98.0
0	0	1	1.3	3	1.4	0	0	2	3.6	1	0.7
73	100.0	77	100.0	216	100.0	10	100.0	56	100.0	150	100.0
11	11.2	10	9.8	24	8.7	0	0	3	4.1	21	10.5
85	86.7	89	87.3	245	89.1	2	100.0	69	94.5	174	87.0
2	2.0	3	2.9	6	2.2	0	0	1	1.4	5	2.5
98	100.0	102	100.0	275	100.0	2	100.0	73	100.0	200	100.0
1	0.2	0	0	2	0.1	0	0	1	0.1	1	0.1
479	99.6	472	100.0	2,632	99.7	681	99.9	1,000	99.6	951	99.8
1	0.2	0	0	5	0.2	1	0.1	3	0.3	1	0.1
481	100.0	472	100.0	2,639	100.0	682	100.0	1,004	100.0	953	100.0
1	0.9	1	0.9	3	0.9	0	0	1	0.9	2	0.9
103	97.2	108	98.2	344	98.3	21	100.0	112	99.1	211	97.7
2	1.9	1	0.9	3	0.9	0	0	0	0	3	1.4
106	100.0	110	100.0	350	100.0	21	100.0	113	100.0	216	100.0
7	9.2	11	14.5	20	9.7	0	0	2	4.1	18	11.8
69	90.8	62	81.6	184	88.9	6	100.0	47	95.9	131	86.2
0	0	3	3.9	3	1.4	0	0	0	0	3	2.0
76	100.0	76	100.0	207	100.0	6	100.0	49	100.0	152	100.0

第54表-2 糖尿病の状況別、糖尿病合併症の状況（網膜症、性・年齢階級別）

	糖尿病の状況	網膜症の状況	20-29歳		30-39歳		40-49歳		50-59歳	
			人数	%	人数	%	人数	%	人数	%
男女計	HbA1c 5.6%未満	網膜症がある								
		はい	1	0.3	0	0	0	0	1	0.1
		いいえ	390	99.5	679	99.9	686	99.7	908	99.7
		不明	1	0.3	1	0.1	2	0.3	2	0.2
		—総数—	392	100.0	680	100.0	688	100.0	911	100.0
	HbA1c 5.6%以上 6.1%未満	網膜症がある								
		はい	0	0	0	0	0	0	1	0.8
		いいえ	4	100.0	27	100.0	48	98.0	118	98.3
		不明	0	0	0	0	1	2.0	1	0.8
		—総数—	4	100.0	27	100.0	49	100.0	120	100.0
	HbA1c 6.1%以上 または 現在治療を受けている	網膜症がある								
		はい	0	0	0	0	0	0	1	1.1
いいえ		2	100.0	6	100.0	30	100.0	89	96.7	
不明		0	0	0	0	0	0	2	2.2	
	—総数—	2	100.0	6	100.0	30	100.0	92	100.0	
男 性	HbA1c 5.6%未満	網膜症がある								
		はい	0	0	0	0	0	0	1	0.3
		いいえ	141	99.3	248	100.0	272	100.0	321	99.4
		不明	1	0.7	0	0	0	0	1	0.3
		—総数—	142	100.0	248	100.0	272	100.0	323	100.0
	HbA1c 5.6%以上 6.1%未満	網膜症がある								
		はい	0	0	0	0	0	0	0	0
		いいえ	3	100.0	7	100.0	9	90.0	45	97.8
		不明	0	0	0	0	1	10.0	1	2.2
		—総数—	3	100.0	7	100.0	10	100.0	46	100.0
	HbA1c 6.1%以上 または 現在治療を受けている	網膜症がある								
		はい	0	0	0	0	0	0	0	0
いいえ		0	0	2	100.0	13	100.0	58	96.7	
不明		0	0	0	0	0	0	2	3.3	
	—総数—	0	0	2	100.0	13	100.0	60	100.0	
女 性	HbA1c 5.6%未満	網膜症がある								
		はい	1	0.4	0	0	0	0	0	0
		いいえ	249	99.6	431	99.8	414	99.5	587	99.8
		不明	0	0	1	0.2	2	0.5	1	0.2
		—総数—	250	100.0	432	100.0	416	100.0	588	100.0
	HbA1c 5.6%以上 6.1%未満	網膜症がある								
		はい	0	0	0	0	0	0	1	1.4
		いいえ	1	100.0	20	100.0	39	100.0	73	98.6
		不明	0	0	0	0	0	0	0	0
		—総数—	1	100.0	20	100.0	39	100.0	74	100.0
	HbA1c 6.1%以上 または 現在治療を受けている	網膜症がある								
		はい	0	0	0	0	0	0	1	3.1
いいえ		2	100.0	4	100.0	17	100.0	31	96.9	
不明		0	0	0	0	0	0	0	0	
	—総数—	2	100.0	4	100.0	17	100.0	32	100.0	

60-69歳		70歳以上		一計		20-39歳(再掲)		40-59歳(再掲)		60歳以上(再掲)	
人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
3	0.4	1	0.1	6	0.1	1	0.1	1	0.1	4	0.2
851	99.4	768	99.6	4,282	99.6	1,069	99.7	1,594	99.7	1,619	99.5
2	0.2	2	0.3	10	0.2	2	0.2	4	0.3	4	0.2
856	100.0	771	100.0	4,298	100.0	1,072	100.0	1,599	100.0	1,627	100.0
2	1.1	2	1.1	5	0.9	0	0	1	0.6	4	1.1
175	97.8	183	97.9	555	98.1	31	100.0	166	98.2	358	97.8
2	1.1	2	1.1	6	1.1	0	0	2	1.2	4	1.1
179	100.0	187	100.0	566	100.0	31	100.0	169	100.0	366	100.0
14	8.0	17	9.6	32	6.6	0	0	1	0.8	31	8.8
158	90.8	155	87.1	440	91.3	8	100.0	119	97.5	313	88.9
2	1.1	6	3.4	10	2.1	0	0	2	1.6	8	2.3
174	100.0	178	100.0	482	100.0	8	100.0	122	100.0	352	100.0
3	0.8	1	0.3	5	0.3	0	0	1	0.2	4	0.6
371	98.9	296	99.0	1,649	99.4	389	99.7	593	99.7	667	99.0
1	0.3	2	0.7	5	0.3	1	0.3	1	0.2	3	0.4
375	100.0	299	100.0	1,659	100.0	390	100.0	595	100.0	674	100.0
2	2.7	2	2.6	4	1.9	0	0	0	0	4	2.7
71	97.3	74	96.1	209	96.8	10	100.0	54	96.4	145	96.7
0	0	1	1.3	3	1.4	0	0	2	3.6	1	0.7
73	100.0	77	100.0	216	100.0	10	100.0	56	100.0	150	100.0
9	9.2	9	8.8	18	6.5	0	0	0	0	18	9.0
87	88.8	90	88.2	250	90.9	2	100.0	71	97.3	177	88.5
2	2.0	3	2.9	7	2.5	0	0	2	2.7	5	2.5
98	100.0	102	100.0	275	100.0	2	100.0	73	100.0	200	100.0
0	0	0	0	1	0.0	1	0.1	0	0	0	0
480	99.8	472	100.0	2,633	99.8	680	99.7	1,001	99.7	952	99.9
1	0.2	0	0	5	0.2	1	0.1	3	0.3	1	0.1
481	100.0	472	100.0	2,639	100.0	682	100.0	1,004	100.0	953	100.0
0	0	0	0	1	0.3	0	0	1	0.9	0	0
104	98.1	109	99.1	346	98.9	21	100.0	112	99.1	213	98.6
2	1.9	1	0.9	3	0.9	0	0	0	0	3	1.4
106	100.0	110	100.0	350	100.0	21	100.0	113	100.0	216	100.0
5	6.6	8	10.5	14	6.8	0	0	1	2.0	13	8.6
71	93.4	65	85.5	190	91.8	6	100.0	48	98.0	136	89.5
0	0	3	3.9	3	1.4	0	0	0	0	3	2.0
76	100.0	76	100.0	207	100.0	6	100.0	49	100.0	152	100.0

第54表-3 糖尿病の状況別、糖尿病合併症の状況（腎症、性・年齢階級別）

	糖尿病の状況	腎症の状況	20-29歳		30-39歳		40-49歳		50-59歳	
			人数	%	人数	%	人数	%	人数	%
男女計	HbA1c 5.6%未満	腎症がある								
		はい	0	0	0	0	2	0.3	5	0.5
		いいえ	391	99.7	679	99.9	684	99.4	904	99.2
		不明	1	0.3	1	0.1	2	0.3	2	0.2
	—総数—	392	100.0	680	100.0	688	100.0	911	100.0	
	HbA1c 5.6%以上 6.1%未満	腎症がある								
		はい	0	0	0	0	0	0	2	1.7
		いいえ	4	100.0	27	100.0	48	98.0	116	96.7
		不明	0	0	0	0	1	2.0	2	1.7
—総数—	4	100.0	27	100.0	49	100.0	120	100.0		
HbA1c 6.1%以上 または 現在治療を受けている	腎症がある									
	はい	0	0	0	0	3	10.0	8	8.7	
	いいえ	2	100.0	6	100.0	27	90.0	82	89.1	
	不明	0	0	0	0	0	0	2	2.2	
—総数—	2	100.0	6	100.0	30	100.0	92	100.0		
男 性	HbA1c 5.6%未満	腎症がある								
		はい	0	0	0	0	2	0.7	4	1.2
		いいえ	141	99.3	248	100.0	270	99.3	318	98.5
		不明	1	0.7	0	0	0	0	1	0.3
	—総数—	142	100.0	248	100.0	272	100.0	323	100.0	
	HbA1c 5.6%以上 6.1%未満	腎症がある								
		はい	0	0	0	0	0	0	1	2.2
		いいえ	3	100.0	7	100.0	9	90.0	43	93.5
		不明	0	0	0	0	1	10.0	2	4.3
—総数—	3	100.0	7	100.0	10	100.0	46	100.0		
HbA1c 6.1%以上 または 現在治療を受けている	腎症がある									
	はい	0	0	0	0	2	15.4	4	6.7	
	いいえ	0	0	2	100.0	11	84.6	54	90.0	
	不明	0	0	0	0	0	0	2	3.3	
—総数—	0	0	2	100.0	13	100.0	60	100.0		
女 性	HbA1c 5.6%未満	腎症がある								
		はい	0	0	0	0	0	0	1	0.2
		いいえ	250	100.0	431	99.8	414	99.5	586	99.7
		不明	0	0	1	0.2	2	0.5	1	0.2
	—総数—	250	100.0	432	100.0	416	100.0	588	100.0	
	HbA1c 5.6%以上 6.1%未満	腎症がある								
		はい	0	0	0	0	0	0	1	1.4
		いいえ	1	100.0	20	100.0	39	100.0	73	98.6
		不明	0	0	0	0	0	0	0	0
—総数—	1	100.0	20	100.0	39	100.0	74	100.0		
HbA1c 6.1%以上 または 現在治療を受けている	腎症がある									
	はい	0	0	0	0	1	5.9	4	12.5	
	いいえ	2	100.0	4	100.0	16	94.1	28	87.5	
	不明	0	0	0	0	0	0	0	0	
—総数—	2	100.0	4	100.0	17	100.0	32	100.0		

60-69歳		70歳以上		— 計 —		20-39歳(再掲)		40-59歳(再掲)		60歳以上(再掲)	
人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
0	0	0	0	7	0.2	0	0	7	0.4	0	0
853	99.6	769	99.7	4,280	99.6	1,070	99.8	1,588	99.3	1,622	99.7
3	0.4	2	0.3	11	0.3	2	0.2	4	0.3	5	0.3
856	100.0	771	100.0	4,298	100.0	1,072	100.0	1,599	100.0	1,627	100.0
1	0.6	3	1.6	6	1.1	0	0	2	1.2	4	1.1
176	98.3	182	97.3	553	97.7	31	100.0	164	97.0	358	97.8
2	1.1	2	1.1	7	1.2	0	0	3	1.8	4	1.1
179	100.0	187	100.0	566	100.0	31	100.0	169	100.0	366	100.0
17	9.8	16	9.0	44	9.1	0	0	11	9.0	33	9.4
154	88.5	157	88.2	428	88.8	8	100.0	109	89.3	311	88.4
3	1.7	5	2.8	10	2.1	0	0	2	1.6	8	2.3
174	100.0	178	100.0	482	100.0	8	100.0	122	100.0	352	100.0
0	0	0	0	6	0.4	0	0	6	1.0	0	0
373	99.5	297	99.3	1,647	99.3	389	99.7	588	98.8	670	99.4
2	0.5	2	0.7	6	0.4	1	0.3	1	0.2	4	0.6
375	100.0	299	100.0	1,659	100.0	390	100.0	595	100.0	674	100.0
1	1.4	3	3.9	5	2.3	0	0	1	1.8	4	2.7
72	98.6	73	94.8	207	95.8	10	100.0	52	92.9	145	96.7
0	0	1	1.3	4	1.9	0	0	3	5.4	1	0.7
73	100.0	77	100.0	216	100.0	10	100.0	56	100.0	150	100.0
10	10.2	11	10.8	27	9.8	0	0	6	8.2	21	10.5
86	87.8	88	86.3	241	87.6	2	100.0	65	89.0	174	87.0
2	2.0	3	2.9	7	2.5	0	0	2	2.7	5	2.5
98	100.0	102	100.0	275	100.0	2	100.0	73	100.0	200	100.0
0	0	0	0	1	0.0	0	0	1	0.1	0	0
480	99.8	472	100.0	2,633	99.8	681	99.9	1,000	99.6	952	99.9
1	0.2	0	0	5	0.2	1	0.1	3	0.3	1	0.1
481	100.0	472	100.0	2,639	100.0	682	100.0	1,004	100.0	953	100.0
0	0	0	0	1	0.3	0	0	1	0.9	0	0
104	98.1	109	99.1	346	98.9	21	100.0	112	99.1	213	98.6
2	1.9	1	0.9	3	0.9	0	0	0	0	3	1.4
106	100.0	110	100.0	350	100.0	21	100.0	113	100.0	216	100.0
7	9.2	5	6.6	17	8.2	0	0	5	10.2	12	7.9
68	89.5	69	90.8	187	90.3	6	100.0	44	89.8	137	90.1
1	1.3	2	2.6	3	1.4	0	0	0	0	3	2.0
76	100.0	76	100.0	207	100.0	6	100.0	49	100.0	152	100.0

第54表-4 糖尿病の状況別、糖尿病合併症の状況（足壊疽、性・年齢階級別）

	糖尿病の状況	足壊疽の状況	20-29歳		30-39歳		40-49歳		50-59歳	
			人数	%	人数	%	人数	%	人数	%
男女計	HbA1c 5.6%未満	足壊疽がある								
		はい	0	0	0	0	0	0	0	0
		いいえ	391	99.7	679	99.9	686	99.7	909	99.8
		不明	1	0.3	1	0.1	2	0.3	2	0.2
	—総数—	392	100.0	680	100.0	688	100.0	911	100.0	
	HbA1c 5.6%以上 6.1%未満	足壊疽がある								
		はい	0	0	0	0	0	0	1	0.8
		いいえ	4	100.0	27	100.0	48	98.0	118	98.3
		不明	0	0	0	0	1	2.0	1	0.8
	—総数—	4	100.0	27	100.0	49	100.0	120	100.0	
	HbA1c 6.1%以上 または 現在治療を受けている	足壊疽がある								
		はい	0	0	0	0	0	0	0	0
いいえ		2	100.0	6	100.0	30	100.0	91	98.9	
不明		0	0	0	0	0	0	1	1.1	
—総数—	2	100.0	6	100.0	30	100.0	92	100.0		
男 性	HbA1c 5.6%未満	足壊疽がある								
		はい	0	0	0	0	0	0	0	0
		いいえ	141	99.3	248	100.0	272	100.0	322	99.7
		不明	1	0.7	0	0	0	0	1	0.3
	—総数—	142	100.0	248	100.0	272	100.0	323	100.0	
	HbA1c 5.6%以上 6.1%未満	足壊疽がある								
		はい	0	0	0	0	0	0	0	0
		いいえ	3	100.0	7	100.0	9	90.0	45	97.8
		不明	0	0	0	0	1	10.0	1	2.2
	—総数—	3	100.0	7	100.0	10	100.0	46	100.0	
	HbA1c 6.1%以上 または 現在治療を受けている	足壊疽がある								
		はい	0	0	0	0	0	0	0	0
いいえ		0	0	2	100.0	13	100.0	59	98.3	
不明		0	0	0	0	0	0	1	1.7	
—総数—	0	0	2	100.0	13	100.0	60	100.0		
女 性	HbA1c 5.6%未満	足壊疽がある								
		はい	0	0	0	0	0	0	0	0
		いいえ	250	100.0	431	99.8	414	99.5	587	99.8
		不明	0	0	1	0.2	2	0.5	1	0.2
	—総数—	250	100.0	432	100.0	416	100.0	588	100.0	
	HbA1c 5.6%以上 6.1%未満	足壊疽がある								
		はい	0	0	0	0	0	0	1	1.4
		いいえ	1	100.0	20	100.0	39	100.0	73	98.6
		不明	0	0	0	0	0	0	0	0
	—総数—	1	100.0	20	100.0	39	100.0	74	100.0	
	HbA1c 6.1%以上 または 現在治療を受けている	足壊疽がある								
		はい	0	0	0	0	0	0	0	0
いいえ		2	100.0	4	100.0	17	100.0	32	100.0	
不明		0	0	0	0	0	0	0	0	
—総数—	2	100.0	4	100.0	17	100.0	32	100.0		

60-69歳		70歳以上		— 計 —		20-39歳(再掲)		40-59歳(再掲)		60歳以上(再掲)	
人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
0	0	0	0	0	0	0	0	0	0	0	0
853	99.6	769	99.7	4,287	99.7	1,070	99.8	1,595	99.7	1,622	99.7
3	0.4	2	0.3	11	0.3	2	0.2	4	0.3	5	0.3
856	100.0	771	100.0	4,298	100.0	1,072	100.0	1,599	100.0	1,627	100.0
0	0	0	0	1	0.2	0	0	1	0.6	0	0
177	98.9	185	98.9	559	98.8	31	100.0	166	98.2	362	98.9
2	1.1	2	1.1	6	1.1	0	0	2	1.2	4	1.1
179	100.0	187	100.0	566	100.0	31	100.0	169	100.0	366	100.0
1	0.6	3	1.7	4	0.8	0	0	0	0	4	1.1
170	97.7	169	94.9	468	97.1	8	100.0	121	99.2	339	96.3
3	1.7	6	3.4	10	2.1	0	0	1	0.8	9	2.6
174	100.0	178	100.0	482	100.0	8	100.0	122	100.0	352	100.0
0	0	0	0	0	0	0	0	0	0	0	0
373	99.5	297	99.3	1,653	99.6	389	99.7	594	99.8	670	99.4
2	0.5	2	0.7	6	0.4	1	0.3	1	0.2	4	0.6
375	100.0	299	100.0	1,659	100.0	390	100.0	595	100.0	674	100.0
0	0	0	0	0	0	0	0	0	0	0	0
73	100.0	76	98.7	213	98.6	10	100.0	54	96.4	149	99.3
0	0	1	1.3	3	1.4	0	0	2	3.6	1	0.7
73	100.0	77	100.0	216	100.0	10	100.0	56	100.0	150	100.0
1	1.0	0	0	1	0.4	0	0	0	0	1	0.5
95	96.9	99	97.1	268	97.5	2	100.0	72	98.6	194	97.0
2	2.0	3	2.9	6	2.2	0	0	1	1.4	5	2.5
98	100.0	102	100.0	275	100.0	2	100.0	73	100.0	200	100.0
0	0	0	0	0	0	0	0	0	0	0	0
480	99.8	472	100.0	2,634	99.8	681	99.9	1,001	99.7	952	99.9
1	0.2	0	0	5	0.2	1	0.1	3	0.3	1	0.1
481	100.0	472	100.0	2,639	100.0	682	100.0	1,004	100.0	953	100.0
0	0	0	0	1	0.3	0	0	1	0.9	0	0
104	98.1	109	99.1	346	98.9	21	100.0	112	99.1	213	98.6
2	1.9	1	0.9	3	0.9	0	0	0	0	3	1.4
106	100.0	110	100.0	350	100.0	21	100.0	113	100.0	216	100.0
0	0	3	3.9	3	1.4	0	0	0	0	3	2.0
75	98.7	70	92.1	200	96.6	6	100.0	49	100.0	145	95.4
1	1.3	3	3.9	4	1.9	0	0	0	0	4	2.6
76	100.0	76	100.0	207	100.0	6	100.0	49	100.0	152	100.0

第55表-1 糖尿病の状況別、心臓病および脳卒中の状況（心臓病の既往、性・年齢階級別）

	糖尿病の状況	心臓病の状況	20-29歳		30-39歳		40-49歳		50-59歳	
			人数	%	人数	%	人数	%	人数	%
男女計	HbA1c 5.6%未満	心臓病の既往がある								
		はい	1	0.3	8	1.2	10	1.5	45	5.0
		いいえ	391	99.7	672	98.8	677	98.5	864	95.0
		—総数—	392	100.0	680	100.0	687	100.0	909	100.0
	HbA1c 5.6%以上 6.1%未満	心臓病の既往がある								
		はい	0	0	0	0	3	6.3	5	4.2
		いいえ	4	100.0	27	100.0	45	93.8	115	95.8
		—総数—	4	100.0	27	100.0	48	100.0	120	100.0
	HbA1c 6.1%以上または 現在治療を受けている	心臓病の既往がある								
はい		0	0	0	0	0	0	4	4.4	
いいえ		2	100.0	6	100.0	30	100.0	87	95.6	
	—総数—	2	100.0	6	100.0	30	100.0	91	100.0	
男 性	HbA1c 5.6%未満	心臓病の既往がある								
		はい	0	0	3	1.2	5	1.8	23	7.1
		いいえ	142	100.0	245	98.8	267	98.2	300	92.9
		—総数—	142	100.0	248	100.0	272	100.0	323	100.0
	HbA1c 5.6%以上 6.1%未満	心臓病の既往がある								
		はい	0	0	0	0	2	20.0	2	4.3
		いいえ	3	100.0	7	100.0	8	80.0	44	95.7
		—総数—	3	100.0	7	100.0	10	100.0	46	100.0
	HbA1c 6.1%以上または 現在治療を受けている	心臓病の既往がある								
はい		0	0	0	0	0	0	2	3.4	
いいえ		0	0	2	100.0	13	100.0	57	96.6	
	—総数—	0	0	2	100.0	13	100.0	59	100.0	
女 性	HbA1c 5.6%未満	心臓病の既往がある								
		はい	1	0.4	5	1.2	5	1.2	22	3.8
		いいえ	249	99.6	427	98.8	410	98.8	564	96.2
		—総数—	250	100.0	432	100.0	415	100.0	586	100.0
	HbA1c 5.6%以上 6.1%未満	心臓病の既往がある								
		はい	0	0	0	0	1	2.6	3	4.1
		いいえ	1	100.0	20	100.0	37	97.4	71	95.9
		—総数—	1	100.0	20	100.0	38	100.0	74	100.0
	HbA1c 6.1%以上または 現在治療を受けている	心臓病の既往がある								
はい		0	0	0	0	0	0	2	6.3	
いいえ		2	100.0	4	100.0	17	100.0	30	93.8	
	—総数—	2	100.0	4	100.0	17	100.0	32	100.0	

60-69歳		70歳以上		— 計 —		20-39歳(再掲)		40-59歳(再掲)		60歳以上(再掲)	
人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
68	8.0	130	17.0	262	6.1	9	0.8	55	3.4	198	12.3
783	92.0	633	83.0	4,020	93.9	1,063	99.2	1,541	96.6	1,416	87.7
851	100.0	763	100.0	4,282	100.0	1,072	100.0	1,596	100.0	1,614	100.0
18	10.1	30	16.3	56	10.0	0	0	8	4.8	48	13.2
161	89.9	154	83.7	506	90.0	31	100.0	160	95.2	315	86.8
179	100.0	184	100.0	562	100.0	31	100.0	168	100.0	363	100.0
33	19.0	39	22.0	76	15.8	0	0	4	3.3	72	20.5
141	81.0	138	78.0	404	84.2	8	100.0	117	96.7	279	79.5
174	100.0	177	100.0	480	100.0	8	100.0	121	100.0	351	100.0
33	8.8	52	17.5	116	7.0	3	0.8	28	4.7	85	12.7
340	91.2	245	82.5	1,539	93.0	387	99.2	567	95.3	585	87.3
373	100.0	297	100.0	1,655	100.0	390	100.0	595	100.0	670	100.0
11	15.1	14	18.4	29	13.5	0	0	4	7.1	25	16.8
62	84.9	62	81.6	186	86.5	10	100.0	52	92.9	124	83.2
73	100.0	76	100.0	215	100.0	10	100.0	56	100.0	149	100.0
16	16.3	19	18.6	37	13.5	0	0	2	2.8	35	17.5
82	83.7	83	81.4	237	86.5	2	100.0	70	97.2	165	82.5
98	100.0	102	100.0	274	100.0	2	100.0	72	100.0	200	100.0
35	7.3	78	16.7	146	5.6	6	0.9	27	2.7	113	12.0
443	92.7	388	83.3	2,481	94.4	676	99.1	974	97.3	831	88.0
478	100.0	466	100.0	2,627	100.0	682	100.0	1,001	100.0	944	100.0
7	6.6	16	14.8	27	7.8	0	0	4	3.6	23	10.7
99	93.4	92	85.2	320	92.2	21	100.0	108	96.4	191	89.3
106	100.0	108	100.0	347	100.0	21	100.0	112	100.0	214	100.0
17	22.4	20	26.7	39	18.9	0	0	2	4.1	37	24.5
59	77.6	55	73.3	167	81.1	6	100.0	47	95.9	114	75.5
76	100.0	75	100.0	206	100.0	6	100.0	49	100.0	151	100.0

第55表-2 糖尿病の状況別、心臓病および脳卒中の状況（脳卒中の既往、性・年齢階級別）

	糖尿病の状況	脳卒中の状況	20-29歳		30-39歳		40-49歳		50-59歳	
			人数	%	人数	%	人数	%	人数	%
男女計	HbA1c 5.6%未満	脳卒中の既往がある								
		はい	0	0	4	0.6	10	1.5	45	5.0
		いいえ	392	100.0	676	99.4	677	98.5	864	95.0
		—総数—	392	100.0	680	100.0	687	100.0	909	100.0
	HbA1c 5.6%以上 6.1%未満	脳卒中の既往がある								
		はい	0	0	0	0	0	0	2	1.7
		いいえ	4	100.0	27	100.0	48	100.0	118	98.3
		—総数—	4	100.0	27	100.0	48	100.0	120	100.0
	HbA1c 6.1%以上または 現在治療を受けている	脳卒中の既往がある								
はい		0	0	0	0	0	0	1	1.1	
いいえ		2	100.0	6	100.0	30	100.0	90	98.9	
	—総数—	2	100.0	6	100.0	30	100.0	91	100.0	
男 性	HbA1c 5.6%未満	脳卒中の既往がある								
		はい	0	0	3	1.2	3	1.1	5	1.5
		いいえ	142	100.0	245	98.8	269	98.9	318	98.5
		—総数—	142	100.0	248	100.0	272	100.0	323	100.0
	HbA1c 5.6%以上 6.1%未満	脳卒中の既往がある								
		はい	0	0	0	0	0	0	1	2.2
		いいえ	3	100.0	7	100.0	10	100.0	45	97.8
		—総数—	3	100.0	7	100.0	10	100.0	46	100.0
	HbA1c 6.1%以上または 現在治療を受けている	脳卒中の既往がある								
はい		0	0	0	0	0	0	0	0	
いいえ		0	0	2	100.0	13	100.0	59	100.0	
	—総数—	0	0	2	100.0	13	100.0	59	100.0	
女 性	HbA1c 5.6%未満	脳卒中の既往がある								
		はい	0	0	1	0.2	1	0.2	11	1.9
		いいえ	250	100.0	431	99.8	413	99.8	573	98.1
		—総数—	250	100.0	432	100.0	414	100.0	584	100.0
	HbA1c 5.6%以上 6.1%未満	脳卒中の既往がある								
		はい	0	0	0	0	0	0	1	1.4
		いいえ	1	100.0	20	100.0	38	100.0	73	98.6
		—総数—	1	100.0	20	100.0	38	100.0	74	100.0
	HbA1c 6.1%以上または 現在治療を受けている	脳卒中の既往がある								
はい		0	0	0	0	0	0	1	3.1	
いいえ		2	100.0	4	100.0	17	100.0	31	96.9	
	—総数—	2	100.0	4	100.0	17	100.0	32	100.0	

60-69歳		70歳以上		一計		20-39歳(再掲)		40-59歳(再掲)		60歳以上(再掲)	
人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
37	4.4	70	9.2	131	3.1	4	0.4	20	1.3	107	6.6
812	95.6	693	90.8	4,146	96.9	1,068	99.6	1,573	98.7	1,505	93.4
849	100.0	763	100.0	4,277	100.0	1,072	100.0	1,593	100.0	1,612	100.0
8	4.5	20	10.9	30	5.3	0	0	2	1.2	28	7.7
171	95.5	163	89.1	531	94.7	31	100.0	166	98.8	334	92.3
179	100.0	183	100.0	561	100.0	31	100.0	168	100.0	362	100.0
17	9.8	20	11.4	38	7.9	0	0	1	0.8	37	10.6
157	90.2	156	88.6	441	92.1	8	100.0	120	99.2	313	89.4
174	100.0	176	100.0	479	100.0	8	100.0	121	100.0	350	100.0
20	5.4	37	12.6	68	4.1	3	0.8	8	1.3	57	8.5
353	94.6	257	87.4	1,584	95.9	387	99.2	587	98.7	610	91.5
373	100.0	294	100.0	1,652	100.0	390	100.0	595	100.0	667	100.0
5	6.8	14	18.7	20	9.3	0	0	1	1.8	19	12.8
68	93.2	61	81.3	194	90.7	10	100.0	55	98.2	129	87.2
73	100.0	75	100.0	214	100.0	10	100.0	56	100.0	148	100.0
14	14.3	13	12.7	27	9.9	0	0	0	0	27	13.5
84	85.7	89	87.3	247	90.1	2	100.0	72	100.0	173	86.5
98	100.0	102	100.0	274	100.0	2	100.0	72	100.0	200	100.0
17	3.6	33	7.0	63	2.4	1	0.1	12	1.2	50	5.3
459	96.4	436	93.0	2,562	97.6	681	99.9	986	98.8	895	94.7
476	100.0	469	100.0	2,625	100.0	682	100.0	998	100.0	945	100.0
3	2.8	6	5.6	10	2.9	0	0	1	0.9	9	4.2
103	97.2	102	94.4	337	97.1	21	100.0	111	99.1	205	95.8
106	100.0	108	100.0	347	100.0	21	100.0	112	100.0	214	100.0
3	3.9	7	9.5	11	5.4	0	0	1	2.0	10	6.7
73	96.1	67	90.5	194	94.6	6	100.0	48	98.0	140	93.3
76	100.0	7475	100.0	205	100.0	6	100.0	49	100.0	150	100.0

第56表-1 糖尿病の状況別、血圧の状況（日本高血圧学会の分類、性・年齢階級別）

	糖尿病の状況	血圧の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男女計	HbA1c 5.6%未満	至適血圧	247	63.7	374	55.4	244	35.6	166	18.3	98	11.6	54	7.0	1,183	27.7
		正常血圧	82	21.1	151	22.4	152	22.2	177	19.6	114	13.5	85	11.1	761	17.8
		正常高値	39	10.1	92	13.6	113	16.5	202	22.3	184	21.7	137	17.9	767	18.0
		軽症高血圧	18	4.6	50	7.4	136	19.9	254	28.1	308	36.4	310	40.4	1,076	25.2
		中等度高血圧	2	0.5	6	0.9	30	4.4	78	8.6	110	13.0	141	18.4	367	8.6
		重症高血圧	0	0	2	0.3	10	1.5	28	3.1	33	3.9	40	5.2	113	2.6
		一総数	388	100.0	675	100.0	685	100.0	905	100.0	847	100.0	767	100.0	4,267	100.0
	HbA1c 5.6%以上 6.1%未満	至適血圧	2	50.0	11	40.7	13	26.5	18	15.3	19	10.6	17	9.2	80	14.2
		正常血圧	1	25.0	5	18.5	12	24.5	25	21.2	32	17.9	16	8.6	91	16.2
		正常高値	0	0	3	11.1	11	22.4	19	16.1	37	20.7	35	18.9	105	18.7
		軽症高血圧	1	25.0	6	22.2	9	18.4	40	33.9	57	31.8	72	38.9	185	32.9
		中等度高血圧	0	0	2	7.4	4	8.2	13	11.0	26	14.5	31	16.8	76	13.5
重症高血圧		0	0	0	0	0	0	3	2.5	8	4.5	14	7.6	25	4.4	
一総数		4	100.0	27	100.0	49	100.0	118	100.0	179	100.0	185	100.0	562	100.0	
HbA1c 6.1%以上 または現在治療を 受けている	至適血圧	2	100.0	1	16.7	6	20.0	8	8.7	12	7.0	4	2.3	33	6.9	
	正常血圧	0	0	1	16.7	4	13.3	12	13.0	16	9.3	12	6.8	45	9.4	
	正常高値	0	0	0	0	7	23.3	15	16.3	33	19.2	31	17.5	86	18.0	
	軽症高血圧	0	0	3	50.0	7	23.3	38	41.3	67	39.0	80	45.2	195	40.7	
	中等度高血圧	0	0	1	16.7	2	6.7	13	14.1	34	19.8	39	22.0	89	18.6	
	重症高血圧	0	0	0	0	4	13.3	6	6.5	10	5.8	11	6.2	31	6.5	
	一総数	2	100.0	6	100.0	30	100.0	92	100.0	172	100.0	177	100.0	479	100.0	
男 性	HbA1c 5.6%未満	至適血圧	57	40.4	68	27.5	59	21.8	33	10.3	29	7.8	20	6.7	266	16.1
		正常血圧	40	28.4	71	28.7	55	20.3	57	17.8	47	12.6	32	10.8	302	18.3
		正常高値	29	20.6	61	24.7	53	19.6	71	22.1	83	22.3	56	18.9	353	21.4
		軽症高血圧	14	9.9	42	17.0	79	29.2	112	34.9	150	40.2	120	40.4	517	31.3
		中等度高血圧	1	0.7	4	1.6	20	7.4	35	10.9	46	12.3	48	16.2	154	9.3
		重症高血圧	0	0	1	0.4	5	1.8	13	4.0	18	4.8	21	7.1	58	3.5
		一総数	141	100.0	247	100.0	271	100.0	321	100.0	373	100.0	297	100.0	1,650	100.0
	HbA1c 5.6%以上 6.1%未満	至適血圧	1	33.3	0	0	0	0	4	8.9	5	6.8	7	9.2	17	7.9
		正常血圧	1	33.3	2	28.6	5	50.0	11	24.4	10	13.7	11	14.5	40	18.7
		正常高値	0	0	1	14.3	2	20.0	8	17.8	13	17.8	16	21.1	40	18.7
		軽症高血圧	1	33.3	3	42.9	3	30.0	16	35.6	31	42.5	24	31.6	78	36.4
		中等度高血圧	0	0	1	14.3	0	0	5	11.1	13	17.8	12	15.8	31	14.5
重症高血圧		0	0	0	0	0	0	1	2.2	1	1.4	6	7.9	8	3.7	
一総数		3	100.0	7	100.0	10	100.0	45	100.0	73	100.0	76	100.0	214	100.0	
HbA1c 6.1%以上 または現在治療を 受けている	至適血圧	0	0	1	50.0	3	23.1	5	8.3	2	2.1	3	2.9	14	5.1	
	正常血圧	0	0	0	0	3	23.1	9	15.0	7	7.2	7	6.9	26	9.5	
	正常高値	0	0	0	0	3	23.1	14	23.3	19	19.6	18	17.6	54	19.7	
	軽症高血圧	0	0	0	0	2	15.4	20	33.3	37	38.1	50	49.0	109	39.8	
	中等度高血圧	0	0	1	50.0	0	0	7	11.7	25	25.8	19	18.6	52	19.0	
	重症高血圧	0	0	0	0	2	15.4	5	8.3	7	7.2	5	4.9	19	6.9	
	一総数	0	0	2	100.0	13	100.0	60	100.0	97	100.0	102	100.0	274	100.0	
女 性	HbA1c 5.6%未満	至適血圧	190	76.9	306	71.5	185	44.7	133	22.8	69	14.6	34	7.2	917	35.0
		正常血圧	42	17.0	80	18.7	97	23.4	120	20.5	67	14.1	53	11.3	459	17.5
		正常高値	10	4.0	31	7.2	60	14.5	131	22.4	101	21.3	81	17.2	414	15.8
		軽症高血圧	4	1.6	8	1.9	57	13.8	142	24.3	158	33.3	190	40.4	559	21.4
		中等度高血圧	1	0.4	2	0.5	10	2.4	43	7.4	64	13.5	93	19.8	213	8.1
		重症高血圧	0	0	1	0.2	5	1.2	15	2.6	15	3.2	19	4.0	55	2.1
		一総数	247	100.0	428	100.0	414	100.0	584	100.0	474	100.0	470	100.0	2,617	100.0
	HbA1c 5.6%以上 6.1%未満	至適血圧	1	100.0	11	55.0	13	33.3	14	19.2	14	13.2	10	9.2	63	18.1
		正常血圧	0	0	3	15.0	7	17.9	14	19.2	22	20.8	5	4.6	51	14.7
		正常高値	0	0	2	10.0	9	23.1	11	15.1	24	22.6	19	17.4	65	18.7
		軽症高血圧	0	0	3	15.0	6	15.4	24	32.9	26	24.5	48	44.0	107	30.7
		中等度高血圧	0	0	1	5.0	4	10.3	8	11.0	13	12.3	19	17.4	45	12.9
重症高血圧		0	0	0	0	0	0	2	2.7	7	6.6	8	7.3	17	4.9	
一総数		1	100.0	20	100.0	39	100.0	73	100.0	106	100.0	109	100.0	348	100.0	
HbA1c 6.1%以上 または現在治療を 受けている	至適血圧	2	100.0	0	0	3	17.6	3	9.4	10	13.3	1	1.3	19	9.3	
	正常血圧	0	0	1	25.0	1	5.9	3	9.4	9	12.0	5	6.7	19	9.3	
	正常高値	0	0	0	0	4	23.5	1	3.1	14	18.7	13	17.3	32	15.6	
	軽症高血圧	0	0	3	75.0	5	29.4	18	56.3	30	40.0	30	40.0	86	42.0	
	中等度高血圧	0	0	0	0	2	11.8	6	18.8	9	12.0	20	26.7	37	18.0	
	重症高血圧	0	0	0	0	2	11.8	1	3.1	3	4.0	6	8.0	12	5.9	
	一総数	2	100.0	4	100.0	17	100.0	32	100.0	75	100.0	75	100.0	205	100.0	

第56表-2 糖尿病の状況別、血圧の状況（WHOの分類、性・年齢階級別）

	糖尿病の状況	血圧の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		— 計 —	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男女計	HbA1c 5.6%未満	正常血圧	368	94.8	617	91.4	509	74.3	545	60.2	396	46.8	276	36.0	2,711	63.5
		境界域血圧	16	4.1	36	5.3	90	13.1	201	22.2	264	31.2	293	38.2	900	21.1
		高血圧	4	1.0	22	3.3	86	12.6	159	17.6	187	22.1	198	25.8	656	15.4
		— 総数 —	388	100.0	675	100.0	685	100.0	905	100.0	847	100.0	767	100.0	4,267	100.0
	HbA1c 5.6%以上 6.1%未満	正常血圧	3	75.0	19	70.4	36	73.5	62	52.5	88	49.2	68	36.8	276	49.1
		境界域血圧	1	25.0	5	18.5	6	12.2	35	29.7	48	26.8	67	36.2	162	28.8
		高血圧	0	0	3	11.1	7	14.3	21	17.8	43	24.0	50	27.0	124	22.1
		— 総数 —	4	100.0	27	100.0	49	100.0	118	100.0	179	100.0	185	100.0	562	100.0
	HbA1c 6.1%以上 または現在治療を 受けている	正常血圧	2	100.0	2	33.3	17	56.7	35	38.0	61	35.5	47	26.6	164	34.2
		境界域血圧	0	0	3	50.0	6	20.0	33	35.9	61	35.5	78	44.1	181	37.8
		高血圧	0	0	1	16.7	7	23.3	24	26.1	50	29.1	52	29.4	134	28.0
		— 総数 —	2	100.0	6	100.0	30	100.0	92	100.0	172	100.0	177	100.0	479	100.0
男 性	HbA1c 5.6%未満	正常血圧	126	89.4	200	81.0	167	61.6	161	50.2	159	42.6	108	36.4	921	55.8
		境界域血圧	12	8.5	32	13.0	49	18.1	88	27.4	128	34.3	113	38.0	422	25.6
		高血圧	3	2.1	15	6.1	55	20.3	72	22.4	86	23.1	76	25.6	307	18.6
		— 総数 —	141	100.0	247	100.0	271	100.0	321	100.0	373	100.0	297	100.0	1,650	100.0
	HbA1c 5.6%以上 6.1%未満	正常血圧	2	66.7	3	42.9	7	70.0	23	51.1	28	38.4	34	44.7	97	45.3
		境界域血圧	1	33.3	2	28.6	2	20.0	13	28.9	26	35.6	22	28.9	66	30.8
		高血圧	0	0	2	28.6	1	10.0	9	20.0	19	26.0	20	26.3	51	23.8
		— 総数 —	3	100.0	7	100.0	10	100.0	45	100.0	73	100.0	76	100.0	214	100.0
	HbA1c 6.1%以上 または現在治療を 受けている	正常血圧	0	0	1	50.0	9	69.2	28	46.7	28	28.9	28	27.5	94	34.3
		境界域血圧	0	0	0	0	2	15.4	16	26.7	35	36.1	48	47.1	101	36.9
		高血圧	0	0	1	50.0	2	15.4	16	26.7	34	35.1	26	25.5	79	28.8
		— 総数 —	0	0	2	100.0	13	100.0	60	100.0	97	100.0	102	100.0	274	100.0
女 性	HbA1c 5.6%未満	正常血圧	242	98.0	417	97.4	342	82.6	384	65.8	237	50.0	168	35.7	1,790	68.4
		境界域血圧	4	1.6	4	0.9	41	9.9	113	19.3	136	28.7	180	38.3	478	18.3
		高血圧	1	0.4	7	1.6	31	7.5	87	14.9	101	21.3	122	26.0	349	13.3
		— 総数 —	247	100.0	428	100.0	414	100.0	584	100.0	474	100.0	470	100.0	2,617	100.0
	HbA1c 5.6%以上 6.1%未満	正常血圧	1	100.0	16	80.0	29	74.4	39	53.4	60	56.6	34	31.2	179	51.4
		境界域血圧	0	0	3	15.0	4	10.3	22	30.1	22	20.8	45	41.3	96	27.6
		高血圧	0	0	1	5.0	6	15.4	12	16.4	24	22.6	30	27.5	73	21.0
		— 総数 —	1	100.0	20	100.0	39	100.0	73	100.0	106	100.0	109	100.0	348	100.0
	HbA1c 6.1%以上 または現在治療を 受けている	正常血圧	2	100.0	1	25.0	8	47.1	7	21.9	33	44.0	19	25.3	70	34.1
		境界域血圧	0	0	3	75.0	4	23.5	17	53.1	26	34.7	30	40.0	80	39.0
		高血圧	0	0	0	0	5	29.4	8	25.0	16	21.3	26	34.7	55	26.8
		— 総数 —	2	100.0	4	100.0	17	100.0	32	100.0	75	100.0	75	100.0	205	100.0

第57表-1 現在の肥満度別、糖尿病の状況（性・年齢階級別）

	現在の肥満度(BMI)	糖尿病の状況	20-29歳		30-39歳		40-49歳		50-59歳		
			人数	%	人数	%	人数	%	人数	%	
男女計	18.5未満	HbA1c 5.6%未満	63	98.4	66	98.5	37	100.0	39	90.7	
		HbA1c 5.6%以上6.1%未満	1	1.6	1	1.5	0	0	4	9.3	
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	0	0	
		—総数—	64	100.0	67	100.0	37	100.0	43	100.0	
	18.5以上22未満	HbA1c 5.6%未満	188	98.4	302	96.2	234	91.4	307	89.2	
		HbA1c 5.6%以上6.1%未満	2	1.0	12	3.8	17	6.6	26	7.6	
		HbA1c 6.1%以上または現在治療を受けている	1	0.5	0	0	5	2.0	11	3.2	
		—総数—	191	100.0	314	100.0	256	100.0	344	100.0	
	22以上25未満	HbA1c 5.6%未満	78	100.0	174	97.2	257	94.1	316	82.3	
		HbA1c 5.6%以上6.1%未満	0	0	5	2.8	11	4.0	40	10.4	
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	5	1.8	28	7.3	
		—総数—	78	100.0	179	100.0	273	100.0	384	100.0	
	25以上30未満	HbA1c 5.6%未満	44	100.0	113	92.6	139	81.8	218	72.2	
		HbA1c 5.6%以上6.1%未満	0	0	5	4.1	16	9.4	43	14.2	
		HbA1c 6.1%以上または現在治療を受けている	0	0	4	3.3	15	8.8	41	13.6	
		—総数—	44	100.0	122	100.0	170	100.0	302	100.0	
	30以上	HbA1c 5.6%未満	15	88.2	21	77.8	18	64.3	25	59.5	
		HbA1c 5.6%以上6.1%未満	1	5.9	4	14.8	5	17.9	5	11.9	
		HbA1c 6.1%以上または現在治療を受けている	1	5.9	2	7.4	5	17.9	12	28.6	
		—総数—	17	100.0	27	100.0	28	100.0	42	100.0	
	男性	18.5未満	HbA1c 5.6%未満	11	100.0	7	100.0	6	100.0	6	75.0
			HbA1c 5.6%以上6.1%未満	0	0	0	0	0	0	2	25.0
			HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	0	0
			—総数—	11	100.0	7	100.0	6	100.0	8	100.0
18.5以上22未満		HbA1c 5.6%未満	55	96.5	68	100.0	53	94.6	96	85.0	
		HbA1c 5.6%以上6.1%未満	2	3.5	0	0	2	3.6	8	7.1	
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	1.8	9	8.0	
		—総数—	57	100.0	68	100.0	56	100.0	113	100.0	
22以上25未満		HbA1c 5.6%未満	40	100.0	90	97.8	124	96.1	111	76.6	
		HbA1c 5.6%以上6.1%未満	0	0	2	2.2	1	0.8	17	11.7	
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	4	3.1	17	11.7	
		—総数—	40	100.0	92	100.0	129	100.0	145	100.0	
25以上30未満		HbA1c 5.6%未満	27	100.0	69	94.5	78	85.7	100	69.4	
		HbA1c 5.6%以上6.1%未満	0	0	3	4.1	7	7.7	16	11.1	
		HbA1c 6.1%以上または現在治療を受けている	0	0	1	1.4	6	6.6	28	19.4	
		—総数—	27	100.0	73	100.0	91	100.0	144	100.0	
30以上		HbA1c 5.6%未満	8	88.9	13	81.3	10	83.3	8	50	
		HbA1c 5.6%以上6.1%未満	1	11.1	2	12.5	0	0	2	12.5	
		HbA1c 6.1%以上または現在治療を受けている	0	0	1	6.3	2	16.7	6	37.5	
		—総数—	9	100.0	16	100.0	12	100.0	16	100.0	
女性		18.5未満	HbA1c 5.6%未満	52	98.1	59	98.3	31	100.0	33	94.3
			HbA1c 5.6%以上6.1%未満	1	1.9	1	1.7	0	0	2	5.7
			HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	0	0
			—総数—	53	100.0	60	100.0	31	100.0	35	100.0
	18.5以上22未満	HbA1c 5.6%未満	133	99.3	234	95.1	181	90.5	211	91.3	
		HbA1c 5.6%以上6.1%未満	0	0	12	4.9	15	7.5	18	7.8	
		HbA1c 6.1%以上または現在治療を受けている	1	0.7	0	0	4	2.0	2	0.9	
		—総数—	134	100.0	246	100.0	200	100.0	231	100.0	
	22以上25未満	HbA1c 5.6%未満	38	100.0	84	96.6	133	92.4	205	85.8	
		HbA1c 5.6%以上6.1%未満	0	0	3	3.4	10	6.9	23	9.6	
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	0.7	11	4.6	
		—総数—	38	100.0	87	100.0	144	100.0	239	100.0	
	25以上30未満	HbA1c 5.6%未満	17	100.0	44	89.8	61	77.2	118	74.7	
		HbA1c 5.6%以上6.1%未満	0	0	2	4.1	9	11.4	27	17.1	
		HbA1c 6.1%以上または現在治療を受けている	0	0	3	6.1	9	11.4	13	8.2	
		—総数—	17	100.0	49	100.0	79	100.0	158	100.0	
	30以上	HbA1c 5.6%未満	7	87.5	8	72.7	8	50.0	17	65.4	
		HbA1c 5.6%以上6.1%未満	0	0	2	18.2	5	31.3	3	11.5	
		HbA1c 6.1%以上または現在治療を受けている	1	12.5	1	9.1	3	18.8	6	23.1	
		—総数—	8	100.0	11	100.0	16	100.0	26	100.0	

60-69歲		70歲以上		—計—		20-39歲(再掲)		40-59歲(再掲)		60歲以上(再掲)	
人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
26	74.3	66	82.5	297	91.1	129	98.5	76	95.0	92	80.0
1	2.9	6	7.5	13	4.0	2	1.5	4	5.0	7	6.1
8	22.9	8	10.0	16	4.9	0	0	0	0	16	13.9
35	100.0	80	100.0	326	100.0	131	100.0	80	100.0	115	100.0
234	77.5	217	72.3	1,482	86.8	490	97.0	541	90.2	451	74.9
35	11.6	41	13.7	133	7.8	14	2.8	43	7.2	76	12.6
33	10.9	42	14.0	92	5.4	1	0.2	16	2.7	75	12.5
302	100.0	300	100.0	1,707	100.0	505	100.0	600	100.0	602	100.0
336	73.2	277	69.9	1,438	81.3	252	98.1	573	87.2	613	71.7
52	11.3	59	14.9	167	9.4	5	1.9	51	7.8	111	13.0
71	15.5	60	15.2	164	9.3	0	0	33	5.0	131	15.3
459	100.0	396	100.0	1,769	100.0	257	100.0	657	100.0	855	100.0
234	65.2	180	58.3	928	71.1	157	94.6	357	75.6	414	62.0
79	22.0	67	21.7	210	16.1	5	3.0	59	12.5	146	21.9
46	12.8	62	20.1	168	12.9	4	2.4	56	11.9	108	16.2
359	100.0	309	100.0	1,306	100.0	166	100.0	472	100.0	668	100.0
17	40.5	25	61.0	121	61.4	36	81.8	43	61.4	42	50.6
12	28.6	12	29.3	39	19.8	5	11.4	10	14.3	24	28.9
13	31.0	4	9.8	37	18.8	3	6.8	17	24.3	17	20.5
42	100.0	41	100.0	197	100.0	44	100.0	70	100.0	83	100.0
9	69.2	28	84.8	67	85.9	18	100.0	12	85.7	37	80.4
1	7.7	1	3.0	4	5.1	0	0	2	14.3	2	4.3
3	23.1	4	12.1	7	9.0	0	0	0	0	7	15.2
13	100.0	33	100.0	78	100.0	18	100.0	14	100.0	46	100.0
95	77.2	88	63.3	455	81.8	123	98.4	149	88.2	183	69.8
10	8.1	21	15.1	43	7.7	2	1.6	10	5.9	31	11.8
18	14.6	30	21.6	58	10.4	0	0	10	5.9	48	18.3
123	100.0	139	100.0	556	100.0	125	100.0	169	100.0	262	100.0
161	69.4	108	62.4	634	78.2	130	98.5	235	85.8	269	66.4
25	10.8	28	16.2	73	9.0	2	1.5	18	6.6	53	13.1
46	19.8	37	21.4	104	12.8	0	0	21	7.7	83	20.5
232	100.0	173	100.0	811	100.0	132	100.0	274	100.0	405	100.0
102	63.0	68	55.7	444	71.7	96	96.0	178	75.7	170	59.9
34	21.0	23	18.9	83	13.4	3	3.0	23	9.8	57	20.1
26	16.0	31	25.4	92	14.9	1	1.0	34	14.5	57	20.1
162	100.0	122	100.0	619	100.0	100	100.0	235	100.0	284	100.0
6	46.2	4	57.1	49	67.1	21	84.0	18	64.3	10	50.0
3	23.1	3	42.9	11	15.1	3	12.0	2	7.1	6	30.0
4	30.8	0	0	13	17.8	1	4.0	8	28.6	4	20.0
13	100.0	7	100.0	73	100.0	25	100.0	28	100.0	20	100.0
17	77.3	38	80.9	230	92.7	111	98.2	64	97.0	55	79.7
0	0	5	10.6	9	3.6	2	1.8	2	3.0	5	7.2
5	22.7	4	8.5	9	3.6	0	0	0	0	9	13.0
22	100.0	47	100.0	248	100.0	113	100.0	66	100.0	69	100.0
139	77.7	129	80.1	1,027	89.2	367	96.6	392	91.0	268	78.8
25	14.0	20	12.4	90	7.8	12	3.2	33	7.7	45	13.2
15	8.4	12	7.5	34	3.0	1	0.3	6	1.4	27	7.9
179	100.0	161	100.0	1,151	100.0	380	100.0	431	100.0	340	100.0
175	77.1	169	75.8	804	83.9	122	97.6	338	88.3	344	76.4
27	11.9	31	13.9	94	9.8	3	2.4	33	8.6	58	12.9
25	11.0	23	10.3	60	6.3	0	0	12	3.1	48	10.7
227	100.0	223	100.0	958	100.0	125	100.0	383	100.0	450	100.0
132	67.0	112	59.9	484	70.5	61	92.4	179	75.5	244	63.5
45	22.8	44	23.5	127	18.5	2	3.0	36	15.2	89	23.2
20	10.2	31	16.6	76	11.1	3	4.5	22	9.3	51	13.3
197	100.0	187	100.0	687	100.0	66	100.0	237	100.0	384	100.0
11	37.9	21	61.8	72	58.1	15	78.9	25	59.5	32	50.8
9	31.0	9	26.5	28	22.6	2	10.5	8	19.0	18	28.6
9	31.0	4	11.8	24	19.4	2	10.5	9	21.4	13	20.6
29	100.0	34	100.0	124	100.0	19	100.0	42	100.0	63	100.0

第57表-2 現在の肥満度別、糖尿病の状況（男女計、年齢階級別）

現在の肥満度	糖尿病の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
-10%未満	HbA1c 5.6%未満	135	98.5	177	97.3	97	96.0	121	92.4	84	77.8	124	76.5	738	89.9
	HbA1c 5.6%以上6.1%未満	2	1.5	5	2.7	4	4.0	10	7.6	8	7.4	19	11.7	48	5.8
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	0	0	16	14.8	19	11.7	35	4.3
	—総数—	137	100.0	182	100.0	101	100.0	131	100.0	108	100.0	162	100.0	821	100.0
-10%以上 0%未満	HbA1c 5.6%未満	116	98.3	191	96.0	174	90.6	225	87.9	176	76.9	159	72.9	1,041	85.9
	HbA1c 5.6%以上6.1%未満	1	0.8	8	4.0	13	6.8	20	7.8	28	12.2	28	12.8	98	8.1
	HbA1c 6.1%以上または現在治療を受けている	1	0.8	0	0	5	2.6	11	4.3	25	10.9	31	14.2	73	6.0
	—総数—	118	100.0	199	100.0	192	100.0	256	100.0	229	100.0	218	100.0	1,212	100.0
0%以上 +10%未満	HbA1c 5.6%未満	65	100.0	137	97.2	202	94.0	248	84.4	248	75.2	203	72.0	1,103	83.1
	HbA1c 5.6%以上6.1%未満	0	0	4	2.8	10	4.7	32	10.9	28	8.5	40	14.2	114	8.6
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	3	1.4	14	4.8	54	16.4	39	13.8	110	8.3
	—総数—	65	100.0	141	100.0	215	100.0	294	100.0	330	100.0	282	100.0	1,327	100.0
+10%以上 +20%未満	HbA1c 5.6%未満	32	100.0	93	97.9	123	88.5	181	79.0	204	67.8	172	65.2	805	75.9
	HbA1c 5.6%以上6.1%未満	0	0	1	1.1	10	7.2	23	10.0	58	19.3	45	17.0	137	12.9
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	1.1	6	4.3	25	10.9	39	13.0	47	17.8	118	11.1
	—総数—	32	100.0	95	100.0	139	100.0	229	100.0	301	100.0	264	100.0	1,060	100.0
+20%以上 +30%未満	HbA1c 5.6%未満	20	100.0	40	85.1	56	86.2	85	67.5	94	61.8	68	55.7	363	68.2
	HbA1c 5.6%以上6.1%未満	0	0	4	8.5	4	6.2	19	15.1	36	23.7	29	23.8	92	17.3
	HbA1c 6.1%以上または現在治療を受けている	0	0	3	6.4	5	7.7	22	17.5	22	14.5	25	20.5	77	14.5
	—総数—	20	100.0	47	100.0	65	100.0	126	100.0	152	100.0	122	100.0	532	100.0
+30%以上 +40%未満	HbA1c 5.6%未満	7	87.5	25	92.6	24	70.6	31	62.0	31	66.0	21	46.7	139	65.9
	HbA1c 5.6%以上6.1%未満	1	12.5	2	7.4	4	11.8	10	20.0	10	21.3	12	26.7	39	18.5
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	6	17.6	9	18.0	6	12.8	12	26.7	33	15.6
	—総数—	8	100.0	27	100.0	34	100.0	50	100.0	47	100.0	45	100.0	211	100.0
+40%以上	HbA1c 5.6%未満	13	92.9	13	72.2	9	50.0	14	48.3	10	33.3	18	54.5	77	54.2
	HbA1c 5.6%以上6.1%未満	0	0	3	16.7	4	22.2	4	13.8	11	36.7	12	36.4	34	23.9
	HbA1c 6.1%以上または現在治療を受けている	1	7.1	2	11.1	5	27.8	11	37.9	9	30.0	3	9.1	31	21.8
	—総数—	14	100.0	18	100.0	18	100.0	29	100.0	30	100.0	33	100.0	142	100.0

第57表-3 現在の肥満度別、糖尿病の状況（男性、年齢階級別）

現在の肥満度	糖尿病の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		—計—	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
-10%未満	HbA1c 5.6%未満	24	96.0	20	100.0	13	100.0	35	92.1	32	72.7	49	69.0	173	82.0
	HbA1c 5.6%以上6.1%未満	1	4.0	0	0	0	0	3	7.9	3	6.8	10	14.1	17	8.1
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	0	0	9	20.5	12	16.9	21	10.0
	—総数—	25	100.0	20	100.0	13	100.0	38	100.0	44	100.0	71	100.0	211	100.0
-10%以上0%未満	HbA1c 5.6%未満	42	97.7	55	100.0	46	93.9	67	80.7	72	78.3	67	66.3	349	82.5
	HbA1c 5.6%以上6.1%未満	1	2.3	0	0	2	4.1	7	8.4	8	8.7	12	11.9	30	7.1
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	2.0	9	10.8	12	13.0	22	21.8	44	10.4
	—総数—	43	100.0	55	100.0	49	100.0	83	100.0	92	100.0	101	100.0	423	100.0
0%以上+10%未満	HbA1c 5.6%未満	31	100.0	68	97.1	93	96.9	86	81.1	120	71.0	77	65.3	475	80.5
	HbA1c 5.6%以上6.1%未満	0	0	2	2.9	1	1.0	14	13.2	14	8.3	18	15.3	49	8.3
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	2	2.1	6	5.7	35	20.7	23	19.5	66	11.2
	—総数—	31	100.0	70	100.0	96	100.0	106	100.0	169	100.0	118	100.0	590	100.0
+10%以上+20%未満	HbA1c 5.6%未満	18	100.0	57	100.0	70	90.9	83	74.8	95	65.1	72	61.0	395	75.0
	HbA1c 5.6%以上6.1%未満	0	0	0	0	4	5.2	10	9.0	24	16.4	18	15.3	56	10.6
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	3	3.9	18	16.2	27	18.5	28	23.7	76	14.4
	—総数—	18	100.0	57	100.0	77	100.0	111	100.0	146	100.0	118	100.0	527	100.0
+20%以上+30%未満	HbA1c 5.6%未満	14	100.0	21	87.5	32	88.9	33	61.1	37	58.7	23	47.9	160	66.9
	HbA1c 5.6%以上6.1%未満	0	0	2	8.3	2	5.6	6	11.1	18	28.6	12	25.0	40	16.7
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	4.2	2	5.6	15	27.8	8	12.7	13	27.1	39	16.3
	—総数—	14	100.0	24	100.0	36	100.0	54	100.0	63	100.0	48	100.0	239	100.0
+30%以上+40%未満	HbA1c 5.6%未満	5	83.3	17	89.5	13	76.5	13	56.5	12	66.7	4	36.4	64	68.1
	HbA1c 5.6%以上6.1%未満	1	16.7	2	10.5	1	5.9	4	17.4	3	16.7	3	27.3	14	14.9
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	3	17.6	6	26.1	3	16.7	4	36.4	16	17.0
	—総数—	6	100.0	19	100.0	17	100.0	23	100.0	18	100.0	11	100.0	94	100.0
+40%以上	HbA1c 5.6%未満	7	100.0	9	81.8	4	66.7	4	36.4	5	45.5	4	57.1	33	62.3
	HbA1c 5.6%以上6.1%未満	0	0	1	9.1	0	0	1	9.1	3	27.3	3	42.9	8	15.1
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	9.1	2	33.3	6	54.5	3	27.3	0	0	12	22.6
	—総数—	7	100.0	11	100.0	6	100.0	11	100.0	11	100.0	7	100.0	53	100.0

第57表-4 現在の肥満度別、糖尿病の状況（女性、年齢階級別）

現在の肥満度	糖尿病の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
-10%未満	HbA1c 5.6%未満	111	99.1	157	96.9	84	95.5	86	92.5	52	81.3	75	82.4	565	92.6
	HbA1c 5.6%以上6.1%未満	1	0.9	5	3.1	4	4.5	7	7.5	5	7.8	9	9.9	31	5.1
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	0	0	7	10.9	7	7.7	14	2.3
	—総数—	112	100.0	162	100.0	88	100.0	93	100.0	64	100.0	91	100.0	610	100.0
-10%以上0%未満	HbA1c 5.6%未満	74	98.7	136	94.4	128	89.5	158	91.3	104	75.9	92	78.6	692	87.7
	HbA1c 5.6%以上6.1%未満	0	0	8	5.6	11	7.7	13	7.5	20	14.6	16	13.7	68	8.6
	HbA1c 6.1%以上または現在治療を受けている	1	1.3	0	0	4	2.8	2	1.2	13	9.5	9	7.7	29	3.7
	—総数—	75	100.0	144	100.0	143	100.0	173	100.0	137	100.0	117	100.0	789	100.0
0%以上+10%未満	HbA1c 5.6%未満	34	100.0	69	97.2	109	91.6	162	86.2	128	79.5	126	76.8	628	85.2
	HbA1c 5.6%以上6.1%未満	0	0	2	2.8	9	7.6	18	9.6	14	8.7	22	13.4	65	8.8
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	0.8	8	4.3	19	11.8	16	9.8	44	6.0
	—総数—	34	100.0	71	100.0	119	100.0	188	100.0	161	100.0	164	100.0	737	100.0
+10%以上+20%未満	HbA1c 5.6%未満	14	100.0	36	94.7	53	85.5	98	83.1	109	70.3	100	68.5	410	76.9
	HbA1c 5.6%以上6.1%未満	0	0	1	2.6	6	9.7	13	11.0	34	21.9	27	18.5	81	15.2
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	2.6	3	4.8	7	5.9	12	7.7	19	13.0	42	7.9
	—総数—	14	100.0	38	100.0	62	100.0	118	100.0	155	100.0	146	100.0	533	100.0
+20%以上+30%未満	HbA1c 5.6%未満	6	100.0	19	82.6	24	82.8	52	72.2	57	64.0	45	60.8	203	69.3
	HbA1c 5.6%以上6.1%未満	0	0	2	8.7	2	6.9	13	18.1	18	20.2	17	23.0	52	17.7
	HbA1c 6.1%以上または現在治療を受けている	0	0	2	8.7	3	10.3	7	9.7	14	15.7	12	16.2	38	13.0
	—総数—	6	100.0	23	100.0	29	100.0	72	100.0	89	100.0	74	100.0	293	100.0
+30%以上+40%未満	HbA1c 5.6%未満	2	100.0	8	100.0	11	64.7	18	66.7	19	65.5	17	50.0	75	64.1
	HbA1c 5.6%以上6.1%未満	0	0	0	0	3	17.6	6	22.2	7	24.1	9	26.5	25	21.4
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	3	17.6	3	11.1	3	10.3	8	23.5	17	14.5
	—総数—	2	100.0	8	100.0	17	100.0	27	100.0	29	100.0	34	100.0	117	100.0
+40%以上	HbA1c 5.6%未満	6	85.7	4	57.1	5	41.7	10	55.6	5	26.3	14	53.8	44	49.4
	HbA1c 5.6%以上6.1%未満	0	0	2	28.6	4	33.3	3	16.7	8	42.1	9	34.6	26	29.2
	HbA1c 6.1%以上または現在治療を受けている	1	14.3	1	14.3	3	25.0	5	27.8	6	31.6	3	11.5	19	21.3
	—総数—	7	100.0	7	100.0	12	100.0	18	100.0	19	100.0	26	100.0	89	100.0

第58表-1 1日の運動量（歩行数）別、糖尿病の状況（男女計、年齢階級別）

1日の運動量 (歩行数)	糖尿病の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		—計—	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
2,000歩未満	HbA1c 5.6%未満	11	100.0	10	100.0	12	75.0	21	75.0	33	62.3	153	63.5	240	66.9
	HbA1c 5.6%以上6.1%未満	0	0	0	0	3	18.8	4	14.3	8	15.1	44	18.3	59	16.4
	HbA1c 6.1%以上または 現在治療を受けている	0	0	0	0	1	6.3	3	10.7	12	22.6	44	18.3	60	16.7
	—総数—	11	100.0	10	100.0	16	100.0	28	100.0	53	100.0	241	100.0	359	100.0
2,000 -3,999歩	HbA1c 5.6%未満	42	97.7	62	91.2	67	94.4	109	74.7	115	70.6	167	64.7	562	75.0
	HbA1c 5.6%以上6.1%未満	0	0	5	7.4	2	2.8	18	12.3	24	14.7	41	15.9	90	12.0
	HbA1c 6.1%以上または 現在治療を受けている	1	2.3	1	1.5	2	2.8	19	13.0	24	14.7	50	19.4	97	13.0
	—総数—	43	100.0	68	100.0	71	100.0	146	100.0	163	100.0	258	100.0	749	100.0
4,000 -5,999歩	HbA1c 5.6%未満	82	98.8	130	95.6	133	88.7	172	85.1	162	68.4	161	69.4	840	80.8
	HbA1c 5.6%以上6.1%未満	1	1.2	6	4.4	10	6.7	15	7.4	43	18.1	43	18.5	118	11.3
	HbA1c 6.1%以上または 現在治療を受けている	0	0	0	0	7	4.7	15	7.4	32	13.5	28	12.1	82	7.9
	—総数—	83	100.0	136	100.0	150	100.0	202	100.0	237	100.0	232	100.0	1,040	100.0
6,000 -7,999歩	HbA1c 5.6%未満	85	100.0	156	97.5	140	88.6	177	81.9	172	73.5	127	75.6	857	83.9
	HbA1c 5.6%以上6.1%未満	0	0	2	1.3	11	7.0	21	9.7	30	12.8	20	11.9	84	8.2
	HbA1c 6.1%以上または 現在治療を受けている	0	0	2	1.3	7	4.4	18	8.3	32	13.7	21	12.5	80	7.8
	—総数—	85	100.0	160	100.0	158	100.0	216	100.0	234	100.0	168	100.0	1,021	100.0
8,000 -9,999歩	HbA1c 5.6%未満	58	98.3	118	95.9	125	89.9	156	80.0	144	75.4	68	69.4	669	83.1
	HbA1c 5.6%以上6.1%未満	0	0	5	4.1	11	7.9	27	13.8	23	12.0	16	16.3	82	10.2
	HbA1c 6.1%以上または 現在治療を受けている	1	1.7	0	0	3	2.2	12	6.2	24	12.6	14	14.3	54	6.7
	—総数—	59	100.0	123	100.0	139	100.0	195	100.0	191	100.0	98	100.0	805	100.0
10,000歩 以上	HbA1c 5.6%未満	98	97.0	176	94.1	197	90.0	249	81.1	210	69.5	65	67.0	995	82.0
	HbA1c 5.6%以上6.1%未満	3	3.0	8	4.3	12	5.5	34	11.1	50	16.6	15	15.5	122	10.1
	HbA1c 6.1%以上または 現在治療を受けている	0	0	3	1.6	10	4.6	24	7.8	42	13.9	17	17.5	96	7.9
	—総数—	101	100.0	187	100.0	219	100.0	307	100.0	302	100.0	97	100.0	1,213	100.0

第58表-2 1日の運動量（歩行数）別、糖尿病の状況（男性、年齢階級別）

1日の運動量 (歩行数)	糖尿病の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
2,000歩未満	HbA1c 5.6%未満	5	100.0	4	100.0	7	100.0	12	66.7	14	51.9	54	56.3	96	61.1
	HbA1c 5.6%以上6.1%未満	0	0	0	0	0	0	3	16.7	3	11.1	17	17.7	23	14.6
	HbA1c 6.1%以上または 現在治療を受けている	0	0	0	0	0	0	3	16.7	10	37.0	25	26.0	38	24.2
	—総数—	5	100.0	4	100.0	7	100.0	18	100.0	27	100.0	96	100.0	157	100.0
2,000 -3,999歩	HbA1c 5.6%未満	16	100.0	18	85.7	28	96.6	39	62.9	52	67.5	62	57.9	215	68.9
	HbA1c 5.6%以上6.1%未満	0	0	2	9.5	1	3.4	9	14.5	15	19.5	19	17.8	46	14.7
	HbA1c 6.1%以上または 現在治療を受けている	0	0	1	4.8	0	0	14	22.6	10	13.0	26	24.3	51	16.3
	—総数—	16	100.0	21	100.0	29	100.0	62	100.0	77	100.0	107	100.0	312	100.0
4,000 -5,999歩	HbA1c 5.6%未満	27	96.4	40	95.2	51	94.4	62	75.6	69	66.3	61	64.9	310	76.7
	HbA1c 5.6%以上6.1%未満	1	3.6	2	4.8	1	1.9	10	12.2	17	16.3	16	17.0	47	11.6
	HbA1c 6.1%以上または 現在治療を受けている	0	0	0	0	2	3.7	10	12.2	18	17.3	17	18.1	47	11.6
	—総数—	28	100.0	42	100.0	54	100.0	82	100.0	104	100.0	94	100.0	404	100.0
6,000 -7,999歩	HbA1c 5.6%未満	24	100.0	52	96.3	50	92.6	65	82.3	62	68.1	51	71.8	304	81.5
	HbA1c 5.6%以上6.1%未満	0	0	1	1.9	2	3.7	5	6.3	11	12.1	9	12.7	28	7.5
	HbA1c 6.1%以上または 現在治療を受けている	0	0	1	1.9	2	3.7	9	11.4	18	19.8	11	15.5	41	11.0
	—総数—	24	100.0	54	100.0	54	100.0	79	100.0	91	100.0	71	100.0	373	100.0
8,000 -9,999歩	HbA1c 5.6%未満	17	100.0	38	100.0	52	91.2	55	78.6	55	70.5	27	62.8	244	80.5
	HbA1c 5.6%以上6.1%未満	0	0	0	0	2	3.5	8	11.4	9	11.5	8	18.6	27	8.9
	HbA1c 6.1%以上または 現在治療を受けている	0	0	0	0	3	5.3	7	10.0	14	17.9	8	18.6	32	10.6
	—総数—	17	100.0	38	100.0	57	100.0	70	100.0	78	100.0	43	100.0	303	100.0
10,000歩 以上	HbA1c 5.6%未満	47	95.9	83	98.8	76	88.4	77	74.0	117	74.1	29	64.4	429	81.6
	HbA1c 5.6%以上6.1%未満	2	4.1	1	1.2	4	4.7	11	10.6	18	11.4	4	8.9	40	7.6
	HbA1c 6.1%以上または 現在治療を受けている	0	0	0	0	6	7.0	16	15.4	23	14.6	12	26.7	57	10.8
	—総数—	49	100.0	84	100.0	86	100.0	104	100.0	158	100.0	45	100.0	526	100.0

第58表-3 1日の運動量(歩行数)別、糖尿病の状況(女性、年齢階級別)

1日の運動量 (歩行数)	糖尿病の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
2,000歩未満	HbA1c 5.6%未満	6	100.0	6	100.0	5	55.6	9	90.0	19	73.1	99	68.3	144	71.3
	HbA1c 5.6%以上6.1%未満	0	0	0	0	3	33.3	1	10.0	5	19.2	27	18.6	36	17.8
	HbA1c 6.1%以上または 現在治療を受けている	0	0	0	0	1	11.1	0	0	2	7.7	19	13.1	22	10.9
	一総数一	6	100.0	6	100.0	9	100.0	10	100.0	26	100.0	145	100.0	202	100.0
2,000 ~3,999歩	HbA1c 5.6%未満	26	96.3	44	93.6	39	92.9	70	83.3	63	73.3	105	69.5	347	79.4
	HbA1c 5.6%以上6.1%未満	0	0	3	6.4	1	2.4	9	10.7	9	10.5	22	14.6	44	10.1
	HbA1c 6.1%以上または 現在治療を受けている	1	3.7	0	0	2	4.8	5	6.0	14	16.3	24	15.9	46	10.5
	一総数一	27	100.0	47	100.0	42	100.0	84	100.0	86	100.0	151	100.0	437	100.0
4,000 ~5,999歩	HbA1c 5.6%未満	55	100.0	90	95.7	82	85.4	110	91.7	93	69.9	100	72.5	530	83.3
	HbA1c 5.6%以上6.1%未満	0	0	4	4.3	9	9.4	5	4.2	26	19.5	27	19.6	71	11.2
	HbA1c 6.1%以上または 現在治療を受けている	0	0	0	0	5	5.2	5	4.2	14	10.5	11	8.0	35	5.5
	一総数一	55	100.0	94	100.0	96	100.0	120	100.0	133	100.0	138	100.0	636	100.0
6,000 ~7,999歩	HbA1c 5.6%未満	61	100.0	104	98.1	90	86.5	112	81.8	110	76.9	76	78.4	553	85.3
	HbA1c 5.6%以上6.1%未満	0	0	1	0.9	9	8.7	16	11.7	19	13.3	11	11.3	56	8.6
	HbA1c 6.1%以上または 現在治療を受けている	0	0	1	0.9	5	4.8	9	6.6	14	9.8	10	10.3	39	6.0
	一総数一	61	100.0	106	100.0	104	100.0	137	100.0	143	100.0	97	100.0	648	100.0
8,000 ~9,999歩	HbA1c 5.6%未満	41	97.6	80	94.1	73	89.0	101	80.8	89	78.8	41	74.5	425	84.7
	HbA1c 5.6%以上6.1%未満	0	0	5	5.9	9	11.0	19	15.2	14	12.4	8	14.5	55	11.0
	HbA1c 6.1%以上または 現在治療を受けている	1	2.4	0	0	0	0	5	4.0	10	8.8	6	10.9	22	4.4
	一総数一	42	100.0	85	100.0	82	100.0	125	100.0	113	100.0	55	100.0	502	100.0
10,000歩 以上	HbA1c 5.6%未満	51	98.1	93	90.3	121	91.0	172	84.7	93	64.6	36	69.2	566	82.4
	HbA1c 5.6%以上6.1%未満	1	1.9	7	6.8	8	6.0	23	11.3	32	22.2	11	21.2	82	11.9
	HbA1c 6.1%以上または 現在治療を受けている	0	0	3	2.9	4	3.0	8	3.9	19	13.2	5	9.6	39	5.7
	一総数一	52	100.0	103	100.0	133	100.0	203	100.0	144	100.0	52	100.0	687	100.0

第58表-4 1日の運動量（歩行数）別、糖尿病の状況

	1日の運動量 (歩行数)	糖尿病の状況	20-39歳		40-59歳		60歳以上		一計	
			人数	%	人数	%	人数	%	人数	%
男女計	4,000歩未満	HbA1c 5.6%未満	125	94.7	209	80.1	468	65.5	802	72.4
		HbA1c 5.6%以上6.1%未満	5	3.8	27	10.3	117	16.4	149	13.4
		HbA1c 6.1%以上または 現在治療を受けている	2	1.5	25	9.6	130	18.2	157	14.2
		—総数—	132	100.0	261	100.0	715	100.0	1,108	100.0
	4,000-7,999歩	HbA1c 5.6%未満	453	97.6	622	85.7	622	71.4	1,697	82.3
		HbA1c 5.6%以上6.1%未満	9	1.9	57	7.9	136	15.6	202	9.8
		HbA1c 6.1%以上または 現在治療を受けている	2	0.4	47	6.5	113	13.0	162	7.9
		—総数—	464	100.0	726	100.0	871	100.0	2,061	100.0
	8,000-11,999歩	HbA1c 5.6%未満	280	94.6	474	84.2	340	71.7	1,094	82.1
		HbA1c 5.6%以上6.1%未満	12	4.1	59	10.5	70	14.8	141	10.6
		HbA1c 6.1%以上または 現在治療を受けている	4	1.4	30	5.3	64	13.5	98	7.4
		—総数—	296	100.0	563	100.0	474	100.0	1,333	100.0
12,000歩以上	HbA1c 5.6%未満	170	97.7	253	85.2	147	68.7	570	83.2	
	HbA1c 5.6%以上6.1%未満	4	2.3	25	8.4	34	15.9	63	9.2	
	HbA1c 6.1%以上または 現在治療を受けている	0	0	19	6.4	33	15.4	52	7.6	
	—総数—	174	100.0	297	100.0	214	100.0	685	100.0	
男性	4,000歩未満	HbA1c 5.6%未満	43	93.5	86	74.1	182	59.3	311	66.3
		HbA1c 5.6%以上6.1%未満	2	4.3	13	11.2	54	17.6	69	14.7
		HbA1c 6.1%以上または 現在治療を受けている	1	2.2	17	14.7	71	23.1	89	19.0
		—総数—	46	100.0	116	100.0	307	100.0	469	100.0
	4,000-7,999歩	HbA1c 5.6%未満	143	96.6	228	84.8	243	67.5	614	79.0
		HbA1c 5.6%以上6.1%未満	4	2.7	18	6.7	53	14.7	75	9.7
		HbA1c 6.1%以上または 現在治療を受けている	1	0.7	23	8.6	64	17.8	88	11.3
		—総数—	148	100.0	269	100.0	360	100.0	777	100.0
	8,000-11,999歩	HbA1c 5.6%未満	94	98.9	171	81.8	148	71.5	413	80.8
		HbA1c 5.6%以上6.1%未満	1	1.1	17	8.1	25	12.1	43	8.4
		HbA1c 6.1%以上または 現在治療を受けている	0	0	21	10.0	34	16.4	55	10.8
		—総数—	95	100.0	209	100.0	207	100.0	511	100.0
12,000歩以上	HbA1c 5.6%未満	91	97.8	89	82.4	80	68.4	260	81.8	
	HbA1c 5.6%以上6.1%未満	2	2.2	8	7.4	14	12.0	24	7.5	
	HbA1c 6.1%以上または 現在治療を受けている	0	0	11	10.2	23	19.7	34	10.7	
	—総数—	93	100.0	108	100.0	117	100.0	318	100.0	
女性	4,000歩未満	HbA1c 5.6%未満	82	95.3	123	84.8	286	70.1	491	76.8
		HbA1c 5.6%以上6.1%未満	3	3.5	14	9.7	63	15.4	80	12.5
		HbA1c 6.1%以上または 現在治療を受けている	1	1.2	8	5.5	59	14.5	68	10.6
		—総数—	86	100.0	145	100.0	408	100.0	639	100.0
	4,000-7,999歩	HbA1c 5.6%未満	310	98.1	394	86.2	379	74.2	1,083	84.3
		HbA1c 5.6%以上6.1%未満	5	1.6	39	8.5	83	16.2	127	9.9
		HbA1c 6.1%以上または 現在治療を受けている	1	0.3	24	5.3	49	9.6	74	5.8
		—総数—	316	100.0	457	100.0	511	100.0	1,284	100.0
	8,000-11,999歩	HbA1c 5.6%未満	186	92.5	303	85.6	192	71.9	681	82.8
		HbA1c 5.6%以上6.1%未満	11	5.5	42	11.9	45	16.9	98	11.9
		HbA1c 6.1%以上または 現在治療を受けている	4	2.0	9	2.5	30	11.2	43	5.2
		—総数—	201	100.0	354	100.0	267	100.0	822	100.0
12,000歩以上	HbA1c 5.6%未満	79	97.5	164	86.8	67	69.1	310	84.5	
	HbA1c 5.6%以上6.1%未満	2	2.5	17	9.0	20	20.6	39	10.6	
	HbA1c 6.1%以上または 現在治療を受けている	0	0	8	4.2	10	10.3	18	4.9	
	—総数—	81	100.0	189	100.0	97	100.0	367	100.0	

第59表 喫煙習慣の状況別、糖尿病の状況（性・年齢階級別）

	喫煙習慣	糖尿病の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男女計	あり	HbA1c 5.6%未満	119	98.3	209	94.6	204	92.3	204	76.7	165	68.2	90	59.6	991	81.1
		HbA1c 5.6%以上6.1%未満	2	1.7	9	4.1	9	4.1	28	10.5	42	17.4	28	18.5	118	9.7
		HbA1c 6.1%以上または現在治療を受けている	0	0	3	1.4	8	3.6	34	12.8	35	14.5	33	21.9	113	9.2
		—総数—	121	100.0	221	100.0	221	100.0	266	100.0	242	100.0	151	100.0	1,222	100.0
	なし	HbA1c 5.6%未満	268	98.5	465	95.7	480	88.6	701	82.6	683	71.4	676	69.2	3,273	80.2
		HbA1c 5.6%以上6.1%未満	2	0.7	18	3.7	40	7.4	90	10.6	137	14.3	157	16.1	444	10.9
		HbA1c 6.1%以上または現在治療を受けている	2	0.7	3	0.6	22	4.1	58	6.8	137	14.3	144	14.7	366	9.0
		—総数—	272	100.0	486	100.0	542	100.0	849	100.0	957	100.0	977	100.0	4,083	100.0
男 性	あり	HbA1c 5.6%未満	75	97.4	136	95.8	141	92.8	149	74.9	135	69.6	77	57.9	713	79.5
		HbA1c 5.6%以上6.1%未満	2	2.6	4	2.8	4	2.6	20	10.1	30	15.5	25	18.8	85	9.5
		HbA1c 6.1%以上または現在治療を受けている	0	0	2	1.4	7	4.6	30	15.1	29	14.9	31	23.3	99	11.0
		—総数—	77	100.0	142	100.0	152	100.0	199	100.0	194	100.0	133	100.0	897	100.0
	なし	HbA1c 5.6%未満	66	98.5	111	97.4	130	91.5	172	75.8	238	68.2	220	64.3	937	75.5
		HbA1c 5.6%以上6.1%未満	1	1.5	3	2.6	6	4.2	25	11.0	43	12.3	51	14.9	129	10.4
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	6	4.2	30	13.2	68	19.5	71	20.8	175	14.1
		—総数—	67	100.0	114	100.0	142	100.0	227	100.0	349	100.0	342	100.0	1,241	100.0
女 性	あり	HbA1c 5.6%未満	44	100.0	73	92.4	63	91.3	55	82.1	30	62.5	13	72.2	278	85.5
		HbA1c 5.6%以上6.1%未満	0	0	5	6.3	5	7.2	8	11.9	12	25.0	3	16.7	33	10.2
		HbA1c 6.1%以上または現在治療を受けている	0	0	1	1.3	1	1.4	4	6.0	6	12.5	2	11.1	14	4.3
		—総数—	44	100.0	79	100.0	69	100.0	67	100.0	48	100.0	18	100.0	325	100.0
	なし	HbA1c 5.6%未満	202	98.5	354	95.2	350	87.5	529	85.0	445	73.2	456	71.8	2,336	82.2
		HbA1c 5.6%以上6.1%未満	1	0.5	15	4.0	34	8.5	65	10.5	94	15.5	106	16.7	315	11.1
		HbA1c 6.1%以上または現在治療を受けている	2	1.0	3	0.8	16	4.0	28	4.5	69	11.3	73	11.5	191	6.7
		—総数—	205	100.0	372	100.0	400	100.0	622	100.0	608	100.0	635	100.0	2,842	100.0

第60表 飲酒習慣の状況別、糖尿病の状況（性・年齢階級別）

	飲酒習慣	糖尿病の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男女計	飲んでいる	HbA1c 5.6%未満	50	96.2	146	97.3	221	94.4	280	81.4	233	74.2	162	68.1	1,092	82.0
		HbA1c 5.6%以上6.1%未満	2	3.8	4	2.7	9	3.8	34	9.9	35	11.1	22	9.2	106	8.0
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	4	1.7	30	8.7	46	14.6	54	22.7	134	10.1
		—総数—	52	100.0	150	100.0	234	100.0	344	100.0	314	100.0	238	100.0	1,332	100.0
	以前飲んでいた	HbA1c 5.6%未満	9	100.0	19	100.0	20	90.9	24	72.7	38	55.9	52	64.2	162	69.8
		HbA1c 5.6%以上6.1%未満	0	0	0	0	1	4.5	5	15.2	10	14.7	11	13.6	27	11.6
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	4.5	4	12.1	20	29.4	18	22.2	43	18.5
		—総数—	9	100.0	19	100.0	22	100.0	33	100.0	68	100.0	81	100.0	232	100.0
	飲んでいない	HbA1c 5.6%未満	328	98.8	508	94.6	443	87.4	601	81.4	577	70.6	552	68.2	3,009	80.5
		HbA1c 5.6%以上6.1%未満	2	0.6	23	4.3	39	7.7	79	10.7	134	16.4	152	18.8	429	11.5
		HbA1c 6.1%以上または現在治療を受けている	2	0.6	6	1.1	25	4.9	58	7.9	106	13.0	105	13.0	302	8.1
		—総数—	332	100.0	537	100.0	507	100.0	738	100.0	817	100.0	809	100.0	3,740	100.0
男性	飲んでいる	HbA1c 5.6%未満	32	94.1	108	97.3	157	94.6	208	80.3	202	72.9	143	66.2	850	80.0
		HbA1c 5.6%以上6.1%未満	2	5.9	3	2.7	5	3.0	25	9.7	30	10.8	21	9.7	86	8.1
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	4	2.4	26	10.0	45	16.2	52	24.1	127	11.9
		—総数—	34	100.0	111	100.0	166	100.0	259	100.0	277	100.0	216	100.0	1,063	100.0
	以前飲んでいた	HbA1c 5.6%未満	1	100.0	12	100.0	15	88.2	13	68.4	30	57.7	46	63.0	117	67.2
		HbA1c 5.6%以上6.1%未満	0	0	0	0	1	5.9	3	15.8	7	13.5	11	15.1	22	12.6
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	5.9	3	15.8	15	28.8	16	21.9	35	20.1
		—総数—	1	100.0	12	100.0	17	100.0	19	100.0	52	100.0	73	100.0	174	100.0
	飲んでいない	HbA1c 5.6%未満	108	99.1	127	95.5	99	89.2	100	67.6	141	65.9	108	58.1	683	75.8
		HbA1c 5.6%以上6.1%未満	1	0.9	4	3.0	4	3.6	17	11.5	36	16.8	44	23.7	106	11.8
		HbA1c 6.1%以上または現在治療を受けている	0	0	2	1.5	8	7.2	31	20.9	37	17.3	34	18.3	112	12.4
		—総数—	109	100.0	133	100.0	111	100.0	148	100.0	214	100.0	186	100.0	901	100.0
女性	飲んでいる	HbA1c 5.6%未満	18	100.0	38	97.4	64	94.1	72	84.7	31	83.8	19	86.4	242	90.0
		HbA1c 5.6%以上6.1%未満	0	0	1	2.6	4	5.9	9	10.6	5	13.5	1	4.5	20	7.4
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	4	4.7	1	2.7	2	9.1	7	2.6
		—総数—	18	100.0	39	100.0	68	100.0	85	100.0	37	100.0	22	100.0	269	100.0
	以前飲んでいた	HbA1c 5.6%未満	8	100.0	7	100.0	5	100.0	11	78.6	8	50.0	6	75.0	45	77.6
		HbA1c 5.6%以上6.1%未満	0	0	0	0	0	0	2	14.3	3	18.8	0	0	5	8.6
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	1	7.1	5	31.3	2	25.0	8	13.8
		—総数—	8	100.0	7	100.0	5	100.0	14	100.0	16	100.0	8	100.0	58	100.0
	飲んでいない	HbA1c 5.6%未満	220	98.7	381	94.3	344	86.9	501	84.9	436	72.3	444	71.3	2,326	81.9
		HbA1c 5.6%以上6.1%未満	1	0.4	19	4.7	35	8.8	62	10.5	98	16.3	108	17.3	323	11.4
		HbA1c 6.1%以上または現在治療を受けている	2	0.9	4	1.0	17	4.3	27	4.6	69	11.4	71	11.4	190	6.7
		—総数—	223	100.0	404	100.0	396	100.0	590	100.0	603	100.0	623	100.0	2,839	100.0

第61表 運動習慣の状況別、糖尿病の状況（性・年齢階級別）

	運動習慣	糖尿病の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		—計—	
			人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
男女計	あり	HbA1c 5.6%未満	62	98.4	133	96.4	152	92.1	253	79.3	337	68.5	279	68.2	1,216	76.7
		HbA1c 5.6%以上6.1%未満	1	1.6	4	2.9	9	5.5	35	11.0	75	15.2	66	16.1	190	12.0
		HbA1c 6.1%以上または現在治療を受けている	0	0	1	0.7	4	2.4	31	9.7	80	16.3	64	15.6	180	11.3
		—総数—	63	100.0	138	100.0	165	100.0	319	100.0	492	100.0	409	100.0	1,586	100.0
	なし (健康上) の理由	HbA1c 5.6%未満	6	100.0	9	100.0	20	80.0	40	75.5	51	63.8	120	60.9	246	66.5
		HbA1c 5.6%以上6.1%未満	0	0	0	0	4	16.0	5	9.4	12	15.0	37	18.8	58	15.7
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	4.0	8	15.1	17	21.3	40	20.3	66	17.8
		—総数—	6	100.0	9	100.0	25	100.0	53	100.0	80	100.0	197	100.0	370	100.0
	なし (その他) の理由	HbA1c 5.6%未満	319	98.5	532	95.0	512	89.4	612	82.4	460	73.4	366	70.2	2,801	83.7
		HbA1c 5.6%以上6.1%未満	3	0.9	23	4.1	36	6.3	78	10.5	92	14.7	82	15.7	314	9.4
		HbA1c 6.1%以上または現在治療を受けている	2	0.6	5	0.9	25	4.4	53	7.1	75	12.0	73	14.0	233	7.0
		—総数—	324	100.0	560	100.0	573	100.0	743	100.0	627	100.0	521	100.0	3,348	100.0
男 性	あり	HbA1c 5.6%未満	32	100.0	54	98.2	56	91.8	77	70.0	161	68.2	119	62.3	499	72.8
		HbA1c 5.6%以上6.1%未満	0	0	1	1.8	3	4.9	14	12.7	30	12.7	27	14.1	75	10.9
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	2	3.3	19	17.3	45	19.1	45	23.6	111	16.2
		—総数—	32	100.0	55	100.0	61	100.0	110	100.0	236	100.0	191	100.0	685	100.0
	なし (健康上) の理由	HbA1c 5.6%未満	2	100.0	0	0	7	100.0	14	77.8	13	48.1	39	57.4	75	61.5
		HbA1c 5.6%以上6.1%未満	0	0	0	0	0	0	1	5.6	6	22.2	14	20.6	21	17.2
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	3	16.7	8	29.6	15	22.1	26	21.3
		—総数—	2	100.0	0	0	7	100.0	18	100.0	27	100.0	68	100.0	122	100.0
	なし (その他) の理由	HbA1c 5.6%未満	107	97.3	193	96.0	208	92.0	230	77.2	199	71.1	139	64.4	1,076	80.8
		HbA1c 5.6%以上6.1%未満	3	2.7	6	3.0	7	3.1	30	10.1	37	13.2	35	16.2	118	8.9
		HbA1c 6.1%以上または現在治療を受けている	0	0	2	1.0	11	4.9	38	12.8	44	15.7	42	19.4	137	10.3
		—総数—	110	100.0	201	100.0	226	100.0	298	100.0	280	100.0	216	100.0	1,331	100.0
女 性	あり	HbA1c 5.6%未満	30	96.8	79	95.2	96	92.3	176	84.2	176	68.8	160	73.4	717	79.6
		HbA1c 5.6%以上6.1%未満	1	3.2	3	3.6	6	5.8	21	10.0	45	17.6	39	17.9	115	12.8
		HbA1c 6.1%以上または現在治療を受けている	0	0	1	1.2	2	1.9	12	5.7	35	13.7	19	8.7	69	7.7
		—総数—	31	100.0	83	100.0	104	100.0	209	100.0	256	100.0	218	100.0	901	100.0
	なし (健康上) の理由	HbA1c 5.6%未満	4	100.0	9	100.0	13	72.2	26	74.3	38	71.7	81	62.8	171	69.0
		HbA1c 5.6%以上6.1%未満	0	0	0	0	4	22.2	4	11.4	6	11.3	23	17.8	37	14.9
		HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	5.6	5	14.3	9	17.0	25	19.4	40	16.1
		—総数—	4	100.0	9	100.0	18	100.0	35	100.0	53	100.0	129	100.0	248	100.0
	なし (その他) の理由	HbA1c 5.6%未満	212	99.1	339	94.4	304	87.6	382	85.8	261	75.2	227	74.4	1,725	85.5
		HbA1c 5.6%以上6.1%未満	0	0	17	4.7	29	8.4	48	10.8	55	15.9	47	15.4	196	9.7
		HbA1c 6.1%以上または現在治療を受けている	2	0.9	3	0.8	14	4.0	15	3.4	31	8.9	31	10.2	96	4.8
		—総数—	214	100.0	359	100.0	347	100.0	445	100.0	347	100.0	305	100.0	2,017	100.0

第62表-1 糖尿病の状況別、血清総コレステロール値の分布（男女計、年齢階級別）

糖尿病の状況	総コレステロール値 (mg/dl)	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計		
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
HbA1c 5.6%未満	150未満	45	11.6	39	5.8	21	3.1	19	2.1	24	2.8	32	4.2	180	4.2	
	150-199	240	61.9	375	55.5	318	46.4	271	29.9	308	36.4	337	43.9	1,849	43.3	
	200-219	54	13.9	122	18.0	149	21.8	219	24.2	203	24.0	174	22.7	921	21.6	
	220-239	26	6.7	81	12.0	112	16.4	182	20.1	158	18.7	125	16.3	684	16.0	
	240-259	11	2.8	38	5.6	52	7.6	111	12.3	84	9.9	67	8.7	363	8.5	
	260以上	12	3.1	21	3.1	33	4.8	103	11.4	70	8.3	32	4.2	271	6.3	
	—総数—	388	100.0	676	100.0	685	100.0	905	100.0	847	100.0	767	100.0	4,268	100.0	
	(再掲)															
	220未満	339	87.4	536	79.3	488	71.2	509	56.2	535	63.2	543	70.8	2,950	69.1	
	220以上240未満	26	6.7	81	12.0	112	16.4	182	20.1	158	18.7	125	16.3	684	16.0	
240以上	23	5.9	59	8.7	85	12.4	214	23.6	154	18.2	99	12.9	634	14.9		
HbA1c 5.6%以上 6.1%未満	150未満	0	0	2	7.4	1	2.0	0	0	1	0.6	6	3.2	10	1.8	
	150-199	3	75.0	10	37.0	19	38.8	34	28.6	54	30.2	83	44.9	203	36.1	
	200-219	1	25.0	5	18.5	15	30.6	15	12.6	43	24.0	49	26.5	128	22.7	
	220-239	0	0	5	18.5	9	18.4	27	22.7	51	28.5	26	14.1	118	21.0	
	240-259	0	0	3	11.1	3	6.1	21	17.6	16	8.9	12	6.5	55	9.8	
	260以上	0	0	2	7.4	2	4.1	22	18.5	14	7.8	9	4.9	49	8.7	
	—総数—	4	100.0	27	100.0	49	100.0	119	100.0	179	100.0	185	100.0	563	100.0	
	(再掲)															
	220未満	4	100.0	17	63.0	35	71.4	49	41.2	98	54.7	138	74.6	341	60.6	
	220以上240未満	0	0	5	18.5	9	18.4	27	22.7	51	28.5	26	14.1	118	21.0	
240以上	0	0	5	18.5	5	10.2	43	36.1	30	16.8	21	11.4	104	18.5		
HbA1c 6.1%以上 または現在治療を 受けている	150未満	0	0	0	0	0	0	0	0	4	2.3	9	5.1	13	2.7	
	150-199	1	50.0	1	16.7	7	23.3	35	38.0	67	38.7	78	44.1	189	39.4	
	200-219	0	0	1	16.7	11	36.7	18	19.6	41	23.7	45	25.4	116	24.2	
	220-239	1	50.0	3	50.0	5	16.7	18	19.6	26	15.0	30	16.9	83	17.3	
	240-259	0	0	0	0	4	13.3	8	8.7	23	13.3	8	4.5	43	9.0	
	260以上	0	0	1	16.7	3	10.0	13	14.1	12	6.9	7	4.0	36	7.5	
	—総数—	2	100.0	6	100.0	30	100.0	92	100.0	173	100.0	177	100.0	480	100.0	
	(再掲)															
	220未満	1	50.0	2	33.3	18	60.0	53	57.6	112	64.7	132	74.6	318	66.3	
	220以上240未満	1	50.0	3	50.0	5	16.7	18	19.6	26	15.0	30	16.9	83	17.3	
240以上	0	0	1	16.7	7	23.3	21	22.8	35	20.2	15	8.5	79	16.5		

第62表-2 糖尿病の状況別、血清総コレステロール値の分布（男性、年齢階級別）

糖尿病の状況	総コレステロール値 (mg/dl)	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計		
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
HbA1c 5.6%未満	150未満	14	9.9	10	4.0	7	2.6	14	4.4	22	5.9	24	8.1	91	5.5	
	150-199	89	63.1	120	48.6	117	43.2	136	42.4	165	44.2	168	56.6	795	48.2	
	200-219	21	14.9	45	18.2	59	21.8	78	24.3	86	23.1	50	16.8	339	20.5	
	220-239	11	7.8	34	13.8	52	19.2	44	13.7	64	17.2	32	10.8	237	14.4	
	240-259	3	2.1	23	9.3	22	8.1	27	8.4	27	7.2	16	5.4	118	7.2	
	260以上	3	2.1	15	6.1	14	5.2	22	6.9	9	2.4	7	2.4	70	4.2	
	一総数一	141	100.0	247	100.0	271	100.0	321	100.0	373	100.0	297	100.0	1,650	100.0	
	(再掲)															
	220未満	124	87.9	175	70.9	183	67.5	228	71.0	273	73.2	242	81.5	1,225	74.2	
	220以上240未満	11	7.8	34	13.8	52	19.2	44	13.7	64	17.2	32	10.8	237	14.4	
240以上	6	4.3	38	15.4	36	13.3	49	15.3	36	9.7	23	7.7	188	11.4		
HbA1c 5.6%以上 6.1%未満	150未満	0	0	0	0	0	0	0	0	1	1.4	5	6.6	6	2.8	
	150-199	3	100.0	2	28.6	2	20.0	18	39.1	27	37.0	39	51.3	91	42.3	
	200-219	0	0	1	14.3	4	40.0	7	15.2	19	26.0	18	23.7	49	22.8	
	220-239	0	0	3	42.9	3	30.0	11	23.9	20	27.4	8	10.5	45	20.9	
	240-259	0	0	0	0	1	10.0	5	10.9	2	2.7	3	3.9	11	5.1	
	260以上	0	0	1	14.3	0	0	5	10.9	4	5.5	3	3.9	13	6.0	
	一総数一	3	100.0	7	100.0	10	100.0	46	100.0	73	100.0	76	100.0	215	100.0	
	(再掲)															
	220未満	3	100.0	3	42.9	6	60.0	25	54.3	47	64.4	62	81.6	146	67.9	
	220以上240未満	0	0	3	42.9	3	30.0	11	23.9	20	27.4	8	10.5	45	20.9	
240以上	0	0	1	14.3	1	10.0	10	21.7	6	8.2	6	7.9	24	11.2		
HbA1c 6.1%以上 または現在治療を 受けている	150未満	0	0	0	0	0	0	0	0	3	3.1	6	5.9	9	3.3	
	150-199	0	0	0	0	5	38.5	26	43.3	48	49.0	54	52.9	133	48.4	
	200-219	0	0	0	0	5	38.5	13	21.7	21	21.4	24	23.5	63	22.9	
	220-239	0	0	1	50.0	2	15.4	10	16.7	10	10.2	15	14.7	38	13.8	
	240-259	0	0	0	0	1	7.7	4	6.7	9	9.2	3	2.9	17	6.2	
	260以上	0	0	1	50.0	0	0	7	11.7	7	7.1	0	0	15	5.5	
	一総数一	0	0	2	100.0	13	100.0	60	100.0	98	100.0	102	100.0	275	100.0	
	(再掲)															
	220未満	0	0	0	0	10	76.9	39	65.0	72	73.5	84	82.4	205	74.5	
	220以上240未満	0	0	1	50.0	2	15.4	10	16.7	10	10.2	15	14.7	38	13.8	
240以上	0	0	1	50.0	1	7.7	11	18.3	16	16.3	3	2.9	32	11.6		

第62表-3 糖尿病の状況別、血清総コレステロール値の分布（女性、年齢階級別）

糖尿病の状況	総コレステロール値 (mg/dl)	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		—計—		
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
HbA1c 5.6%未満	150未満	31	12.6	29	6.8	14	3.4	5	0.9	2	0.4	8	1.7	89	3.4	
	150-199	151	61.1	255	59.4	201	48.6	135	23.1	143	30.2	169	36.0	1,054	40.3	
	200-219	33	13.4	77	17.9	90	21.7	141	24.1	117	24.7	124	26.4	582	22.2	
	220-239	15	6.1	47	11.0	60	14.5	138	23.6	94	19.8	93	19.8	447	17.1	
	240-259	8	3.2	15	3.5	30	7.2	84	14.4	57	12.0	51	10.9	245	9.4	
	260以上	9	3.6	6	1.4	19	4.6	81	13.9	61	12.9	25	5.3	201	7.7	
	—総数—	247	100.0	429	100.0	414	100.0	584	100.0	474	100.0	470	100.0	2,618	100.0	
	(再掲)															
	220未満	215	87.0	361	84.1	305	73.7	281	48.1	262	55.3	301	64.0	1,725	65.9	
	220以上240未満	15	6.1	47	11.0	60	14.5	138	23.6	94	19.8	93	19.8	447	17.1	
240以上	17	6.9	21	4.9	49	11.8	165	28.3	118	24.9	76	16.2	446	17.0		
HbA1c 5.6%以上 6.1%未満	150未満	0	0	2	10.0	1	2.6	0	0	0	0	1	0.9	4	1.1	
	150-199	0	0	8	40.0	17	43.6	16	21.9	27	25.5	44	40.4	112	32.2	
	200-219	1	100.0	4	20.0	11	28.2	8	11.0	24	22.6	31	28.4	79	22.7	
	220-239	0	0	2	10.0	6	15.4	16	21.9	31	29.2	18	16.5	73	21.0	
	240-259	0	0	3	15.0	2	5.1	16	21.9	14	13.2	9	8.3	44	12.6	
	260以上	0	0	1	5.0	2	5.1	17	23.3	10	9.4	6	5.5	36	10.3	
	—総数—	1	100.0	20	100.0	39	100.0	73	100.0	106	100.0	109	100.0	348	100.0	
	(再掲)															
	220未満	1	100.0	14	70.0	29	74.4	24	32.9	51	48.1	76	69.7	195	56.0	
	220以上240未満	0	0	2	10.0	6	15.4	16	21.9	31	29.2	18	16.5	73	21.0	
240以上	0	0	4	20.0	4	10.3	33	45.2	24	22.6	15	13.8	80	23.0		
HbA1c 6.1%以上 または現在治療を 受けている	150未満	0	0	0	0	0	0	0	0	1	1.3	3	4.0	4	2.0	
	150-199	1	50.0	1	25.0	2	11.8	9	28.1	19	25.3	24	32.0	56	27.3	
	200-219	0	0	1	25.0	6	35.3	5	15.6	20	26.7	21	28.0	53	25.9	
	220-239	1	50.0	2	50.0	3	17.6	8	25.0	16	21.3	15	20.0	45	22.0	
	240-259	0	0	0	0	3	17.6	4	12.5	14	18.7	5	6.7	26	12.7	
	260以上	0	0	0	0	3	17.6	6	18.8	5	6.7	7	9.3	21	10.2	
	—総数—	2	100.0	4	100.0	17	100.0	32	100.0	75	100.0	75	100.0	205	100.0	
	(再掲)															
	220未満	1	50.0	2	50.0	8	47.1	14	43.8	40	53.3	48	64.0	113	55.1	
	220以上240未満	1	50.0	2	50.0	3	17.6	8	25.0	16	21.3	15	20.0	45	22.0	
240以上	0	0	0	0	6	35.3	10	31.3	19	25.3	12	16.0	47	22.9		

第63表-1 糖尿病の状況別、血清HDLコレステロール値の分布（男女計、年齢階級別）

糖尿病の状況	HDLコレステロール値(mg/dl)	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		— 計 —	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
HbA1c 5.6%未満	30未満	3	0.8	1	0.1	3	0.4	3	0.3	5	0.6	5	0.7	20	0.5
	30-39	15	3.9	23	3.4	25	3.6	43	4.8	55	6.5	62	8.1	223	5.2
	40-49	46	11.9	99	14.6	104	15.2	142	15.7	167	19.7	131	17.1	689	16.1
	50-59	93	24.0	159	23.5	139	20.3	209	23.1	227	26.8	225	29.3	1,052	24.6
	60-69	99	25.5	178	26.3	169	24.7	215	23.8	196	23.1	167	21.8	1,024	24.0
	70-79	68	17.5	115	17.0	134	19.6	150	16.6	96	11.3	103	13.4	666	15.6
	80-89	34	8.8	66	9.8	66	9.6	83	9.2	60	7.1	49	6.4	358	8.4
	90-99	19	4.9	20	3.0	28	4.1	34	3.8	28	3.3	15	2.0	144	3.4
	100以上	11	2.8	15	2.2	17	2.5	26	2.9	13	1.5	10	1.3	92	2.2
	— 総数 —	388	100.0	676	100.0	685	100.0	905	100.0	847	100.0	767	100.0	4,268	100.0
	(再掲)														
40未満	18	4.6	24	3.6	28	4.1	46	5.1	60	7.1	67	8.7	243	5.7	
40以上	370	95.4	652	96.4	657	95.9	859	94.9	787	92.9	700	91.3	4,025	94.3	
HbA1c 5.6%以上 6.1%未満	30未満	0	0	2	7.4	0	0	0	0	1	0.6	3	1.6	6	1.1
	30-39	0	0	0	0	0	0	8	6.7	18	10.1	23	12.4	49	8.7
	40-49	0	0	1	3.7	10	20.4	25	21.0	36	20.1	49	26.5	121	21.5
	50-59	1	25.0	10	37.0	16	32.7	28	23.5	45	25.1	51	27.6	151	26.8
	60-69	0	0	6	22.2	12	24.5	26	21.8	43	24.0	29	15.7	116	20.6
	70-79	3	75.0	3	11.1	6	12.2	16	13.4	21	11.7	19	10.3	68	12.1
	80-89	0	0	2	7.4	3	6.1	9	7.6	10	5.6	5	2.7	29	5.2
	90-99	0	0	3	11.1	2	4.1	5	4.2	2	1.1	2	1.1	14	2.5
	100以上	0	0	0	0	0	0	2	1.7	3	1.7	4	2.2	9	1.6
	— 総数 —	4	100.0	27	100.0	49	100.0	119	100.0	179	100.0	185	100.0	563	100.0
	(再掲)														
40未満	0	0	2	7.4	0	0	8	6.7	19	10.6	26	14.1	55	9.8	
40以上	4	100.0	25	92.6	49	100.0	111	93.3	160	89.4	159	85.9	508	90.2	
HbA1c 6.1%以上 または現在治療を 受けている	30未満	0	0	0	0	0	0	1	1.1	4	2.3	4	2.3	9	1.9
	30-39	0	0	1	16.7	6	20.0	16	17.4	25	14.5	18	10.2	66	13.8
	40-49	0	0	2	33.3	6	20.0	27	29.3	37	21.4	44	24.9	116	24.2
	50-59	1	50.0	2	33.3	7	23.3	24	26.1	53	30.6	58	32.8	145	30.2
	60-69	1	50.0	1	16.7	5	16.7	13	14.1	24	13.9	28	15.8	72	15.0
	70-79	0	0	0	0	4	13.3	7	7.6	13	7.5	13	7.3	37	7.7
	80-89	0	0	0	0	1	3.3	2	2.2	9	5.2	8	4.5	20	4.2
	90-99	0	0	0	0	1	3.3	2	2.2	5	2.9	3	1.7	11	2.3
	100以上	0	0	0	0	0	0	0	0	3	1.7	1	0.6	4	0.8
	— 総数 —	2	100.0	6	100.0	30	100.0	92	100.0	173	100.0	177	100.0	480	100.0
	(再掲)														
40未満	0	0	1	16.7	6	20.0	17	18.5	29	16.8	22	12.4	75	15.6	
40以上	2	100.0	5	83.3	24	80.0	75	81.5	144	83.2	155	87.6	405	84.4	

第63表-2 糖尿病の状況別、血清HDLコレステロール値の分布（男性、年齢階級別）

糖尿病の状況	HDLコレステロール値(mg/dl)	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
HbA1c 5.6%未満	30未満	3	2.1	1	0.4	3	1.1	3	0.9	4	1.1	5	1.7	19	1.2
	30-39	13	9.2	20	8.1	21	7.7	30	9.3	42	11.3	38	12.8	164	9.9
	40-49	30	21.3	64	25.9	73	26.9	74	23.1	88	23.6	63	21.2	392	23.8
	50-59	35	24.8	76	30.8	66	24.4	94	29.3	103	27.6	98	33.0	472	28.6
	60-69	34	24.1	46	18.6	53	19.6	62	19.3	79	21.2	52	17.5	326	19.8
	70-79	17	12.1	24	9.7	37	13.7	34	10.6	27	7.2	23	7.7	162	9.8
	80-89	4	2.8	11	4.5	12	4.4	15	4.7	15	4.0	13	4.4	70	4.2
	90-99	4	2.8	2	0.8	4	1.5	7	2.2	9	2.4	3	1.0	29	1.8
	100以上	1	0.7	3	1.2	2	0.7	2	0.6	6	1.6	2	0.7	16	1.0
	—総数—		141	100.0	247	100.0	271	100.0	321	100.0	373	100.0	297	100.0	1,650
(再掲)	40未満	16	11.3	21	8.5	24	8.9	33	10.3	46	12.3	43	14.5	183	11.1
	40以上	125	88.7	226	91.5	247	91.1	288	89.7	327	87.7	254	85.5	1,467	88.9
HbA1c 5.6%以上 6.1%未満	30未満	0	0	1	14.3	0	0	0	0	1	1.4	2	2.6	4	1.9
	30-39	0	0	0	0	0	0	8	17.4	13	17.8	10	13.2	31	14.4
	40-49	0	0	0	0	5	50.0	13	28.3	21	28.8	21	27.6	60	27.9
	50-59	1	33.3	3	42.9	4	40.0	10	21.7	13	17.8	24	31.6	55	25.6
	60-69	0	0	2	28.6	0	0	9	19.6	14	19.2	13	17.1	38	17.7
	70-79	2	66.7	1	14.3	0	0	3	6.5	9	12.3	2	2.6	17	7.9
	80-89	0	0	0	0	1	10.0	3	6.5	1	1.4	1	1.3	6	2.8
	90-99	0	0	0	0	0	0	0	0	0	0	1	1.3	1	0.5
	100以上	0	0	0	0	0	0	0	0	1	1.4	2	2.6	3	1.4
	—総数—		3	100.0	7	100.0	10	100.0	46	100.0	73	100.0	76	100.0	215
(再掲)	40未満	0	0	1	14.3	0	0	8	17.4	14	19.2	12	15.8	35	16.3
	40以上	3	100.0	6	85.7	10	100.0	38	82.6	59	80.8	64	84.2	180	83.7
HbA1c 6.1%以上 または現在治療を受けている	30未満	0	0	0	0	0	0	1	1.7	4	4.1	3	2.9	8	2.9
	30-39	0	0	1	50.0	6	46.2	14	23.3	21	21.4	12	11.8	54	19.6
	40-49	0	0	0	0	3	23.1	20	33.3	24	24.5	26	25.5	73	26.5
	50-59	0	0	1	50.0	3	23.1	16	26.7	26	26.5	33	32.4	79	28.7
	60-69	0	0	0	0	1	7.7	5	8.3	8	8.2	14	13.7	28	10.2
	70-79	0	0	0	0	0	0	2	3.3	6	6.1	7	6.9	15	5.5
	80-89	0	0	0	0	0	0	1	1.7	4	4.1	4	3.9	9	3.3
	90-99	0	0	0	0	0	0	1	1.7	2	2.0	2	2.0	5	1.8
	100以上	0	0	0	0	0	0	0	0	3	3.1	1	1.0	4	1.5
	—総数—		0	0	2	100.0	13	100.0	60	100.0	98	100.0	102	100.0	275
(再掲)	40未満	0	0	1	50.0	6	46.2	15	25.0	25	25.5	15	14.7	62	22.5
	40以上	0	0	1	50.0	7	53.8	45	75.0	73	74.5	87	85.3	213	77.5

第63表-3 糖尿病の状況別、血清HDLコレステロール値の分布（女性、年齢階級別）

糖尿病の状況	HDLコレステロール値 (mg/dl)	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
HbA1c 5.6%未満	30未満	0	0	0	0	0	0	0	0	1	0.2	0	0	1	0.0
	30-39	2	0.8	3	0.7	4	1.0	13	2.2	13	2.7	24	5.1	59	2.3
	40-49	16	6.5	35	8.2	31	7.5	68	11.6	79	16.7	68	14.5	297	11.3
	50-59	58	23.5	83	19.3	73	17.6	115	19.7	124	26.2	127	27.0	580	22.2
	60-69	65	26.3	132	30.8	116	28.0	153	26.2	117	24.7	115	24.5	698	26.7
	70-79	51	20.6	91	21.2	97	23.4	116	19.9	69	14.6	80	17.0	504	19.3
	80-89	30	12.1	55	12.8	54	13.0	68	11.6	45	9.5	36	7.7	288	11.0
	90-99	15	6.1	18	4.2	24	5.8	27	4.6	19	4.0	12	2.6	115	4.4
	100以上	10	4.0	12	2.8	15	3.6	24	4.1	7	1.5	8	1.7	76	2.9
	—総数—		247	100.0	429	100.0	414	100.0	584	100.0	474	100.0	470	100.0	2,618
(再掲)	40未満	2	0.8	3	0.7	4	1.0	13	2.2	14	3.0	24	5.1	60	2.3
	40以上	245	99.2	426	99.3	410	99.0	571	97.8	460	97.0	446	94.9	2,558	97.7
HbA1c 5.6%以上 6.1%未満	30未満	0	0	1	5.0	0	0	0	0	0	0	1	0.9	2	0.6
	30-39	0	0	0	0	0	0	0	0	5	4.7	13	11.9	18	5.2
	40-49	0	0	1	5.0	5	12.8	12	16.4	15	14.2	28	25.7	61	17.5
	50-59	0	0	7	35.0	12	30.8	18	24.7	32	30.2	27	24.8	96	27.6
	60-69	0	0	4	20.0	12	30.8	17	23.3	29	27.4	16	14.7	78	22.4
	70-79	1	100.0	2	10.0	6	15.4	13	17.8	12	11.3	17	15.6	51	14.7
	80-89	0	0	2	10.0	2	5.1	6	8.2	9	8.5	4	3.7	23	6.6
	90-99	0	0	3	15.0	2	5.1	5	6.8	2	1.9	1	0.9	13	3.7
	100以上	0	0	0	0	0	0	2	2.7	2	1.9	2	1.8	6	1.7
	—総数—		1	100.0	20	100.0	39	100.0	73	100.0	106	100.0	109	100.0	348
(再掲)	40未満	0	0	1	5.0	0	0	0	0	5	4.7	14	12.8	20	5.7
	40以上	1	100.0	19	95.0	39	100.0	73	100.0	101	95.3	95	87.2	328	94.3
HbA1c 6.1%以上 または現在治療を受けている	30未満	0	0	0	0	0	0	0	0	0	0	1	1.3	1	0.5
	30-39	0	0	0	0	0	0	2	6.3	4	5.3	6	8.0	12	5.9
	40-49	0	0	2	50.0	3	17.6	7	21.9	13	17.3	18	24.0	43	21.0
	50-59	1	50.0	1	25.0	4	23.5	8	25.0	27	36.0	25	33.3	66	32.2
	60-69	1	50.0	1	25.0	4	23.5	8	25.0	16	21.3	14	18.7	44	21.5
	70-79	0	0	0	0	4	23.5	5	15.6	7	9.3	6	8.0	22	10.7
	80-89	0	0	0	0	1	5.9	1	3.1	5	6.7	4	5.3	11	5.4
	90-99	0	0	0	0	1	5.9	1	3.1	3	4.0	1	1.3	6	2.9
	100以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	—総数—		2	100.0	4	100.0	17	100.0	32	100.0	75	100.0	75	100.0	205
(再掲)	40未満	0	0	0	0	0	0	2	6.3	4	5.3	7	9.3	13	6.3
	40以上	2	100.0	4	100.0	17	100.0	30	93.8	71	94.7	68	90.7	192	93.7

第64表-1 糖尿病の状況別、血清トリグリセリド値（食後3時間以上）の分布（男女計、年齢階級別）

糖尿病の状況	血清トリグリセリド値(mg/dl)	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		— 計 —	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
HbA1c 5.6%未満	150未満	219	87.3	399	80.3	365	74.5	467	68.5	426	67.3	410	72.3	2,286	73.3
	150-199	17	6.8	43	8.7	58	11.8	94	13.8	93	14.7	86	15.2	391	12.5
	200-249	6	2.4	25	5.0	29	5.9	54	7.9	60	9.5	39	6.9	213	6.8
	250-299	3	1.2	11	2.2	12	2.4	31	4.5	22	3.5	21	3.7	100	3.2
	300以上	6	2.4	19	3.8	26	5.3	36	5.3	32	5.1	11	1.9	130	4.2
	—総数—	251	100.0	497	100.0	490	100.0	682	100.0	633	100.0	567	100.0	3,120	100.0
	(再掲)														
150未満	219	87.3	399	80.3	365	74.5	467	68.5	426	67.3	410	72.3	2,286	73.3	
150以上	32	12.7	98	19.7	125	25.5	215	31.5	207	32.7	157	27.7	834	26.7	
HbA1c 5.6%以上 6.1%未満	150未満	4	100.0	14	66.7	23	69.7	58	66.7	77	59.2	89	70.6	265	66.1
	150-199	0	0	4	19.0	5	15.2	13	14.9	22	16.9	20	15.9	64	16.0
	200-249	0	0	0	0	3	9.1	8	9.2	14	10.8	10	7.9	35	8.7
	250-299	0	0	1	4.8	0	0	2	2.3	5	3.8	5	4.0	13	3.2
	300以上	0	0	2	9.5	2	6.1	6	6.9	12	9.2	2	1.6	24	6.0
	—総数—	4	100.0	21	100.0	33	100.0	87	100.0	130	100.0	126	100.0	401	100.0
	(再掲)														
150未満	4	100.0	14	66.7	23	69.7	58	66.7	77	59.2	89	70.6	265	66.1	
150以上	0	0	7	33.3	10	30.3	29	33.3	53	40.8	37	29.4	136	33.9	
HbA1c 6.1%以上 または現在治療を 受けている	150未満	1	50.0	2	40.0	12	57.1	37	50.7	78	61.9	71	55.9	201	56.8
	150-199	0	0	1	20.0	2	9.5	15	20.5	20	15.9	30	23.6	68	19.2
	200-249	1	50.0	0	0	4	19.0	7	9.6	10	7.9	9	7.1	31	8.8
	250-299	0	0	0	0	1	4.8	1	1.4	8	6.3	8	6.3	18	5.1
	300以上	0	0	2	40.0	2	9.5	13	17.8	10	7.9	9	7.1	36	10.2
	—総数—	2	100.0	5	100.0	21	100.0	73	100.0	126	100.0	127	100.0	354	100.0
	(再掲)														
150未満	1	50.0	2	40.0	12	57.1	37	50.7	78	61.9	71	55.9	201	56.8	
150以上	1	50.0	3	60.0	9	42.9	36	49.3	48	38.1	56	44.1	153	43.2	

第64表-2 糖尿病の状況別、血清トリグリセリド値（食後3時間以上）の分布（男性、年齢階級別）

糖尿病の状況	血清トリグリセリド値(mg/dl)	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
HbA1c 5.6%未満	150未満	70	77.8	125	65.1	118	59.0	155	59.8	193	66.3	157	69.2	818	65.0
	150-199	11	12.2	29	15.1	34	17.0	44	17.0	44	15.1	38	16.7	200	15.9
	200-249	3	3.3	17	8.9	18	9.0	27	10.4	25	8.6	17	7.5	107	8.5
	250-299	2	2.2	6	3.1	7	3.5	14	5.4	12	4.1	12	5.3	53	4.2
	300以上	4	4.4	15	7.8	23	11.5	19	7.3	17	5.8	3	1.3	81	6.4
	—総数—	90	100.0	192	100.0	200	100.0	259	100.0	291	100.0	227	100.0	1,259	100.0
	(再掲)														
150未満	70	77.8	125	65.1	118	59.0	155	59.8	193	66.3	157	69.2	818	65.0	
150以上	20	22.2	67	34.9	82	41.0	104	40.2	98	33.7	70	30.8	441	35.0	
HbA1c 5.6%以上 6.1%未満	150未満	3	100.0	2	40.0	1	14.3	19	54.3	26	48.1	38	70.4	89	56.3
	150-199	0	0	1	20.0	2	28.6	7	20.0	11	20.4	9	16.7	30	19.0
	200-249	0	0	0	0	2	28.6	4	11.4	7	13.0	5	9.3	18	11.4
	250-299	0	0	0	0	0	0	2	5.7	1	1.9	1	1.9	4	2.5
	300以上	0	0	2	40.0	2	28.6	3	8.6	9	16.7	1	1.9	17	10.8
	—総数—	3	100.0	5	100.0	7	100.0	35	100.0	54	100.0	54	100.0	158	100.0
	(再掲)														
150未満	3	100.0	2	40.0	1	14.3	19	54.3	26	48.1	38	70.4	89	56.3	
150以上	0	0	3	60.0	6	85.7	16	45.7	28	51.9	16	29.6	69	43.7	
HbA1c 6.1%以上 または現在治療を 受けている	150未満	0	0	0	0	4	33.3	26	54.2	46	59.7	53	65.4	129	58.6
	150-199	0	0	1	50.0	1	8.3	10	20.8	10	13.0	13	16.0	35	15.9
	200-249	0	0	0	0	4	33.3	4	8.3	7	9.1	5	6.2	20	9.1
	250-299	0	0	0	0	1	8.3	0	0	6	7.8	5	6.2	12	5.5
	300以上	0	0	1	50.0	2	16.7	8	16.7	8	10.4	5	6.2	24	10.9
	—総数—	0	0	2	100.0	12	100.0	48	100.0	77	100.0	81	100.0	220	100.0
	(再掲)														
150未満	0	0	0	0	4	33.3	26	54.2	46	59.7	53	65.4	129	58.6	
150以上	0	0	2	100.0	8	66.7	22	45.8	31	40.3	28	34.6	91	41.4	

第64表-3 糖尿病の状況別、血清トリグリセリド値（食後3時間以上）の分布（女性、年齢階級別）

糖尿病の状況	血清トリグリセリド値 (mg/dl)	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		— 計 —	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
HbA1c 5.6%未満	150未満	149	92.5	274	89.8	247	85.2	312	73.8	233	68.1	253	74.4	1,468	78.9
	150-199	6	3.7	14	4.6	24	8.3	50	11.8	49	14.3	48	14.1	191	10.3
	200-249	3	1.9	8	2.6	11	3.8	27	6.4	35	10.2	22	6.5	106	5.7
	250-299	1	0.6	5	1.6	5	1.7	17	4.0	10	2.9	9	2.6	47	2.5
	300以上	2	1.2	4	1.3	3	1.0	17	4.0	15	4.4	8	2.4	49	2.6
	— 総数 —	161	100.0	305	100.0	290	100.0	423	100.0	342	100.0	340	100.0	1,861	100.0
	(再掲)														
150未満	149	92.5	274	89.8	247	85.2	312	73.8	233	68.1	253	74.4	1,468	78.9	
150以上	12	7.5	31	10.2	43	14.8	111	26.2	109	31.9	87	25.6	393	21.1	
HbA1c 5.6%以上 6.1%未満	150未満	1	100.0	12	75.0	22	84.6	39	75.0	51	67.1	51	70.8	176	72.4
	150-199	0	0	3	18.8	3	11.5	6	11.5	11	14.5	11	15.3	34	14.0
	200-249	0	0	0	0	1	3.8	4	7.7	7	9.2	5	6.9	17	7.0
	250-299	0	0	1	6.3	0	0	0	0	4	5.3	4	5.6	9	3.7
	300以上	0	0	0	0	0	0	3	5.8	3	3.9	1	1.4	7	2.9
	— 総数 —	1	100.0	16	100.0	26	100.0	52	100.0	76	100.0	72	100.0	243	100.0
	(再掲)														
150未満	1	100.0	12	75.0	22	84.6	39	75.0	51	67.1	51	70.8	176	72.4	
150以上	0	0	4	25.0	4	15.4	13	25.0	25	32.9	21	29.2	67	27.6	
HbA1c 6.1%以上 または現在治療を 受けている	150未満	1	50.0	2	66.7	8	88.9	11	44.0	32	65.3	18	39.1	72	53.7
	150-199	0	0	0	0	1	11.1	5	20.0	10	20.4	17	37.0	33	24.6
	200-249	1	50.0	0	0	0	0	3	12.0	3	6.1	4	8.7	11	8.2
	250-299	0	0	0	0	0	0	1	4.0	2	4.1	3	6.5	6	4.5
	300以上	0	0	1	33.3	0	0	5	20.0	2	4.1	4	8.7	12	9.0
	— 総数 —	2	100.0	3	100.0	9	100.0	25	100.0	49	100.0	46	100.0	134	100.0
	(再掲)														
150未満	1	50.0	2	66.7	8	88.9	11	44.0	32	65.3	18	39.1	72	53.7	
150以上	1	50.0	1	33.3	1	11.1	14	56.0	17	34.7	28	60.9	62	46.3	

第65表-1 糖尿病の状況別、血漿血糖値（食後3時間以上）の分布（男女計、年齢階級別）

糖尿病の状況	血糖値 (mg/dl)	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		—計—		
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
HbA1c 5.6%未満	110未満	241	96.0	476	96.0	456	93.3	603	88.7	506	80.1	420	74.1	2,702	86.7	
	110-125	7	2.8	17	3.4	31	6.3	63	9.3	89	14.1	110	19.4	317	10.2	
	126-139	1	0.4	3	0.6	1	0.2	10	1.5	21	3.3	27	4.8	63	2.0	
	140-159	2	0.8	0	0	1	0.2	3	0.4	13	2.1	8	1.4	27	0.9	
	160-199	0	0	0	0	0	0	1	0.1	3	0.5	2	0.4	6	0.2	
	200以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	—総数—	251	100.0	496	100.0	489	100.0	680	100.0	632	100.0	567	100.0	3,115	100.0	
	(再掲)															
	110未満	241	96.0	476	96.0	456	93.3	603	88.7	506	80.1	420	74.1	2,702	86.7	
	110以上	10	4.0	20	4.0	33	6.7	77	11.3	126	19.9	147	25.9	413	13.3	
HbA1c 5.6%以上 6.1%未満	110未満	4	100.0	19	90.5	25	75.8	60	69.0	90	69.2	59	47.2	257	64.3	
	110-125	0	0	1	4.8	6	18.2	17	19.5	28	21.5	37	29.6	89	22.3	
	126-139	0	0	1	4.8	2	6.1	5	5.7	5	3.8	15	12.0	28	7.0	
	140-159	0	0	0	0	0	0	3	3.4	5	3.8	10	8.0	18	4.5	
	160-199	0	0	0	0	0	0	2	2.3	1	0.8	4	3.2	7	1.8	
	200以上	0	0	0	0	0	0	0	0	1	0.8	0	0	1	0.3	
	—総数—	4	100.0	21	100.0	33	100.0	87	100.0	130	100.0	125	100.0	400	100.0	
	(再掲)															
	110未満	4	100.0	19	90.5	25	75.8	60	69.0	90	69.2	59	47.2	257	64.3	
	110以上	0	0	2	9.5	8	24.2	27	31.0	40	30.8	66	52.8	143	35.8	
HbA1c 6.1%以上 または現在治療を 受けている	110未満	2	100.0	0	0	11	52.4	19	26.0	29	23.2	39	31.5	100	28.6	
	110-125	0	0	0	0	4	19.0	15	20.5	27	21.6	15	12.1	61	17.4	
	126-139	0	0	0	0	3	14.3	8	11.0	11	8.8	18	14.5	40	11.4	
	140-159	0	0	0	0	0	0	11	15.1	17	13.6	18	14.5	46	13.1	
	160-199	0	0	3	60.0	3	14.3	9	12.3	16	12.8	20	16.1	51	14.6	
	200以上	0	0	2	40.0	0	0	11	15.1	25	20.0	14	11.3	52	14.9	
	—総数—	2	100.0	5	100.0	21	100.0	73	100.0	125	100.0	124	100.0	350	100.0	
	(再掲)															
	110未満	2	100.0	0	0	11	52.4	19	26.0	29	23.2	39	31.5	100	28.6	
	110以上	0	0	5	100.0	10	47.6	54	74.0	96	76.8	85	68.5	250	71.4	

第65表-2 糖尿病の状況別、血漿血糖値（食後3時間以上）の分布（男性、年齢階級別）

糖尿病の状況	血糖値 (mg/dl)	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計		
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
HbA1c 5.6%未満	110未満	86	95.6	181	94.3	187	93.5	235	90.7	245	84.2	179	78.9	1,113	88.4	
	110-125	3	3.3	9	4.7	12	6.0	20	7.7	34	11.7	29	12.8	107	8.5	
	126-139	1	1.1	2	1.0	1	0.5	4	1.5	7	2.4	14	6.2	29	2.3	
	140-159	0	0	0	0	0	0	0	0	4	1.4	4	1.8	8	0.6	
	160-199	0	0	0	0	0	0	0	0	1	0.3	1	0.4	2	0.2	
	200以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	—総数—	90	100.0	192	100.0	200	100.0	259	100.0	291	100.0	227	100.0	1,259	100.0	
	(再掲)															
	110未満	86	95.6	181	94.3	187	93.5	235	90.7	245	84.2	179	78.9	1,113	88.4	
	110以上	4	4.4	11	5.7	13	6.5	24	9.3	46	15.8	48	21.1	146	11.6	
HbA1c 5.6%以上 6.1%未満	110未満	3	100.0	4	80.0	5	71.4	23	65.7	37	68.5	34	63.0	106	67.1	
	110-125	0	0	1	20.0	1	14.3	8	22.9	12	22.2	11	20.4	33	20.9	
	126-139	0	0	0	0	1	14.3	3	8.6	2	3.7	6	11.1	12	7.6	
	140-159	0	0	0	0	0	0	1	2.9	3	5.6	2	3.7	6	3.8	
	160-199	0	0	0	0	0	0	0	0	0	0	1	1.9	1	0.6	
	200以上	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	—総数—	3	100.0	5	100.0	7	100.0	35	100.0	54	100.0	54	100.0	158	100.0	
	(再掲)															
	110未満	3	100.0	4	80.0	5	71.4	23	65.7	37	68.5	34	63.0	106	67.1	
	110以上	0	0	1	20.0	2	28.6	12	34.3	17	31.5	20	37.0	52	32.9	
HbA1c 6.1%以上 または現在治療を 受けている	110未満	0	0	0	0	7	58.3	12	25.0	20	26.0	28	35.9	67	30.9	
	110-125	0	0	0	0	2	16.7	10	20.8	15	19.5	8	10.3	35	16.1	
	126-139	0	0	0	0	1	8.3	6	12.5	8	10.4	10	12.8	25	11.5	
	140-159	0	0	0	0	0	0	7	14.6	9	11.7	10	12.8	26	12.0	
	160-199	0	0	1	50.0	2	16.7	3	6.3	9	11.7	13	16.7	28	12.9	
	200以上	0	0	1	50.0	0	0	10	20.8	16	20.8	9	11.5	36	16.6	
	—総数—	0	0	2	100.0	12	100.0	48	100.0	77	100.0	78	100.0	217	100.0	
	(再掲)															
	110未満	0	0	0	0	7	58.3	12	25.0	20	26.0	28	35.9	67	30.9	
	110以上	0	0	2	100.0	5	41.7	36	75.0	57	74.0	50	64.1	150	69.1	

第65表-3 糖尿病の状況別、血漿血糖値（食後3時間以上）の分布（女性、年齢階級別）

糖尿病の状況	血糖値 (mg/dl)	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計一		
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
HbA1c 5.6%未満	110未満	155	96.3	295	97.0	269	93.1	368	87.4	261	76.5	241	70.9	1,589	85.6	
	110-125	4	2.5	8	2.6	19	6.6	43	10.2	55	16.1	81	23.8	210	11.3	
	126-139	0	0	1	0.3	0	0	6	1.4	14	4.1	13	3.8	34	1.8	
	140-159	2	1.2	0	0	1	0.3	3	0.7	9	2.6	4	1.2	19	1.0	
	160-199	0	0	0	0	0	0	1	0.2	2	0.6	1	0.3	4	0.2	
	200以上															
	一総数一		161	100.0	304	100.0	289	100.0	421	100.0	341	100.0	340	100.0	1,856	100.0
(再掲)	110未満	155	96.3	295	97.0	269	93.1	368	87.4	261	76.5	241	70.9	1,589	85.6	
	110以上	6	3.7	9	3.0	20	6.9	53	12.6	80	23.5	99	29.1	267	14.4	
HbA1c 5.6%以上 6.1%未満	110未満	1	100.0	15	93.8	20	76.9	37	71.2	53	69.7	25	35.2	151	62.4	
	110-125	0	0	0	0	5	19.2	9	17.3	16	21.1	26	36.6	56	23.1	
	126-139	0	0	1	6.3	1	3.8	2	3.8	3	3.9	9	12.7	16	6.6	
	140-159	0	0	0	0	0	0	2	3.8	2	2.6	8	11.3	12	5.0	
	160-199	0	0	0	0	0	0	2	3.8	1	1.3	3	4.2	6	2.5	
	200以上	0	0	0	0	0	0	0	0	1	1.3	0	0	1	0.4	
	一総数一		1	100.0	16	100.0	26	100.0	52	100.0	76	100.0	71	100.0	242	100.0
(再掲)	110未満	1	100.0	15	93.8	20	76.9	37	71.2	53	69.7	25	35.2	151	62.4	
	110以上	0	0	1	6.3	6	23.1	15	28.8	23	30.3	46	64.8	91	37.6	
HbA1c 6.1%以上 または現在治療を 受けている	110未満	2	100.0	0	0	4	44.4	7	28.0	9	18.8	11	23.9	33	24.8	
	110-125	0	0	0	0	2	22.2	5	20.0	12	25.0	7	15.2	26	19.5	
	126-139	0	0	0	0	2	22.2	2	8.0	3	6.3	8	17.4	15	11.3	
	140-159	0	0	0	0	0	0	4	16.0	8	16.7	8	17.4	20	15.0	
	160-199	0	0	2	66.7	1	11.1	6	24.0	7	14.6	7	15.2	23	17.3	
	200以上	0	0	1	33.3	0	0	1	4.0	9	18.8	5	10.9	16	12.0	
	一総数一		2	100.0	3	100.0	9	100.0	25	100.0	48	100.0	46	100.0	133	100.0
(再掲)	110未満	2	100.0	0	0	4	44.4	7	28.0	9	18.8	11	23.9	33	24.8	
	110以上	0	0	3	100.0	5	55.6	18	72.0	39	81.3	35	76.1	100	75.2	

第66表-1 地域ブロック別、糖尿病の状況（男女計、年齢階級別）

地域 ブロック	糖尿病の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
北海道	HbA1c 5.6%未満	18	100.0	23	88.5	36	100.0	36	83.7	30	62.5	36	70.6	179	80.6
	HbA1c 5.6%以上6.1%未満	0	0	3	11.5	0	0	2	4.7	7	14.6	6	11.8	18	8.1
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	5	11.6	11	22.9	9	17.6	25	11.3
	—総数—	18	100.0	26	100.0	36	100.0	43	100.0	48	100.0	51	100.0	222	100.0
東北	HbA1c 5.6%未満	38	100.0	49	92.5	63	94.0	84	83.2	83	76.9	72	63.7	389	81.0
	HbA1c 5.6%以上6.1%未満	0	0	3	5.7	2	3.0	8	7.9	16	14.8	19	16.8	48	10.0
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	1.9	2	3.0	9	8.9	9	8.3	22	19.5	43	9.0
	—総数—	38	100.0	53	100.0	67	100.0	101	100.0	108	100.0	113	100.0	480	100.0
関東Ⅰ	HbA1c 5.6%未満	70	100.0	157	95.7	108	90.8	163	85.3	152	72.0	123	71.9	773	83.5
	HbA1c 5.6%以上6.1%未満	0	0	6	3.7	7	5.9	22	11.5	28	13.3	29	17.0	92	9.9
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	0.6	4	3.4	6	3.1	31	14.7	19	11.1	61	6.6
	—総数—	70	100.0	164	100.0	119	100.0	191	100.0	211	100.0	171	100.0	926	100.0
関東Ⅱ	HbA1c 5.6%未満	34	100.0	69	93.2	84	91.3	81	79.4	62	61.4	55	59.8	385	77.8
	HbA1c 5.6%以上6.1%未満	0	0	4	5.4	5	5.4	13	12.7	24	23.8	21	22.8	67	13.5
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	1.4	3	3.3	8	7.8	15	14.9	16	17.4	43	8.7
	—総数—	34	100.0	74	100.0	92	100.0	102	100.0	101	100.0	92	100.0	495	100.0
北陸	HbA1c 5.6%未満	24	96.0	32	94.1	53	89.8	53	86.9	51	66.2	51	72.9	264	81.0
	HbA1c 5.6%以上6.1%未満	1	4.0	2	5.9	6	10.2	2	3.3	15	19.5	10	14.3	36	11.0
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	6	9.8	11	14.3	9	12.9	26	8.0
	—総数—	25	100.0	34	100.0	59	100.0	61	100.0	77	100.0	70	100.0	326	100.0
東海	HbA1c 5.6%未満	57	95.0	94	97.9	103	90.4	149	78.4	131	73.2	81	65.9	615	80.7
	HbA1c 5.6%以上6.1%未満	2	3.3	2	2.1	8	7.0	22	11.6	24	13.4	21	17.1	79	10.4
	HbA1c 6.1%以上または現在治療を受けている	1	1.7	0	0	3	2.6	19	10.0	24	13.4	21	17.1	68	8.9
	—総数—	60	100.0	96	100.0	114	100.0	190	100.0	179	100.0	123	100.0	762	100.0
近畿Ⅰ	HbA1c 5.6%未満	40	97.6	80	98.8	52	85.2	92	78.6	108	67.9	87	66.9	459	77.9
	HbA1c 5.6%以上6.1%未満	1	2.4	1	1.2	6	9.8	15	12.8	25	15.7	24	18.5	72	12.2
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	3	4.9	10	8.5	26	16.4	19	14.6	58	9.8
	—総数—	41	100.0	81	100.0	61	100.0	117	100.0	159	100.0	130	100.0	589	100.0
近畿Ⅱ	HbA1c 5.6%未満	12	100.0	7	100.0	19	79.2	30	81.1	24	75.0	26	66.7	118	78.1
	HbA1c 5.6%以上6.1%未満	0	0	0	0	2	8.3	4	10.8	1	3.1	7	17.9	14	9.3
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	3	12.5	3	8.1	7	21.9	6	15.4	19	12.6
	—総数—	12	100.0	7	100.0	24	100.0	37	100.0	32	100.0	39	100.0	151	100.0
中国	HbA1c 5.6%未満	34	100.0	54	96.4	47	94.0	76	76.0	71	69.6	55	59.8	337	77.6
	HbA1c 5.6%以上6.1%未満	0	0	1	1.8	2	4.0	13	13.0	15	14.7	15	16.3	46	10.6
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	1.8	1	2.0	11	11.0	16	15.7	22	23.9	51	11.8
	—総数—	34	100.0	56	100.0	50	100.0	100	100.0	102	100.0	92	100.0	434	100.0
四国	HbA1c 5.6%未満	13	100.0	19	90.5	30	93.8	30	78.9	29	80.6	38	74.5	159	83.2
	HbA1c 5.6%以上6.1%未満	0	0	1	4.8	0	0	5	13.2	5	13.9	6	11.8	17	8.9
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	4.8	2	6.3	3	7.9	2	5.6	7	13.7	15	7.9
	—総数—	13	100.0	21	100.0	32	100.0	38	100.0	36	100.0	51	100.0	191	100.0
北九州	HbA1c 5.6%未満	22	95.7	47	92.2	59	89.4	61	83.6	51	65.4	65	69.9	305	79.4
	HbA1c 5.6%以上6.1%未満	0	0	4	7.8	3	4.5	7	9.6	13	16.7	12	12.9	39	10.2
	HbA1c 6.1%以上または現在治療を受けている	1	4.3	0	0	4	6.1	5	6.8	14	17.9	16	17.2	40	10.4
	—総数—	23	100.0	51	100.0	66	100.0	73	100.0	78	100.0	93	100.0	384	100.0
南九州	HbA1c 5.6%未満	30	100.0	49	98.0	34	72.3	56	80.0	64	82.1	82	73.9	315	81.6
	HbA1c 5.6%以上6.1%未満	0	0	0	0	8	17.0	7	10.0	6	7.7	17	15.3	38	9.8
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	2.0	5	10.6	7	10.0	8	10.3	12	10.8	33	8.5
	—総数—	30	100.0	50	100.0	47	100.0	70	100.0	78	100.0	111	100.0	386	100.0

第66表-2 地域ブロック別、糖尿病の状況 (男性、年齢階級別)

地域 ブロック	糖尿病の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
北海道	HbA1c 5.6%未満	8	100.0	7	100.0	16	100.0	11	64.7	15	65.2	12	63.2	69	76.7
	HbA1c5.6%以上6.1%未満	0	0	0	0	0	0	1	5.9	2	8.7	3	15.8	6	6.7
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	5	29.4	6	26.1	4	21.1	15	16.7
	一総数	8	100.0	7	100.0	16	100.0	17	100.0	23	100.0	19	100.0	90	100.0
東北	HbA1c 5.6%未満	14	100.0	22	100.0	20	95.2	39	81.3	33	73.3	27	56.3	155	78.3
	HbA1c5.6%以上6.1%未満	0	0	0	0	0	0	4	8.3	6	13.3	10	20.8	20	10.1
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	4.8	5	10.4	6	13.3	11	22.9	23	11.6
	一総数	14	100.0	22	100.0	21	100.0	48	100.0	45	100.0	48	100.0	198	100.0
関東Ⅰ	HbA1c 5.6%未満	25	100.0	54	94.7	30	100.0	44	77.2	74	70.5	63	75.9	290	81.2
	HbA1c5.6%以上6.1%未満	0	0	3	5.3	0	0	10	17.5	11	10.5	10	12.0	34	9.5
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	3	5.3	20	19.0	10	12.0	33	9.2
	一総数	25	100.0	57	100.0	30	100.0	57	100.0	105	100.0	83	100.0	357	100.0
関東Ⅱ	HbA1c 5.6%未満	17	100.0	23	88.5	40	90.9	30	69.8	28	62.2	23	53.5	161	73.9
	HbA1c5.6%以上6.1%未満	0	0	2	7.7	2	4.5	8	18.6	9	20.0	7	16.3	28	12.8
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	3.8	2	4.5	5	11.6	8	17.8	13	30.2	29	13.3
	一総数	17	100.0	26	100.0	44	100.0	43	100.0	45	100.0	43	100.0	218	100.0
北陸	HbA1c 5.6%未満	6	85.7	16	94.1	22	91.7	20	87.0	21	61.8	19	65.5	104	77.6
	HbA1c5.6%以上6.1%未満	1	14.3	1	5.9	2	8.3	0	0	6	17.6	5	17.2	15	11.2
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	3	13.0	7	20.6	5	17.2	15	11.2
	一総数	7	100.0	17	100.0	24	100.0	23	100.0	34	100.0	29	100.0	134	100.0
東海	HbA1c 5.6%未満	27	96.4	31	100.0	39	86.7	52	69.3	61	74.4	34	65.4	244	78.0
	HbA1c5.6%以上6.1%未満	1	3.6	0	0	4	8.9	8	10.7	12	14.6	8	15.4	33	10.5
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	2	4.4	15	20.0	9	11.0	10	19.2	36	11.5
	一総数	28	100.0	31	100.0	45	100.0	75	100.0	82	100.0	52	100.0	313	100.0
近畿Ⅰ	HbA1c 5.6%未満	10	90.9	25	100.0	16	84.2	34	79.1	44	66.7	30	56.6	159	73.3
	HbA1c5.6%以上6.1%未満	1	9.1	0	0	2	10.5	5	11.6	9	13.6	11	20.8	28	12.9
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	5.3	4	9.3	13	19.7	12	22.6	30	13.8
	一総数	11	100.0	25	100.0	19	100.0	43	100.0	66	100.0	53	100.0	217	100.0
近畿Ⅱ	HbA1c 5.6%未満	3	100.0	3	100.0	6	66.7	11	78.6	10	66.7	7	58.3	40	71.4
	HbA1c5.6%以上6.1%未満	0	0	0	0	0	0	1	7.1	0	0	2	16.7	3	5.4
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	3	33.3	2	14.3	5	33.3	3	25.0	13	23.2
	一総数	3	100.0	3	100.0	9	100.0	14	100.0	15	100.0	12	100.0	56	100.0
中国	HbA1c 5.6%未満	10	100.0	27	100.0	22	95.7	24	64.9	26	56.5	19	50.0	128	70.7
	HbA1c5.6%以上6.1%未満	0	0	0	0	0	0	4	10.8	9	19.6	6	15.8	19	10.5
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	4.3	9	24.3	11	23.9	13	34.2	34	18.8
	一総数	10	100.0	27	100.0	23	100.0	37	100.0	46	100.0	38	100.0	181	100.0
四国	HbA1c 5.6%未満	7	100.0	8	100.0	19	95.0	8	66.7	14	87.5	14	70.0	70	84.3
	HbA1c5.6%以上6.1%未満	0	0	0	0	0	0	2	16.7	2	12.5	4	20.0	8	9.6
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	5.0	2	16.7	0	0	2	10.0	5	6.0
	一総数	7	100.0	8	100.0	20	100.0	12	100.0	16	100.0	20	100.0	83	100.0
北九州	HbA1c 5.6%未満	4	100.0	15	93.8	24	96.0	21	84.0	23	65.7	21	56.8	108	76.1
	HbA1c5.6%以上6.1%未満	0	0	1	6.3	0	0	2	8.0	5	14.3	5	13.5	13	9.2
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	4.0	2	8.0	7	20.0	11	29.7	21	14.8
	一総数	4	100.0	16	100.0	25	100.0	25	100.0	35	100.0	37	100.0	142	100.0
南九州	HbA1c 5.6%未満	11	100.0	17	94.4	18	94.7	29	82.9	26	76.5	30	68.2	131	81.4
	HbA1c5.6%以上6.1%未満	0	0	0	0	0	0	1	2.9	2	5.9	6	13.6	9	5.6
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	5.6	1	5.3	5	14.3	6	17.6	8	18.2	21	13.0
	一総数	11	100.0	18	100.0	19	100.0	35	100.0	34	100.0	44	100.0	161	100.0

第66表-3 地域ブロック別、糖尿病の状況（女性、年齢階級別）

地域 ブロック	糖尿病の状況	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		一計	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
北海道	HbA1c 5.6%未満	10	100.0	16	84.2	20	100.0	25	96.2	15	60.0	24	75.0	110	83.3
	HbA1c 5.6%以上 6.1%未満	0	0	3	15.8	0	0	1	3.8	5	20.0	3	9.4	12	9.1
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	0	0	5	20.0	5	15.6	10	7.6
	—総数—	10	100.0	19	100.0	20	100.0	26	100.0	25	100.0	32	100.0	132	100.0
東北	HbA1c 5.6%未満	24	100.0	27	87.1	43	93.5	45	84.9	50	79.4	45	69.2	234	83.0
	HbA1c 5.6%以上 6.1%未満	0	0	3	9.7	2	4.3	4	7.5	10	15.9	9	13.8	28	9.9
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	3.2	1	2.2	4	7.5	3	4.8	11	16.9	20	7.1
	—総数—	24	100.0	31	100.0	46	100.0	53	100.0	63	100.0	65	100.0	282	100.0
関東Ⅰ	HbA1c 5.6%未満	45	100.0	103	96.3	78	87.6	119	88.8	78	73.6	60	68.2	483	84.9
	HbA1c 5.6%以上 6.1%未満	0	0	3	2.8	7	7.9	12	9.0	17	16.0	19	21.6	58	10.2
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	0.9	4	4.5	3	2.2	11	10.4	9	10.2	28	4.9
	—総数—	45	100.0	107	100.0	89	100.0	134	100.0	106	100.0	88	100.0	569	100.0
関東Ⅱ	HbA1c 5.6%未満	17	100.0	46	95.8	44	91.7	51	86.4	34	60.7	32	65.3	224	80.9
	HbA1c 5.6%以上 6.1%未満	0	0	2	4.2	3	6.3	5	8.5	15	26.8	14	28.6	39	14.1
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	1	2.1	3	5.1	7	12.5	3	6.1	14	5.1
	—総数—	17	100.0	48	100.0	48	100.0	59	100.0	56	100.0	49	100.0	277	100.0
北陸	HbA1c 5.6%未満	18	100.0	16	94.1	31	88.6	33	86.8	30	69.8	32	78.0	160	83.3
	HbA1c 5.6%以上 6.1%未満	0	0	1	5.9	4	11.4	2	5.3	9	20.9	5	12.2	21	10.9
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	3	7.9	4	9.3	4	9.8	11	5.7
	—総数—	18	100.0	17	100.0	35	100.0	38	100.0	43	100.0	41	100.0	192	100.0
東海	HbA1c 5.6%未満	30	93.8	63	96.9	64	92.8	97	84.3	70	72.2	47	66.2	371	82.6
	HbA1c 5.6%以上 6.1%未満	1	3.1	2	3.1	4	5.8	14	12.2	12	12.4	13	18.3	46	10.2
	HbA1c 6.1%以上または現在治療を受けている	1	3.1	0	0	1	1.4	4	3.5	15	15.5	11	15.5	32	7.1
	—総数—	32	100.0	65	100.0	69	100.0	115	100.0	97	100.0	71	100.0	449	100.0
近畿Ⅰ	HbA1c 5.6%未満	30	100.0	55	98.2	36	85.7	58	78.4	64	68.8	57	74.0	300	80.6
	HbA1c 5.6%以上 6.1%未満	0	0	1	1.8	4	9.5	10	13.5	16	17.2	13	16.9	44	11.8
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	2	4.8	6	8.1	13	14.0	7	9.1	28	7.5
	—総数—	30	100.0	56	100.0	42	100.0	74	100.0	93	100.0	77	100.0	372	100.0
近畿Ⅱ	HbA1c 5.6%未満	9	100.0	4	100.0	13	86.7	19	82.6	14	82.4	19	70.4	78	82.1
	HbA1c 5.6%以上 6.1%未満	0	0	0	0	2	13.3	3	13.0	1	5.9	5	18.5	11	11.6
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	0	0	1	4.3	2	11.8	3	11.1	6	6.3
	—総数—	9	100.0	4	100.0	15	100.0	23	100.0	17	100.0	27	100.0	95	100.0
中国	HbA1c 5.6%未満	24	100.0	27	93.1	25	92.6	52	82.5	45	80.4	36	66.7	209	82.6
	HbA1c 5.6%以上 6.1%未満	0	0	1	3.4	2	7.4	9	14.3	6	10.7	9	16.7	27	10.7
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	3.4	0	0	2	3.2	5	8.9	9	16.7	17	6.7
	—総数—	24	100.0	29	100.0	27	100.0	63	100.0	56	100.0	54	100.0	253	100.0
四国	HbA1c 5.6%未満	6	100.0	11	84.6	11	91.7	22	84.6	15	75.0	24	77.4	89	82.4
	HbA1c 5.6%以上 6.1%未満	0	0	1	7.7	0	0	3	11.5	3	15.0	2	6.5	9	8.3
	HbA1c 6.1%以上または現在治療を受けている	0	0	1	7.7	1	8.3	1	3.8	2	10.0	5	16.1	10	9.3
	—総数—	6	100.0	13	100.0	12	100.0	26	100.0	20	100.0	31	100.0	108	100.0
北九州	HbA1c 5.6%未満	18	94.7	32	91.4	35	85.4	40	83.3	28	65.1	44	78.6	197	81.4
	HbA1c 5.6%以上 6.1%未満	0	0	3	8.6	3	7.3	5	10.4	8	18.6	7	12.5	26	10.7
	HbA1c 6.1%以上または現在治療を受けている	1	5.3	0	0	3	7.3	3	6.3	7	16.3	5	8.9	19	7.9
	—総数—	19	100.0	35	100.0	41	100.0	48	100.0	43	100.0	56	100.0	242	100.0
南九州	HbA1c 5.6%未満	19	100.0	32	100.0	16	57.1	27	77.1	38	86.4	52	77.6	184	81.8
	HbA1c 5.6%以上 6.1%未満	0	0	0	0	8	28.6	6	17.1	4	9.1	11	16.4	29	12.9
	HbA1c 6.1%以上または現在治療を受けている	0	0	0	0	4	14.3	2	5.7	2	4.5	4	6.0	12	5.3
	—総数—	19	100.0	32	100.0	28	100.0	35	100.0	44	100.0	67	100.0	225	100.0

平成14年度糖尿病実態調査企画・解析検討会

*は座長

赤沼安夫* 社団法人 日本糖尿病学会理事長

門脇 孝 東京大学大学院医学系研究科
糖尿病・代謝内科助教授

岸本泰子 島根県雲南保健所長

櫻井秀也 社団法人 日本医師会常任理事

田嶋尚子 東京慈恵会医科大学内科学講座
糖尿病・代謝・内分泌内科教授

伊達ちぐさ 武庫川女子大学生生活環境学部教授

田中平三 独立行政法人 国立健康・栄養研究所理事長

松岡健平 社団法人 日本糖尿病協会副理事長

山本光昭 茨城県保健福祉部長